

1 **Title:**

2 Electromyographic Analysis of Shoulder Girdle Muscles during Common
3 Internal Rotation Exercises

4 **Authors:**

5 Omid Alizadehkhayat (MD, PhD)¹, David H Hawkes (MD)², Graham J Kemp (DM,
6 FRCPath)³, Simon P Frostick (DM, FRCS)²

7 1. School of Health Sciences (Sport and Exercise Science), Liverpool Hope University,
8 Liverpool, UK

9 2. Musculoskeletal Science Research Group, Institute of Translational Medicine, University
10 of Liverpool, Liverpool, UK

11 3. Department of Musculoskeletal Biology II, Institute of Ageing & Chronic Disease,
12 Faculty of Health & Life Sciences, University of Liverpool, UK

13 **First Author and Corresponding Author:**

14 Dr Omid Alizadehkhayat

15 School of Health Sciences, Liverpool Hope University

16 Hope Park, Liverpool L16 9JD, UK

17 Tel: 0151 291 3262

18 Email: alizado@hope.ac.uk

19 **Abstract**

20 **Background:** High level throwing performance requires the development of effective muscle
21 activation within shoulder girdle muscles particularly during forceful internal rotation (IR)
22 motions.

23 **Study Design:** Controlled Laboratory Descriptive Study

24 **Purpose:** To investigate activation pattern of 16 shoulder girdle muscles/muscle sub-regions
25 during three common shoulder IR exercises.

26 **Method:** EMG was recorded in 30 healthy subjects from 16 shoulder girdle muscles/muscle
27 sub-regions (surface electrode: anterior, middle and posterior deltoid, upper, middle and
28 lower trapezius, serratus anterior, teres major, upper and lower latissimus dorsi, upper and
29 lower pectoralis major; fine wire electrodes: supraspinatus, infraspinatus, subscapularis and
30 rhomboid major) using a telemetric EMG system. Three IR exercises (standing IR at 0° and
31 90° of Abduction, and IR at Zero-Position) were studied. EMG amplitudes were normalized
32 to EMG_{max} (EMG at maximal IR force in a standard position) and compared using one-way
33 repeated-measures analysis of variance (ANOVA).

34 **Results:** There were significant differences in muscles' activation across IR exercises
35 ($p < 0.05$ – $p < 0.001$). Rotator cuff and deltoid muscles were highly activated during IR at 90° of
36 Abduction. Latissimus dorsi exhibited markedly higher activation during IR at Zero-Position.
37 While upper trapezius had the highest activation during IR at Zero-Position, middle and
38 lower trapezius were activated at highest during IR at 90° of Abduction. The highest
39 activation of serratus anterior and rhomboid major occurred in IR at Zero-Position and IR at
40 90° of Abduction, respectively.

41 **Conclusions:** Studied exercises have the potential to effectively activate glenohumeral and
42 scapular muscles involved in throwing motions. Results provide further evidence for
43 developing rehabilitation, injury prevention, and training strategies.

44 **Key Words:** Electromyography; Internal Rotation Exercises; Rehabilitation; Shoulder

45 Muscle Activation

46 Level of Evidence: 4, Controlled laboratory study

47 **INTRODUCTION**

48 The glenohumeral joint (GHJ) is the most mobile joint in the human body due to its bony
49 structure which requires the coordinated activation of shoulder complex musculature to
50 achieve functional stability during movements.¹ The activation of key rotator cuff (RC)
51 muscles is a fundamental contributor to shoulder joint stability (centring the humeral head
52 into the glenoid) and efficient force development during arm elevation and overhead
53 activities such as throwing.²⁻⁴ The parts of the deltoid work along with the RC to develop
54 force couples required for arm motion during elevation and rotation. Pectoralis major,
55 latissimus dorsi, and teres major produce coordinated adduction moments during GHJ
56 elevation and abduction. Concurrent activation of these muscles and the subscapularis
57 stabilize the GHJ inferiorly.⁵

58 A synchronised contribution from scapular musculature is also critical for optimal
59 positioning, stability, and functioning of the shoulder complex. In addition to linking the
60 upper extremity and trunk, the scapula provides insertion points for several muscles involved
61 in scapulohumeral and scapulothoracic motions.^{6,7} Scapular stabilizers play substantial roles
62 in maintaining the center of glenohumeral rotation during arm-scapula-trunk motion, raising
63 the acromion during glenohumeral rotation to increase subacromial space, and transition of
64 forces from the feet to the hand by kinetically linking the upper extremity to the trunk.

65 During rotational motions, a coordinated balance between mobility and functional stability is
66 essential for the safe transmission of the high forces placed on the shoulder complex. Yet,
67 repetitive forceful movements may impose stress on the GHJ beyond the physiologic limits
68 of composing tissues and lead to injury. For example, cadaveric studies have shown that
69 vigorous abduction and external rotation (e.g. late cocking phase of throwing motion) in the
70 presence of decreased subscapularis muscle force can lead to forceful internal impingement

71 due to significant increase in GHJ contact pressure.⁸ Furthermore, biceps pulley lesions
72 caused by repetitive forceful IR above the horizontal plane can potentially lead to internal
73 impingement by causing frictional impairment between the pulley system and the
74 subscapularis tendon and the anterior superior glenoid rim.^{9, 10} Earlier electromyography
75 EMG studies have documented shoulder girdle muscle activation during common internal
76 rotation (IR) exercises to support the development of evidence-based rehabilitation and injury
77 prevention programs.^{2, 6, 11} The results, however, remain inconclusive and uncertainty exists
78 regarding optimal IR exercises that elicit optimal activation and strengthening of key
79 shoulder girdle muscles. Furthermore, the majority of previous studies compared the EMG
80 activity of a limited number of muscles during exercises.

81 There is, thus a lack of comprehensive data regarding shoulder musculature activation
82 strategies during common internal rotation exercises. This knowledge would guide the
83 planning of effective training programs, and establish a base of evidence for developing
84 optimal rehabilitation and training programs for overhead athletes with and without shoulder
85 pathology. The purpose of this study was to provide such a knowledge base by
86 comprehensive measurement of the EMG activity of 16 shoulder girdle muscles/muscle
87 segments during commonly prescribed shoulder IR exercises.

88 **METHODS**

89 **Participants**

90 Thirty healthy volunteers (15 male; 15 female) with normal upper limb clinical examination
91 and no history of upper limb painful conditions were recruited for participation in the study.
92 The mean (\pm SD) age, height and weight for the whole group was 33.1 ± 9.9 y, 1.71 ± 0.08 m,
93 and 70.5 ± 12.7 kg, respectively. This study received approval from local research ethics
94 committee and written informed consent was obtained from participants. The data were
95 collected in a university laboratory setting.

96 **EMG Measurements**

97 Signal acquisition, processing and analysis were performed using a TeleMyo 2400 G2
98 Telemetry System (Noraxon Inc., Arizona; USA). Signals were differentially amplified
99 (CMRR $>$ 100 dB; input impedance $>$ 100 Mohm; gain 500 dB), digitized at a sampling rate of
100 3000 Hz and band-pass filtered at 10-500 Hz and 10-1500 Hz for surface and fine-wire
101 electrodes, respectively. A cancellation algorithm was applied to remove ECG signal
102 contamination.

103 Disposable Ag/AgCl bipolar surface electrodes with 10mm conducting area and 20mm inter-
104 electrode distance (Noraxon Inc., Arizona, USA) were used to record the EMG from anterior,
105 middle, and posterior deltoid (AD, MD, PD, respectively), upper, middle and lower trapezius
106 (UT, MT, LT, respectively), upper and lower latissimus dorsi (ULD, LLD, respectively),
107 upper and lower pectoralis major pectoralis major (UPM, LPM, respectively), serratus
108 anterior (SA), and teres major (TM), consistent with established guidelines (SENIAM).^{12,13}
109 Bipolar hooked fine-wire electrodes (Nicolet Biomedical, Division of VIASYS, Madison,
110 USA) were used to record signals from supraspinatus (SSP), infraspinatus (ISP),
111 subscapularis (SUBS), and rhomboid major (RHOM) according to Basmajian and DeLuca.¹⁴

112 The dominant shoulder was tested in all participants. Figure 1 demonstrates the relative
113 locations of surface and fine-wire EMG electrodes.

114 Raw EMG signals from ten IR exercise cycles (the first and last IR exercise cycles were
115 omitted) were full-wave rectified and smoothed (100 ms root mean square [RMS]). For
116 normalization purpose, EMG_{max} was recorded during a standardized production of maximal
117 IR force (MVC) using a shoulder Nottingham Mecmesin Myometer with an accuracy of ± 0.1
118 % of full-scale and 1,000 N capacity (Mecmesin Ltd., Slinfold, UK) while seated, shoulder in
119 a neutral position, elbow in 90° flexion tucked to the side of body, and forearm in neutral
120 position. Data were collected during three 5-second contractions, and the average of three
121 trials was taken as EMG_{max} which was used as a reference value for EMG amplitude
122 normalization during IR exercises.

123 **Exercises**

124 Exercises are demonstrated in Figure 2. Participants were tested for three shoulder IR
125 exercises in a random order: isotonic standing IR at 0° and 90° of abduction (IR at 0° ABD
126 and IR at 90° ABD) and IR at Zero-Position (Zero rotation of the humerus with arm elevated
127 155° in the scapular plane and elastic resistance applied against IR as described by Saha).¹⁵
128 This particular exercise was chosen as during the cocking phase of throwing motion, the arm
129 is moved into external rotation past the zero position; and then during the acceleration the
130 arm is moved into forward internal rotation past the zero position again.¹⁶ Each exercise was
131 accurately demonstrated and participants were allowed time to familiarize themselves with
132 the exercise. Participants performed 12 cycles of each exercise using either a 1 kg dumbbell
133 in hand (IR at 0° ABD and IR at 90° ABD) or an elastic band (IR at Zero-Position) according
134 to a metronome set at 60 beats per minute (each concentric and eccentric phase was
135 performed during 1 beat). All participants were given a period of three-minute rest between
136 each set of exercises to minimise the impact of fatigue on measurements.

137 **Data analyses**

138 Descriptive statistics are presented as mean \pm standard deviation (SD) or standard error of the
139 mean (SEM), as appropriate. A one-way repeated-measures analysis of variance (ANOVA)
140 was used to determine the main effect of IR exercises on each muscle's activity. A
141 Bonferroni post-hoc test was then applied for the comparative pair-wise analysis of mean
142 normalized EMG (%EMG_{max}) to detect significant differences in the activation of muscles
143 across three exercises. The alpha level for statistical significance was set at $p < 0.05$. SPSS
144 release 20.0 for Windows (Armonk, NY: IBM Corp.) was used for statistical analysis.

145 **RESULTS**

146 Table 1 and Figure 3 present and compare the activation of muscles during IR exercises.

147 **Deltoids**: The highest activation of AD, MD, and PD occurred in IR at 90°ABD followed by
148 IR at Zero-Position; both significantly higher than IR at 0°ABD ($p < 0.001$). Collective deltoid
149 (AD+MD+PD) activation in IR at 90°ABD and IR at Zero-Position was also markedly higher
150 than IR at 0° ABD (346.4% vs. 252.2% vs. 49.7%; $p = 0.006 - < 0.001$).

151 **Rotator Cuff**: The activity of SSP, ISP, and SUBS in IR at 90°ABD was significantly higher
152 than IR at 0°ABD ($p < 0.05 - < 0.001$). They also showed a similar trend towards higher muscle
153 activity higher activation in IR at Zero-Position, but this difference was not statistically
154 significantly different. As a group (SSP+ISP+SUBS), higher activation occurred in IR at
155 90°ABD compared to other exercises (325.0% vs. 94.0-188.3%; $p < 0.05$).

156 **Pectoralis Major**: UPM and LPM activation did not vary across exercises. Both segments
157 showed a trend towards higher muscle activity during IR at Zero-Position, but were not
158 statistically significantly different.

159 **Latissimus Dorsi:** ULD had the highest activation in IR at Zero-Position, significantly
160 higher than IR at 0°ABD ($p<0.05$) followed by IR at 90°ABD. The activity of LLD and
161 combined segments (ULD+LLD) was similar across exercises.

162 **Teres Major:** There was no significant difference across exercises.

163 **Trapezius:** Highest UT activation occurred in IR at Zero-Position followed IR at 90°ABD,
164 both significantly higher than IR at 0°ABD ($p<0.001$). MT and LT were activated
165 considerably more in IR at 90°ABD compared to other two exercises ($p<0.001$). MT
166 activation was also higher in IR at Zero-Position than IR at 0°ABD ($p<0.05$). Collective
167 activation of the trapezius muscles (UT+MT+LT) was markedly higher in both IR at 90°ABD
168 and IR at Zero-Position compared to IR at 0°ABD (230.2% vs. 64.3-158.8%; $p<0.001$).

169 **Serratus Anterior:** The highest SA activation occurred in IR at Zero-Position which was
170 markedly higher than IR at 0°ABD ($p<0.05$).

171 **Rhomboid Major:** RM had the highest activation in IR at 90°ABD compared to other IR
172 exercises ($p<0.001$). The activity was also markedly higher in IR at Zero-Position compared
173 to IR at 0°ABD ($p<0.05$).

174 **DISCUSSION**

175 The results of the present study provide additional support for the use of these common IR
176 exercises. Furthermore, the results illustrate novel strategies for the selective activation of
177 shoulder complex muscles during specific exercises, which may be helpful during
178 implementation in training, injury prevention, and rehabilitation programs.

179 Optimal performance of shoulder complex during both daily activities and sporting
180 movements necessitates appropriately balanced activation of muscles responsible for
181 shoulder mobility and functional stability.^{1,3,7,17} The high occurrence of shoulder complex

182 injuries highlights the need for implementation of sound evidence in developing
183 rehabilitation, injury prevention, and training strategies.^{1,2,6,15}

184 Current shoulder rehabilitation strategies give emphasis to correcting muscle imbalances and
185 strength deficiencies through selectively activating dysfunctional muscles. Considering that a
186 low ER/IR ratio has been suggested as a key risk factor for shoulder injuries,^{18,19} several
187 investigators have studied muscle activation patterns during shoulder rotational exercises,
188 with inconsistent results.^{11,20-22} EMG studies of IR exercises have mainly focused on the
189 principal internal rotators such as SUBS and pectoralis muscles.²²⁻²⁴ Moreover, there is
190 growing interest in applying exercises in sport-specific positions that reflect capsular strain
191 and muscular length-tension relationships throughout the shoulder complex during sport
192 competition (e.g. ER and IR at 90°ABD) in order to facilitate enhanced functional
193 rehabilitation.^{23, 25}

194 **Glenohumeral Muscles**

195 In the present study, the highest activation of all deltoid sub-regions was found in IR at
196 90°ABD followed by IR at Zero-Position. This is consistent with the role of MD and AD
197 during dynamic arm abduction and with role of PD as humeral abductor and compressor in
198 higher degrees (>80°) of abduction.⁵ This high activation of PD is contradictory to the reports
199 of its ineffectiveness in generating abduction forces.^{26,27} Hughes and An²⁸ reported a minimal
200 force generation of 2 N for PD compared to 434 N for MD and 323 N for AD when the arm
201 is positioned at 90°ABD. It is generally suggested that exercises producing high levels of
202 deltoid activity (MD in particular) are disadvantageous for majority of patients and athletes
203 with shoulder injury due to significant impact on superior humeral head migration.^{17, 23}

204 Similar to deltoids, RC muscles including SSP, ISP, and SUBS had their highest activation in
205 IR at 90°ABD followed by IR at Zero-Position. Jenp et al²⁹ reported substantial activity in the

206 SSP during shoulder IR. The activation patterns in the deltoids and RC demonstrated in the
207 current study indicate a balanced motor strategy with similar contribution from both muscle
208 groups for both stability (maintaining central position of the humeral head within the glenoid)
209 and dynamic mobility of the GHJ in abducted positions. In order to counterbalance the
210 impact of AD and MD activation on superior translation of the humeral head during shoulder
211 abduction,⁵ SUBS and ISP activation generates an inferior force which serves to minimize the
212 risk of subacromial impingement.³⁰

213 While standing IR at 90°ABD effectively activated both deltoid and RC muscles and may
214 have functional advantages by replicating overhead and sport-specific positions,³¹ the blend
215 of abduction and rotation can impose high levels of stress on shoulder's ligaments and
216 capsulolabral complex.²⁵ In the presence of RC pathology it is important to select exercises
217 that generate high RC activation with minimal deltoid involvement. Hence, IR at 0°ABD
218 with low-to-moderate activation of muscles may be considered in individuals who are at risk
219 or suffering from shoulder complex injuries particularly impingement syndrome.

220 Previous researchers have placed an emphasis on SUBS activity during IR exercises
221 particularly in relation to other large muscles involved in glenohumeral IR such as PM and
222 LD.^{22,23} It has been suggested that SUBS action during IR at 0°ABD is assisted by PM, LD,
223 and TM. While EMG activation differences between high- and low skill pitchers has
224 demonstrated the importance of SUBS conditioning (strength and endurance) in enhancing
225 pitching ability and preventing injury,³² the optimal position for selective activation of SUBS
226 for muscle strengthening and strength testing remains unclear.³³ In addition to its role as
227 internal rotator of humerus,²⁷ according to EMG studies of sport-specific activities SUBS also
228 acts as shoulder abductor, anterior stabiliser, and humeral head depressor.^{26,28,33,34} While
229 some authors reported greater SUBS activity in IR at 90°ABD,³⁵ others found greater
230 activation at 0°ABD.²² Based on three dimensional (3-D) biomechanical studies, SUBS

231 maximal force generation during IR at 90°ABD and 0°ABD is 1725N and 1297N,
232 respectively²⁸ which is consistent with the current finding of higher SUBS activation at
233 90°ABD compared to 0°ABD.

234 While previous authors have recommended SUBS strengthening exercises in adducted
235 positions,³⁶ significantly higher activation of SUBS along with low-to-moderate activation of
236 PM, LD, and, TM in IR at 90°ABD as demonstrated in the present study, suggest the
237 preference of this exercise for selective SUBS activation. In an EMG study of IR at various
238 positions, Suenaga et al²⁴ demonstrated high activation of LPM and UPM during resistive IR
239 at 0°ABD compared to other positions. Decker et al²² also demonstrated higher levels of PM
240 and LD activation IR at 0°ABD compared to 90°ABD and suggested that IR at 90°ABD may
241 be beneficial in strengthening the SUBS due to minimizing the contributions of larger muscle
242 groups.

243 **Scapular Muscles**

244 Effective scapular muscle function is fundamental for maximized performance in both daily
245 activities and overhead sports such as the volleyball serve and spike, the tennis serve, and
246 baseball pitching.^{17,34} Furthermore, current suggestions regarding the role of impaired
247 scapular motions (e.g. aberrant muscle activation patterns and fatigue) in developing a
248 dysfunctional shoulder complex and subsequent injury highlights the importance of
249 integrating scapulothoracic musculature into shoulder complex rehabilitation programs.^{6,37}
250 Amongst scapular muscles that predominantly control synchronized scapular motion during
251 arm movements, the present study assessed three parts of trapezius (UT, MT, and LT), SA,
252 and RHOM major.

253 The main functions of the trapezius include upward rotation and elevation (UT), retraction
254 (MT), and upward rotation and depression (LT) of the scapula. Importantly, LT activation

255 supports posterior tilt and ER of the scapula during arm elevation which consequently
256 decreases the risk of subacromial impingement.³⁸ The main body of existing literature
257 focuses on trapezius activity during ER and sparse data are available regarding activity
258 during IR exercises. While UT activation was found to be highest in IR at Zero-Position, MT
259 and LT had their highest activation in IR at 90°ABD. It is clinically important to enhance the
260 LT/UT and MT/UT activation ratios as a dominant UT (as compared to the other portions of
261 the trapezius) has been linked to shoulder pathologies due to contributions of poor posture
262 and muscle imbalances.⁶ Hence, the current findings support IR at 90°ABD as the more
263 advantageous exercise to enhance the LT/UT and MT/UT activation ratios over the other two
264 studied exercises. This recommendation is in agreement with other authors who have
265 reported relatively high MT activity during arm positions of 90° abduction and higher^{2, 22} but
266 not with those of Moseley et al¹¹ who reported low EMG activity of the MT during IR at
267 90°ABD. Higher LT activation in IR at 90°ABD is also consistent with previous reports of its
268 increased activity from 90° to 180°.^{2, 11}

269 Contribution of the SA to upward rotation, posterior tilt, and ER rotation of the scapula
270 during arm elevation is important for preserving a healthy scapulohumeral rhythm.^{2, 39} In the
271 presence of a dysfunctional SA, an overactive UT may cause abnormal scapular motion
272 (extreme scapular elevation and anterior tilt) and lead to muscle imbalance and functional
273 shoulder impairment.^{2, 6, 7, 39} In the presence of scapular muscle imbalances such as
274 disproportionate UT/SA activation/strength ratio, emphasis has been placed upon the
275 selective activation of underactive muscles with the minimal involvement of hyperactive
276 muscles for balance restoration.⁶ The authors' observed noticeably higher activation of SA in
277 IR at Zero-Position followed by IR at 90°ABD which represent a similar activation pattern to
278 UT during the same exercises. While IR at Zero-Position may enhance scapular function in
279 healthy athletes by mirroring shoulder positioning and motion patterns occurring during

280 overhead and throwing performance,⁴⁰ it may need to be avoided in those with or at risk of
281 subacromial impingement due to increased UT/SA activation ratio. While higher activation of
282 SA during IR at elevated and abducted arm positions has been reported by previous authors²³,
283 ⁴¹ there is a lack of information regarding IR at Zero-Position.

284 RHOM contributes to scapular retraction, downward rotation, and elevation of scapula. In
285 general, there is limited information on RHOM activation during shoulder exercises mainly
286 because of technical complications in positioning intramuscular electrodes. It is suggested
287 that several exercises used for the training and strengthening RC and other scapular muscles
288 such as ER at 0°- and 90°ABD and prone horizontal abduction at 90°ABD with IR also
289 efficiently provoke RHOM activity.^{6,23} The results of the present study demonstrated
290 markedly higher activation of RHOM activation in IR 90°ABD when compared to the other
291 exercises. This is in agreement with findings of Myers et al⁴¹ who reported relatively high
292 RHOM activity during the same exercise.¹¹

293 **Technical Considerations and Study Limitations**

294 The authors of the current study attempted to overcome inherent limitations of EMG and
295 maximize the reliability of findings. Broad experience with shoulder girdle EMG informed
296 accurate electrode positioning for optimal electrode positioning and EMG recording. EMG
297 studies have employed alternative normalization methods such as the use of MVC to study
298 muscle activation, however, use of an isometric contraction remains questionable particularly
299 in relation to studying dynamic movements.⁴²⁻⁴⁵ Hence, in view of conflicting opinions and
300 uncertainties surrounding the reliability of MMTs and related MVC for EMG amplitude
301 normalization,⁴² the present study reported each muscle's EMG activity (mean RMS) during
302 each IR exercise as a percentage of a reference value, i.e. EMGmax in a standard IR position,
303 allowing appropriate assessment and comparison of each muscles' contribution across the
304 exercises. A similar method has been applied by previous authors (e.g. maximum sprinting
305 for normalizing the EMG during walking, maximum sprint cycling for normalizing the EMG
306 during cycling).⁴³⁻⁴⁵ This normalization method may have advantages for the examination of
307 relative muscle function around the shoulder complex by minimizing intrinsic limitations in
308 reliability and validity associated with communal reference to MVC as there is no consensus
309 as to which test generates maximal activation in all individuals in any given muscle.⁴⁶⁻⁴⁸
310 While this normalization approach produced large EMG % values for some of the muscles, it
311 was deemed appropriate for comparing activity of each individual muscle across the IR
312 exercises (between-exercise comparison) as the reference value is task dependent. However,
313 it may not be the preferred method for comparing activations between the muscles (between-
314 muscle comparison) as maximum force production during the task used for normalization
315 does not necessarily produce a maximum activation in the muscles under investigation.

316 Muscle activations during IR exercises were examined using a single load (1kg) in hand or
317 against resistance from an elastic band in order to gain further insight into functional roles of
318 the muscles contributing to glenohumeral stability. According to studies by other authors,
319 increasing load does not alter shoulder muscle recruitment patterns and the functional role of
320 muscles does not change with higher muscle activity levels associated with increased
321 loads.^{21,49,50} Considering the task-specific nature of shoulder muscle function, muscle
322 recruitment strategy for a particular task such as IR is not expected to change with increasing
323 resistance/load due to a systematic increase in the activity of all shoulder muscles involved in
324 generating IR torque.^{21, 49} However, applying different loads might have provided a greater
325 information regarding the contribution of each muscle to maintaining glenohumeral stability
326 when performing exercises. The clinical implications of current study findings with regard to
327 symptomatic subjects are limited as this study included only asymptomatic participants.
328 Finally, the use of arm support or placement of a rolled towel in the axilla for isolating or
329 certain muscles without simultaneous deltoid activation was not considered in this study. This
330 is particularly important for the focused rehabilitation of RC where minimal activation of the
331 deltoid is desirable.

332 **Conclusion**

333 Activation patterns of 16 muscles/muscle sub-regions were reported during three common IR
334 exercises in order to provide descriptive data regarding their activation. Despite the fact that
335 coactivation of deltoid and RC muscles standing IR at 90°ABD may provide a functional
336 advantage by mirroring shoulder position and soft tissue mechanics (e.g. capsular strain and
337 muscle fiber length-tension relationships) during overhead activities and sports, it can place
338 high levels of stress on shoulder's tissues. Hence, IR at 0°ABD which generates low-to-
339 moderate activation of muscles may be preferred in the rehabilitation of the individuals at risk
340 or affected by shoulder injuries. Considering the current emphasis on the SUBS activity

341 during IR exercises, findings of markedly higher activation of SUBS along with low-to-
342 moderate activation of PM, LD, and, TM in IR at 90°ABD support the use of this exercise for
343 selective SUBS activation. Considering the significance of incorporating scapular muscles
344 into training and rehabilitation programs by means of enhanced LT/UT and MT/UT activity
345 ratios, the current findings support the use of IR at 90°ABD for such purposes.

346 **REFERENCES**

- 347 1. Cain PR, Mutschler TA, Fu FH, et al. Anterior stability of the glenohumeral joint. A dynamic
348 model. *Am J Sports Med.* 1987;15:144-148.
- 349 2. Ekstrom RA, Donatelli RA, Soderberg GL. Surface electromyographic analysis of exercises for
350 the trapezius and serratus anterior muscles. *J Orthop Sports Phys Ther.* 2003;33:247-258
- 351 3. Reinold MM, Gill TJ, Wilk KE, et al. Current concepts in the evaluation and treatment of the
352 shoulder in overhead throwing athletes, part 2: injury prevention and treatment. *Sports Health.*
353 2010;2:101-115.
- 354 4. Wilk KE, Andrews JR, Arrigo CA. Preventive and Rehabilitative Exercises for the Shoulder &
355 Elbow. Edited by Birmingham, AL, American Sports Medicine Institute, 2001.
- 356 5. Ackland DC, Pak P, Richardson M, et al. Moment arms of the muscles crossing the anatomical
357 shoulder. *J Anat.* 2008;213:383-390.
- 358 6. Cools AM, Dewitte V, Lanszweert F, et al. Rehabilitation of scapular muscle balance: which
359 exercises to prescribe? *Am J Sports Med.* 2007;35:1744-1751.
- 360 7. Cricchio M, Frazer C. Scapulothoracic and scapulohumeral exercises: a narrative review of
361 electromyographic studies. *J Hand Ther.* 2011;24:322-333.
- 362 8. Mihata T, Gates J, MCGarry MH, Lee J, et al. Effect of rotator cuff muscle imbalance on forceful
363 internal impingement and peel-back of the superior labrum: a cadaveric study. *Am J Sports Med.*
364 2009;37, 2222-7.
- 365 9. Gerber C, Sebesta A. Impingement of the deep surface of the subscapularis tendon and the
366 reflection pulley on the anterosuperior glenoid rim: a preliminary report. *J Shoulder Elbow Surg.*
367 2000;9, 483-90.
- 368 10. Habermeyer, P, Magosch, P, Pritsch, M, et al. Anterosuperior impingement of the shoulder as a
369 result of pulley lesions: a prospective arthroscopic study. *J Shoulder Elbow Surg.* 2004;13, 5-12.
- 370 11. Moseley JB, Jobe FW, Pink M, et al. EMG analysis of the scapular muscles during a shoulder
371 rehabilitation program. *Am J Sports Med.* 1992;20:128-134.

- 372 12.ISEK. Standards for Reporting EMG Data. *Journal of Electromyography and Kinesiology*.
373 2014;24:I-II.
- 374 13.SENIAM. Surface ElectroMyoGraphy for the Non-Invasive Assessment of Muscles. Available at:
375 <http://www.seniam.org> Accessed August 2015.
- 376 14.Basmajian JV, De Luca CJ. Muscles alive: their functions revealed by electromyography. Edited
377 by Baltimore, Williams & Wilkins, 1985.
- 378 15.Saha AK. Dynamic stability of the glenohumeral joint. *Acta Orthop Scand*. 1971;42:491-505.
- 379 16.Liotard JP, Walch G. Rehabilitation of the unstable shoulder pp. 61-72 Edited by Puddu G,
380 Giombini A, Selvanetti A. Berlin Springer-Verlag, 2001, p.62.
- 381 17.Escamilla RF, Andrews JR. Shoulder muscle recruitment patterns and related biomechanics during
382 upper extremity sports. *Sports Med*. 2009;39:569-590.
- 383 18.Ellenbecker TS, Mattalino AJ. Concentric isokinetic shoulder internal and external rotation
384 strength in professional baseball pitchers. *J Orthop Sports Phys Ther*. 1997;25:323-328.
- 385 19.Wilk KE, Andrews JR, Arrigo CA, et al. The strength characteristics of internal and external
386 rotator muscles in professional baseball pitchers. *Am J Sports Med*. 1993;21:61-66.
- 387 20.Blackburn TA, McLeod WD, White B, et al. EMG analysis of posterior rotator cuff exercises. *Athl*
388 *Train*. 1990;25:40-45.
- 389 21.Dark A, Ginn KA, Halaki M. Shoulder muscle recruitment patterns during commonly used rotator
390 cuff exercises: an electromyographic study. *Phys Ther*. 2007;87:1039-1046
- 391 22.Decker MJ, Tokish JM, Ellis HB, et al. Subscapularis muscle activity during selected rehabilitation
392 exercises. *Am J Sports Med*. 2003;31:126-134
- 393 23.Reinold MM, Escamilla RF, Wilk KE. Current concepts in the scientific and clinical rationale
394 behind exercises for glenohumeral and scapulothoracic musculature. *J Orthop Sports Phys Ther*.
395 2009;39:105-117
- 396 24.Suenaga N, Minami A, Fujisawa H. Electromyographic analysis of internal rotational motion of
397 the shoulder in various arm positions. *J Shoulder Elbow Surg*. 2003;12:501-505.
- 398 25.Wilk KE, Arrigo CA, Andrews JR. Current concepts: the stabilizing structures of the glenohumeral
399 joint. *J Orthop Sports Phys Ther*. 1997;25:364-379.

400 26.Liu J, Hughes RE, Smutz WP, et al. Roles of deltoid and rotator cuff muscles in shoulder
401 elevation. *Clin Biomech (Bristol, Avon)*. 1997;12:32-38.

402 27.Otis JC, Jiang CC, Wickiewicz TL, et al. Changes in the moment arms of the rotator cuff and
403 deltoid muscles with abduction and rotation. *J Bone Joint Surg Am*. 1994;76:667-676.

404 28.Hughes RE, An KN. Force analysis of rotator cuff muscles. *Clin Orthop Relat Res*. 1996;75-83.

405 29.Jenp YN, Malanga GA, Growney ES, An KN. Activation of the rotator cuff in generating
406 isometric shoulder rotation torque. *Am J Sports Med*. 1996;24:477-485.

407 30.Sharkey NA, Marder RA. The rotator cuff opposes superior translation of the humeral head. *Am J*
408 *Sports Med*. 1995;23:270-275.

409 31.Fleisig GS, Barrentine SW, Escamilla RF, et al. Biomechanics of overhand throwing with
410 implications for injuries. *Sports Med*. 1996, 21:421-437.

411 32.Gowan ID, Jobe FW, Tibone JE, et al. A comparative electromyographic analysis of the shoulder
412 during pitching. Professional versus amateur pitchers. *Am J Sports Med*. 1987;15:586-590.

413 33.Jobe FW, Moynes DR. Delineation of diagnostic criteria and a rehabilitation program for rotator
414 cuff injuries. *Am J Sports Med*. 1982;10:336-339.

415 34.Jobe FW, Moynes DR, Tibone JE, et al. An EMG analysis of the shoulder in pitching. A second
416 report. *Am J Sports Med*. 1984;12:218-220.

417 35.Kadaba MP, Cole A, Wootten ME, et al. Intramuscular wire electromyography of the
418 subscapularis. *J Orthop Res*. 1992;10:394-397.

419 36.Matsen FA, Thomas SC, Rockwood CA. Glenohumeral instability. Edited by Rockwood CA,
420 Matsen FA. Philadelphia, Saunders. 1990,;pp. 526-622.

421 37.Ebaugh DD, McClure PW, Karduna AR. Scapulothoracic and glenohumeral kinematics following
422 an external rotation fatigue protocol. *J Orthop Sports Phys Ther*. 2006;36:557-571.

423 38.Graichen H, Bonel H, Stammberger T, et al. Three-dimensional analysis of the width of the
424 subacromial space in healthy subjects and patients with impingement syndrome. *Am J Roentgenol*.
425 1999;172:1081-1086.

426 39.Ludewig PM, Reynolds JF. The association of scapular kinematics and glenohumeral joint
427 pathologies. *J Orthop Sports Phys Ther*. 2009;39:90-104.

- 428 40. Puddu G, Giombini A, Selvanetti A. Rehabilitation of Sports Injuries: Current Concepts. Berlin,
429 Springer, 2001.
- 430 41. Myers JB, Pasquale MR, Laudner KG, et al. On-the-Field Resistance-Tubing Exercises for
431 Throwers: An Electromyographic Analysis. *J Athl Train.* 2005;40:15-22.
- 432 42. Halaki M, Ginn K. Normalization of EMG Signals: To Normalize or Not to Normalize and What
433 to Normalize to? Computational Intelligence in Electromyography Analysis - A Perspective on
434 Current Applications and Future Challenges 2012.
- 435 43. Albertus-Kajee Y, Tucker R, Derman W, et al. Alternative methods of normalising EMG during
436 running. *J Electromyogr Kinesiol.* 2011;21:579-586.
- 437 44. Fernández-Peña E, Lucertini F, Ditroilo M. A maximal isokinetic pedalling exercise for EMG
438 normalization in cycling. *J Electromyogr Kinesiol.* 2009;19:e162-e170.
- 439 45. Rouffet DM, Hautier CA. EMG normalization to study muscle activation in cycling. *J*
440 *Electromyogr Kinesiol.* 2008;18:866-878.
- 441 46. Cram JR, Kasman GS, Holtz J. Electrode placement. Edited by Cram JR, Kasman GS, Holtz J.
442 Gaithersburg, MD, Aspen Publishers, 1998.
- 443 47. Ha SM, Cynn HS, Kwon OY, et al. A reliability of electromyographic normalization methods for
444 the infraspinatus muscle in healthy subjects. *J Hum Kinet,* 2013;36:69-76
- 445 48. Marras WS, Davis KG. A non-MVC EMG normalization technique for the trunk musculature: Part
446 1. Method development. *J Electromyogr Kinesiol.* 2001.11:1-9
- 447 49. Boettcher CE, Cathers I, Ginn KA. The role of shoulder muscles is task specific. *J Sci Med Sport.*
448 2010;13:651-656.
- 449 50. Reed D, Halaki M, Ginn K. The rotator cuff muscles are activated at low levels during shoulder
450 adduction: an experimental study. *J Physiother.* 2010;56:259-264.