First report of an adelgid from Malta (Hemiptera, Aphidoidea, Adelgidae)

David MIFSUD1 & Nicolás PÉREZ HIDALGO2

In recent years, the aphid fauna of the Maltese Islands has received considerable attention. MIFSUD et al. (2011) reviewed earlier works on the subject and provided information on previously reported species and on several new records. An updated list of aphids provided therein included 98 Aphididae and a single species of Phylloxeridae. The only other family currently included with aphids, the Adelgidae was not represented in the list. Samples of young shoots of *Pinus halepensis* Mill, were observed infested with a whitish flocculent material (Fig. 1A) in several localities in Malta and examination of this material proved it to belong to an adelgid which we report here under.

Pineus pini (Macquart, 1819)

Material examined. MALTA: Attard, St. Anton Garden, 3.v.2012, numerous apterous colonies on shoots of *Pinus halepensis*, leg. D. Mifsud; Zejtun, 30.xi.2011/1.v.2012, numerous apterous colonies on shoots of Pinus halepensis, leg. D. Mifsud.

Using the dichotomous keys of BLACKMAN & EASTOP (2010), the adelgid was identified as Pineus pini (Figs. 1B, 1C, 1D), commonly referred to as the pine woolly aphid. This species is anholocyclic on Pinus sylvestris and Pinus mugo in western and central Europe, with an overwintering (hiemosistens) generation on the twigs and two or more overlapping generations of progredientes attacking the current year's shoots. The second progrediens generation in May-June includes alatae (Blackman & Eastop, 2010). According to Blackman & Eastop (2010) the morphs of P. pini are almost indistinguishable from the equivalent secondary host generations of *P. orientalis* (Dreyfus), and it seems probable that the anholocyclic P. pini arose from the holocyclic P. orientalis by loss of a functional sexual generation (see references in Blackman & Eastop, 2010). Pineus orientalis is host-alternating between certain *Picea* spp. and various *Pinus* spp. of the subsection sylvestris (densiflora, mugo, sylvestris, thunbergii) from Western Europe across Asia to Japan, but there are no species of *Picea* in Malta. *Pineus pini* is present in Italy (including Sicily) and in many other countries in Europe and the Mediterranean area (Nieto Nafría et al., 2004).

A revision of this species complex is required because in the Australian literature populations of P. pini group have in the past been given the name of P. laevis Maskell and confused with P. boerneri Annand, which also occurs on a range of pines in both Australia and New Zealand.

ACKNOWLEDGEMENTS

We would like to thank Robert G. Foottit of the Canadian National Collection of Insects, Ontario for reviewing the present work and for all suggestions.

¹ Department of Biology, University of Malta, Junior College, Msida MSD 1251, Malta. E-mail: david.a.mifsud@ um.edu.mt

² Department of Biodiversity and Environmental Management, University of Leon, E-24071, León, Spain. E-mail: nperh@unileon.es




Figure 1: *Pineus pini*. **A**: Colony on *Pinus halepensis*; **B**: habitus of adult (from microscopic preparation); **C**: detail of cephaloprothoracic shield; **D**: detail of wax glands comprising groups of 2-10 facets.

REFERENCES

BLACKMAN, R.L. & EASTOP, V.F. (2010) Aphids on the World's Trees - An identification and information guide. Second edition. Published online at http://www.aphidsonworldsplants.info/x APHIDS P.htm#Pineus [accessed 5th August 2012].

MIFSUD, D., MANGION, M., AZZOPARDI, E., ESPADALER, X., CUESTA-SEGURA, D., WATSON, G.W. & PÉREZ HIDALGO, N. (2011) Aphids associated with shrubs, herbaceous plants and crops in the Maltese Archipelago (Hemiptera: Aphidoidea). *Bulletin of the Entomological Society of Malta*, 4: 5-53.

Nieto Nafría, J.M., Andreev, A.V., Binazzi, A., Mier Durante, M.P., Pérez Hidalgo, N., Rakauskas, R. & Stekolshchikov, A.V. (2004) Aphidoidea. Fauna Europaea version 2.4. http://www.faunaeur.org [accessed 20th August 2012].

Received: August 10, 2012 Accepted: September 20, 2012