

Review Article

Molecular mechanisms in haematological malignancies

Godfrey Grech, Roberto Avellino, Pierre Schembri Wismayer

Abstract

Haematopoiesis requires the constant production of large numbers of peripheral blood cells. This process is under tight control of transcription factor networks as well as cytokines, growth factors and hormones. We will review the importance of transcription factors in programming the haematopoietic lineage commitment and the role of the microenvironment and the corresponding cellular sensitivity to ensure production of mature functional cells in response to the physiological demand. Understanding the molecular mechanism of this complex process gives the opportunity to identify the underlying molecular deregulation in haematopoietic malignancies. The different levels of deregulation include hyperproliferation, block in differentiation and sensitivity to growth factors. In this review, leukaemic transformation is selected to give evidence of cell signalling deregulation. The clinical implications will be reviewed in the context of the potential opportunities in the future to identify specific therapeutic patient groups that can be defined using prognostic and predictive biomarkers.

Keywords

Haematopoiesis, haematological malignancies, leukaemia, cytogenetic aberrations, transcription factors, molecular genetics

Godfrey Grech* PhD, MMCPATH
Department of Pathology, Medical School, University of Malta
Email: godfrey.grech@um.edu.mt

Roberto Avellino BSc (MLS), MRes
Department of Pathology, Mater Dei Hospital, Malta

Pierre Schembri Wismayer MD, PhD
Department of Anatomy, Biomedical Sciences Building,
University of Malta, Malta

* corresponding author

Introduction

The quiescent haematopoietic stem cell (HSC) compartment maintains a multipotent cell population and gives rise to oligolineage progenitors. These progenitors expand to maintain the haematopoietic compartment and differentiate into various blood lineage progenitors. Lineage positive progenitors are committed for differentiation into mature functional blood cells.

Lineage commitment in haematopoiesis

Transcription factors have a pivotal role in haematopoiesis and regulate HSC early development, survival, proliferation and lineage commitment. Transcription factors, including Runx1, SCL, Gata-2, ALL-1 and Evi-1 maintain a gene expression program unique to HSCs.¹ The maintenance of HSC self-renewal requires specific transcription factors among which are HoxB4², Notch1³ and Bmi-1.⁴

Oligolineage progenitors originate from the HSC and develop into either the common myeloid precursor (CMP) or the common lymphoid precursor (CLP), in the latter case through the down regulation of PU.1.⁵ The CMPs undergo further lineage divergence into megakaryocytic/erythroid progenitors (MEPs) and granulocytic/monocytic progenitors (GMPs) upon Gata-1 and PU.1 mutually exclusive expression, respectively. C/EBPs, including C/EBP α and C/EBP ϵ play an important role in the granulocytic branch of myelomonocytic commitment both through cell-cycle arrest⁶ and the up-regulation of tissue specific genes.⁷ Under the influence of PU.1, GATA-3 and Ikaros transcription factors, HSCs give rise to CLPs.⁸ Upregulation of such transcription factors determine the commitment of the lymphoid lineage and subsequently the downregulation of myeloid lineage transcription factors such as GATA-1, C/EBP α , and NFE2.⁹

Differentiation of lineage committed progenitors

Upon commitment to the lymphoid lineage, lymphocyte progenitors start their maturation in the bone marrow and differentiate either as mature T cells in the thymus gland or as mature B cells in germinal centres (GCs) of secondary lymphoid organs. Lymphocyte fate is under tight regulation of mutually exclusive transcription factors and signalling pathways, which force CLP differentiation into a pro-B cell or a pro-T cell. Increased expression levels of E2A, EBF, Bcl11a¹⁰ and PAX5^{11,12} promote CLP differentiation to pro-B cells, pre-B cells and further to 'naive' B cells by the participation of the

SOX4¹³ and LEF1¹⁴ transcription factors. Consistently, PAX5 acts as a repressor by recruiting co-repressors to genes encoding NOTCH1¹⁵, therefore direct differentiation towards the B cell lineage.

The microenvironment and the cellular sensitivity to the surrounding cytokines and growth factors allows appropriate differentiation to various mature cells in response to the physiological demand. Upon commitment, progenitors migrate to the specific microenvironment and are programmed to express lineage specific receptors, as governed by the transcription program, altering the sensitivity to the microenvironment. Lineage specific cytokines and growth factors regulate maintenance and differentiation of the committed progenitor cells. For instance, the differentiation towards the T cell lineage is promoted via the migration of CLPs to the thymus gland. The developmental progression of T cells is initiated by the downregulation of genes encoding KIT, CD44 and CD25.¹⁶ This proceeds further to the upregulation of genes encoding the NOTCH proteins^{17,18} and the response to cytokines and growth factors, such as IL-7 and morphogens, such as the Sonic Hedgehog proteins.¹⁹

Stem Cell Factor (SCF) supports proliferation of various haematopoietic compartments. In erythropoiesis the lineage-specific cytokine erythropoietin (Epo) works in concert with SCF to regulate the balance between proliferation and differentiation of erythroid progenitors. Epo and SCF transduce signals via multiple cooperating pathways in erythroid progenitors.^{20,21} Activation of Protein Kinase B (PKB) results in phosphorylation of transcription factor Foxo3a, which results in its cytoplasmic retention and inhibition of transcriptional activation of cell cycle progression inhibitors, such as p27, p130Rb2, Btg1 and cyclin G2.²² PKB, also activates the mTOR/eIF4E pathway²³ resulting in enhanced translation efficiency of structured mRNAs representing growth promoting transcripts (Figure 1). Decreased SCF signalling results in downregulation of PI3K, releasing active Foxo3a that promotes expression of cell cycle inhibitors initiating signals that execute terminal differentiation.

Dysregulation of haematopoiesis

The balance between proliferation and differentiation of committed progenitors is under tight control, to maintain the progenitor pool and ensure maturation in response to physiological demand. The production of increased numbers of mature blood cells during stress, requires higher progenitor proliferation rates. Concurrently, feedback mechanisms must be closely coordinated to repress progenitor proliferation and to restore physiological cell numbers when the stress is over.²⁴ Deregulation of this balance will result in disease.

Hyperproliferation

Myelo-Proliferative Disorders (MPD) and anaemia originate from hyperproliferative potential that can be sustained or can result in bone marrow exhaustion, respectively. Constitutive active tyrosine kinase receptors (FLT-3, cKit), activated kinases like activated Jak2 and mutant phosphatase PTEN promote proliferation and/or survival in committed progenitors that are

still capable of differentiation. This results in a hyperproliferative phenotype.^{9,25,26}

A common mutational event in lymphoid malignancies is the juxtaposition of genes to the BCR or TCR enhancer/promoter elements. One classical example is the BCL1 protein or CyclinD1 in the t(11;14) anomaly, where the BCL1 protein is juxtaposed to the Ig gene enhancer. The BCL1 protein is a cell cycle regulator required for the cellular maintenance of G1 progression and G1/S transition of the cell cycle. The enhancers of the Ig and BCL1 gene are exchanged leading to the possibility of overexpression of BCL1 gene with an accelerated passage through the G1 phase of the cell cycle.²⁸ Other similar examples are seen in mature B cell malignancies such as the t(14;18) in Follicular lymphomas²⁹, the t(1;14) in MALT lymphomas³⁰ and the t(8;14) in Burkitt lymphomas.³¹ Table 1 summarises the translocations resulting in dysregulation of proteins involved in cell cycle regulation and apoptosis. Dysregulation is a result of promoter exchange upon juxtaposition of one gene to another.

Differentiation block

In addition to hyperproliferation, a perturbed program blocking terminal differentiation plays a role in the development of leukaemia. Experimental reduction of the terminal differentiation transcription factor PU.1, results in leukaemia following failed differentiation.⁵ Translocations t(8;21) and t(15;17) give rise to the fusion proteins AML1/RUNX1 and PML-RAR α , both of which inhibit haematopoietic differentiation by recruiting repression complexes to target genes of the AML1 and RAR α transcription factors respectively.^{32, 33} This is also observed in early transformed lymphoid cells where a block in differentiation occurs either by the involvement of a transcription factor in the generation of a fusion gene or by promoter exchange with the BCR or TCR enhancer loci. The translocation t(1;19), encoding the E2A-PBX fusion protein, blocks the activity of the transcription factor E2A contributing to the onset of pre-B ALL.³⁴ In a recurrent chromosomal abnormality detected in T-ALL patients, t(1;14), the TAL1 gene is brought under the influence of the TCRA/D enhancer, deregulating the expression of the transcription factor TAL1 and blocks the T cell differentiation pathway giving rise to the onset of T-ALL.³⁵ Table 2 summarises recurrent chromosomal translocations found in leukemias that involve transcription factors.

Expression of fusion proteins in haematopoietic progenitors confers propagation in serial murine transplantation models^{36,37} but is not sufficient to induce leukaemia. Cooperation events exemplified by the complementation of Flt3-ITD mutant to PML-RAR α transgenic bone marrow cells results in 100% penetrance of an APL-like disease when transplanted to secondary recipients.³² Moreover, chromosomal translocations such as the t(9;22)³⁸, t(14;18)³⁹ that are detected in CML and NHL respectively, have been also detected in normal and healthy individuals without any evidence of malignancy. Supporting this evidence is the t(12;21) in monozygotic twins. Studies have shown that although both twins may carry the t(12;21) anomaly, only the one with the additional mutation, 12p deletion, develops

precursor B acute lymphoblastic leukaemia.⁴⁰ This provides strong evidence that a second hit mutation, in addition to any other present mutation, is required for leukaemogenesis. It is in fact this combination of a proliferative switch together with a differentiation-blocking event which usually results in hematopoietic malignancy.

Growth factor sensitivity in disease

The high demand of erythrocytes in circulation is satisfied by continuous high level of erythroid progenitor expansion and differentiation in the bone marrow. Hence, negative feedback of EpoR signalling is required to prevent erythrocytosis. EpoR mutations resulting in truncations in the C-terminal region lack recruitment of the phosphatase SHP-1 resulting in constitutive proliferative signals and hypersensitivity to Epo.⁴¹ These mutants are associated with primary familial polycythemia⁴² with normal cell maturation and increased blood cell production. Hypersensitivity can be a result of loss of negative feedback but

also be caused by activating mutations resulting in enhanced proliferation and/or survival. For instance the V617F *JAK2* mutation⁴³ is found predominantly in Polycythemia Vera (PV) patients, a myeloproliferative disorder (MPD) characterized by massive erythrocytosis. The mutation resides in the kinase inhibitory domain, resulting in constitutive *JAK2* kinase activity. Pro-proliferative signal transduction pathway mutants have a central role in the pathogenesis of MPD and require cooperative events that lead to leukaemia. In human disease, constitutive activating mutations in *FLT3* are found in 30-35% of adult AML, *N-RAS* and *K-RAS* mutations in 20% and *cKIT* mutations account for 5% of cases,^{27,44} supporting the hypothesis that tyrosine kinase receptor mutations represent collaborative events in leukaemogenesis together with loss or gain of function mutations in haematopoietic transcription factors such as AML1, GATA-1, C/EBP α and PU1. In addition to mutations that deregulate kinases, a number of translocations involving kinase molecules, are summarised in Table 3.

Table 1: Dysregulation of proteins involved in cell cycle regulation and apoptosis. In these translocations, fusion genes are not generated but dysregulation is brought about by promoter exchange upon juxtaposition of one gene to another

Chromosomal Translocation	Gene involved	Function of deregulated gene	Associated Phenotype/disease	References
t(14;18)(q32;q21)	BCL2	Apoptotic inhibition	Follicular lymphoma	29
t(11;14)(q13;q32)	BCL1	Overexpression accelerates passage through the G1 phase	Mantle cell lymphoma	28
t(1;14)(q21;q32)	BCL9	Apoptosis inhibition through NF- κ B	Mucosa associated lymphoid tissue	30
3q27 rearrangements	BCL6	Bind to sequence specific DNA and repress its transcription in addition to recruiting other protein repressors; Cell cycle control; Apoptosis inhibition	Diffuse Large B cell Lymphoma	59
t(8;14)(q24;q32)	c-MYC	Constitutive activation of c-myc	Burkitt lymphoma	31

Table 2: Transcription factors commonly involved in recurrent chromosomal translocations in leukemias

Chromosomal Translocation	Fusion gene	Fusion Protein – oncogenic function	Associated phenotype	References
T(8;21)(q22;q22)	AML1-ETO	Recruitment of repression complex; negative dominance of ETO on AML1	AML - FAB M2 subtype	53
T(12;21)(p12;q22)	TEL-AML1	Recruitment of repression complex; negative dominance of TEL on AML1	B-ALL	54
T(15;17)(q22;q21)	PML-RARA	Dysregulation of retinoid-inducible genes involved in myeloid differentiation; Higher binding affinity for HDACs	AML – FAB M3 subtype	55
T(16;21)(p11;q22)	TLS-ERG	Seems to act as a transcriptional activator	de novo acute myeloid leukemia	56
T(1;19)(q23;p13)	E2A-PBX1	Potent transcriptional activator with pleiotropic transforming activity	pre B ALL	57
T(3;21)(q26;q22)	AML1-EVI1	Chimeric transcription factor with the dual functions of AML1 and EVI1: differentiation block (due to Runt) and stimulation of proliferation (from the zn fingers)	CML-BC of myeloid type; AML and MDS	58

Conversely, in myelodysplastic syndrome (MDS) lack of circulating erythrocytes occurs due to impaired responsiveness to Epo⁴⁵ or aberrant response to inhibitory cytokines. Epo stimulation of erythroid progenitors derived from MDS bone marrow fail to induce DNA-binding of transcription factor Stat5,⁴⁵ suggesting that dysplastic cells result from maturation commitment without the capacity to drive the terminal differentiation program.

Signalling deregulation in leukaemia transformation

Mutations that enhance the translation machinery also play a central role in enhanced aggressiveness of various human cancers including AML.⁴⁶ The D816V mutation in the kinase domain of cKit activates the PI3K/PKB/mTOR pathway conferring sensitivity to rapamycin.⁴⁷ Interestingly, rapamycin induces cell cycle arrest and apoptosis in patient-derived neoplastic mast cells harbouring the D816V *cKIT*, but not in

Table 3: Table showing deregulated kinases involved in chromosomal translocations contributing to the onset of leukemia

Chromosomal Translocation	Protein kinases	Fusion gene	Associated Phenotype	Reference
t(9;22)(q34;q11)	ABL	BCR - ABL	CML; ALL; AML	60
t(9;12)(q34;p13)	ABL	TEL - ABL	CML; ALL; AML	61
t(9;12)(p24;p13)	JAK2	TEL - JAK2	T-ALL; CD10+-ALL; 'CML-like' disease in transformation	62
t(9;22)(p24;q11.2)	JAK2	BCR-JAK2	Typical CML	63
t(2;5)(p23;q35)	ALK	NPM-ALK	Anaplastic large cell lymphoma	64
t(5;12)(q33;p13)	PDGFR	TEL-PDGFR	CMML	27
t(12;15)(p13;q25)	NTRK3	TEL-NTRK3	AML - FAB M2 subtype	65
				66




Figure 1: Stem cell factor (SCF) signalling pathway. SCF binds to the tyrosine kinase receptor, cKit, activating the PI3K/PKB signal transduction pathway. PKB phosphorylates and redistributes Fox3a, sequestering the molecule from the nucleus, inhibiting transcription of cell cycle inhibitors. In addition, activation of mTOR results in the translation of growth factors (eIF4E sensitive mRNA transcripts) that are otherwise not recruited to polysomes

normal human cord-blood derived mast cells.⁴⁷ The frequency of active PI3K is higher than the incidence of mutations in *RAS* or in the receptor tyrosine kinases (RTK) *FLT3* and *cKIT*,⁴⁸ suggesting the need for targeting PI3K downstream effectors. Phosphatases that attenuate kinase activity are eligible for targeting. The phosphatases, PTEN and pp2a attenuate the PI3K/mTOR pathway, directly affecting ribosome biosynthesis and translation initiation.^{23,49} The leukaemic potential of BCR/ABL-expressing cells can be inhibited by pharmacological activation of the phosphatase pp2a.⁵⁰ This suggests a central role of deregulated PI3K/mTOR/translation machinery in chronic myeloid leukemia (CML).

Conclusion and Further Directions

The knowledge of the molecular mechanism underlying specific groups of patients gives the opportunity to identify specific therapeutic targets that can be treated with known, or offer the opportunity to develop, specific pharmaceutical agents. To achieve this goal, patients are required to be classified into therapeutic groups. As mentioned above, activation of the PI3K pathway by various molecular aberrations classifies patients into a risk group that is sensitive to rapamycin treatment. Patients with cytogenetic abnormalities *inv(16)* (p13q22), and *t(8;21)* (q22;q22) are classified as core binding factor (CBF) leukemias (Table 2). These patients are represented into both the good and poor prognostic groups. Poor prognosis is driven by mutations in the *cKit* gene.⁵¹ The identification of *cKit* mutations in CBF leukaemia cases is today used as a predictive biomarker strategy that will allow the use of specific tyrosine kinase inhibitors, such as Imatinib, in the treatment of such cases. Specific tyrosine kinase inhibitors are also available for *Jak2* positive myeloproliferative disorders (MPD); and ABL translocations (Table 3). Although still being tested in clinical trials, FLT3 inhibitors are showing promising results in AML patients that carry FLT3 mutations (approximately 30% of AML patients).⁵² The BCR/ABL product of the translocation *t(9;22)*(q34;q11) is efficiently and directly targeted by the tyrosine kinase inhibitor, Imatinib. Molecular studies of the acquired mutations in the BCR/ABL molecule give predictive information on resistance to therapy. Knowledge of these mutations has been used to develop new therapeutic drugs to minimise the burden of resistance to therapy in CML patients.

Hence, patients will benefit from the identification of additional biomarkers that could predict clinical outcome at diagnosis. In addition, prognostic markers that can potentially serve as targets for novel agents are continuously investigated. Taken together, this knowledge will ensure informed therapeutic decisions and allow clinicians to use specific targeted therapies.

Acknowledgements

We would like to acknowledge Dr Marieke von Lindern (Hematology Department, Erasmus Medical Centre, Rotterdam, the Netherlands) for her constructive review of main parts of this manuscript. We are also thankful to Prof Denis Alexander (Haematology Department, Belfast City Hospital) for the exposure to current research in lymphoid malignancies.

References

- Phillips RL, Ernst RE, Brunk B, Ivanova N, Mahan MA, Deanehan JK, et al. The genetic program of hematopoietic stem cells. *Science*. 2000;288:1635-40.
- Sauvageau G, Thorsteinsdottir U, Eaves CJ, Lawrence HJ, Largman C, Lansdorp PM, et al. Overexpression of HOXB4 in hematopoietic cells causes the selective expansion of more primitive populations in vitro and in vivo. *Genes Dev*. 1995;9:1753-65.
- Varnum-Finney B, Xu L, Brashem-Stein C, Nourigat C, Flowers D, Bakkour S, et al. Pluripotent, cytokine-dependent, hematopoietic stem cells are immortalized by constitutive Notch1 signaling. *Nat Med*. 2000;6:1278-81.
- Park IK, Qian D, Kiel M, Becker MW, Pihalja M, Weissman IL, et al. Bmi-1 is required for maintenance of adult self-renewing haematopoietic stem cells. *Nature*. 2003;423:302-5.
- DeKoter RP, Kamath MB, Houston IB. Analysis of concentration-dependent functions of PU.1 in hematopoiesis using mouse models. *Blood Cells Mol Dis*. 2007;39:316-20.
- Johnson PF. Molecular stop signs: regulation of cell-cycle arrest by C/EBP transcription factors. *J Cell Sci*. 2005;118(Pt 12):2545-55.
- Friedman AD. Transcriptional regulation of myelopoiesis. *Int J Hematol*. 2002;75:466-72.
- Busslinger M. Transcriptional control of early B cell development. *Annu Rev Immunol*. 2004;22:55-79.
- Akashi K, He X, Chen J, Iwasaki H, Niu C, Steenhard B, et al. Transcriptional accessibility for genes of multiple tissues and hematopoietic lineages is hierarchically controlled during early hematopoiesis. *Blood*. 2003;101:383-9.
- Liu P, Keller JR, Ortiz M, Tessarollo L, Rachel RA, Nakamura T, et al. Bcl11a is essential for normal lymphoid development. *Nat Immunol*. 2003;4:525-32.
- Nutt SL, Heavey B, Rolink AG, Busslinger M. Commitment to the B-lymphoid lineage depends on the transcription factor Pax5. *Nature*. 1999;401:556-62.
- Nutt SL, Morrison AM, Dorfler P, Rolink A, Busslinger M. Identification of BSAP (Pax-5) target genes in early B-cell development by loss- and gain-of-function experiments. *Embo J*. 1998;17:2319-33.
- Schilham MW, Oosterwegel MA, Moerer P, Ya J, de Boer PA, van de Wetering M, et al. Defects in cardiac outflow tract formation and pro-B-lymphocyte expansion in mice lacking Sox-4. *Nature*. 1996;380:711-4.
- Schilham MW, Clevers H. HMG box containing transcription factors in lymphocyte differentiation. *Semin Immunol*. 1998;10:127-32.
- Souabni A, Cobaleda C, Schebesta M, Busslinger M. Pax5 promotes B lymphopoiesis and blocks T cell development by repressing Notch1. *Immunity*. 2002;17:781-93.
- Scimone ML, Aifantis I, Apostolou I, von Boehmer H, von Andrian UH. A multistep adhesion cascade for lymphoid progenitor cell homing to the thymus. *Proc Natl Acad Sci U S A*. 2006;103:7006-11.
- Barrick D, Kopan R. The Notch transcription activation complex makes its move. *Cell*. 2006;124:883-5.
- Robey E, Chang D, Itano A, Cado D, Alexander H, Lans D, et al. An activated form of Notch influences the choice between CD4 and CD8 T cell lineages. *Cell*. 1996;87:483-92.
- von Boehmer H, Aifantis I, Gounari F, Azogui O, Haughn L, Apostolou I, et al. Thymic selection revisited: how essential is it? *Immunol Rev*. 2003;191:62-78.
- von Lindern M, Zauner W, Mellitzer G, Steinlein P, Fritsch G, Huber K, et al. The glucocorticoid receptor cooperates with the erythropoietin receptor and c-Kit to enhance and sustain proliferation of erythroid progenitors in vitro. *Blood*. 1999;94:550-9.
- Wessely O, Bauer A, Quang CT, Deiner EM, von Lindern M, Mellitzer G, et al. A novel way to induce erythroid progenitor self renewal: cooperation of c-Kit with the erythropoietin receptor. *Biol Chem*. 1999;380:187-202.
- Bakker WJ, Blazquez-Domingo M, Kolbus A, Besooyen J, Steinlein P, Beug H, et al. FoxO3a regulates erythroid differentiation and induces BTG1, an activator of protein arginine methyl transferase 1. *J Cell Biol*. 2004;164:175-84.
- Blazquez-Domingo M, Grech G, von Lindern M. Translation initiation factor 4E inhibits differentiation of erythroid progenitors. *Mol Cell Biol*. 2005;25:8496-506.

24. Vattem KM, Wek RC. Reinitiation involving upstream ORFs regulates ATF4 mRNA translation in mammalian cells. *Proc Natl Acad Sci U S A*. 2004;101:11269-74.
25. Vainchenker W, Constantinescu SN. A Unique Activating Mutation in JAK2 (V617F) Is at the Origin of Polycythemia Vera and Allows a New Classification of Myeloproliferative Diseases. *Hematology Am Soc Hematol Educ Program*. 2005:195-200.
26. Kelly LM, Liu Q, Kutok JL, Williams IR, Boulton CL, Gilliland DG. FLT3 internal tandem duplication mutations associated with human acute myeloid leukemias induce myeloproliferative disease in a murine bone marrow transplant model. *Blood*. 2002;99:310-8.
27. Golub TR, Barker GF, Lovett M, Gilliland DG. Fusion of PDGF receptor beta to a novel ets-like gene, tel, in chronic myelomonocytic leukemia with t(5;12) chromosomal translocation. *Cell*. 1994; 77:307-16.
28. Bosch F, Jares P, Campo E, Lopez-Guillermo A, Piris MA, Villamor N, et al. PRAD-1/cyclin D1 gene overexpression in chronic lymphoproliferative disorders: a highly specific marker of mantle cell lymphoma. *Blood*. 1994;84:2726-32.
29. Yunis JJ, Oken MM, Kaplan ME, Ensrud KM, Howe RR, Theologides A. Distinctive chromosomal abnormalities in histologic subtypes of non-Hodgkin's lymphoma. *NEJM*. 1982;307:1231-6.
30. Willis TG, Zalberg IR, Coignet LJ, Wlodarska I, Stul M, Jadayel DM, et al. Molecular cloning of translocation t(1;14)(q21;q32) defines a novel gene (BCL9) at chromosome 1q21. *Blood*. 1998;91:1873-81.
31. Kornblau SM, Goodacre A, Cabanillas F. Chromosomal abnormalities in adult non-endemic Burkitt's lymphoma and leukemia: 22 new reports and a review of 148 cases from the literature. *Hematol Oncol* 1991;9:63-78.
32. Kelly LM, Kutok JL, Williams IR, Boulton CL, Amaral SM, Curley DP, et al. PML/RARalpha and FLT3-ITD induce an APL-like disease in a mouse model. *Proc Natl Acad Sci U S A*. 2002;99:8283-8.
33. Tallman MS, Nabhan C, Feusner JH, Rowe JM. Acute promyelocytic leukemia: evolving therapeutic strategies. *Blood*. 2002;99:759-67.
34. Dederda DA, Waller EK, LeBrun DP, Sen-Majumdar A, Stevens ME, Barsh GS, et al. Chimeric homeobox gene E2A-PBX1 induces proliferation, apoptosis, and malignant lymphomas in transgenic mice. *Cell*. 1993;74:833-43.
35. Bash RO, Crist WM, Shuster JJ, Link MP, Amylon M, Pullen J, et al. Clinical features and outcome of T-cell acute lymphoblastic leukemia in childhood with respect to alterations at the TAL1 locus: a Pediatric Oncology Group study. *Blood*. 1993;81:2110-7.
36. Higuchi M, O'Brien D, Kumaravelu P, Lenny N, Yeoh EJ, Downing JR. Expression of a conditional AML1-ETO oncogene bypasses embryonic lethality and establishes a murine model of human t(8;21) acute myeloid leukemia. *Cancer Cell*. 2002;1:63-74.
37. Grisolan JL, Wesselschmidt RL, Pelicci PG, Ley TJ. Altered myeloid development and acute leukemia in transgenic mice expressing PML-RAR alpha under control of cathepsin G regulatory sequences. *Blood*. 1997;89:376-87.
38. Bose S, Deininger M, Gora-Tybor J, Goldman JM, Melo JV. The presence of typical and atypical BCR-ABL fusion genes in leukocytes of normal individuals: biologic significance and implications for the assessment of minimal residual disease. *Blood*. 1998;92:3362-7.
39. Schuler F, Hirt C, Dolken G. Chromosomal translocation t(14;18) in healthy individuals. *Semin Cancer Biol*. 2003;13:203-9.
40. Ford AM, Bennett CA, Price CM, Bruin MC, Van Wering ER, Greaves M. Fetal origins of the TEL-AML1 fusion gene in identical twins with leukemia. *Proc Natl Acad Sci U S A*. 1998;95:4584-8.
41. Klingmuller U, Lorenz U, Cantley LC, Neel BG, Lodish HF. Specific recruitment of SH-PTP1 to the erythropoietin receptor causes inactivation of JAK2 and termination of proliferative signals. *Cell*. 1995;80:729-38.
42. Furukawa T, Narita M, Sakaue M, Otsuka T, Kuroha T, Masuko M, et al. Primary familial polycythaemia associated with a novel point mutation in the erythropoietin receptor. *Br J Haematol* 1997; 99:222-7.
43. James C, Ugo V, Le Couedic JP, Staerk J, Delhommeau F, Lacout C, et al. A unique clonal JAK2 mutation leading to constitutive signalling causes polycythaemia vera. *Nature*. 2005;434:1144-8.
44. Beghini A, Peterlongo P, Ripamonti CB, Larizza L, Cairoli R, Morra E, et al. C-kit mutations in core binding factor leukemias. *Blood*. 2000;95:726-7.
45. Hoefsloot LH, van Amelsvoort MP, Broeders LC, van der Plas DC, van Lom K, Hoogerbrugge H, et al. Erythropoietin-induced activation of STAT5 is impaired in the myelodysplastic syndrome. *Blood*. 1997;89:1690-700.
46. Longley BJ, Reguera MJ, Ma Y. Classes of c-KIT activating mutations: proposed mechanisms of action and implications for disease classification and therapy. *Leuk Res*. 2001;25:571-6.
47. Gabillot-Carre M, Lepelletier Y, Humbert M, de Sepuvela P, Ben Hamouda N, Zappulla JP, et al. Rapamycin inhibits growth and survival of D-816-V-mutated c-kit mast cells. *Blood*. 2006. 108:1065-72.
48. Cornillet-Lefebvre P, Cuccuini W, Bardet V, Tamburini J, Gillot L, Ifrah N, et al. Constitutive phosphoinositide 3-kinase activation in acute myeloid leukemia is not due to p110delta mutations. *Leukemia*. 2006;20:374-6.
49. Podsypanina K, Lee RT, Politis C, Hennessy I, Crane A, Puc J, et al. An inhibitor of mTOR reduces neoplasia and normalizes p70/S6 kinase activity in Pten +/- mice. *Proc Natl Acad Sci U S A*. 2001;98:10320-5.
50. Perrotti D, Turturro F, Neviani P. BCR/ABL, mRNA translation and apoptosis. *Cell Death Differ*. 2005;12:534-40.
51. Paschka P, Marcucci G, Ruppert AS, Mro'zek K, Chen H, Kittles RA, et al. Adverse Prognostic Significance of KIT Mutations in Adult Acute Myeloid Leukemia With inv(16) and t(8;21): A Cancer and Leukemia Group B Study. *Journal of clinical investigation*. 2006;24:3904-3911.
52. Kottaridis PD, Gale RE, Frew ME, Harrison G, Langabeer SE, Belton AA, et al. The presence of a FLT3 internal tandem duplication in patients with acute myeloid leukemia (AML) adds important prognostic information to cytogenetic risk group and response to the first cycle of chemotherapy: analysis of 854 patients from the United Kingdom Medical Research Council AML 10 and 12 trials. *Blood*. 2001;98:1752-1759.
53. Ohki M. Molecular basis of the t(8;21) translocation in acute myeloid leukaemia. *Semin Cancer Biol*. 1993;4:369-75.
54. Romana SP, Poirel H, Leconiat M, Flexor MA, Mauchauffe M, Jonveaux P, et al. High frequency of t(12;21) in childhood B-lineage acute lymphoblastic leukemia. *Blood*. 1995;86:4263-9.
55. Borrow J, Goddard AD, Sheer D, Solomon E. Molecular analysis of acute promyelocytic leukemia breakpoint cluster region on chromosome 17. *Science*. 1990;249:1577-80.
56. Berkowicz M, Rosner E, Resnitzky P, Mamon Z, Ben-Bassat I, Ramot B. Acute nonlymphocytic leukemia with t(16;21). *Cancer Genet Cytogenet*. 1990;47:139-40.
57. Kamps MP, Murre C, Sun XH, Baltimore D. A new homeobox gene contributes the DNA binding domain of the t(1;19) translocation protein in pre-B ALL. *Cell*. 1990;60:547-55.
58. Secker-Walker LM, Mehta A, Bain B. Abnormalities of 3q21 and 3q26 in myeloid malignancy: a United Kingdom Cancer Cytogenetic Group study. *Br J Haematol*. 1995;91:490-501.
59. Ohno H. Pathogenetic and clinical implications of non-immunoglobulin ; BCL6 translocations in B-cell non-Hodgkin's lymphoma. *J Clin Exp Hematol*. 2006;46:43-53.
60. Heisterkamp N, Groffen J. Molecular insights into the Philadelphia translocation. *Hematol Pathol*. 1991;5:1-10.
61. Papadopoulos P, Ridge SA, Boucher CA, Stocking C, Wiedemann LM. The novel activation of ABL by fusion to an ets-related gene, TEL. *Cancer Res*. 1995;55:34-8.
62. Lacroque V, Boureux A, Valle VD, Poirel H, Quang CT, Mauchauffe M, et al. A TEL-JAK2 fusion protein with constitutive kinase activity in human leukemia. *Science*. 1997;278:1309-12.
63. Griesinger F, Hennig H, Hillmer F, Podleschny M, Steffens R, Pies A, et al. A BCR-JAK2 fusion gene as the result of a t(9;22)(p24;q11.2) translocation in a patient with a clinically typical chronic myeloid leukemia. *Genes Chromosomes Cancer*. 2005;44:329-33.
64. Morris SW, Kirstein MN, Valentine MB, Dittmer KG, Shapiro DN, Saltman DL, et al. Fusion of a kinase gene, ALK, to a nucleolar protein gene, NPM, in non-Hodgkin's lymphoma. *Science*. 1994;263:1281-4.
65. Eguchi M, Eguchi-Ishimae M, Tojo A, Morishita K, Suzuki K, Sato Y, et al. Fusion of ETV6 to neurotrophin-3 receptor TRKC in acute myeloid leukemia with t(12;15)(p13;q25). *Blood*. 1999;93:1355-63.
66. Lannon CL, Sorensen PH. ETV6-NTRK3: a chimeric protein tyrosine kinase with transformation activity in multiple cell lineages. *Semin Cancer Biol*. 2005;15:215-23.