

Marchetti M, Sierecki M, Oriot D, Ghazali A. Brugada-type ECG associated with pectus excavatum. *Images Paediatr Cardiol* 2015;17(3):1-2.

IMAGES in PAEDIATRIC CARDIOLOGY

Marchetti M,^a Sierecki M,^a Oriot D,^b Ghazali A.^a Brugada-type ECG associated with pectus excavatum. *Images Paediatr Cardiol* 2015;17(3):1-2.

^a Department of Emergency Medicine, University Hospital Center of Poitiers, Poitiers, France

^b Department of Paediatrics Emergencies, University Hospital Center of Poitiers, Poitiers, France

MeSH: pectus excavatum ; Brugada phenocopie

Brugada-type ECG associated with pectus excavatum

Introduction

Brugada phenocopies (BrP) are new clinical entities characterized by an ECG pattern that is identical to type 1 or type 2 Brugada pattern, despite the absence of the true congenital Brugada syndrome (BrS).¹ BrP are caused by various factors such as mechanical mediastinal compression, myocardial ischemia, pulmonary embolism, pericardial diseases and metabolic conditions.² However, only few cases have been reported of patients with pectus excavatum and BrP. They have an ECG showing right bundle branch block, but also mild ST-segment elevation in the right precordial leads, mimicking the ECG patterns of type 2 Brugada syndrome.³

Case


A 13-year old adolescent with pectus excavatum of the thorax (Figure 1) was admitted to the emergency department with an electrocardiogram (ECG) on admission showing a Brugada type 2 pattern (Figure 2). The patient had no personal or familial history of Brugada syndrome, sudden cardiac death, cardiac arrest, or non-vagal syncope. Moreover, he was known to paediatricians for a history of tachycardia and had several ECGs before his adolescence, without ST-segment modification.

Figure 1: Pectus excavatum



© Images Paediatr Cardiol

Figure 2: Brugada type 2 ECG pattern



Discussion

This case illustrates with typical images a rare association of symptoms. We speculate that the Brugada-type ECG observed in our patient was probably caused by long-term mechanical injury to the right ventricular free wall, as a result of chronic compression by the pectus excavatum (as it appeared only after his adolescence).⁴ Our patient was discharged from the emergency department without any evidence of true Brugada syndrome and a normal 24h Holter ECG recording. No additional investigation was done.

References

1. Baranchuk A, Nguyen T, Ryu MH, Femeni'a F, Zareba W, Wilde AA, et al. Brugada phenocopy: new terminology and proposed classification. *Ann Noninvasive Electrocardiol.* 2012;17:299–314.
2. Anselm DD, Baranchuk A. Brugada Phenocopy: redefinition and updated classification. *Am J Cardiol.* 2013;111:453-456.
3. Kataoka H. Electrocardiographic patterns of the Brugada syndrome in 2 young patients with pectus excavatum. *J Electrocardiol.* 2002;35:169-171.
4. Cartoski MJ, Nuss D, Goretsky MJ, Proud VK, Croitoru DP, Gustin T, et al. Classification of the dysmorphology of pectus excavatum. *J Pediatr Surg.* 2006;9:1573-1581.

Contact Information

Matthieu Marchetti,
Department of Emergency Medicine,
University Hospital Center of Poitiers, 2 Rue
de la Milétrie, 86000 Poitiers, France.
Tel: +33 5 49 44 33 10 / +33 6 88 46 03 81
matthieu.marchetti@gmail.com

© Images in
Paediatric Cardiology
(1999-2015)

