

The Antiplatelet Activity of Aspirin

Claude A. Farrugia B.Pharm. (Hons.), Ph.D. (UIC)

Completion of Assessment Questions following this article entitles members to one credit towards C.E. requirement.

The synthesis of aspirin in the late nineteenth century marked the development of what was to become the most widely used household analgesic in the twentieth century. However, with the increased understanding of the major role of platelets in vascular occlusion over the last ten years, the importance of aspirin as an antiplatelet drug and, consequently, its applications in reducing the risks of thrombotic vascular events, have received major attention.

Pharmacology and Mechanism of Action

Aspirin, otherwise known as acetylsalicylic acid, acts by acetylation of the two isoenzyme forms of prostaglandin G/H synthase. This results in the irreversible inactivation of the cyclooxygenase activity of this enzyme, preventing the conversion of arachidonic acid to prostaglandin G_2 , and consequently decreasing biosynthesis of prostaglandin H_2 and thromboxane A_2 . The type one isoenzyme of prostaglandin G/H synthase is constitutively expressed in platelets, where synthesis and release

of thromboxane A_2 occurs in response to a variety of stimuli, resulting in irreversible platelet aggregation. Platelets, platelet products and thrombosis play important roles in the occurrence of acute occlusive vascular events, including myocardial infarction (MI) and ischemic stroke, since the disruption of platelet- and fibrin-rich atherosclerotic plaque may be followed by aggressive platelet deposition and, ultimately the development of a thrombus that can precipitate an acute occlusive event. The decreased platelet aggregation caused by aspirin is the most plausible mechanism for the cardioprotective effects of this drug.

Dose-dependant inhibition of platelet cyclooxygenase activity occurs with single aspirin oral doses of 5 to 100 mg, with the latter dose resulting in practically total suppression of thromboxane A_2 biosynthesis. The onset of activity of the drug is extremely rapid and unrelated to systemic bioavailability, probably due to platelet prostaglandin synthase suppression in the portal circulation. Furthermore, since platelets lack the intracellular

machinery necessary to regenerate prostaglandin synthase, recovery from the effects of aspirin is related to platelet turnover (7 to 10 days), with repeated doses exerting a cumulative effect. This accounts for the fact that a drug with a half-life of 20 minutes is effective even when administered once daily.

Primary Prevention of Occlusive Cardiovascular Disease

The efficacy of aspirin at preventing important cardiovascular events, chiefly myocardial infarction, stroke and cardiovascular mortality has been assessed in healthy subjects and subjects with stable chronic angina. Also, since patients with insulin-dependent and non-insulin-dependent diabetes mellitus, and hypertensive individuals have an increased risk of cardiovascular complications, including myocardial infarction and stroke, these categories of patients have also received attention in studies assessing the antiplatelet activity of aspirin. The results of these studies are summarised in Table 1.

Despite the significant reduction in MI incidence noted in the Physicians' Health Study, the benefit of aspirin in healthy individuals appears to be minor, given the very low absolute risk of cardiovascular events in healthy individuals (only 5% had an event during approximately 5 years of follow-up). However, in patients with an existing medical condition predisposing to a cardiovascular event, the use of aspirin appeared to have significant benefits. In particular, in the medium-risk population investigated in the Swedish Angina Pectoris Aspirin Trial, the results suggested the potential to prevent 51 important cardiovascular events in 1000 patients over four years, an absolute benefit at least tenfold greater than that obtained in the British Doctors' Trial and the Physicians' Health Study. Similar benefits were also observed in hypertensive patients.

Secondary Prevention of Occlusive Vascular Disease

The benefits afforded by aspirin in the prevention of secondary cardiovascular events supercede those

Table 1: Trials of Aspirin in the Primary Prevention of Occlusive Cardiovascular Disease.

Study	Type of Subjects	Number of Patients	Aspirin Dose	Outcome
British Doctors' Trial	Healthy males, 50-78 yrs old	5,139	500 mg daily vs placebo	No significant difference
Physicians' Health Study (USA)	Healthy males, 40-84 yrs old	22,071	325 mg every other day vs placebo	39% decrease in MI; 18% decrease in important cardiovascular events
Swedish Angina Pectoris Aspirin Trial	Patients with chronic stable angina without previous MI	2,035	75 mg daily vs placebo	34% decrease in MI and sudden death occurrence; 22 - 32% decrease in secondary outcomes
Early Treatment Diabetic Retinopathy Study	Patients with diabetes, 49% of whom with cardiovascular disease history	3,711	650 mg daily vs placebo	28% decrease in MI, 16% increase in stroke, 18% decrease in important cardiovascular events
Hypertension Optimal Treatment Trial	Hypertensive patients (diastolic BP100-115 mmHg), 50-80 yrs old	18,790	75 mg daily vs placebo	36% decrease in MI, 15% decrease in major cardiovascular events

for primary events. The drug remains the standard antiplatelet reference compound for secondary prevention of myocardial infarction, cardiovascular-associated death, and stroke. To this date, no other antiplatelet agent can compete with aspirin in these indications. Three main areas, in increasing order of severity, have been identified where the antiplatelet activity of aspirin plays a significant role in preventing secondary cardiovascular events. These are:

- a) unstable angina and nonacute myocardial infarction
- b) suspected acute evolving myocardial infarction and acute nonfatal myocardial infarction, and
- c) transient cerebral ischaemia and stroke.

Three major studies all established that the administration of aspirin, both short- and long-term, was particularly effective at reducing the risk of acute myocardial infarction and/or death (Table 2). Subgroup analysis in the RISC study also showed that the risk of myocardial infarction was reduced both in patients with silent ischaemia as well as in those with asymptomatic ischaemia detected during a pre-discharge exercise test. However, doubt remains as to whether aspirin can affect the incidence or severity of myocardial ischaemia after an episode of unstable angina. However, the drug did reduce incidence of death and myocardial infarction in patients with this syndrome, regardless of any influence on episodes of transient ischaemia.

Although the results of most early studies in the benefits of aspirin against secondary cardiovascular events in patients with suspected acute evolving myocardial infarction and acute nonfatal myocardial infarction were consistent with the benefits of antiplatelet therapy, most trials were too small individually in sample size. In 1988, an overview was published by the Antiplatelets Trialists' Collaboration of the 25 completed trials in secondary prevention, followed by an updated overview in 1994. The results are summarised in Table 3 and, together with those of the Second International Study of Infarct Survival, demonstrate

the effectiveness of antiplatelet treatment in promoting survival following myocardial infarction. Moreover, in the latter study, the combination of aspirin with intravenous streptokinase proved to be more effective than either agent alone. In absolute terms, the percentage reductions observed imply that the treatment of 1000 patients with acute myocardial infarction with aspirin for one month would prevent 38 fatal and nonfatal cardiovascular events.

The efficacy of aspirin in reducing the risk of recurrent vascular complications in patients with a history of transient ischaemic attacks or stroke was evaluated in various trials. An overview of these trials carried out by the Antiplatelets Trialists' Collaboration concluded that the administration of antiplatelet therapy for three years was effective at reducing the risk of nonfatal stroke, nonfatal myocardial infarction and cardiovascular death by about 25%, respectively, in absolute terms, the prevention of 37 cardiovascular events per 1000 patients over three years. Some doubt remains over the magnitude of the aspirin dose which is effective; while some investigators claim that larger doses are more effective, patients receiving larger doses in those studies reporting a beneficial effect were also receiving other antiplatelet drugs, in particular dipyridamole and sulfinpyrazone.

Maintenance of Vascular Grafts or Arterial Patency

A further outcome of the overview conducted by the Antiplatelets Trialists' Collaboration was the evidence of the benefits provided by antiplatelet therapy in reducing the incidence of arterial or graft occlusion after coronary-artery surgery (30% decrease) or angioplasty (50% decrease), and after the formation of a haemodialysis shunt or fistula (56% decrease). The data provided represented absolute benefits of 90 patients protected per 1000 over 7 months following coronary artery surgery, 40 patients per 1000 over 6 months following angioplasty, and 200 patients per 1000 over two months following formation of a haemodialysis shunt. The same group also concluded that antiplatelet therapy

was effective at reducing the incidence of deep vein thrombosis and pulmonary embolism by 26% and 64% respectively, representing absolute prevention in 90 and 17 patients, respectively, per 1000 treated. While effective prevention was observed in a wide range of medical and surgical patients, the highest benefits were seen in general and orthopaedic surgery.

Side Effects and Safety of Antiplatelet Therapy with Aspirin

The regulation of several homeostatic mechanisms, including haemostasis, renal function, gastric acid secretion and blood pressure control, is modulated locally through eicosanoid synthesis. Inhibition of the constitutive prostaglandin G/H synthase pathway responsible for the production of these mediators is essentially responsible for the side effects associated with the long term administration of aspirin.

Gastrointestinal complications, such as bleeding and perforation, are observed with greater frequency in individuals on nonsteroidal anti-inflammatory therapy. The incidence and severity of these complications are affected by the dosage regimen, the duration of treatment and the type of formulation used (plain vs enteric-coated). Comparison of daily doses of 900 to 1300 mg daily (300 or 325 mg three times daily) as against placebo for various studies exhibited a 40 to 60% increased incidence of stomach pain, heartburn and nausea. On the other hand, both the Veterans Administration Cooperative Study and the Research Group on Instability in Coronary Artery Disease in Southeast Sweden (Table 2) noted only minor increases in the frequency of gastrointestinal symptoms with daily doses of 324 mg and 75 mg aspirin, respectively. Similar dose-dependant results were also obtained by the United Kingdom Transient Ischaemic Attack (UK-TIA) Trial, which noted a 19% increase in upper gastrointestinal symptoms in patients given 300 mg aspirin daily over placebo, and a 58% increase in patients given 1200 mg daily.

The incidence of haemorrhagic

Table 2: Percent reductions in risk of death or acute myocardial infarction in trials among patients following episodes of unstable angina or nonacute myocardial infarction.

Study	Type of Subjects	Aspirin Regimen	Percent reduction
Research Group on Instability in Coronary Artery Disease in Southeast Sweden (RISC)	796 male patients	75 mg daily for 5 days	57-69%
		75 mg daily for 3 months	64%
		75 mg daily for 1 year	48%
Veterans Administration Cooperative Study	1,266 male patients	324 mg daily for 3 months	51%
Canadian Multicenter Trial	555 patients	325 mg 4 times daily for 2 years	30%

Table 3: Percent reductions in cardiovascular events among patients with suspected acute evolving myocardial infarction and acute nonfatal myocardial infarction.

Endpoint	Percent reductions in patients assigned antiplatelet therapy	
	Antiplatelet Trialists' Collaboration (long-term therapy; 1-3 years)	2nd International Study of Infarct Survival (short-term therapy; 5 weeks)
Nonfatal myocardial infarction	31	49
Nonfatal stroke	39	46
Cardiovascular mortality	15	23
Cardiovascular events	25	28

stroke amongst patients involved in antiplatelet studies was too small to permit any comparison amongst different aspirin dosage regimens. However, the UK-TIA Trial did observe a dose-response relation for extracranial haemorrhagic events, with an increased incidence of gastrointestinal haemorrhage at higher doses (5% incidence for patients taking 1200 mg aspirin daily, as against 3% for those taking 300 mg daily), possibly reflecting the presence of more severe gastric mucosal damage and thus a higher possibility of gastric bleeding.

Long-term administration of aspirin may also be associated with an increased risk of chronic renal disease and interference with blood pressure control in hypertensive individuals, since prostaglandin G/H synthase is involved in the renal synthesis of vasodilatory prostaglandins. While the use of aspirin at sub-antiinflammatory doses is unlikely to exert any effect (aspirin is only a weak inhibitor of renal prostaglandin synthase), the UK-TIA trial noted elevations in systolic blood pressures in patients on 300 mg and 1200 mg aspirin daily. Furthermore, this effect of aspirin in both renal and other peripheral tissues interferes with the action of angiotensin-converting-enzyme (ACE) inhibitors in controlling hypertension.

Conclusions

The benefits to be derived from long-term aspirin therapy are evident, particularly in secondary prevention of cardiovascular events. The progress in the understanding of the underlying mechanism of action of the drug and its dose-response relationship have led to the current recommendations of an initial loading dose of about 300 mg followed by a daily dose of 75 mg, the latter serving to decrease the possible incidence of side effects.

While the role of prescribing aspirin antiplatelet therapy and monitoring of its cardiovascular effects remain strictly that of the physician, the pharmacist, too, has a contribution to make. There exists the danger of the idea spreading among the general population that "an aspirin (or baby aspirin) tablet a day is good for one's health". Such an idea is, at the very least, questionable in the

absence of predisposing health conditions and becomes certainly more so if the possibility of side effects and drug interactions discussed above is prevalent.

Nonetheless, the pharmacist remains in an ideal position to identify candidates in whom initiation of such

therapy appears feasible, and should take it upon himself/herself to consult with and make recommendations to the patient's physician, particularly where administration of an aspirin antiplatelet dosage regimen might be effective in the prevention of cardiovascular events. ★

Bibliography

- Antiplatelet Trialists' Collaboration (1994) Collaborative overview of randomised trials of antiplatelet therapy - I: Prevention of death, myocardial infarction, and stroke by prolonged antiplatelet therapy in various categories of patients. *British Medical Journal*, 308: 81-106.
- Antiplatelet Trialists' Collaboration (1994) Collaborative overview of randomised trials of antiplatelet therapy - II: Maintenance of vascular graft or arterial patency by antiplatelet therapy. *British Medical Journal*, 308: 159-168.
- Antiplatelet Trialists' Collaboration (1994) Collaborative overview of randomised trials of antiplatelet therapy - III: Reduction in venous thrombosis and pulmonary embolism by antiplatelet prophylaxis among surgical and medical patients. *British Medical Journal*, 308: 235-246.
- Cairns, J.A., Gent, M., Singer, J., Finnie, K.J., Froggatt, G.M., Holder, D.A., Jablonsky, G., Kostuk, W.J., Melendez, L.J., Myers, M.G., et al. (1985) Aspirin, sulfinpyrazone, or both in unstable angina. Results of a Canadian multicenter trial. *The New England Journal of Medicine*, 313: 1369-1375.
- EDTRS Investigators (1992) Aspirin effects on mortality and morbidity in patients with diabetes mellitus. Early Treatment Diabetic Retinopathy Study report 14. *Journal of the American Medical Association* 268: 1292-1300.
- Farrell, B., Godwin, J., Richards, S., and Warlow, C. (1991) The United Kingdom transient ischaemic attack (UK-TIA) aspirin trial: final results. *Journal of Neurology, Neurosurgery and Psychiatry*, 54: 1044-1054.
- Hansson, L., Zanchetti, A., Carruthers, S.G., Dahlof, B., Elmfeldt, D., Julius, S., Menard, J., Rahn, K.H., Wedel, H., and Westerling, S. (1998) Effects of intensive blood-pressure lowering and low-dose aspirin in patients with hypertension: principle results of the Hypertension Optimal Treatment (HOT) randomised trial. HOT Study Group. *Lancet*, 351: 1755-1762.
- Hennekens, C.H. (1997) Aspirin in the treatment and prevention of cardiovascular disease. *Annual Review of Public Health*, 18: 37-49.
- ISIS-2 (Second International Study of Infarct Survival) Collaborative Group. (1988) Randomised trial of intravenous streptokinase, oral aspirin, both, or neither among 17,187 cases of suspected acute myocardial infarction: ISIS-2. *Lancet*, 2: 349-360.
- Juul-Möller, S., Edvardsson, N., Jahnmatz, B., Rosen, A., Sorensen, S., and Omblus, R. (1992) Double-blind trial of aspirin in primary prevention of myocardial infarction in patients with stable chronic angina pectoris. *Lancet*, 340: 1421-1425.
- Lewis, H.D. Jr, Davis, J.W., Archibald, D.G., Steinke, W.E., Smitherman, T.C., Doherty, J.E., Schnaper, H.W., LeWinter, M.M., Linares, E., Pouget, J.M., Sabharwal, S.C., Chesler, E., and DeMots, H. (1983) Protective effects of aspirin against acute myocardial infarction and death in men with unstable angina. Results of a Veterans Administration Cooperative Study. *The New England Journal of Medicine*, 309: 396-403.
- Patrono, C. (1994) Aspirin as an antiplatelet drug. *The New England Journal of Medicine*, 330: 1287-1294.
- Peto, R., Gray, R., Collins, R., Wheatley, K., Hennekens, C., Jamrozik, K., Warlow, C., Hafner, B., Thompson, E., Norton, S., et al. (1988) Randomised trial of prophylactic daily aspirin in British male doctors. *British Medical Journal*, 296: 313-316.
- Schrör, K. (1995) Antiplatelet drugs: A comparative review. *Drugs*, 50: 7-28.
- Steering Committee of the Physicians' Health Study Research Group (1989) Final report on the aspirin component of the ongoing Physicians' Health Study. *The New England Journal of Medicine*, 321: 129-135.
- The RISC Group (1990) Risk of myocardial infarction and death during treatment with low dose aspirin and intravenous heparin in men with unstable coronary heart disease. *Lancet*, 336: 827-830.

THE **chronic**★**ill**

Pharmacists wishing to publish in the next issue are invited to submit their articles to the Editor, Malta College of Pharmacy Practice, c/o Department of Pharmacy, University of Malta, Msida; or fax on 340427. Tel: 3290 2899

Assessment Questions:

1. Platelet aggregation is caused by release of:
 - a) prostaglandin G₂
 - b) prostaglandin synthase
 - c) thromboxane A₂
 - d) acetylsalicylic acid
2. A once daily aspirin antiplatelet regimen is effective because:
 - a) very high doses of aspirin are used
 - b) the drug accumulates intracellularly from where it is gradually released
 - c) sustained release preparations are used
 - d) recovery from the effects of aspirin is related to platelet turnover
3. Aspirin is effective as a primary preventive agent in:
 - a) hypertensive patients
 - b) perfectly healthy individuals
 - c) patients with chronic stable angina
 - d) both (a) and (c)
4. The minimum aspirin dose effective in secondary prevention in patients with unstable angina is:
 - a) 324 mg daily
 - b) 75 mg daily
 - c) 1200 mg daily
 - d) subject to debate
5. The minimum aspirin dose effective in secondary prevention in patients with a history of transient ischaemic attacks is:
 - a) 324 mg daily
 - b) 75 mg daily
 - c) 1200 mg daily
 - d) subject to debate
6. In order for a beneficial effect to be observed with aspirin therapy in patients with acute MI, therapy must last for at least:
 - a) about 3 years
 - b) about 2 years
 - c) about 1 year
 - d) slightly more than 1 month
7. Aspirin can prevent arterial or graft occlusion after:
 - a) coronary artery surgery
 - b) coronary artery angioplasty
 - c) formation of a haemodialysis shunt
 - d) all of the above
8. The highest benefit in reduction of deep vein thrombosis and pulmonary embolism is seen in:
 - a) all medical and surgical patients
 - b) all surgical patients
 - c) general and orthopaedic surgery patients
 - d) all medical patients
9. The major underlying mechanism responsible for the side effects of aspirin is:
 - a) direct erosion of the intestinal lining, since aspirin is an acid
 - b) inhibition of the constitutive prostaglandin synthesis pathway
 - c) metabolism by the liver to toxic metabolites
 - d) none of the above
10. Gastrointestinal complications with aspirin:
 - a) occur more frequently at higher doses
 - b) occur less frequently at lower doses
 - c) exhibit a dose-dependent response
 - d) all of the above
11. Aspirin-mediated inhibition of prostaglandin synthesis:
 - a) has no effect on blood pressure control in hypertensive individuals
 - b) can result in undesirable drug interactions with captopril and enalapril
 - c) has been observed to result in elevations in diastolic blood pressure
 - d) probably occurs even at maintenance doses of 75 mg daily
12. The recommended aspirin dosage regimen for antiplatelet therapy is:
 - a) an Alka-Seltzer tablet daily
 - b) 300 mg aspirin q.i.d.
 - c) an initial loading dose of about 300 mg followed by a daily dose of 75 mg
 - d) none of the above
13. The minimal aspirin dose which will result in practically total suppression of platelet thromboxane A₂ biosynthesis is:
 - a) 75 mg
 - b) 100 mg
 - c) 300 mg
 - d) 1200 mg
14. The gastrointestinal side effects associated with prolonged use of high dose aspirin include:
 - a) stomach pain
 - b) heartburn
 - c) diarrhoea
 - d) both (a) and (b)
15. If a patient asks a pharmacist whether it is recommendable to take an aspirin tablet a day for good health, the pharmacist should:
 - a) answer with an unequivocal yes
 - b) check if the patient is hypertensive or has a history of cardiovascular events
 - c) refer to a physician
 - d) both (b) and (c)
16. Aspirin plays a major role in preventing secondary cardiovascular events in patients with:
 - a) unstable angina
 - b) a history of acute myocardial infarction but not nonacute MI
 - c) a history of any form of MI
 - d) both (a) and (c)
17. The main indicators in the effectiveness of aspirin in both primary and secondary prevention of occlusive cardiovascular disease are:
 - a) decrease in mortality
 - b) decrease in incidence of MI
 - c) both of the above
 - d) none of the above
18. How does the combination of aspirin with intravenous streptokinase compare with the use of either agent alone in promoting survival following MI?
 - a) The combination is more effective
 - b) The combination is less effective
 - c) The combination is equally effective
 - d) There is no clear evidence for a conclusion to be reached.
19. Other drugs which are sometimes combined with aspirin for antiplatelet activity include:
 - a) dipyridamole
 - b) enalapril
 - c) sulfipyrazone
 - d) both (a) and (c)
20. The incidence and severity of gastrointestinal complications associated with long-term aspirin use are effected by:
 - a) dosage regimen
 - b) duration of treatment
 - c) the type of formulation used
 - d) all of the above