



Title	Molecular weight-dependent degradation of D-lactate-containing polyesters by polyhydroxyalkanoate depolymerases from <i>Variovorax</i> sp C34 and <i>Alcaligenes faecalis</i> T1
Author(s)	Sun, Jian; Matsumoto, Ken'ichiro; Tabata, Yuta; Kadoya, Ryosuke; Ooi, Toshihiko; Abe, Hideki; Taguchi, Seiichi
Citation	Applied microbiology and biotechnology, 99(22), 9555-9563 https://doi.org/10.1007/s00253-015-6756-1
Issue Date	2015-11
Doc URL	http://hdl.handle.net/2115/63400
Rights	The final publication is available at link.springer.com
Type	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	Matsumo-AMB(99)suppl. file.pdf



[Instructions for use](#)

Supporting information

Molecular weight-dependent degradation of D-lactate-containing polyesters by polyhydroxyalkanoate depolymerases from *Variovorax* sp. C34 and *Alcaligenes faecalis* T1

Jian Sun¹, Ken'ichiro Matsumoto¹, Yuta Tabata^{2,3}, Ryosuke Kadoya¹, Toshihiko Ooi^{1,4}, Hideki Abe^{2,3} and Seiichi Taguchi^{1,4,*}

¹Division of Biotechnology and Macromolecular Chemistry, Graduate School of Engineering, Hokkaido University, N13-W8, Kita-ku, Sapporo 060-8628, Japan,

²Department of Innovative and Engineering Materials, Tokyo Institute of Technology, 4259 Nagatsuta, Midori-ku, Yokohama, Kanagawa 226-8502, Japan

³Bioplastic Research Team, Biomass Engineering Program Cooperation Division, RIKEN Center for Sustainable Resource Science, 2-1, Hirosawa, Wako-shi, Saitama 351-0198, Japan

⁴CREST, JST, 4-1-8 Honcho, Kawaguchi, Saitama, 332-0012, Japan

*Corresponding author

staguchi@eng.hokudai.ac.jp

TEL/FAX +81-117066610

Figure S1 ^1H NMR analysis of chemically synthesized trimers. (a) 3HB-LA-LA-methyl, (b) LA-3HB-LA-methyl, (c) LA-LA-3HB-methyl.

