

# Journal of Applied Research on Children: Informing Policy for Children at Risk

Volume 1

Issue 1 *Latino Children*

Article 5

2010

## Latino Teen Pregnancy in Texas: Prevalence, Prevention, and Policy

Susan R. Tortolero

*Center for Health Promotion and Prevention Research, University of Texas Health Science Center at Houston, TX,*  
[susan.tortolero@uth.tmc.edu](mailto:susan.tortolero@uth.tmc.edu)

Belinda F. Hernandez

*Center for Health Promotion and Prevention Research, University of Texas Health Science Center at Houston, TX,*  
[belinda.flores@uth.tmc.edu](mailto:belinda.flores@uth.tmc.edu)

Paula M. Cuccaro

*Center for Health Promotion and Prevention Research, University of Texas Health Science Center at Houston, TX,*  
[Paula.M.Cuccaro@uth.tmc.edu](mailto:Paula.M.Cuccaro@uth.tmc.edu)

Melissa F. Peskin

*Center for Health Promotion and Prevention Research, University of Texas Health Science Center at Houston, TX,*  
[Melissa.F.Peskin@uth.tmc.edu](mailto:Melissa.F.Peskin@uth.tmc.edu)

Christine M. Markham

*Center for Health Promotion and Prevention Research, University of Texas Health Science Center at Houston, TX,*  
[Christine.Markham@uth.tmc.edu](mailto:Christine.Markham@uth.tmc.edu)

*See next page for additional authors*

Follow this and additional works at: <http://digitalcommons.library.tmc.edu/childrenatrisk>

### Recommended Citation

Tortolero, Susan R.; Hernandez, Belinda F.; Cuccaro, Paula M.; Peskin, Melissa F.; Markham, Christine M.; and Shegog, Ross (2010) "Latino Teen Pregnancy in Texas: Prevalence, Prevention, and Policy," *Journal of Applied Research on Children: Informing Policy for Children at Risk*: Vol. 1: Iss. 1, Article 5.

Available at: <http://digitalcommons.library.tmc.edu/childrenatrisk/vol1/iss1/5>

The *Journal of Applied Research on Children* is brought to you for free and open access by CHILDREN AT RISK at DigitalCommons@The Texas Medical Center. It has a "cc by-nc-nd" Creative Commons license" (Attribution Non-Commercial No Derivatives) For more information, please contact [digitalcommons@exch.library.tmc.edu](mailto:digitalcommons@exch.library.tmc.edu)



---

# Latino Teen Pregnancy in Texas: Prevalence, Prevention, and Policy

## **Acknowledgements**

Belinda F. Hernandez is a recipient of an Association of Schools of Public Health/Centers for Disease Control/Prevention Research Center Minority Public Health Fellowship. The content is solely the responsibility of the authors and does not necessarily represent the official views of the Association of Schools of Public Health, Centers for Disease Control, or the Prevention Research Centers.

## **Authors**

Susan R. Tortolero, Belinda F. Hernandez, Paula M. Cuccaro, Melissa F. Peskin, Christine M. Markham, and Ross Shegog

## Introduction

Latinos (defined as a person of Cuban, Mexican, Puerto Rican, South or Central American, or other Spanish culture or origin regardless of race) are the largest minority group in the United States, with almost half living in Texas or California. Texas is home to approximately 9 million Latinos comprising 37% of the population.<sup>1</sup> In the U.S., the Latino population is growing at twice the rate of the total population, and estimates indicate that by 2025 one in two teens living in Texas will be Latino.<sup>2, 3</sup> More than one-third of Texas Latinos are 18 years or younger, and half are under the age of 26.

Compared to other youth, Latino youth are more likely to experience high levels of poverty (32%)<sup>4</sup>, high dropout rates<sup>5</sup>, high unemployment rates<sup>6</sup>, and low rates of health insurance.<sup>7</sup> Additional challenges that Texas Latinos face include negative reproductive health outcomes, such as teen pregnancy and sexually transmitted infections (STIs), which disproportionately impact the health of Latino teens in Texas. Given that high rates of teen pregnancy fuel high rates of poverty, school dropout, and unemployment<sup>8</sup>, it is vital that we understand the factors that contribute to this health burden and potential solutions. By tackling teen pregnancy, we can positively impact the future and well-being of not only Latinos, but of all Texans, and subsequently can contribute to the social and economic success of Texas. This paper describes the current status of teen pregnancy, STIs, sexual behaviors, and contraceptive use among Latino youth in Texas. Potential statewide solutions to impact the teen pregnancy rate among this growing group in Texas are discussed.

## Teen Pregnancies and Births

Pregnancy is a serious public health issue for young people living in Texas, which has the 4<sup>th</sup> highest teen pregnancy rate in the nation (in women aged 15-19, the pregnancy rate in Texas is 88 per 1000 vs. 70 per 1000 in the U.S.).<sup>9</sup> In 2005, more than 73,000 girls aged 15-19 years got pregnant in Texas, and 58% (42,000) were Latina teens.<sup>9</sup> More than half of all Latina girls experience a pregnancy before they turn twenty years of age.<sup>10</sup> It is projected that by 2015 the Texas teen pregnancy rate will increase by 13%, resulting in a projected teen pregnancy rate of 127 per 1000 Texas women aged 15-19 years.<sup>11</sup> This increase is primarily due to the changing demographics and projected growth in the number of Latino adolescents in Texas.

There are over 53,000 teen births in Texas every year, accounting for 12% of all teen births in the U.S.<sup>12</sup> The Texas teen birth rate is the third highest in the U.S. (63.1 per 1000 Texas women aged 15-19 vs. 43 per 1000 U.S. women aged 15-19), and it is the second highest when examining school-aged females (15-17 years). Texas has the highest

repeat teen birth rate in the U.S.: 23% in Texas vs. 19% in the U.S.<sup>13</sup> The statistics among Latino girls in Texas are even more startling: 98 per 1000 Latino girls aged 15-19 are giving birth and one in four are repeat births. This translates to over 32,000 births each year among Latina teens in Texas, over 8,000 of which are to Latina teens who already have at least one child.<sup>14</sup>

While birth rates have declined over the past 15 years between 1990 and 2006, the overall Texas teen birth rate declined at a slower pace than the U.S. teen birth rate (-20% vs. -32%). This slower decline is especially pronounced among Latina teens living in Texas compared to U.S. Latina teens (-6% vs. -17%).

Since half of all Latinos in the U.S. live in California and Texas<sup>15</sup> with about a third of the population in each of these states being of Latino origin, it is helpful to compare Texas' and California's teen birth rates. In 2000, the Latina teen birth rate in both Texas and California was 103 per 1000 Latina females aged 15-19; however, 2006 data indicate that the teen birth rate for California Latinas dramatically decreased by 30% to 68 per 1000 Latina teens, while the teen birth rate for Texas Latinas has decreased only 4% to 98 per 1000 Latina teens. These data indicate that factors beyond population proportions affect teen birth rates.<sup>16</sup>


The estimated annual direct medical costs of all teen births in Texas were almost \$159 million. Direct medical costs per year among Texas Latina teens totaled almost \$98 million. The estimated total costs for Latina births including public sector health care, child welfare costs, costs of incarceration, and lost revenue due to lower taxes paid by the children of teen mothers over their lifetimes as a result of decreased education and earnings was well over \$638 million.<sup>17</sup> These costs are largely paid by taxpayers.<sup>18</sup> If Texas could decrease its Latina teen birth rate to that of California (68 per 1000 Latina teens), there would be almost 10,000 fewer Latina teen births each year, resulting in annual savings of over \$29 million. These costs are likely to be understated because it is assumed that medical costs of unintended pregnancies are no greater than the average costs of all births. An Institute of Medicine report concludes that the costs of an unwanted pregnancy are greater because the mother is less likely to seek prenatal care, more likely to expose the fetus to harmful substances by smoking and using alcohol, and the child is at greater risk of being low birth weight, dying in the first year, being abused, and of not receiving sufficient resources for healthy development.<sup>19</sup>

### **Sexual Behavior**

Texas teens are becoming sexually active at an early age, with as many as 14% of 7<sup>th</sup> grade students reporting ever having sex.<sup>20, 21</sup> In a study conducted by Tortolero and colleagues among a large sample of urban

middle school youth, 7% of Latino students reported having had sex by 7<sup>th</sup> grade (unpublished data, 2010). The proportion of Latino students engaging in sex more than doubled by 8<sup>th</sup> grade and quadrupled by 9<sup>th</sup> grade. When ethnic differences in the initiation of sex were examined (See Figure 1), however, Latino students were less likely to report initiating sex than black students. In that study, Latino students had greater intentions to abstain from sex until after high school and greater intentions to abstain from sex until marriage than black students. However, Latino students had lower knowledge about STIs and HIV and lower self-efficacy to use a condom than black students (unpublished data, 2010).

**Figure 1. Initiation of Sexual Activity in Middle School by Grade**




The 2009 Youth Risk Behavior Surveillance System (YRBS) data for high school students in Texas show that Latino students (54%) were less likely to report ever having had sexual intercourse compared to black students (63%), but were more likely to report ever having sex than white students (45%).<sup>22</sup> Among Latino students, over one-third (39%) of 9<sup>th</sup> grade students, half (52%) of 10<sup>th</sup> grade students, two-thirds (65%) of 11<sup>th</sup> grade students, and 69% of high school seniors report initiating sex.

Figure 2 displays the proportion of high school students initiating sex by grade for white and Latino students. The difference between Latino and white high school students is particularly pronounced among males: 58% vs. 45%, respectively; however by 12<sup>th</sup> grade, differences in initiation of sex between Latino and white students is not significantly different (69% vs. 65%) and is still lower than black students (77%). According to the YRBS, an estimated 15% of Texas Latino high school students report having sex with four or more partners in their lifetime, and 40% of Texas Latino high school students report being sexually active in the last three months.<sup>22</sup>

In summary, Latino teens are at greater risk of pregnancy than other teens even when fewer Latino teens are sexually experienced and they have greater intentions to wait until marriage to become sexually active. Most teens become sexually experienced while they are of school age, which translates to an estimated 622,015 sexually experienced students attending Texas public schools, an estimated 414,583 of which are Latino. Of these Latino youth, 237,466 report being currently sexually active, and 89,000 report having four or more sexual partners in their lifetime.<sup>22, 23</sup> These numbers are alarming and point to the need to understand why Texas is doing so poorly while other states such as California have been so successful in implementing programs and policies that work.

**Figure 2. Initiation of Sexual Activity in High School by Grade**



## Contraception

Rising rates of unintended pregnancies and STIs also suggest that sexually experienced youth do not protect themselves with effective methods of contraception. In Texas only 58% of sexually active teens reported using a condom during their last sexual encounter, and only 20% of females reported using an effective method of birth control (oral contraceptive pill or Depo Provera injectable contraceptive) compared to 27% of U.S. female teens.<sup>22</sup> Latino students are less likely to use birth control or condoms than their white and black peers. For example, a 1995 study reported that almost half of Latina teens did not use any type of birth control the first time they had sex, compared to 29% of all teens.<sup>24</sup> Further, according to the 2009 YRBS, only 64% of Texas Latino high school students reported using a condom, and 11% reported using birth control pills or Depo Provera during their last sexual encounter.<sup>22</sup> Studies show that generational and language status are associated with less frequent contraceptive use among Latino teens. For example, first or second generation Latino teens are less likely to consistently use contraception than third generation teens, and Latino teens interviewed in Spanish and those who speak Spanish at home are less likely to use contraception the first time they have sex than Latino teens who speak only English at home.<sup>25</sup>

## Impact

Latino teens are twice as likely as black teens and four times as likely as white teens to drop out of high school; teen pregnancy is cited as a major reason for school dropout. In a national study, almost 40% of dropouts either had a child or were expecting one. This was the case for 54% of females and 26% of males.<sup>26</sup> Over half (54%) of Latina teen mothers do not complete high school, compared to 34% of all teen mothers.<sup>27</sup> One study reported that of all Latina teens who drop out of high school between sophomore and senior year, 36% cite being pregnant or becoming a mother as a reason for dropping out.<sup>28</sup>

Teen mothers are more likely to live in poverty, to come from poor or low-income families, and to depend on welfare. According to the National Campaign to Prevent Teen and Unplanned Pregnancy, over 80% of teen mothers receive welfare during the 10 years following the birth of their first child, and 44% of them for a period of more than 5 years.<sup>24</sup>

The consequences for the child of the teen mother include poverty, welfare dependence, school failure, drop out, increased risk of abuse, time spent in the foster care system, being unemployed, and if female, becoming a teen mother herself. Children of teenage mothers are also at greater risk for low birth weight, premature birth, lack of father

involvement, receiving insufficient health care, incarceration and mental retardation.<sup>29</sup>

## **Prevention**

We know what works to prevent teen pregnancy: implementation of evidence-based sexual health education programs in middle and high school; improving access to community and school-based reproductive health services; providing training on adolescent sexual health to teachers, administrators, parents, and school nurses; and providing clear state-wide policies and leadership regarding effective strategies to prevent pregnancies, and to increase parental involvement.<sup>30, 31</sup>

**Implementation of evidence-based sexual health education programs:** Research confirms that some sexual health education curricula can delay the initial sexual encounter, reduce the frequency of sex, reduce the number of sexual partners, and increase the use of condoms and contraceptives.<sup>29</sup> Implementation of evidence-based sexual health education programming is essential in preventing teen pregnancy. However, Texas is not systematically implementing effective programs and practices, and great disparities exist in who even receives the most basic instruction related to sexual health. A recent report indicated that the vast majority (96%) of Texas students did not receive medically accurate or evidence-based information on responsible pregnancy and disease prevention, and 41% of school districts used sexual health education materials that contained factual errors about condoms and STIs.<sup>32</sup>

Texas laws regarding health education and sexual health education are confusing and generally unsupportive of school-based implementation of effective sexual health programs. For example, Texas recently passed a law indicating that health education is no longer a required course for high school students.<sup>33</sup> While the provision of health education in and of itself does not ensure that Texas students are exposed to effective sexual health curricula, the probability that Texas students are exposed to even basic sexual health information will be greatly impacted by this new law.

Prior to Texas law not mandating health education as a required course for high school students, the CDC reported that only 19% of high school students were required to take two or more health education courses. While the majority (82%) of Texas schools taught eight key topics related to pregnancy, HIV, or other STIs in a required course in high school, only 46% taught key topics related to condom use in a required course. Texas teachers are also poorly trained to deliver information on HIV prevention: only 45% of secondary schools reported having a lead health education teacher who received professional development on HIV prevention in the prior two years.<sup>34</sup> In a survey of professional staff (e.g., teachers and principals) from a large urban school district in Texas, 93%



of respondents strongly believed that sexual health education should be taught in middle schools; however only half perceived the Texas Education Agency as supportive of teaching sexual health in middle school.<sup>35</sup>

In Texas, Latino students are less likely than white or black students to receive the most basic sexual health information on HIV and AIDS. One-fourth of Latino high school students reported never learning about HIV in school compared to 11% of white students and 11% of black students. Unfortunately, simply providing information and increasing knowledge is not enough to decrease risky sexual behavior and to impact the teen pregnancy rate. Implementing a program rigorously tested and shown to impact behavior is essential in reducing risky sexual behavior and teen pregnancy.<sup>29</sup> An effective pregnancy prevention strategy follows two principles: 1) choosing not to have sex is the surest way to avoid STIs and unintended pregnancies, and 2) if teens are sexually active, young men should use condoms and young women should use hormonal/long-acting reversible birth control every time they have sex.

While few programs have been developed specifically for Latino youth, many programs have been tested and found to be effective in reducing risky sexual behaviors among this population (See Table 1).

- The evaluation of *¡Cuidate!/Take Care of Yourself: The Hispanic Youth Health Promotion Program* found that one year after baseline, youth exposed to *¡Cuidate!* were less likely to report having had sex recently, having multiple sexual partners, having had sex without a condom and more likely to report consistent condom use compared to youth in the control condition.<sup>36</sup>
- *It's Your Game: Keep It Real (IYG)* was implemented in middle schools and found that over 24 months, Latino students who did *not* receive the intervention were 64% more likely to initiate sex than Latino students who received the *IYG* curriculum.<sup>21</sup>
- *Draw the Line/Respect the Line* was evaluated over a 36-month follow-up period and was found to delay sexual initiation (among boys only) in a sample which was made up of 59% Latinos.<sup>37</sup>
- *Positive Prevention* was implemented in the 9<sup>th</sup> grade and was found to delay sexual initiation at the six-month follow-up in a sample of youth that was 60% Latino.<sup>38</sup>
- *Safer Choices* is a school-based multi-component intervention implemented in high school over two consecutive years. As with *IYG*, *Safer Choices* had its greatest effect on Latino youth.<sup>39</sup> At the 31-month follow-up, Latino teens in the program were 43% less likely to have initiated sex when compared to Latino teens in the control group. Additionally, Latinos in the program who

were sexually active were 65% more likely than Latinos in the control group to have used a condom during their last sexual encounter.<sup>39</sup>

- *Reach for Health Community Youth Service (RFH-CYS)* is a multi-component program that combines service learning with skills-based health instruction in middle school. At the six-month follow-up, *RFH-CYS* was found to delay sexual initiation among a sample of youth that were 16% Latino.<sup>40</sup>
- The *Children's Aid Society-Carrera Program* is a multi-component long term, after school, youth development program implemented with youth aged 13-15 years. Tested in a population that was 39% Latino, the *Carrera* program significantly delayed sexual initiation, increased condom use, and reduced pregnancy and birth rates among females exposed to the program.<sup>41</sup>

While Texas has relied on abstinence-only programming upon which there is no evidence of impact on sexual behavior, California's strong sex education policy has been credited as a major factor in the state's success in decreasing teen birth rates. For the last several decades, California has had strong bipartisan commitment to implementing sex education programming that includes information about abstinence, while also providing medically accurate information on other methods of preventing pregnancy and STDs.<sup>30</sup> These programs are monitored by the California Department of Education by visiting school districts every four years to interview parents, students, and teachers, and to review classroom instruction.

**Access to contraception services:** Research indicates that once a teen is sexually active, he or she is not likely to revert to sexual abstinence<sup>29</sup> and that access to contraceptive services is critical to preventing teen pregnancy.<sup>42, 43</sup> Studies show that a sexually active young girl who does not use contraceptives has a 90% chance of becoming pregnant within a year.<sup>44</sup> In 2008 there were an estimated 1.2 million teen females of reproductive age (13 to 19 years) living in Texas, of which over half a million were Latina.<sup>45</sup> About half of all teen females, 43% of which are Latina,<sup>45</sup> are in need of contraceptive services.<sup>46</sup> An estimated one-third of Latina females aged 13-19 years live below the federal poverty level<sup>47</sup> and may be in need of public-supported contraceptive services. In 2006 the 426 publicly-funded family planning centers provided services to 100,200 teenagers of all racial and ethnic backgrounds.<sup>48</sup> Guttmacher reports that in 2008, family planning clinics in Texas served 32% of all women (ages 13-44) in need of publicly-supported contraceptive services. In Texas, 158 family planning centers received some support through the

Federal Title X family planning program and provided contraceptives to 47,100 teenagers in 2006. Publicly-funded family planning services can help teens avoid pregnancy that could result in unintended births and abortions. The data indicate that for every dollar spent providing contraceptive care, \$3.74 in public funds are saved.<sup>46</sup>

Texas has some of the most restrictive policies regarding teen access to contraceptive services. While 21 states and the District of Columbia explicitly allow minors to consent for contraceptive services, Texas requires parental consent for contraceptive services in state-funded family planning programs.<sup>49</sup> While some policymakers are opposed to increasing teen access to contraceptives, one study found positive effects of teen-friendly family planning policies in reducing teen birth rates; specifically, Medicaid Family Planning was shown to be effective in reducing teen birth rates among all racial and ethnic groups.<sup>16</sup> The majority (90%) of family planning clinics counsel clients younger than 18 about abstinence and the importance of talking to parents about sex.<sup>16, 50</sup> Laws limiting teen access to family planning services fail to reduce sexual activity and increase the risk of unintended pregnancy.<sup>16, 51</sup> Improving access to culturally-appropriate, youth-friendly reproductive health services and increasing support for comprehensive coverage of contraceptive products are cost-effective steps to reduce the personal and social burden of Latina teen pregnancies.

**Parental involvement:** Parental involvement and training are additional strategies that may impact the teen pregnancy rate among Latinas in Texas. Parents play an important role in the lives of Latino youth; however, many have limited resources, knowledge, and skills necessary to address sexual health for their teens.<sup>52</sup> Studies have shown that relationships with engaged and communicative parents help some Latinas delay pregnancy. Research shows that over 90% of parents support implementation of sex education programs in schools and that the curriculum should include information about contraception.<sup>53</sup> Parents often are unprepared and lack the knowledge and skills to discuss human sexuality with their teen, but want to do a better job of informing their children.<sup>52</sup>

## Conclusion

Texas is home to over one million Latino teens, and estimates indicate that by 2025 one in two teens living in Texas will be Latino.<sup>2,3</sup> Texas Latinos face enormous negative reproductive health outcomes, such as teen pregnancy and STIs, which disproportionately impact the health of Latino teens in Texas and place them at risk of continued high rates of poverty, school dropout, and unemployment<sup>8</sup> unless Texas makes a concerted effort to reduce its teen pregnancy rate. There are over 53,000

teen births in Texas every year, accounting for 12% of all teen births in the U.S.<sup>12</sup> The birth rate among Latina girls is astonishing: 98 per 1000 Latinas (aged 15-19 years) are giving birth. This translates to over 32,000 births each year among Latina teens, costing almost \$98 million in direct medical expenditures and well over \$638 million if other costs are included, such as public sector health care, child welfare costs, costs of incarceration, and lost revenue due to lower taxes paid by the children of teen mothers over their lifetimes as a result of decreased education and earnings.<sup>17</sup>

Most teens become sexually experienced while they are of school age, which translates to an estimated 622,015 sexually experienced students attending Texas public schools, an estimated 414,583 of which are Latino. Of these Latino youth, 237,466 report being currently sexually active, and 89,000 report having had four or more sexual partners in their lifetime.<sup>22, 23</sup>

While causes of teen pregnancy are complex, the solutions to teen pregnancy are known.<sup>30,31</sup> Texas needs an effective comprehensive approach to address the sexual health needs of Texas Latino youth that includes: statewide implementation and monitoring of evidence-based sex education for middle school and high school students, access to reproductive health services for students who are already sexually active, and widespread training on adolescent sexual health for teachers, service providers, and parents. By tackling teen pregnancy, we can positively impact the future and well-being of not only Latinos, but of all Texans, and subsequently can contribute to the social and economic success of Texas.

## References

- (1) U.S. Bureau of the Census. State and county quickfacts: Texas. Available at: <http://quickfacts.census.gov/qfd/states/48000.html>. Accessed on June 4, 2010.
- (2) Sabatiuk L, Flores R. *Toward a Common Future: Latino Teens and Adults Speak Out About Teen Pregnancy*. Washington, DC: National Campaign to Prevent Teen and Unplanned Pregnancy; 2009.
- (3) Vexler EJ, Suellentrop K. *Bridging Two Worlds: How Teen Pregnancy Prevention Programs Can Better Serve Latino Youth*. Washington, DC: National Campaign to Prevent Teen and Unplanned Pregnancy; 2006.
- (4) Federal Interagency Forum on Child and Family Statistics. *America's Children in Brief: Key National Indicators of Well-Being, 2008*. Washington, DC: U.S. Government Printing Office; 2008.
- (5) Aud S, Hussar W, Planty M et al. *The Condition of Education 2010 (NCES 2010-028)*. Washington, DC: National Center of Education Statistics, Institute of Education Sciences, US Department of Education; 2010.
- (6) U.S. Department of Labor. Labor force statistics from the current population survey. Available at: <http://www.bls.gov/data/>. Accessed on July 8, 2010.
- (7) DeNavas-Walt C, Proctor BD, Smith JC. *Income Poverty and Health Insurance Coverage in the United States: 2008*. Washington, DC: U.S. Government Printing Office; 2009.
- (8) Hoffman SD. *By the Numbers: The Public Costs of Teen Childbearing*. Washington, D.C.: National Campaign to Prevent Teen Pregnancy; 2006.
- (9) Kost K, Henshaw S, Carlin L. U.S. teenage pregnancies, births, and abortions: national and state trends and trends by race and ethnicity. Available at: <http://www.guttmacher.org/pubs/USTPtrends.pdf>. Accessed on July 10, 2010.
- (10) National Campaign to Prevent Teen and Unplanned Pregnancy. A look at Latinos: an overview of Latina teen pregnancy and birth rates. Available at: [http://www.thenationalcampaign.org/espanol/PDF/latino\\_overview.pdf](http://www.thenationalcampaign.org/espanol/PDF/latino_overview.pdf). Accessed on July 8, 2010.
- (11) Sayegh MA, Castrucci BC, Lewis K, Hobbs-Lopez A. Teen pregnancy in Texas: 2005 to 2015. *Matern Child Health J*. January 2010;14(1):94-101.

- (12) Centers for Disease Control and Prevention, National Center for Health Statistics. Vital stats: birth data files. Available at: <http://www.cdc.gov/nchs/vitalstats.htm>. Accessed on July 8, 2010.
- (13) Child Trends. Facts at a glance: a fact sheet reporting national, state, and city trends in teen childbearing. Available at: [http://www.childtrends.org/Files//Child\\_Trends-2009\\_08\\_31\\_FG\\_Edition.pdf](http://www.childtrends.org/Files//Child_Trends-2009_08_31_FG_Edition.pdf). Accessed on July 8, 2010.
- (14) Schelar E, Franzetta K, Manlove J. Repeat teen childbearing: differences across states and by race and ethnicity. Available at: [http://www.childtrends.org/Files//Child\\_Trends-2007\\_11\\_27\\_RB\\_RepeatCB.pdf](http://www.childtrends.org/Files//Child_Trends-2007_11_27_RB_RepeatCB.pdf). Accessed on July 15, 2010.
- (15) U.S. Bureau of the Census. Population estimates for states by race and Hispanic origin, July 2009. Available at: <http://www.census.gov/popest/states/asrh/stasrh.html>. Accessed on July 8, 2010.
- (16) Yang Z, Gaydos LM. Reasons for and challenges of recent increases in teen birth rates: a study of family planning service policies and demographic changes at the state level. *J Adolesc Health*. June 2010;46(6):517-24.
- (17) The University of Texas Prevention Research Center (UTPRC). *Special calculations tabulated by the UTPRC*. Houston, TX: UTPRC. 2010.
- (18) Texas Medical Association. The uninsured in Texas. Available at: <http://www.texmed.org/Template.aspx?id=5517>. Accessed on September 10, 2010.
- (19) Brown S, Eisenberg L. *The Best Intentions: Unintended Pregnancy and the well-being of Children and Families*. Washington, DC: National Academy Press; 1995.
- (20) Markham CM, Peskin MF, Addy RC, Baumler E, Tortolero SR. Patterns of vaginal, oral, and anal sexual intercourse in an urban seventh grade population. *J Sch Health*. 2009;79:193-200.
- (21) Tortolero SR, Markham CM, Peskin MF et al. It's Your Game: Keep It Real: delaying sexual behavior with an effective middle school program. *J Adolesc Health*. February 2010;46(2):169-79.
- (22) Centers for Disease Control and Prevention. 2009 Youth Risk Behavior Survey. Available at: [www.cdc.gov/yrbss](http://www.cdc.gov/yrbss). Accessed on July 8, 2010.
- (23) Texas Education Agency, Division of Accountability Research. Enrollment in Texas public schools. Available at: <http://ritter.tea.state.tx.us/research/>. Accessed on July 8, 2010.
- (24) Population Resource Center. Executive summary. Latina teen pregnancy: problems and prevention. Available at:

- [http://www.clients.prc.sprocketlogic.com/files/Latina\\_Teen\\_Pregnancy.pdf](http://www.clients.prc.sprocketlogic.com/files/Latina_Teen_Pregnancy.pdf). Accessed on July 8, 2010.
- (25) National Campaign to Prevent Teen and Unplanned Pregnancy. Science Says. Acculturation and sexual behaviors among Latino youth: findings from the national longitudinal survey of youth 1997-2003. Available at: [http://www.thenationalcampaign.org/resources/pdf/SS/SS35\\_Acculturation.pdf](http://www.thenationalcampaign.org/resources/pdf/SS/SS35_Acculturation.pdf). Accessed on July 8, 2010.
  - (26) Coley RJ. *Dreams Deferred: High School Dropouts in the United States. Policy Information Report*. Princeton, NJ: Policy Information Center; 1995.
  - (27) Perper K, Peterson K, Manlove J. Diploma attainment among teen mothers. Available at: [http://www.childtrends.org/Files/Child\\_Trends-2010\\_01\\_22\\_FS\\_DiplomaAttainment.pdf](http://www.childtrends.org/Files/Child_Trends-2010_01_22_FS_DiplomaAttainment.pdf). Access on September 10, 2010.
  - (28) National Campaign to Prevent Teen and Unplanned Pregnancy. A look at Latinos: Latina teen pregnancy and educational attainment. Available at: [http://www.thenationalcampaign.org/espanol/PDF/latino\\_education.pdf](http://www.thenationalcampaign.org/espanol/PDF/latino_education.pdf). Accessed on September 10, 2010.
  - (29) Kirby D. *Emerging answers 2007: Research Findings on Programs to Reduce Teen Pregnancy and Sexually Transmitted Diseases*. Washington, D.C.: National Campaign to Prevent Teen and Unplanned Pregnancy; 2007.
  - (30) Boonstra H. Winning campaign: California's concerted effort to reduce its teen pregnancy rate. *Guttmacher Policy Review*. 2010;13(2):18-24.
  - (31) Donovan P. School-based sexuality education: the issues and challenges. *Fam Plann Perspect*. 1998;30(4):188-93.
  - (32) Wiley D, Wilson K. *Just Say Don't Know: Sexuality Education in Texas Public Schools*. Austin, TX: Texas Freedom Network; no date.
  - (33) Section 28.025 of the Texas Education Code: Texas Education Code §28.025.
  - (34) U.S. Department of Health and Human Services. HIV, other STD, and teen pregnancy prevention and Texas students. Available at: [http://www.cdc.gov/HealthyYouth/yrbs/pdf/hiv/tx\\_hiv\\_combo.pdf](http://www.cdc.gov/HealthyYouth/yrbs/pdf/hiv/tx_hiv_combo.pdf). Accessed on September 10, 2010.
  - (35) Peskin MF, Markham C, Shegog R et al.; Sexual health education from the perspective of school personnel: implications for dissemination of effective middle school programs. Oral presentation presented at the 137<sup>th</sup> Annual Meeting and Exposition

- of the American Public Health Association: November 2009; Philadelphia, PA.
- (36) Villarruel AM, Jemmott JB, III, Jemmott LS. A randomized controlled trial testing an HIV prevention intervention for Latino youth. *Arch Pediatr Adolesc Med.* August 2006;160(8):772-7.
  - (37) Coyle KK, Kirby DB, Marin BV, Gomez CA, Gregorich SE. Draw the line/respect the line: a randomized trial of a middle school intervention to reduce sexual risk behaviors. *Am J Public Health.* May 2004;94(5):843-51.
  - (38) LaChausse R. Evaluation of the positive prevention HIV/ STD curriculum for students grades 9-12. *American Journal of Health Education* 2006;37(4):203-9.
  - (39) Kirby DB, Baumler E, Coyle KK et al. The "Safer choices" intervention: its impact on the sexual behaviors of different subgroups of high school students. *Journal of Adolescent Health.* December 2004;35(6):442-52.
  - (40) O'Donnell L, Stueve A, San DA et al. The effectiveness of the reach for health community youth service learning program in reducing early and unprotected sex among urban middle school students. *Am J Public Health.* February 1999;89(2):176-81.
  - (41) Philliber S, Kaye JW, Herrling S, West E. Preventing pregnancy and improving health care access among teenagers: an evaluation of the children's aid society-Carrera program. *Perspect Sex Reprod Health.* September 2002;34(5):244-51.
  - (42) Franzini L, Marks E, Cromwell PF et al. Projected economic costs due to health consequences of teenagers' loss of confidentiality in obtaining reproductive health care services in Texas. *Arch Pediatr Adolesc Med.* December 2004;158(12):1140-6.
  - (43) Kahn JG, Brindis CD, Gleit DA. Pregnancies averted among U.S. teenagers by the use of contraceptives. *Fam Plann Perspect.* January 1999;31(1):29-34.
  - (44) Harlap S, Kost K, Forrest JD. Preventing Pregnancy, Protecting Health: *A New Look at Birth Control Choices in the United States.* New York: Alan Guttmacher Institute; 1991.
  - (45) Texas State Data Center and Office of the State Demographer. Texas population estimates program. 2008 age, sex and race/ethnicity estimates for the state of Texas and all Texas counties. Available at: <http://txsdc.utsa.edu/tpepp/txpopest.php>. Accessed on July 8, 2010.
  - (46) Frost JJ, Henshaw SK, Sonfield A. Contraceptive Needs and Services: *National and State Data, 2008 Update.* New York: Guttmacher Institute; 2010.



- (47) Annie E. Casey Foundation, Kids Count Data Center. Data across states: children in poverty by race, Available at: <http://datacenter.kidscount.org/data/acrossstates/Rankings.aspx?id=44>. Accessed on September 10, 2010.
- (48) Guttmacher Institute. *Contraceptive Needs and Services, 2006*. Available at: <http://www.guttmacher.org/pubs/win/allstates2006.pdf>. Accessed on September 10, 2010.
- (49) Guttmacher Institute. Minor's access to contraceptive services. *State Policies in Brief*. Available at: [http://www.guttmacher.org/statecenter/spibs/spib\\_MACS.pdf](http://www.guttmacher.org/statecenter/spibs/spib_MACS.pdf). Accessed on July 8, 2010.
- (50) Lindberg LD, Frost JJ, Sten C, Dailard C. Provision of contraceptive and related services by publicly funded family planning clinics, 2003. *Perspect Sex Reprod Health*. September 2006;38(3):139-47.
- (51) Jones RK, Purcell A, Singh S, Finer LB. Adolescents' reports of parental knowledge of adolescents' use of sexual health services and their reactions to mandated parental notification for prescription contraception. *JAMA*. January 2005;293(3):340-8.
- (52) Whitaker DJ, Miller KS, May DC, Levin ML. Teenage partners' communication about sexual risk and condom use: the importance of parent-teenager discussions. *Fam Plann Perspect*. May 1999;31(3):117-21.
- (53) Kaiser Family Foundation. Sex education in America. An NPR/Kaiser/Kennedy school poll. Available at: <http://www.npr.org/templates/story/story.php?storyId=1622610>. Accessed on March 17, 2010.

**Table 1: Effective pregnancy/HIV/STD programs for Latino youth**

<b>Program</b>	<b>Goal of Program</b>	<b>Population Served</b>	<b>Outcomes</b>
<b>Stand Alone Programs</b>			
<i>¡Cuidate!/Take Care of Yourself: The Hispanic Youth Health Promotion Program</i> <sup>36</sup>	Reduce risk behaviors for HIV	Latino teens ages 13-18 years, boys and girls	Reduced frequency of sex, number of sexual partners, frequency of unprotected sex, increased consistent condom use
<i>Draw the Line/Respect the Line</i> <sup>37</sup>	Delay sexual initiation, reduce sexual activity and increase condom use among youth who are sexually active	Youth in 6 <sup>th</sup> -8 <sup>th</sup> grades, boys and girls, 59% were Latino, 17% White, and 16% Asian	Found effective for boys only: Delayed sexual initiation, reduced frequency of sex and number of sexual partners
<i>It's Your Game: Keep It Real (IYG)</i> <sup>22</sup>	Delay sexual initiation, reduce sexual activity and increase condom use among youth who are sexually active	Youth in 7 <sup>th</sup> -8 <sup>th</sup> grades, Boys and girls, 44% were Latino, and 42% were African American	Delayed sexual initiation, reduced frequency of sex
<i>Positive Prevention</i> <sup>38</sup>	Delay sexual initiation, reduce frequency of sex, increase condom use	Youth in high school, boys and girls, 59% were Latino, 17% were African American, 10% were White, 4% were Asian, and 1% Native American	Delayed sexual initiation
<b>Multi-Component Programs</b>			
<i>Children's Aid Society-Carrera Program</i> <sup>41</sup>	Reduce sexual activity, increase contraceptive use, reduce teen pregnancy, and increase access to health care using a youth development approach	Youth 13-15 years, boys and girls, 60% were African American and 39% were Latino	Found effective for girls only: Delayed sexual initiation, increased contraceptive use, decreased pregnancy rate
<i>Reach for Health Community Youth Service (RFH-CYS)</i> <sup>40</sup>	Prevent risky sexual behaviors and other detrimental behaviors among at-risk youth through a service learning approach	Youth in 7 <sup>th</sup> -8 <sup>th</sup> grades, boys and girls, 16% were Latino, 79% were African American	Delayed sexual initiation, reduced frequency of sex, decreased unprotected sex
<i>Safer Choices</i> <sup>39</sup>	Delay sexual initiation and increase condom use among youth sexually experienced	Youth in high school, boys and girls, 28% were Latino, 20% were African American, 30% were White, and 14% were Asian in the intervention condition	Delayed sexual initiation (among Latino youth only), increased condom use, increased contraceptive use, decreased unprotected sex