

EXCLI Journal 2007;6:10-22 – ISSN 1611-2156

Received: 7. November 2006, accepted: 16. November 2006, published: 26. January 2007

Original article:

Escherichia coli K88 Interaction with IgA Oligosaccharides

Gabriela Ramos-Clamont M¹, Evelia Acedo-Felix¹, Joy Winzerling² and Luz Vázquez-Moreno^{1*}

¹Laboratorio de Bioquímica de Proteínas de la Coordinación de Ciencia de los Alimentos, Centro de Investigación en Alimentación y Desarrollo A.C., Carretera a la Victoria Km. 0.6, Hermosillo, CP 83000, México, Phone: + 52 (662) 289-2400, Fax: + 52 (662) 280-0058, e-mail lvazquez@ciad.mx (*Corresponding author); ²Department of Nutritional Sciences, 309 Shantz Building, University of Arizona, Tucson, A.Z., 85721, United States.

ABSTRACT

Diarrhea from enterotoxigenic *Escherichia coli* (*E. coli*) expressing the K88 fimbrial adhesin causes high morbidity and mortality among newborn and weaned piglets. K88 fimbrial adhesins are surface filaments with lectin activity that recognize specific glycoconjugates (glycoproteins or glycolipids) on the surface of intestinal cells. Carbohydrates that compete for adhesion attachment could serve as an alternative for disease prevention. In this study, IgA, IgG and IgM oligosaccharides were tested to inhibit the adhesion of *E. coli* K88 to piglet mucins. Immunoglobulins were isolated from porcine serum by hydrophobic interaction chromatography (HIC) and purified by affinity chromatography. *In vitro* K88 adhesin interacts specifically with IgA oligosaccharides, but not with carbohydrates of IgG or IgM. Also IgA oligosaccharides partially inhibit the adherence of K88 strain to porcine intestinal mucins.

Keywords: Affinity purification, IgA oligosaccharides, *E. coli* K88 adhesin, piglet mucins, inhibition of adhesion.

INTRODUCTION

Protein-carbohydrate interactions appear to play a critical role in the adherence of pathogens to eukaryotic cells (Karlsson, 2001). Many microbial pathogens including those responsible for enteric infections, exploit oligosaccharides that are displayed on the surface of host cells as receptor for toxin and adhesin (Paton et al., 2006). *E. coli* expressing the K88 adhesin on their surface

is a common cause of diarrhea in newborn and weaned piglets (Wilson and Francis, 1986). K88 fimbrial adhesins are filamentous surface appendages with lectin activity that allows *E. coli* to attach to specific glycoconjugates (receptors) on porcine intestinal mucins and epithelial cells (Grange et al, 1998). Adhesion appears to prevent bacterium removal by intestinal peristalsis, facilitating colonization of the small intestine. K88 fimbriae can specifically

adhere to sialoglycoproteins, intestinal mucin-type glycoproteins, and neutral glycosphingolipids via α and β -linked galactose structures (Grange et al, 1999; Grange et al., 1998; Blomberg et al., 1993; Payne et al., 1993; Neeser et al., 1986; Gibbons et al., 1975). Thus, oligosaccharides that compete for microorganism-adhesins could provide an alternative for disease prevention.

The importance of immunoglobulins (Igs) in protecting new borne against diarrhea reflects not only antigen-antibody interactions but also possible interactions with the complex carbohydrates associated to immunoglobulin heavy chains (Owusu et al, 2002; Araujo et al, 2000; Gomez et al, 1998; Cravioto et al, 1991; Wold et al., 1990; Cruz et al, 1988; Rutter et al., 1976). In fact, the carbohydrate portion of human IgA binds to pathogens to prevent their adhesion to intestinal cells (Cravioto et al, 1991) and suggests that carbohydrates of IgA could serve as a possible target for prevention of infection (Rajan et al, 1999). Carbohydrates that specifically interfere with the initial microorganism interaction could hinder the disease process. This concept is supported by studies in which sugars or other synthetic receptor analogs that block the binding of microorganisms to cells and prevent microbial invasion (Karlsson et al., 2001; Rajan et al, 1999; Rosenstein, 1994; Karlsson et al, 1992; Schaeffer et al, 1984). The objective of the present study was to determine whether K88 adhesin could interact with porcine serum IgA oligosaccharides and compete for mucin sites.

MATERIALS AND METHODS

Materials

Sepharose CL 6B was purchased from GE Healthcare (Uppsala, Sweden). Radial

immunodiffusion VET-RID kits, goat anti-pig for IgA, IgG, and IgM were from Bethyl Labs (Montgomery, TX, USA). Divinylsulfone (DVS) activated agarose was from Kem-En-Tek, (Copenhagen, Denmark). BSA-Gal α (1-3)Gal was from Glycorex, (Sweden). All other reagents were analytical grade from Sigma Aldrich (St. Louis, Mo, USA).

Sample preparation

Porcine blood was collected in sterile containers at the bleeding line of a certified slaughterhouse located in Hermosillo, México. Blood was allowed to coagulate at 25°C and serum separated by decanting, followed by centrifugation at 24 000 g for 15 min at 4°C. The fat layer was removed at this time and aliquots of serum stored at -40 °C until use.

IgA, IgM and IgG purification

Porcine immunoglobulins were isolated in a single step by hydrophobic interaction chromatography (HIC) using a highly acetylated Sepharose (HA-Sepharose) synthesized according to Vázquez-Moreno et al. (1992). HA-Sepharose was packed at 1.5-mL/min flow rate, to obtain a 2 mL bed volume (10 x 0.5 cm) column. HA-Sepharose was equilibrated with five bed volumes of 0.5 M Na₂SO₄, 10 mM MOPS, pH 7.6 (Buffer A) and samples of 5 mL of porcine serum (adjusted to 0.5 M Na₂SO₄, 10 mM MOPS, pH 7.6) loaded onto column. The gel was washed with Buffer A to remove unadsorbed proteins that correspond to albumin fraction. Igs were eluted with 10 mM MOPS Buffer, pH 7.6 (Buffer B). Following elution, HA-Sepharose was cleaned with two bed volumes of 4 M guanidine hydrochloride (guanidine-HCl), pH 7.6; washed with five bed volumes of distilled water and re-equilibrated with Buffer A (Ramos-Clamont et al., 2006). All chromatographic procedures were monitored

by absorbance at 280 nm (Spectronic 21 spectrophotometer, Milton Roy, USA) and analyzed for protein content by Bradford (1976) using bovine serum albumin (BSA) as a standard.

Each type of immunoglobulin (IgA, IgM or IgG) was purified by affinity chromatography using an anti-IgA, anti-IgM or anti-IgG agarose matrix according to Vázquez-Moreno et al. (1993). Forty mg of anti-pig for IgA, IgM, or IgG coupled individually to 2 g of DVS activated agarose (Kem-En-Tek, Copenhagen, Denmark). Gels were packed in columns (4 x 1 cm), equilibrated with PBS, pH 7.2 and loaded with 10 mg of isolated Igs fraction (in PBS, pH 7.2) from hydrophobic interaction chromatography. Unbound proteins were washed with PBS and the specific immunoglobulin eluted from the column with 0.02 M glycine HCl, pH 3.6. Elution fractions were immediately neutralized with 0.1 M Tris buffer, pH 8.0.

Quantitative radial immunodiffusion

The relative amount of IgA, IgM or IgG in unabsorbed and adsorbed fractions, as well as in serum samples were determined by radial immunodiffusion (RID) according to Fahey and McKelvey (1965). Quantitative RID kits containing goat anti-pig for IgA, IgM or IgG were used according to vendor (Bethyl Labs, Montgomery TX, USA). Plate wells were loaded with reference standards of Igs or with 160 µg of protein from each chromatographic fraction. Immunoglobulin concentration of unknown samples was determined by locating their precipitation diameter on a semi log plot. Different concentrations of pig Igs were used as standard. Precipitation diameters were measured with a VET-RID reader and an ocular device Finescale (Horscale, Labconco, USA). Duplicate samples were analyzed for immunoglobulin content at three independent times.

Polyacrylamide gel electrophoresis

Protein fractions were analyzed by electrophoresis using 8 % sodium dodecyl sulfate polyacrylamide gel (SDS-PAGE) according to Laemmli (1970). Gels were stained with 1 % Coomassie blue in 30% methanol and 10% acetic acid.

Immunoblotting assay

Assays were performed as described by Towbin et al. (1979). Unadsorbed and elution fractions (25 µg) were separated on 8 % SDS-PAGE gels, under reducing conditions (Laemmli, 1970) and transferred to nitrocellulose membranes at 2.5 mA/cm² for 40 min (Semi-dry blotter, Bluchler, Labconco, USA). IgA was detected with anti-pig IgA (α chain specific, 1:1000 dilution) raised in goat, followed by incubation with anti-goat IgG peroxidase (as secondary antibody, 1:3000 dilution). Procedures for detection of IgG and IgM were similar but using the corresponding antibody. Color reactions were developed by H₂O₂ addition and 3,3'-diaminobenzidine.

Piglet mucin isolation

Intestinal sections (duodenum, jejunum and ileum) were obtained from 9 healthy newborn piglets (17 days-old) according to the method of Miller and Hoskins (1981). Briefly, small intestines were taken immediately after slaughter, flushed with phosphate buffered saline A (PBS-A; 10 mM phosphate, pH 7.2) containing 1 mM phenylmethylsulfonyl fluoride, 0.5 mg/mL NaN₃, 10 mM EDTA and 2 mM iodoacetamide), intestinal sections were opened longitudinally and cut into 20 cm segments. Mucus was gently removed with a glass slide and suspended in four volumes of ice-cold PBS-A. The suspension was shaken for 1 h at 4°C and centrifuged for 30 min at 15,000 g to remove particulate matter. Mucus

was isolated from the extracts by dual precipitation with ice-cold ethanol, freeze-dried and suspended in 10 mM HEPES-Hanks buffer at a concentration of 0.5 mg/ml. Mucin content was determined by the alcian blue method of Hall et al., (1980). Bovine submaxillary mucins were used as standard. All assays were done in duplicate.

Adhesion assays

The *E. coli* K88 strain, kindly donated by Dr. Carlos Eslava from Universidad Nacional Autonoma de Mexico, was maintained as frozen stocks. Overnight cultures were grown in nutritive broth on a shaker at 37°C. After harvesting, bacteria cells were washed twice in Hanks balanced salt solution (HBSS, Sigma, St. Louis, MO. USA) and suspensions adjusted to $\approx 2 \times 10^9$ CFU ml⁻¹ by optical density equal to 1.0 at 600 nm (Jin et al., 2000).

E. coli K88 was biotin labeled by mixing 0.9 mL of culture (2×10^9 CFU ml⁻¹) with 0.1 ml of dimethyl sulfoxide containing 0.2 mg of NHS-LC-Biotin (Pierce, Rockford, Ill. USA). Labeling of bacteria was performed in the presence of 100 mM lactose to protect lectin-binding sites (Ruhl et al., 1996). After 3 h of incubation, at room temperature with occasional mixing, labeled bacteria were washed 4 times in HBSS to remove free NHS-LC-Biotin.

For blotting assays, serial two fold dilutions containing from 1 µg/µL to 1 ng/µL of protein were prepared for IgA, IgM, IgG, piglet mucins and control glycoproteins. Aliquots of 1 µL were applied to nitrocellulose (Ruhl et al., 1996). Glycoproteins used as positive controls included fetuin (F), asialofetuin (AF), ovalbumin (OV), porcine gastric mucin (PGM), asialo porcine gastric mucin (APGM), bovine submaxillary mucin (BSM)

and a neoglycoprotein BSA-Gal α(1-3)Gal prepared by conjugation of Galα(1-3)Gal to BSA (Glycorex, Sweden). Untreated BSA was used as negative control.

Membranes spotted with glycoproteins were blocked overnight in TTBS (0.15 M NaCl, 20 mM Tris HCl, 1 mM CaCl₂, 1 mM MgCl₂ and 0.02% sodium azide and 0.05 % Tween 20, pH 7.8) containing 3% defatted milk and 1 µg/mL of avidin (Wood and Warnke, 1981). After washing with TTBS, membranes were overlaid with biotinylated bacteria to a final concentration of approximately 5×10^8 CFU per ml in the same buffer. Following overnight incubation at 4°C, membranes were washed to remove unbound bacteria and developed with avidin-alkaline phosphatase to detect bacterial adhesion. Similarly, bacterial adhesion was also tested using denatured and dissociated IgA, IgM or IgG. Igs samples (25 µg) were reduced, separated by SDS-PAGE and transferred to nitrocellulose (Ruhl et al., 1996).

To test the inhibition of lectin-mediated adhesion of *E. coli* K88 to piglet mucin by IgA, samples containing 2 µg of duodenum jejunum and ileum, mucins, were spotted on nitrocellulose and incubated with $\approx 5 \times 10^8$ CFU/ml of labeled *E. coli* K88 (positive control). For inhibition studies, bacteria was mixed with different concentrations IgA (0.3-5 µg/ml) and pre-incubated for 45 min before overlaid to nitrocellulose membranes as described before.

Hemagglutination activity and carbohydrate inhibition were performed by the serial double-dilution method using piglet erythrocytes (Jaffe, 1980). Bacteria cells were washed and suspended in PBS pH 7.2, containing 1 mM CaCl₂ and 1 mM MgCl₂. Suspensions were adjusted to 10⁹ CFU/mL. Piglet erythrocytes were suspended to 2% in

a PBS solution, pH 7.2. For agglutination activity, 25 μ L of bacterial suspension were deposited on an agglutination plate, serially diluted and followed by the addition of 25 μ L of erythrocytes. HT was defined as the inverse maximal dilution showing visible agglutination.

Inhibition assays were done by mixing 25 μ L of appropriately diluted glycoprotein solutions with 25 μ L of bacterial suspension. After 40 min incubation, 25 μ L of 2 % erythrocyte suspension were added. Inhibition was determined as the concentration of glycoproteins that prevent 50% of the HT (Jaffe, 1980). Glycoproteins included F, AF, OV, PGM, APM, BSM, porcine Igs (IgA, IgG and IgM), and piglet duodenum mucin. The initial stock concentration of glycoproteins was 1 mg/mL.

RESULTS

IgA purification

The immunoglobulin fraction (containing IgA, IgM, and IgG) was isolated from porcine serum by HIC as described by Ramos-Clamont et al., (2006), and IgA was purified from this fraction by affinity chromatography (Fig. 1). Fractions containing 10 mg of immunoglobulins were applied to anti-IgA column; unadsorbed proteins were washed with PBS pH 7.6 (Fig. 1, fraction A), while those interacting with the matrix were eluted with 0.02 M Glycine-HCl, pH 3.5 (Fig. 1 fraction B).

The IgA content in serum and chromatographic fractions was estimated by radial immunodiffusion (Table 1). Serum contained 2.2 ± 0.1 mg/mL of IgA; also $2.6 \pm$

0.1 and 20.2 ± 0.1 mg/ml of IgM and IgG respectively (data not shown). IgA remained in the adsorbed HIC fraction. IgA (2.2 ± 0.1 mg/mL) was purified in the adsorbed fraction of anti-IgA agarose. As expected the IgA content corresponded to 9-10% of total immunoglobulin fraction (Butler and Brown, 1996).




Figure 1: Elution of IgA purified from porcine immunoglobulins. Column containing immobilized anti-IgA agarose was equilibrated with PBS, pH 7.6 and porcine immunoglobulins applied. Unbound proteins remained in the flow through (A). Adsorbed proteins eluted (B) with Glycine-HCl, pH 3.5; flow rate 0.5 ml/min; fractions, 3.0 ml/tube.

Table 1. Immunoglobulin A content in chromatography fractions

| | HIC | | Affinity Chromatography | | |
|-------------------------------|-----------------------|------------------------|--------------------------------|-------------|-----------|
| | Total Protein (mg/mL) | IgA (mg/mL) | Total Protein (mg/mL) | IgA (mg/mL) | |
| Serum | 40.0 ± 0.3 | 2.2 ± 0.1 ^a | Ig fraction | 10 ± 0.1 | 2.7 ± 0.1 |
| Unadsorbed (albumin fraction) | 32.4 ± 0.2 | ND | Unadsorbed (IgM, IgA, IgG) | 9 ± 0.2 | 1.7 ± 0.2 |
| Adsorbed (Ig fraction) | 7.6 ± 0.3 | 2.2 ± 0.1 | Adsorbed (IgA) | 1 ± 0.1 | 1 ± 0.1 |

^aConcentrations were estimated by RID according to Fahey and McKelvey (1965). Average value and standard deviation of six determinations

Ig: Immunoglobulin

ND: Not Detected

Adhesion assays

Serial two dilution samples containing purified IgA, IgM and IgG, piglet mucins and glycoproteins were immobilized on nitrocellulose and overlaid with biotin-labeled *E. coli* K88 for adhesion assays. K88 fimbrial adhesin bind to IgA at concentrations from 1 µg to 0.25 µg (Fig. 2A). In contrast, no bacterial adhesion was observed when IgG and IgM were spotted (data not shown). Bacterial adhesion was greater to duodenum mucins than to jejunum or ileum mucins (Fig. 2B). Also bovine submaxillary mucin and asialofetuin were recognized (Fig 2C). No adhesion was observed with fetuin, porcine gastric mucin or ovalbumin (data not shown). In contrast adhesion of *E. coli* K88 was observed to as little as 1.9 ng of BSA-Gal α(1-3)Gal

neoglycan. These results suggest that the galactosyl residue containing glycoproteins may serve as receptors for the lectin-like components of *E. coli* K88.

Bacterial adhesion to denatured IgA was also examined. IgA was reduced separated by SDS-PAGE (Fig. 3A) and transferred to nitrocellulose membranes (Fig. 3B). Membranes were incubated with *E. coli* K88 labeled with biotin. *E. coli* interacted with both BSA-Gal(α 1-3)Gal neoglycan and with the heavy chain of porcine IgA (Fig. 3B). Immunoglobulin heavy chain was confirmed with anti-pig IgA antiserum (Bethyl Labs, Montgomery, TX) as shown in figure 3C. Mucins were spotted in nitrocellulose and overlaid with *E. coli* K88 bacteria that was pretreated with purified IgA (0.3 -5 µg/ml).




Figure 2: Lectin-mediated adhesion of *E. coli* K88 to immobilized glycoproteins. Proteins were applied in serial dilution (1 µg to 1 ng) on nitrocellulose. Membranes were then overlaid with biotinylated *E. coli* K88 incubated for 24 h and washed. Interactions were detected using avidin-alkaline phosphatase. A). Purified porcine IgA from serum (1); B). Piglet intestinal mucins from duodenum (2), jejunum (3), and ileum (4); C). Glycoproteins: asialofetuin (6), bovine submaxillary mucin (7) and BSA-Gal $\alpha(1-3)$ Gal neoglycan (8).




Figure 3: Detection of IgA heavy chain by bacterial overlay and immunoblotting. (A) SDS-PAGE of porcine serum IgA. Lane 1. Molecular weight standards; lane 2. BSA-Gal $\alpha(1-3)$ Gal; lane 3. purified IgA. (B). Western blot of porcine IgA. Lane 1; BSA-Gal $\alpha(1-3)$ Gal and lane 2 reduced and denatured IgA. Samples containing 25 µg of purified IgA were reduced, separated by SDS-PAGE, and transferred to nitrocellulose. Membranes were overlaid with biotinylated *E. coli* K88 and the interaction developed with avidin-alkaline phosphatase to detect adherent bacteria. (C) Western blot of the IgA heavy chain (α). IgA α was detected using antiserum raised against anti-pig α chain.

IgA was able to reduce bacterial adhesion to mucins in a dose-dependent manner (Figure 4).




Figure 4. Partial inhibition of lectin-mediated adhesion of *E. coli* K88 to piglet mucin by porcine serum IgA. Samples containing 2 µg of piglet mucin were spotted on nitrocellulose and incubated with $\approx 5 \times 10^8$ CFU/ml of biotin labeled *E. coli* K88 (Positive control). For inhibition studies, bacteria was mixed with the indicated concentrations of porcine serum IgA and incubated for 45 min before overlaying the nitrocellulose membranes. Interactions were revealed with avidin-alkaline phosphatase to detect adherent bacteria.

The bacterial adherence to the intestinal mucosa is a prerequisite in colonization, and therefore, considerable attention has been given to the factors related to adherence. The adhesive ability of an enteropathogen is usually assessed by determining the hemagglutinating ability, because the erythrocyte membrane is believed to possess similar receptors as those of the mucosal involved in bacterial adherence (Anantha et al., 2004; Nagayama et al., 1994, Cravioto et al., 1982). In addition, high correlation has been observed between agglutination of erythrocytes from different species caused by pili and production of enterotoxin by porcine, bovine, and human strains of *E. coli* (Meng et al., 1998; Evans et al., 1979, 1977; Jones and

Rutter, 1974, 1972). Thus piglet erythrocytes were used to test of hemagglutination caused by *E. coli* K88 (Table 2). Fimbrial bacterial agglutination was inhibited by piglet duodenum mucin (0.001 mg/mL) and porcine purified IgA (0.008 mg/mL), while Immunoglobulin fraction required 0.010 mg/mL). This strongly suggests that inhibition is due to IgA present in the total immunoglobulin fraction. Bovine submaxillary mucin was as effective as Igs. On the contrary, asialofetuin was required in greater concentration (8x) to inhibit agglutination.

Table 2. Inhibition of hemagglutination by IgA containing fractions

| Inhibition of Hemagglutination | mg/mL ^a |
|---------------------------------|--------------------|
| <i>Study fractions</i> | |
| Piglet duodenum mucin | 0.001 |
| Porcine IgA | 0.008 |
| Porcine Igs (IgA,IgG,IgM) | 0.010 |
| <i>Commercial glycoproteins</i> | |
| Bovine submaxillary mucin | 0.008 |
| Asialofetuin | 0.063 |
| Porcine gastric mucin | N.I |
| Bovine serum albumin | N.I |
| Ovalbumin | N.I |

(n = 9)

^a Minimum concentration (mg/mL) required to inhibit a hemmagglutination titer of 8

N.I. No inhibition

DISCUSSION

Pathogen Gram-negative bacteria adhesion is mediated mainly by adhesins. Bacterial adhesins bind to host cell complex carbohydrates located at the mucosal epithelial cells leading to an infection (Sharon and Ofek, 2001). K88 fimbriae can specifically adhere to gut receptors via α and

β -linked galactose structures (Grange et al., 1998; Blomberg et al., 1993; Payne et al., 1993; Neeser et al., 1986; Gibbons et al., 1975).

Among the strategies included to prevent diarrhea in animals is the use of probiotic bacteria or specific monoclonal antibodies directed against adhesins. The mechanisms, by which probiotics exert beneficial health effects, are not well understood, but appear to involve immune modulation of the host, production of antimicrobial compounds and competitive exclusion of probiotics against pathogens (Kirjavainen et al., 1998). Recent studies with *Lactobacillus gasseri* K7 found that this probiotic was effective in reducing *E. coli* adhesion to Caco-2 enterocytes, however was not able to do so in *ex vivo* conditions tested for pig jejunal tissue (Bogovic et al., 2006). Alternatively, products from fermented food, such as soybean Tempe have high inhibitory activity of K88 hemagglutination as a result of its ability to aggregate the ETEC K88 cells (Kiers et al., 2002). This inhibition of aggregation was also observed by Van der Broeck et al. (2000) with anti-K88 specific monoclonal antibodies.

In this study, we used non-specific IgA for the inhibition of *E. coli* K88 agglutination. Grange and Mouricout (1996) and Grange et al. (2002) reported that K88 adhesin binds to porcine serum transferrin (pST) which contains a biantennary complex N-glycan with the structures Gal β (1-4)GlcNAc β (1-2)Man-- linked to Asn-497 (Sharma et al., 1994). This glycan moiety is also found in human IgA (Mattu et al., 1998). Furthermore, we show that the oligosaccharide moiety of purified porcine serum IgA was recognized by *E. coli* K88 as well as the epitope Gal α (1-3)Gal attached to BSA. *E. coli* K88

interaction was specific to IgA oligosaccharides, since no bacterial attachment was observed to the major Igs (IgG or IgM) that are also glycosylated with complex carbohydrates (Butler and Brown, 1994).

Bacteria can carry different adhesins with distinct specificities giving cells flexibility to attach to different receptors (Karlsson, 2001; Guinee and Jensen, 1979). The three serologically distinguishable variants of K88 (K88ab, K88ac, and K88ad) present lectins with different but related carbohydrate specificity, as reflected in their hemagglutination patterns. K88ab and K88ac agglutinate porcine erythrocytes (Bijlsma and Frik, 1987) while K88ad did not (Guinee et al., 1979; Orskov, 1964).

Further studies are required to define the specificities of adhesins from pathogens in order to propose treatments with well-defined glycoproteins that possess structurally known glycans.

ACKNOWLEDGMENT

This research was supported financially by the National Council of Science and Technology of Mexico, CONACYT, under project 31458-B.

REFERENCES

Anantha RP, McVeigh AL, Lee LH, Agnew MK, Cassels FJ, Scott DA, Whittam TS, Savarino SJ, Evolutionary and Functional Relationships of Colonization Factor Antigen I and Other Class 5 Adhesive Fimbriae of Enterotoxigenic *Escherichia coli*, *Infect. Immun.*, 2000; 72: 7190-7201.

- Araujo NA, Giugliano LG, Human milk fractions inhibit the adherence of diffusely adherent *Escherichia coli* (DAEC) and enteroaggregative *E. coli* (EAEC) to HeLa cells, *FEMS Microbiol. Lett.*, 2000; 184: 91–94.
- Bijlsma I, Frik JF, Hemagglutination patterns of the different variants of *Escherichia coli* K88 antigen with porcine, bovine, guinea pig, chicken, ovine and equine erythrocytes, *Res. Vet. Sci.*, 1987; 43: 122-123.
- Blomberg L, Krivan HC, Cohen PS, Conway PL, Piglet ileal mucus contains protein and glycolipid (galactosylceramide) receptors specific for *Escherichia coli* K88 fimbriae, *Infect. Immun.*, 1993; 61: 2526-2531.
- Bogovic Matijacic B, Narat M, Zori Peternel M, Rogelj I, Ability of *Lactobacillus gasseri* K7 to inhibit *Escherichia coli* adhesion in vitro on Caco-2 cells and ex vivo on pigs' jejunal tissue *Int. J. Food Microbiol.*, 2006; 107: 92-96.
- Butler JE, Brown WR, The immunoglobulins and immunoglobulin genes of swine, *Vet. Immunol. Immunopath.*, 1994; 43: 5-12.
- Coffey RD, Cromwell GL, The impact of environment and antimicrobial agents on the growth response of early-weaned pigs to spray-dried porcine plasma, *J. Anim. Sci.*, 1995; 73: 2532–2539.
- Cravioto AS, Scotland M, Rowe B, Hemagglutination activity and colonization factor antigens I and II in enterotoxigenic and non-enterotoxigenic strains of *Escherichia coli* isolated from humans, *Infect. Immun.*, 1982; 36:189-197.
- Cravioto AS, Tello A, Villafan H, Ruiz J, Vedovo S, Neeser JR, Inhibition of localized adhesion of enteropathogenic *Escherichia coli* to HEP-2 cells by immunoglobulin and oligosaccharide fractions of human colostrum and breast milk, *J Infect. Dis.*, 1991; 163: 1247–1255.
- Cruz CR, Gil L, Cano F, Caceres P, Pareja G, Breast milk anti *Escherichia coli* heat labile toxin: IgA antibodies protect against toxin induced infantile diarrhea, *Acta Paediatr. Scand.*, 1998; 77: 658–662.
- Evans DG, Evans DJ, DuPont HL, Hemagglutination patterns of enterotoxigenic and enteropathogenic *Escherichia coli* determined with human, bovine, chicken, and guinea-pig erythrocytes in the presence and absence of mannose, *Infect. Immun.*, 1979; 21:336-346.
- Evans DG, Evans DJ, Tjoa W, Hemagglutination of human group A erythrocytes by enterotoxigenic *Escherichia coli* isolated from adults with diarrhea: correlation with colonization factor. *Infect. Immun.*, 1977; 18:330-337.
- Fahey JL, McKelvey EM, Quantitative determination of serum immunoglobulins in antibody agar plates, *J. Immunol.*, 1965; 94: 84-90.
- Gomez GG, Phillips O, Goforth RA, Effect of immunoglobulin source on survival, growth and hematological and immunological variables in pigs, *J. Anim. Sci.*, 1998; 76: 1-7.
- Gibbons RA, Jones GW, Sellwood R, An attempt to identify the intestinal receptor for the K88 adhesin by means of a hemagglutination inhibition test using glycoproteins and fractions from sow colostrum, *J. Gen. Microbiol.*, 1975; 86: 228–240.

- Guinee PAM, Jansen WH, Behavior of *Escherichia coli* K antigens K88ab, K88ac, and K88ad in immunoelectrophoresis, double diffusion, and hemagglutination, *Infect. Immun.*, 1979; 23: 700–705.
- Grange PA, Mouricout MA, Transferrin associated with the porcine intestinal mucosa is a receptor specific for K88ab fimbriae of *Escherichia coli*, *Infect. Immun.*, 1996; 64: 606-610.
- Grange PA, Erickson AK, Anderson TJ, Francis DH, Characterization of the carbohydrate moiety of intestinal mucin-type sialoglycoprotein receptors for the K88ac fimbrial adhesin of *Escherichia coli*, *Infect. Immun.*, 1998; 66: 1613-1621.
- Grange PA, Erickson AK, Lavery SB, Francis DH, Identification of an intestinal neutral glycosphingolipid as a phenotype-specific receptor for the K88ad fimbrial adhesin of *Escherichia coli*, *Infect. Immun.*, 1999; 67: 165-172.
- Grange PA, Mouricout MA, Lavery SB, Francis DH, Erickson AK, Evaluation of receptor binding specificity of *Escherichia coli* K88 (F4) fimbrial adhesin variants using porcine serum transferrin and glycosphingolipids as model receptors. *Infect. Immun.*, 2002; 70: 2336-2343.
- Hall RL, Miller RJ, Peatfield AC, Richardson PS, Williams I, Lampert I, A colorimetric assay for mucous glycoproteins using alcian blue, *Biochem. Soc. Trans.*, 1980; 8: 72-76.
- Jaffe WG, Lectins, in M. Liener (Ed.), *Toxic constituents of plant foodstuffs* Academic Press, London, 1980, 73-98.
- Jin LZ, Marquardt RR, Zhao X, Strain of *Enterococcus faecium* (18C23) inhibits adhesion of enterotoxigenic *Escherichia coli* K88 to porcine small intestine mucus, *Appl. Environ. Microbiol.*, 2000; 66: 4200–4204.
- Jones G.W, Rutter JM, The association of K88 antigen with haemagglutinating activity in porcine strains of *Escherichia coli*, *J. Gen. Microbiol.*, 1974; 84, 135–144.
- Jones GW, Rutter JM, Role of K88 antigen in the pathogenesis of neonatal diarrhea caused by *Escherichia coli* in piglets, *Infect. Immun.*, 1972; 6: 918-927.
- Karlsson KA, Pathogen-host protein-carbohydrate interactions as the basis of important infections, *Adv. Exp. Med. Biol.*, 2001; 491: 431-43.
- Karlsson KA, Angstrom J, Bergstrom J, Lanne B, Microbial interaction with animal cell surface carbohydrates, *APMIS Suppl.*, 1992; 27: 71-83.
- Kiers JL, Nout MJR, Rombouts FM, Nabuurs MJA, van der Meulen J, Inhibition of adhesion of enterotoxigenic *Escherichia coli* K88 by soya bean tempe, *Letters in Appl. Microbiol.*, 2002; 35: 311–315.
- Kirjavainen PV, Ouwehand AC, Isolauri E, Salminen SJ, The ability of probiotic bacteria to bind to human intestinal mucus. *FEMS Microbiol Lett.* 1998; 167:185-189.
- Laemmli UK, Cleavage of structural proteins during the assembly of head of bacteriophage T4. *Nature*, 1970; 227: 680-685.
- Mattu TS, Pleass RJ, Willis AC, Kilian M, Wormald MR, Lellouch AC, Rudd PM, Woof JM, Dwek RA, The glycosylation and structure of human serum IgA1, Fab, and Fc regions and the role of N-glycosylation on Fc alpha-receptor interactions, *J. Biol. Chem.*, 1998; 273: 2260-2272.

- Meng Q, Kerley MS, Russel TJ, Allee GL, Lectin-like activity of *Escherichia coli* K88, *Salmonella choleraesuis*, and *Bifidobacteria pseudolongum* of porcine gastrointestinal origin, *J. Anim. Sci.*, 1998; 76, 551–556.
- Miller RS, Hoskins LC, Mucin degradation in human colon ecosystems. Fecal population densities of mucin-degrading bacteria estimated by a “most probable number” method, *Gastroenterology*, 1981; 81: 759–765.
- Nagayama K, Oguchi T, Arita M, Honda T, Correlation between cell-associated mannose-sensitive hemagglutination by *Vibrio parahaemolyticus* and adherence to human colonic cell line Caco-2, *FEMS Microbiol. Lett.*, 1994; 120:207–210.
- Neeser JR, Koellreutter B, Wuersch P, Oligomannoside-type glycopeptides inhibiting adhesion of *Escherichia coli* strains mediated by type 1 pili: preparation of potent inhibitors from plant glycoproteins, *Infect Immun.*, 1986; 52: 428–436.
- Orskov I, Orskov F, Sojka WJ, Wittig W, K antigens K88ab(L) and K88ac(L) in *E. coli*, *Acta Pathol. Microbiol. Scand.*, 1964; 62: 439–447.
- Owusu-Asiedu A, Baidoo SK, Nyachoti CM, Marquardt RR, Response of early-weaned pigs to spray-dried porcine or animal plasma-based diets supplemented with egg-yolk antibodies against enterotoxigenic *Escherichia coli*, *J. Anim. Sci.*, 2002; 80: 2895–2903.
- Paton A, Morona N, Paton JC, Designer probiotics for prevention of enteric infections, *Nat. Rev. Microbiol.*, 2006; 4:193-200.
- Payne D, O'Reilly M, Williamson D, The K88 fimbrial adhesin of enterotoxigenic *Escherichia coli* binds to β 1-linked galactosyl residues in glycosphingolipids, *Infect. Immun.*, 1993; 61: 3673-3677.
- Rajan N, Cao Q, Anderson BE, Pruden DL, Sensibar J, Duncan JL, Schaeffer AJ, Roles of glycoproteins and oligosaccharides found in human vaginal fluid in bacterial adherence, *Infect Immun.*, 1999; 67: 5027-5032.
- Rosenstein IJ, The use of lipid-linked oligosaccharides (neoglycolipids) in the identification of carbohydrate receptors for microbial pathogens, *Biomed Pharmacother.*, 1994; 48: 319-326.
- Ruhl S, Sandberg AL, Cole MF, Cisar JO, Recognition of Immunoglobulin A by oral *Actinomyces* and *Streptococcal* lectins, *Infect. Immun.*, 1996; 64: 5421- 5424.
- Rutter JM, Jones GW, Brown GTH, Burrows MR, Luther PD, Antibacterial activity in colostrum and milk associated with protection against enteric disease caused by K88-positive *Escherichia coli*, *Infect. Immun.*, 1976; 13: 667-676.
- Schaeffer AJ, Chmiel JS, Duncan JL, Falkowski WS, Mannose-sensitive adherence of *Escherichia coli* to epithelial cells from women with recurrent urinary tract infections, *J. Urol.*, 1984; 131: 906-910.
- Sharma N, Evans R, Patel K, Gorinsky B, Mallet A, Aitken A, Evidence for the glycosylation of porcine serum transferrin at a single site located with the C-terminal lobe, *Biochem. Biophys. Acta*, 1994; 1206: 286-288.
- Stromberg N, Berén T, *Actinomyces* tissue specificity may depend on differences in

- receptor specificity for GaLNac3-containing glycoconjugates, *Infect Immun.*, 1992; 60: 3268-3277.
- Towbin H, Staehelin T, Gordon J, Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc. Natl. Acad. Sci. USA*, 1979; 76: 4350-4354.
- Van den Broeck W, Cox E, Oudega B, Goddeeris BM, The F4 fimbrial antigen of *Escherichia coli* and its receptors, *Vet. Microbiol.*, 2000; 71: 223–244.
- Vázquez-Moreno L, Porath J, Schluter SF, Marchalonis JJ, Purification of a novel heterodimer from shark (*Carcharhinus plumbeus*) serum by gel-immobilized metal chromatography. *Com. Biochem. Physiol.*, 1992; 103B: 563-568.
- Vázquez-Moreno L, Calderon de la Barca AM, Robles-Burgueño R, Comparison of Divinylsulphone activated agarose with different matrix materials to purify lectins by affinity chromatography, in E, Van Driessche, H. Franz, S. Beeckmans, U. Pfiller, A. Allikorm, T.C. Bog-Hansen TC (Eds), *Lectins: Biology, biochemistry, clinical biochemistry*, Vol. 5. Helleurp, Denmark, 1993, 110-114.
- Wilson RA, Francis DH, Fimbriae and enterotoxins associated with *E. coli* serotypes isolated from clinical cases of porcine colibacillosis. *Am. J. Vet. Res.*, 1986; 47: 213-217.
- Wold AE, Mestecky J, Tomana M, Kobata A, Ohbayashi H, Endo T, Edén CS, Secretory immunoglobulin A carries oligosaccharide receptors for *Escherichia coli* type 1 fimbrial lectin, *Infect Immun.*, 1990; 58: 3073–3077.
- Wood G, Warnke R, Suppression of endogenous avidin binding in tissues and its relevance to biotin-avidin detection systems, *J. Histoche and Cytochem.*, 1981; 29: 1196-1204.