

---

## Editorial:

### CARCINOGENESIS AND GENOTOXICITY

R. Marchan

Leibniz Research Centre for Working Environment and Human Factors (IfADo), Ardeystrasse 67, 44139 Dortmund, Germany

E-mail: [marchan@ifado.de](mailto:marchan@ifado.de), Telephone: +49 231-1084-213, Fax: +49 231-1084-403

Carcinogenesis (Table 1A) and genotoxicity (Table 1B) are two of the most popular topics published in our partner journal *Archives of Toxicology*. A particular highlight was an article published in 2008 by Köhle et al. on hepatocarcinogenesis in humans and rodents. It is fairly well-accepted that major interspecies differences between human and rodents can compromise interspecies extrapolation in hepatocarcinogenesis. In their publication, Köhle et al. (2008) addressed the topic of interspecies extrapolation and concluded that a relatively high degree of similarity exists between humans and rodent for genotoxic-initiating carcinogens. However, remarkable species differences have been observed for nongenotoxic tumor promoters. Köhle and colleagues differentiated between two modes of action: (1) chronic cytotoxicity leading to replacement proliferation, which often occurs in both humans and rodents, and (2) sustained activation of orphan receptors such as CAR, PPAR alpha and the Ah receptor. Köhle et al. report that these mechanisms are much more relevant in rodents than in humans which may explain a considerable part of interspecies differences.

A second highlight is the novel classification of carcinogens by the SCOEL (Scientific Committee on Occupational Exposure Limits) reported by Bolt and Huici-Montagud (2008), where carcinogens are classified according to the possibility to establish either “practical” or “true” thresholds. Class A represents non-threshold genotoxic carcinogens where the linear non-threshold model appears appropriate. Class B comprises genotoxic carcinogens for which the existence of a threshold cannot be sufficiently supported. Therefore because of the scientific uncertainty, a linear non-threshold model should be used. Class C includes carcinogens with a practical threshold (as described by Hengstler et al., 2003) where exposure limits may be based on an established NOAEL. Class D are non-genotoxic and non-DNA-reactive carcinogens where a “true” or “perfect” threshold can be established.

An overview of the publications on carcinogenesis and genotoxicity is given in Table 1.

**Table 1A:** Recent studies on **carcinogenesis**

Key message	Reference
Troglitazone, an insulin-sensitizing agent, shows a weak tumor-promoting effect in the rasH2/urethane mouse model.	Jin et al., 2008
Piperonyl butoxide activates c-Jun and ATF-2 in mouse hepatocytes, which may play a role in early stages of hepatocarcinogenesis.	Muguruma et al., 2008
This article focuses on interspecies differences of hepatocarcinogenesis in humans and rodents. While a high degree of similarity is often observed for genotoxic carcinogens, considerable species differences have been observed for the nongenotoxic carcinogens. Two modes of action are of particular relevance: (i) sustained activation of receptors such as CAR, PPARalpha and the Ah receptor, (ii) chronic cytotoxicity leading to replacement proliferation	Köhle et al., 2008
Dietary diphenyl diselenide delays chemically induced breast cancer in rats.	de Vargas Barbosa et al., 2008
2-Methylimidazole, which has been identified in food and is used in the manufacture of dyes and rubber, induces thyroid and liver tumors in rats and mice.	Chan et al., 2008a
The size of GST-P liver foci is associated with the rates of cell division and death.	Lu et al., 2008
The Scientific Committee on Occupational Exposure Limits (SCOEL) recommends consideration of four groups of carcinogens: (A) non-threshold genotoxic carcinogens, (B) genotoxic carcinogens for which the existence of a threshold cannot be sufficiently supported at present, (C) genotoxic carcinogens with a practical threshold, (D) non-genotoxic and non-DNA reactive carcinogens for which a true threshold can be derived.	Bolt and Huici-Montagud, 2008
4-Methylimidazole causes alveolar and bronchiolar adenoma and carcinoma in rats.	Chan et al., 2008b

**Table 1B:** Recent progress in **genotoxicity research**

Key message	Reference
The Fast Micromethod is a 96-well microplate assay for identification of DNA strand breaks. The technique was further improved by omission of the cell lysis step and the time-consuming cell-counting.	Ullmann et al., 2008
Wastewater from a petrochemical plant, a petroleum refinery and a coke plant of the steel industry caused chromosomal aberrations in mononuclear blood cells.	Krishnamurthi et al., 2008
Vitamin B6 antagonizes the clastogenic effect of doxorubicin in rat bone marrow cells.	Takeuchi et al., 2008
The Ty1 assay detects a wide spectrum of genotoxic carcinogens based on induction of the transposition of a Ty1 retrotransposon in yeast cells.	Pesheva et al., 2008
Inhibition of poly (ADP-ribose) polymerase (PARP) can shift balances between apoptotic and necrotic cell death in HaCaT cells.	Kehe et al., 2008
Non-specific chromosomal genotoxicity can be predicted by physico-chemical properties, such as the relation of polar surface to the total molecular surface of compounds.	Dorn et al., 2008a
Anabolic doping steroids induce micronuclei in V79 cells.	Dorn et al., 2008b
This review discusses the mechanisms responsible for chromosomal instability in urothelial carcinomas.	Flori and Schulz, 2008

Key message	Reference
Panax ginseng extract reduces the genotoxicity of ethylenediamine-tetraacetic acid in rats.	Khalil et al., 2008
Phosphate fluoride is used by dental practitioners to prevent caries in children. However, this study has shown that phosphate fluoride induces DNA strand breaks in epithelial cells.	Tsai et al., 2008
Ginsenoside Rg(3) protects against cyclophosphamide-induced DNA damage in bone marrow cells of mice.	Zhang et al., 2008

## REFERENCES

- Bolt HM, Huici-Montagud A. Strategy of the scientific committee on occupational exposure limits (SCOEL) in the derivation of occupational exposure limits for carcinogens and mutagens. *Arch Toxicol* 2008;82:61-4.
- Chan PC, Sills RC, Kissling GE, Nyska A, Richter W. Induction of thyroid and liver tumors by chronic exposure to 2-methylimidazole in F344/N rats and B6C3F1 mice. *Arch Toxicol* 2008a;82:399-412.
- Chan PC, Hill GD, Kissling GE, Nyska A. Toxicity and carcinogenicity studies of 4-methylimidazole in F344/N rats and B6C3F1 mice. *Arch Toxicol*. 2008b;82:45-53.
- de Vargas Barbosa NB, Nogueira CW, Guecheva TN, Bellinaso Mde L, Rocha JB. Diphenyl diselenide supplementation delays the development of N-nitroso-N-methyl-urea-induced mammary tumors. *Arch Toxicol* 2008;82:655-63.
- Dorn SB, Degen GH, Bolt HM, van der Louw J, van Acker FA, van den Dobbelen DJ, Lommerse JP. Some molecular descriptors for non-specific chromosomal genotoxicity based on hydrophobic interactions. *Arch Toxicol* 2008a;82:333-8.
- Dorn SB, Bolt HM, Thevis M, Diel P, Degen GH. Induction of micronuclei in V79 cells by the anabolic doping steroids tetrahydrogestrinone and trenbolone. *Arch Toxicol* 2008b;82:257-63.
- Florl AR, Schulz WA. Chromosomal instability in bladder cancer. *Arch Toxicol* 2008;82:173-82.
- Hengstler JG, Bogdanffy MS, Bolt HM, Oesch F: Challenging dogma: thresholds for genotoxic carcinogens? The case of vinyl acetate. *Annu Rev Pharmacol Toxicol* 2003;43:485-520.
- Jin M, Matsumoto S, Dewa Y, Nishimura J, Saekusa Y, Hasumi K, Mitsumori K. Extremely weak tumor-promoting effect of troglitazone on splenic hemangiosarcomas in rasH2 mice induced by urethane. *Arch Toxicol* 2008;82:771-7.
- Kehe K, Raithel K, Kreppel H, Jochum M, Worek F, Thiermann H. Inhibition of poly(ADP-ribose) polymerase (PARP) influences the mode of sulfur mustard (SM)-induced cell death in HaCaT cells. *Arch Toxicol* 2008;82:461-70.
- Khalil WK, Ahmed KA, Park MH, Kim YT, Park HH, Abdel-Wahhab MA. The inhibitory effects of garlic and Panax ginseng extract standardized with ginsenoside Rg3 on the genotoxicity, biochemical, and histological changes induced by ethylenediaminetetraacetic acid in male rats. *Arch Toxicol* 2008;82:183-95.

Köhle C, Schwarz M, Bock KW: Promotion of hepatocarcinogenesis in humans and animal models. *Arch Toxicol* 2008;82:623-31.

Krishnamurthi K, Saravana Devi S, Hengstler JG, Hermes M, Kumar K, Dutta D, Muhil Vannan S, Subin TS, Yadav RR, Chakrabarti T. Genotoxicity of sludges, wastewater and effluents from three different industries. *Arch Toxicol* 2008;82:965-71.

Lu Y, Lohitnavy M, Reddy M, Lohitnavy O, Eickman E, Ashley A, Gerjevic L, Xu Y, Conolly RB, Yang RS. Quantitative analysis of liver GST-P foci promoted by a chemical mixture of hexachlorobenzene and PCB 126: implication of size-dependent cellular growth kinetics. *Arch Toxicol* 2008;82:103-16.

Muguruma M, Arai K, Moto M, Nishimura J, Dewa Y, Mitsumori K. Piperonyl butoxide activates c-Jun and ATF-2 in the hepatocytes of mice. *Arch Toxicol* 2008;82:749-53.

Pesheva M, Krastanova O, Stamenova R, Kantardjiev D, Venkov P. The response of Ty1 test to genotoxins. *Arch Toxicol* 2008; 82:779-85.

Takeuchi PL, Antunes LM, Takahashi CS. Modulation of doxorubicin-induced clastogenesis in Wistar rat bone marrow cells by vitamin B(6). *Arch Toxicol* 2008;82:869-73.

Tsai CL, Lin JW, Kuo HK, Tai MH, Wu YC, Shyr CR, Wu PC. Induction of apoptosis in rabbit oral mucosa by 1.23% acidulated phosphate fluoride gel. *Arch Toxicol* 2008;82:81-7.

Ullmann K, Müller C, Steinberg P. Two essential modifications strongly improve the performance of the Fast Micromethod to identify DNA single- and double-strand breaks. *Arch Toxicol* 2008;82:861-7.

Zhang QH, Wu CF, Duan L, Yang JY. Protective effects of ginsenoside Rg(3) against cyclophosphamide-induced DNA damage and cell apoptosis in mice. *Arch Toxicol* 2008;82:117-23.