

Guest editorial:

HIGHLIGHT REPORT: NEW METHODS FOR QUANTIFICATION OF BILE CANALICULAR DYNAMICS

Ahmed Ghallab

Forensic Medicine and Toxicology Department, Faculty of Veterinary Medicine,
South Valley University, Qena, Egypt, E-mail: ghallab@vet.svu.edu.eg

<http://dx.doi.org/10.17179/excli2015-763>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).

Cholestasis represents one of the main causes of hepatotoxicity. It is defined by impaired bile flux from hepatocytes to the gall bladder. Cholestasis can be caused by inhibition of the transport of bile acids from hepatocytes into bile canaliculi. Moreover, physical obstructions or impaired flux in the canaliculi or bile ducts may be responsible. Since decades cultivated hepatocytes have been used to study the cholestatic properties of chemicals (Dunn et al., 1989). Tests in this field are based on the release and subsequent analysis of the canalicular content (Liu et al., 1999a-c; Bi et al., 2006; Kostrubsky et al., 2003). However, relatively little is known whether compromised bile canalicular dynamics, the rhythmic contractions of bile canaliculi, play a role in cholestasis. Therefore, Raymond Reif and colleagues have established an *in vitro* system that based on time-lapse microscopy allows the quantification of bile canalicular dynamics (Reif et al., 2015). This technique measures the repetitive swelling and collapsing of bile canaliculi. This period typically lasts for approximately 30 h after plating of mouse hepatocytes. Afterwards, the luminal diameter of the bile canaliculi becomes narrower and more constant (Reif et al., 2015). In this state it resembles morphology and dynamics of bile canalicular structures *in vivo* (Reif et al., 2015). An interesting observation is that exposure to dexamethasone/insulin increased

both, the amplitude of the initial contractions during the first 30 h and also the diameter of the mature canaliculi between 30 and 75 h after plating of the hepatocytes. In conclusion, Reif and colleagues have established a relatively easy to handle technique to quantify bile canalicular dynamics and they have shown that this parameter can be influenced by chemicals. However, whether compromised bile canalicular dynamics play a key role in pathogenesis of cholestasis still remains to be studied in future.

Prediction of hepatotoxicity represents a cutting edge topic in toxicology (Ghallab et al., 2015; Godoy et al., 2009, 2013, 2015; Kim et al., 2015; Hengstler et al., 2014; Drasdo et al., 2014a, b; Frey et al., 2014; Mazzanti et al., 2015; Tolosa et al., 2015; Chen et al., 2014; Vitins et al., 2014; Liu et al., 2014). An intensively studied field are *in vitro* systems for hepatotoxicity, because they may improve the throughput and avoid difficulties due to interspecies extrapolation, because human cells can be used (Schyschka et al., 2013; Messner et al., 2013; Wobus and Löser, 2011; O'Brien et al., 2006). However, because of their limitations, *in vitro* systems have also been critically discussed (Reif, 2014; Ghallab, 2013, 2014a, b; Stöber, 2014; Hammad, 2013). Many *in vitro* systems in the field of hepatotoxicity focus on alterations in gene expression caused by the test compounds (Shinde et al., 2015a, b; Campos

et al., 2014; Rodrigues et al., 2013; Driessen et al., 2013; Tolosa et al., 2013; Grinberg et al., 2014). Other endpoints, such as bile canalicular dynamics have been studied only rarely, possibly due to technical challenges. Therefore, it is fortunate that techniques for analysis of bile canalicular dynamics are now available. However, if - and if yes to which degree - compromised motility of bile canaliculi contributes to hepatotoxicity still has to be elucidated.

REFERENCES

- Bi YA, Kazolias D, Duignan DB. Use of cryopreserved human hepatocytes in sandwich culture to measure hepatobiliary transport. *Drug Metab Dispos.* 2006;34:1658-65.
- Campos G, Schmidt-Heck W, Ghallab A, Rochlitz K, Pütter L, Medinas DB, et al. The transcription factor CHOP, a central component of the transcriptional regulatory network induced upon CCl4 intoxication in mouse liver, is not a critical mediator of hepatotoxicity. *Arch Toxicol.* 2014;88:1267-80.
- Chen M, Tung CW, Shi Q, Guo L, Shi L, Fang H, et al. A testing strategy to predict risk for drug-induced liver injury in humans using high-content screen assays and the 'rule-of-two' model. *Arch Toxicol.* 2014;88:1439-49.
- Drasdo D, Bode J, Dahmen U, Dirsch O, Dooley S, Gebhardt R, et al. The virtual liver: state of the art and future perspectives. *Arch Toxicol.* 2014a;88:2071-5.
- Drasdo D, Hoehme S, Hengstler JG. How predictive quantitative modelling of tissue organisation can inform liver disease pathogenesis. *J Hepatol.* 2014b;61:951-6.
- Driessen M, Kienhuis AS, Pennings JL, Pronk TE, van de Brandhof EJ, Roodbergen M, et al. Exploring the zebrafish embryo as an alternative model for the evaluation of liver toxicity by histopathology and expression profiling. *Arch Toxicol.* 2013;87:807-23.
- Dunn JC, Yarmush ML, Koebe HG, Tompkins RG. Hepatocyte function and extracellular matrix geometry: long-term culture in a sandwich configuration. *FASEB J.* 1989;3:174-7.
- Frey O, Misun PM, Fluri DA, Hengstler JG, Hierlemann A. Reconfigurable microfluidic hanging drop network for multi-tissue interaction and analysis. *Nat Commun.* 2014;5:4250.
- Ghallab A. In vitro test systems and their limitations. *EXCLI J.* 2013;12:1024-6.
- Ghallab A. The rediscovery of HepG2 cells for prediction of drug induced liver injury (DILI). *EXCLI J.* 2014a;13:1286-8.
- Ghallab A. Human non-parenchymal liver cells for co-cultivation systems. *EXCLI J.* 2014b;13:1295-6.
- Ghallab A, Cellière G, Henkel SG, Driesch D, Hoehme S, Hofmann U, et al. Model guided identification and therapeutic implications of an ammonia sink mechanism. *J Hepatol.* 2015; pii: S0168-8278(15)00776-X. doi: 10.1016/j.jhep.2015.11.018. [Epub ahead of print].
- Godoy P, Hengstler JG, Ilkavets I, Meyer C, Bachmann A, Müller A, et al. Extracellular matrix modulates sensitivity of hepatocytes to fibroblastoid dedifferentiation and transforming growth factor beta-induced apoptosis. *Hepatology.* 2009;49:2031-43.
- Godoy P, Hewitt NJ, Albrecht U, Andersen ME, Ansari N, Bhattacharya S, et al. Recent advances in 2D and 3D in vitro systems using primary hepatocytes, alternative hepatocyte sources and non-parenchymal liver cells and their use in investigating mechanisms of hepatotoxicity, cell signaling and ADME. *Arch Toxicol.* 2013;87:1315-530.
- Godoy P, Schmidt-Heck W, Natarajan K, Lucendo-Villarin B, Szkolnicka D, Asplund A, et al. Gene networks and transcription factor motifs defining the differentiation of stem cells into hepatocyte-like cells. *J Hepatol.* 2015;63:934-42.
- Grinberg M, Stöber RM, Edlund K, Rempel E, Godoy P, Reif R, et al. Toxicogenomics directory of chemically exposed human hepatocytes. *Arch Toxicol.* 2014;88:2261-87.
- Hammad S. Advances in 2D and 3D in vitro systems for hepatotoxicity testing. *EXCLI J.* 2013;12:993-6.
- Hengstler JG, Marchan R, Bolt HM. Standard compounds for establishment of in vitro test systems. *Arch Toxicol.* 2014;88:2083-4.
- Kim JY, Fluri DA, Marchan R, Boonen K, Mohanty S, Singh P, et al. 3D spherical microtissues and microfluidic technology for multi-tissue experiments and analysis. *J Biotechnol.* 2015;205:24-35.
- Kostrubsky VE, Strom SC, Hanson J, Urda E, Rose K, Burliegh J, et al. Evaluation of hepatotoxic potential of drugs by inhibition of bile-acid transport in cultured primary human hepatocytes and intact rats. *Toxicol Sci.* 2003;76:220-8.
- Liu A, Krausz KW, Fang ZZ, Brocker C, Qu A, Gonzalez FJ. Gemfibrozil disrupts lysophosphatidylcholine and bile acid homeostasis via PPAR α and its relevance to hepatotoxicity. *Arch Toxicol.* 2014;88:983-96.

- Liu X, Chism JP, LeCluyse EL, Brouwer KR, Brouwer KL. Correlation of biliary excretion in sandwich-cultured rat hepatocytes and in vivo in rats. *Drug Metab Dispos.* 1999a;27:637-44.
- Liu X, LeCluyse EL, Brouwer KR, Gan LS, Lemasters JJ, Stieger B, et al. Biliary excretion in primary rat hepatocytes cultured in a collagen-sandwich configuration. *Am J Physiol.* 1999b;277:G12-21.
- Liu X, LeCluyse EL, Brouwer KR, Lightfoot RM, Lee JJ, Brouwer KL. Use of Ca²⁺ modulation to evaluate biliary excretion in sandwich-cultured rat hepatocytes. *J Pharmacol Exp Ther.* 1999c;289:1592-9.
- Mazzanti G, Di Sotto A, Vitalone A. Hepatotoxicity of green tea: an update. *Arch Toxicol.* 2015;89:1175-91.
- Messner S, Agarkova I, Moritz W, Kelm JM. Multi-cell type human liver microtissues for hepatotoxicity testing. *Arch Toxicol.* 2013;87:209-13.
- O'Brien PJ, Irwin W, Diaz D, Howard-Coffield E, Krejsa CM, Slaughter MR, et al. High concordance of drug-induced human hepatotoxicity with in vitro cytotoxicity measured in a novel cell-based model using high content screening. *Arch Toxicol.* 2006;80:580-604.
- Reif R. The body-on-a-chip concept: possibilities and limitations. *EXCLI J.* 2014;13:1283-5.
- Reif R, Karlsson J, Günther G, Beattie L, Wrangborg D, Hammad S, et al. Bile canalicular dynamics in hepatocyte sandwich cultures. *Arch Toxicol.* 2015;89:1861-70.
- Rodrigues AV, Rollison HE, Martin S, Sarda S, Schulz-Utermoehl T, Stahl S, et al. In vitro exploration of potential mechanisms of toxicity of the human hepatotoxic drug fenclozic acid. *Arch Toxicol.* 2013;87:1569-79.
- Schyschka L, Sánchez JJ, Wang Z, Burkhardt B, Müller-Vieira U, Zeilinger K, et al. Hepatic 3D cultures but not 2D cultures preserve specific transporter activity for acetaminophen-induced hepatotoxicity. *Arch Toxicol.* 2013;87:1581-93.
- Shinde V, Klima S, Sureshkumar PS, Meganathan K, Jagtap S, Rempel E, et al. Human pluripotent stem cell based developmental toxicity assays for chemical safety screening and systems biology data generation. *J Vis Exp.* 2015a;17;(100):e52333.
- Shinde V, Stöber R, Nemade H, Sotiriadou I, Hescheler J, Hengstler J, et al. Transcriptomics of hepatocytes treated with toxicants for investigating molecular mechanisms underlying hepatotoxicity. *Methods Mol Biol.* 2015b;1250:225-40.
- Stöber R. Transcriptome based differentiation of harmless, teratogenic and cytotoxic concentration ranges of valproic acid. *EXCLI J.* 2014;13:1281-2.
- Tolosa L, Gómez-Lechón MJ, Pérez-Cataldo G, Castell JV, Donato MT. HepG2 cells simultaneously expressing five P450 enzymes for the screening of hepatotoxicity: identification of bioactivable drugs and the potential mechanism of toxicity involved. *Arch Toxicol.* 2013;87:1115-27.
- Tolosa L, Gómez-Lechón MJ, Donato MT. High-content screening technology for studying drug-induced hepatotoxicity in cell models. *Arch Toxicol.* 2015;89:1007-22.
- Vitins AP, Kienhuis AS, Speksnijder EN, Roodbergen M, Luijten M, van der Ven LT. Mechanisms of amiodarone and valproic acid induced liver steatosis in mouse in vivo act as a template for other hepatotoxicity models. *Arch Toxicol.* 2014;88:1573-88.
- Wobus AM, Löser P. Present state and future perspectives of using pluripotent stem cells in toxicology research. *Arch Toxicol.* 2011;85:79-117.