

3-4-2009

# Animal pigment bilirubin discovered in plants

Cary Pirone

*Department of Biological Sciences, Florida International University*

J. Martin E. Quirke

*Department of Chemistry and Biochemistry, Florida International University*

Horacio A. Priestap

*Department of Biological Sciences, Florida International University*

David W. Lee

*Department of Biological Sciences, Florida International University*

Follow this and additional works at: [https://digitalcommons.fiu.edu/cas\\_bio](https://digitalcommons.fiu.edu/cas_bio)



Part of the [Biology Commons](#)

---

## Recommended Citation

Pirone, C., Quirke, J. M., Priestap, H. A., & Lee, D. W. (January 01, 2009). Animal pigment bilirubin discovered in plants. *Journal of the American Chemical Society*, 131, 8.)

This work is brought to you for free and open access by the College of Arts, Sciences & Education at FIU Digital Commons. It has been accepted for inclusion in Department of Biological Sciences by an authorized administrator of FIU Digital Commons. For more information, please contact [dcc@fiu.edu](mailto:dcc@fiu.edu).



Published in final edited form as:

*J Am Chem Soc.* 2009 March 4; 131(8): 2830. doi:10.1021/ja809065g.

## The Animal Pigment Bilirubin Discovered in Plants

Cary Pirone<sup>†,\*</sup>, J. Martin E. Quirke<sup>‡</sup>, Horacio A. Priestap<sup>†</sup>, and David W. Lee<sup>†</sup>

<sup>†</sup> Department of Biological Sciences, Florida International University, Miami, 33199, USA

<sup>‡</sup> Department of Chemistry and Biochemistry, Florida International University, Miami, 33199, USA

Cyclic tetrapyrroles occur throughout the plant kingdom and include vital biosynthetic products such as chlorophyll and heme. In plants, oxidative degradation of heme forms first biliverdin-IX $\alpha$  and subsequently phytychromobilin, the precursor of the phytochrome chromophore, an essential light sensing molecule<sup>1</sup>. In animals, oxidative degradation of heme also leads to the formation of biliverdin-IX $\alpha$ , but it is transformed into the yellow-orange pigment bilirubin-IX $\alpha$ . Here, we present spectroscopic and chromatographic evidence that bilirubin (Figure 1) is the major pigment of the orange aril of *Strelitzia nicolai* Regel & Koern. (Strelitziaceae, order Zingiberales), the white bird of paradise tree.

This is the first example of bilirubin in a plant<sup>2</sup>, a finding which likely necessitates the revision of the plant tetrapyrrole pathway since there is currently no known mechanism of bilirubin production in the plant kingdom.

*S. nicolai* is native to South Africa and widely cultivated in the tropics. It produces woody capsular fruits which contain orange arillate seeds. Analytical high-performance liquid chromatography (HPLC) of the aril extract<sup>3</sup> revealed one major peak, which had a UV-visible spectrum with a maximum absorbance at 444 nm. After purification using preparative scale HPLC, the isolated pigment was analyzed by <sup>1</sup>H NMR, <sup>13</sup>C NMR (Bruker, 400 MHz, (CD<sub>3</sub>)<sub>2</sub>S=O), and liquid chromatography-positive ion electrospray mass spectrometry (LC-ESI), (Thermo-Finnigan LCQ).

The <sup>1</sup>H NMR and <sup>13</sup>C NMR spectra of the isolated pigment matched published values of authentic bilirubin<sup>4,5,10</sup> (Tables S1 and S2). Identification was further supported by <sup>1</sup>H NMR analysis of bilirubin standard (Aldrich). This yielded a spectrum which matched that of the *S. nicolai* pigment (Figure S1). Both the positive ion ES mass spectrum and the product ion spectrum matched those of authentic bilirubin standard and previous published data<sup>6</sup> (molecular ion, *m/z* 585 (M + H)<sup>+</sup>, product ion *m/z* 299).

Given the unexpected discovery of bilirubin in plants, it was essential to confirm the identity of the pigment as bilirubin-IX $\alpha$ , and not other isomers. Previous chromatographic studies have demonstrated that the ability of bilirubin-IX $\alpha$  to undergo intramolecular hydrogen bonding makes it significantly less polar than bilirubin-IX $\beta,\gamma$  or  $\delta$ <sup>7,8</sup> (Figure 1).

In our HPLC analyses, a single peak was observed when bilirubin-IX $\alpha$  standard was co-injected with the isolated pigment, thereby eliminating the possibility that the pigment was bilirubin-IX $\beta,\gamma$  or  $\delta$ . Furthermore, the visible spectrum of bilirubin-IX $\alpha$  has an intense peak at 458 nm in dimethylsulfoxide (DMSO), which is approximately 50 nm longer than bilirubin-

Email: cary.pirone@fiu.edu.

Supporting Information Available: Methods, UV-VIS spectrum, and <sup>1</sup>H and <sup>13</sup>C NMR data. This material is available free of charge on the internet at <http://pubs.acs.org>.

IX  $\beta,\gamma$  or  $\delta^7$ . Other bilirubin isomers, including bilirubin-III $\alpha$  and bilirubin-XIII $\alpha$ , were eliminated because their  $^1\text{H}$  NMR spectra are substantially different<sup>9</sup>.

The occurrence of bilirubin is not restricted to *S. nicolai*. Two other species in the Strelitziaceae, *Phenakospermum guyanense* Endl., and *S. reginae* Aiton, the bird of paradise, contain aril pigments which co-eluted with authentic bilirubin in HPLC and had similar UV-visible spectra. We are currently examining species in related families. This information, in combination with studies on the synthesis of bilirubin-IX $\alpha$  in *S. nicolai*, will provide the basis for a more thorough understanding of the evolutionary origin of this pigment in plants.

## Supplementary Material


Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

This research has been funded in part by the United States Environmental Protection Agency (EPA) under the Greater Research Opportunities (GRO) Graduate Program, the McBryde Science Program at the National Tropical Botanical Garden, and the RISE Biomedical Research Initiative (NIH). We thank the Center for Ethnobiology and Natural Products (CeNAP) at FIU for lab facilities.

## References

1. Tanaka R, Tanaka A. *Ann Rev Plant Biol* 2007;58:321–346. [PubMed: 17227226]
2. Willows, RD.; Cuttriss, A.; Pogson, B.; Schwinn, KE.; Davies, KM.; Zryd, JP.; Christinet, L. *Plant Pigments and Their Manipulation*. Davies, KM., editor. Vol. Chapters 2–7. Blackwell; Oxford, UK: 2004.
3. Spivak W, Yuey W. *Biochem J* 1986;234:101–109. [PubMed: 3707537]
4. Kuenzle C. *Biochem J* 1970;119:395–409. [PubMed: 5500301]
5. Kaplan D, Navon G. *Org Mag Res* 1980;13:59–62.
6. Lightner, DA. *Bilirubin*. Heirwegh, KPM.; Brown, SB., editors. Vol. Chapter 1. CRC Press; Boca Raton, FL: 1982. p. 1
7. Blanckart N, Heirwegh KPM, Compennolle F. *Biochem J* 1976;155:405–417. [PubMed: 938489]
8. Bonnett R, Davies JE, Hursthouse MB, Sheldrich GM. *Proc R Soc Lond B* 1978;202:249–268. [PubMed: 28528]
9. McDonagh AF, Palma LA, Lightner DA. *J Am Chem Soc* 1982;104:6865–6867.
10. Muller N. *Magn Reson Chem* 1985;23:688–689.



**Figure 1.**  
Bilirubin-IX $\alpha$  demonstrating intramolecular H-bonding.