

## Florida International University FIU Digital Commons

FCE LTER Journal Articles

FCE LTER

2009

# Roseate spoonbill reproduction as an indicator for restoration of the Everglades and the Everglades estuaries

Jerome J. Lorenz

*Audubon of Florida, Tavernier Science Center*

Brynne Langan-Mulrooney

*Audubon of Florida, Tavernier Science Center*

Peter E. Frezza

*Audubon of Florida, Tavernier Science Center*

Rebecca G. Harvey

*University of Florida, Ft. Lauderdale Research and Education Center*

Frank J. Mazzotti

*University of Florida, Ft. Lauderdale Research and Education Center*

Follow this and additional works at: [http://digitalcommons.fiu.edu/fce\\_lter\\_journal\\_articles](http://digitalcommons.fiu.edu/fce_lter_journal_articles)

 Part of the [Life Sciences Commons](#)

### Recommended Citation

Lorenz, J.J., B. Langan-Mulrooney, P. Frezza, R.G. Harvey, F.J. Mazzotti. 2009. Roseate spoonbill reproduction as an indicator for restoration of the Everglades and the Everglades estuaries. *Ecological Indicators* 9(6): S96-S107.

This material is based upon work supported by the National Science Foundation through the Florida Coastal Everglades Long-Term Ecological Research program under Cooperative Agreements #DBI-0620409 and #DEB-9910514. Any opinions, findings, conclusions, or recommendations expressed in the material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

This work is brought to you for free and open access by the FCE LTER at FIU Digital Commons. It has been accepted for inclusion in FCE LTER Journal Articles by an authorized administrator of FIU Digital Commons. For more information, please contact [dcc@fiu.edu](mailto:dcc@fiu.edu), [jkrefft@fiu.edu](mailto:jkrefft@fiu.edu).

Elsevier Editorial System(tm) for Ecological Indicators  
Manuscript Draft

Manuscript Number:

Title: Roseate Spoonbills as an Indicator for Restoration of the Everglades and Florida Bay

Article Type: Indicators for Restoration

Section/Category:

Keywords: ecological indicators, Everglades restoration, Roseate Spoonbill, Wading Birds, restoration assessment

Corresponding Author: Dr. Jerome J Lorenz, Ph.D.

Corresponding Author's Institution: Audubon of Florida

First Author: Jerome J Lorenz, Ph.D.

Order of Authors: Jerome J Lorenz, Ph.D.; Brynne Langan-Mulrooney, B.S.; Peter E Frezza, MS; Frank J Mazzotti, Ph.D.

Manuscript Region of Origin:

Abstract: Ecological monitoring is a key part of adaptive management and successful restoration. Not everything within an ecosystem can be monitored so it is important to select indicators that are representative of the system, integrate system responses, show clear responses to system change, can be effectively and efficiently monitored, and are easily communicated. Roseate Spoonbills are one of the indicators that meet these criteria within the Everglades ecosystem. Monitoring of Roseate Spoonbills in Florida Bay over the past 70 years has shown that this species responds to changes in hydrology and corresponding changes in prey abundance and availability. This indicator uses nesting location, nest numbers and nesting success in response to food abundance and availability. In turn, prey abundance is a function of hydrological conditions including depth, and salinity. These relationships have been well documented such that spoonbills responses can be directly related to changes in hydrology and salinity. The spoonbill indicator uses performance measures that have been shown to be both effective and efficient

in tracking trends. They are: nesting success, nest number, locations of nests, and prey fish community composition. Targets for these performance measures we established based on previous findings. The performance measures are then reported as suitability indices identified as stoplight colors with green indicating that targets have been met, yellow indicating that conditions are below the target but within a suitable range of it and red indicating the measure is performing poorly in relation to the target.

1            **Roseate Spoonbills as an Indicator for Restoration of the Everglades and**  
2            **Florida Bay**

3

4            Jerome J. Lorenz\*, Audubon of Florida, Tavernier Science Center, 115 Indian  
5            Mound Trail, Tavernier FL 33070. Ph: 305-852-5318, Fax: 305-852-8012,  
6            [jlorenz@audubon.org](mailto:jlorenz@audubon.org)

7

8            Brynne Langan-Mulrooney, Audubon of Florida, Tavernier Science Center, 115  
9            Indian Mound Trail, Tavernier FL 33070. Ph: 305-852-5318, Fax: 305-852-8012,  
10           [blangan@audubon.org](mailto:blangan@audubon.org)

11

12           Peter E. Frezza, Audubon of Florida, Tavernier Science Center, 115 Indian  
13           Mound Trail, Tavernier FL 33070. Ph: 305-852-5318, Fax: 305-852-8012,  
14           [pfrezza@audubon.org](mailto:pfrezza@audubon.org)

15

16           Frank J. Mazzotti, University of Florida, Ft. Lauderdale Research and Education  
17           Center, 3205 College Avenue, Davie Florida 33314. [fjmaAufl.edu](mailto:fjmaAufl.edu)

18

19

20

21

22

23           \*Corresponding author

24           **Abstract**

25           Ecological monitoring is a key part of adaptive management and successful  
26 restoration. Not everything within an ecosystem can be monitored so it is important to  
27 select indicators that are representative of the system, integrate system responses, show  
28 clear responses to system change, can be effectively and efficiently monitored, and are  
29 easily communicated. Roseate Spoonbills are one of the indicators that meet these  
30 criteria within the Everglades ecosystem. Monitoring of Roseate Spoonbills in Florida  
31 Bay over the past 70 years has shown that this species responds to changes in hydrology  
32 and corresponding changes in prey abundance and availability. This indicator uses  
33 nesting location, nest numbers and nesting success in response to food abundance and  
34 availability. In turn, prey abundance is a function of hydrological conditions including  
35 depth, and salinity. These relationships have been well documented such that spoonbills  
36 responses can be directly related to changes in hydrology and salinity. The spoonbill  
37 indicator uses performance measures that have been shown to be both effective and  
38 efficient in tracking trends. They are: nesting success, nest number, locations of nests,  
39 and prey fish community composition. Targets for these performance measures were  
40 established based on previous findings. The performance measures are then reported as  
41 suitability indices identified as stoplight colors with green indicating that targets have  
42 been met, yellow indicating that conditions are below the target but within a suitable  
43 range of it and red indicating the measure is performing poorly in relation to the target.

44

45           Key words: ecological indicators, Everglades restoration, Roseate Spoonbill,  
46 Wading Birds, restoration assessment

47 **1. Introduction and Background**

48

49 Ecological monitoring is a key part of adaptive management (Williams et al.,  
50 2007, Lovett et al., 2007) and successful restoration. Not everything within an ecosystem  
51 can be monitored so it is important to select indicators that are representative of the  
52 system, integrate system responses, show clear responses to system change, can be  
53 effectively and efficiently monitored, and are easily communicated (Doren, 2006, Doren  
54 et al., intro chapter, Schiller et al., 2001).

55

56 Roseate Spoonbills are one of the indicators that meet these criteria within the  
57 Everglades ecosystem. Restoration of hydrology is a major part of the Comprehensive  
58 Everglades Restoration Plan (CERP, U.S. Army Corps of Engineers, 1999), and  
59 indicators used for tracking progress of Everglades restoration should have clear  
60 relationships to hydrologic conditions (Doren et al., intro. chapter, U. S. Army Corps of  
61 Engineers, 2004).

62

63 Monitoring of Roseate Spoonbills (*Platalea ajaja*) in Florida Bay over the past 70  
64 years has shown that this species responds to changes in hydrology and corresponding  
65 changes in prey abundance and availability (Powell et al., 1989, Lorenz et al., 2002).  
66 This indicator uses nesting location, nest numbers and nesting success in response to food  
67 abundance and availability. In turn, prey abundance is a function of hydrological  
68 conditions including depth, and salinity (Lorenz and Serafy, 2006). These relationships

69 have been well documented such that spoonbills responses can be directly related to  
70 changes in hydrology and salinity (Lorenz and Serafy, 2006).

71

72 Spoonbill nesting success is dependent on suitable environmental conditions.  
73 Correlations between biological responses and environmental conditions contribute to an  
74 understanding of the species' status and trends over time (Lorenz, 2000, Lorenz and  
75 Serafy, 2006). The positive or negative trends of this indicator relative to hydrological  
76 changes (Lorenz, 2000, Lorenz et al., 2002, Bartell et al., 2005) permit an assessment of  
77 positive or negative trends in restoration. Restoration success or failure would be  
78 evaluated by comparing recent and future trends and status of spoonbills with historical  
79 population data and model predictions, as stated in the CERP hypotheses related to the  
80 food web (CERP Monitoring and Assessment Plan section 3.1.2.4; U. S. Army Corps of  
81 Engineers, 2004).

82

83 The spoonbill indicator uses performance measures that have been shown to be  
84 both effective and efficient in tracking trends. They include: nesting success, nest  
85 number, locations of nests, and prey fish community composition. These parameters  
86 have been correlated with hydrologic conditions including water depth, hydroperiod,  
87 timing, spatial extent and salinity, which are influenced by water management practices.

88

89 Roseate Spoonbills are one of several charismatic megafauna found in the  
90 Everglades. They are both umbrella and flagship species to which the public can relate.  
91 In addition, the parameters used to track trends are easy to understand: How have the

92 number of spoonbills changed through time? Are they as productive as they were  
93 historically? Are the animals in the places where they should be? Are their prey as  
94 abundant as under natural conditions?

95

### 96 ***1.1. Indicator History***

97       There is a seventy year intermittent database of spoonbill nesting activity in  
98 Florida Bay (Figure 1). Lorenz et al., (2002) demonstrated that nesting patterns are  
99 highly dependant on hydrologic conditions on the foraging ground most proximal to the  
100 nesting colonies (Figure 2). Spoonbills primarily feed on wetland fishes (Dumas, 2000)  
101 and time their nesting with low water levels which result in the prey base fishes  
102 becoming highly concentrated into the remaining wetted areas (Loftus and Kushlan,  
103 1987, DeAngelis et al., 1997, Lorenz, 2000). Studies suggest that tactile feeding wading  
104 birds, such as the Roseate Spoonbill, are particularly dependent on high prey density in  
105 order to successfully forage, probably more so than the visually oriented avian predators  
106 (Kahl, 1964, Frederick and Spalding, 1994, Gawlik, 2002). Tactile feeders are more  
107 efficient when prey density is very high and visual predators are more efficient at lower  
108 prey densities (Kahl, 1965). Gawlik (2002) experimentally demonstrated that two  
109 species of tactile feeders (wood storks and white ibis) abandoned foraging sites while  
110 prey was still abundant enough to attract visually oriented wading birds in high numbers.  
111 Although no spoonbills visited the study site, Gawlik's (2002) experimental approach  
112 lends empirical evidence to the idea that tactile feeders are more sensitive to prey  
113 availability. Because tactile foraging birds in general and roseate spoonbill in particular  
114 are more dependant on high prey concentration than other wading bird species (Kahl,



115 1964, Gawlik, 2002), they are more sensitive to changes in environmental conditions that  
116 determine fish concentrations, specifically water levels (Gawlik, 2002). The requirement  
117 for highly concentrated prey is exacerbated during nesting cycles when the high-energy  
118 demands of their offspring require a consistently available high density of prey items  
119 (Kahl, 1964, Lorenz, 2000, Dumas, 2000).

120

121           Beginning with the completion of a series of canals and water-control structures  
122 known as the South Dade Conveyance System (SDCS) in the early 1980's, water  
123 deliveries to Taylor Slough and northeastern Florida Bay (Figure 2) changed dramatically  
124 (Light and Dineen, 1994, McIvor et al., 1994, Lorenz, 2000). This canal system is  
125 immediately adjacent to Taylor Slough and just upstream from where the majority of  
126 spoonbills nested in Florida Bay at the time (Figure 2; Powell et al., 1989) and heavily  
127 impacted the coastal wetlands that were the primary feeding grounds for the spoonbill  
128 nesting population (Bjork and Powell, 1994). In 1979, 1,250 Roseate Spoonbill nests  
129 were located in Florida Bay, with more than half the nests located in the northeastern bay  
130 (Figure 1, Powell et al., 1989, Lorenz et al., 2002). Today, the number of nests is less  
131 than a third of that in 1979 and distribution of nesting by roseate spoonbills has shifted  
132 from northeastern Florida Bay to the northwestern region (Figure 2, Lorenz et al., 2002).  
133 The shift is attributed to the lack of nest production following the completion of the  
134 SDCS: Lorenz et al., (2002) calculated that prior to the SDCS northeastern Florida Bay  
135 produced an average of 1.38 chicks per nest attempt but dropped to 0.67 chicks per nest  
136 following its' completion. Lorenz (2000) demonstrated that this decline was the result of

137 the SDCS causing changes in hydrology and salinity that affected the production (Figure  
138 3) and availability of the spoonbill prey base.

139

140 In addition to a large nesting population in Florida Bay, spoonbills “nested in the  
141 thousands” along the southwest coast south of Cape Romano (Scott, 1889). Restoration  
142 of more historic hydrological conditions should promote greater prey abundance and  
143 availability in both Florida Bay and the southwestern estuaries of the Everglades, leading  
144 to an increase in the number of years spoonbills can successfully nest, defined as the  
145 survival of offspring to fledging. Therefore, roseate spoonbills are good indicators for  
146 evaluating the CERP’s effectiveness at restoring estuarine conditions (Lorenz et al.,  
147 2002).

148

149 The major anthropogenic perturbations to spoonbill foraging grounds have been  
150 the filling of wetlands for urban development in the upper Florida Keys and the alteration  
151 of wetland type and function along the northeast coast of Florida Bay by water  
152 management practices (Lorenz et al., 2002). A striking implication of these findings is  
153 that current water management practices in the southern Everglades have resulted in the  
154 ecological degradation of the coastal wetlands in northeastern Florida Bay.

155

## 156 ***1.2 CERP Hypotheses for Spoonbills***

157

158 A system-wide Monitoring and Assessment Plan (MAP) has been developed that  
159 describes the monitoring necessary to track ecological responses to Everglades

160 restoration (U.S. Army Corps of Engineers, 2004). Included in that plan are descriptions  
161 of selected indicators, how those indicators are linked to key aspects of restoration  
162 (hypotheses), and performance measures (monitoring parameters). MAP hypotheses for  
163 Roseate Spoonbills are:

164

165           • Spoonbill's should experience successful nesting (defined as an  
166 average production of >1chick/nest) in 7 of 10 years and average 1.5 chicks/nest  
167 overall (initially using a five year running average for nest production and a ten  
168 year running average successful years).

169           • Restore nest numbers to pre-SDCS levels of 1250 nests with at  
170 least half in the northeastern region (as defined by Lorenz et al., 2002) of Florida  
171 Bay. Although specific numbers for the pre-plume hunting era are unknown for  
172 Florida Bay, anecdotal evidence suggests that the long term target should be in  
173 excess of 2000 nests bay wide.

174           • A return of significant nesting activity along the southwestern  
175 coast of Florida in the estuarine areas of Shark River and Lostman's sloughs  
176 (Figure 2).

177

### 178 ***1.3. Areas of the Everglades this Indicator Covers***

179

180 Spoonbills are found throughout the Everglades landscape, however, the species  
181 is predominantly an indicator for the Florida Bay estuary (Figure 2) and cover the Greater  
182 Everglades and Southern Estuaries region. Spoonbills are included as attributes in the

183 Total System, Everglades Mangrove Estuaries, and Florida Bay conceptual ecological  
184 models. A monitoring and assessment plan has been developed for spoonbills nesting in  
185 Florida Bay. We perform a complete nest count of the entire bay, monitor nesting success  
186 at focal colonies in five regions of Florida Bay and perform quantitative assessments of  
187 the mangrove fish community which makes up the bulk of the spoonbill's diet while  
188 nesting in Florida Bay.

189

#### 190 **1.4. Significance of the Indicator to Everglades Restoration**

191 *1.4.1. The indicator is relevant to the Everglades ecosystem and responds to*  
192 *variability at a scale that makes it applicable to a large or portion of the ecosystem.*

193

194 Spoonbills were abundant in Florida Bay and throughout the Southern Estuaries  
195 region prior to Everglades drainage activities and have responded negatively to water  
196 management activities. They are top predators that share a common prey base (small  
197 demersal fishes) and foraging habitat with myriad other species. Spoonbills feed by  
198 tactolocation rather than visual hunting; this makes them more sensitive to perturbations  
199 than the other species dependant on the same resource (i.e., they are an early warning  
200 indicator). Spoonbill nesting productivity is directly linked to hydrologic conditions  
201 within the Southern Estuaries and nest production is linked to hydrology through the  
202 impact of water management on primary producers (e.g. periphyton, submerged aquatic  
203 vegetation) and lower trophic level consumers (i.e., prey base fishes).

204

205 *1.4.2. The indicator is feasible to implement and is scientifically defensible.*

206

207        Research on Roseate Spoonbills has been conducted for over 70 years, providing a  
208 remarkable long-term data base. Currently, there are funded cooperative research and  
209 monitoring programs with U.S. Fish and Wildlife Service, Everglades National Park,  
210 U.S. Geological Service-Biological Resources Division, U.S. Army Corps of Engineers  
211 and the South Florida Water Management District. Reliable models from such research  
212 are available that determine the impacts of water management on nesting patterns. Pattern  
213 metrics (e.g. nest numbers and nesting success) are statistically correlated to Ecosystem  
214 Drivers, and a Spatially Explicit Species Index model is being developed as part of the  
215 Across Trophic Level System Simulation modeling effort. This research has provided  
216 numerous peer reviewed journal articles. This indicator is already part of the CERP  
217 RECOVER interim goals and Food-Web Monitoring Component of the CERP MAP.

218

219            ***1.4.3. The indicator is sensitive to system drivers (stressors).***

220

221        Key environment drivers, such as water depth, hydroperiod and salinity, are  
222 statistically correlated to spoonbill nesting success (Lorenz, 2000, Lorenz et al., 2002). A  
223 causal link exists between hydropatterns, prey abundance and availability, and nesting  
224 success (Lorenz, 2000, Lorenz and Serafy, 2006). Nesting failure has been statistically  
225 linked to nest number and location in a given region such that persistent nesting failure  
226 results in a decline in nesting effort and a concurrent increase in other regions.

227

228            ***1.4.4. The indicator is integrative.***

229

230 Spoonbill nesting success is linked to fish production and in turn, fish production  
231 is linked to periphyton and SAV production. Spoonbill nesting responses are  
232 representative of hydrological improvement (i.e. Water Management). Spoonbills are  
233 also included in the CERP Food-Web Monitoring Component that includes an index  
234 of food-web function and landscape connectivity (“intactness”).

235

## 236 **2. The Spoonbill Indicator Performance Measures**

237

### 238 ***2.1. Indicator Metrics***

239

240 The spoonbill indicator consists of four performance measures:

- 241 • Nesting success (average number of chicks fledged per nesting attempt and  
242 number of years out of the last ten in which production exceeded 1.0 chicks per  
243 nest fledged)
- 244 • Number of nests
- 245 • Distribution of nests (number of nests in northeastern Florida Bay and 10,000  
246 islands area)
- 247 • Prey community structure (percent of total community that are considered  
248 freshwater species as defined by Lorenz and Serafy, 2006)

249

250 In addition there will be a metric for spoonbills nesting in the northwestern region  
251 of Florida Bay to act as a control metric for restoration efforts that will affect the  
252 northeastern region.

253

## 254 ***2.2. The Stoplight Restoration Report Card System Applied to Spoonbills***

255

256 This communication tool is based on MAP performance measures (either by  
257 module or system-wide) and is expected to be able to distinguish between responses to  
258 restoration and natural patterns. A set of parameters (Table 1) has been developed for  
259 each performance measure. Answers are translated as suitability indices identified as  
260 stoplight colors with green indicating that targets have been met, yellow indicating that  
261 conditions are below the target but within a suitable range of it and red indicating the  
262 measure is performing poorly in relation to the target. Two questions are addressed using  
263 suitability indices: 1) have we reached the restoration target, or if not, 2) are we making  
264 progress toward targets?

265

266 Methods for producing suitability curves vary among performance measures. For  
267 example, a ten-year running average was used for percentage of years that spoonbills  
268 were successful. A five-year running average was used for average annual nest  
269 production and nest numbers. Fish community structure changes to a greater percentage  
270 of freshwater species only when salinity conditions have been favorable to these species  
271 for a two to three year period, therefore this parameter will be reported as an annual  
272 metric that covers a three year period. Nesting success will be reported annually because

273 short-term water depth conditions dominate this parameter. By using this suite of  
274 performance measures this indicator covers time scales from annual to three, five and ten  
275 year cycles.

276

### 277 ***2.3. Calculation of Metrics and Thresholds for the Spoonbill Stoplight Restoration***

#### 278 ***Report Card***

279

280 *2.3.1. Spoonbill nesting success.* Lorenz et al., (2002) divided Florida Bay into  
281 five regions based on the primary foraging grounds for each of the colonies within each  
282 region (Figure 2). They also demonstrated that, under the SDCS operations, the nest  
283 productivity and nest number in the northeastern region have experienced a significant  
284 decline. The method used to calculate this metric is based on surveys of focal colonies  
285 (defined as the two largest colonies within the region) . These surveys entailed marking  
286 up to 50 nests shortly after full clutches had been laid and re-visiting the nests on an  
287 approximate 7-10d cycle to monitor chick development. The metric is the number of  
288 chicks per nest to survive to twenty-one days. After twenty-one days, the chicks become  
289 very active and move throughout the colony precluding accurate accounting of individual  
290 nest production. Since 2003, chicks have also been leg-banded so that individual chicks  
291 can be identified. By resighting these individuals later in the nesting cycle, we are able to  
292 use a second method to estimate nest production. Preliminary analysis of this mark-  
293 resighting technique generally confirms that the twenty-one day survival is an accurate  
294 method to calculate nest production..

295



296 This stoplight uses two metrics for nest production. The number of successful  
297 nesting years out of ten with success being defined as an average nest production of  
298 greater than one chick per nest (c/n) for all nest starts. This metric uses only the  
299 northeastern region of the Bay (Figure 2) as this has been demonstrated to be the region  
300 most impacted by water management practices (Lorenz et al., 2002). Prior to the  
301 establishment of the SDCS, spoonbills nesting in the northeastern region averaged 71%  
302 successful years (Lorenz et al., 2002). Stoplight colors were based on this threshold  
303 (Table 1, Figure 4).

304

305 The second metric of nest production is the five year mean of nest production in  
306 the northeastern region. Lorenz et al., (2002) demonstrated that prior to the SDCS annual  
307 mean spoonbill production in the northeast region was 1.38c/n and that this dropped to  
308 0.67 post-SDCS. Initially we set this as the target for the stoplight metric where annual  
309 production was divided by 1.5 c/n with greater than 67% set as the threshold for a green  
310 rating. However, as can be seen in Figure 5, there are no trends in the data with rapid  
311 changes occurring from one year to the next. This is due to the interannual differences in  
312 hydrologic conditions that affect the ability of spoonbills to capture enough prey to  
313 successfully raise young. Simply put, some years are naturally better than others. Taking  
314 a multi-year running average smoothes this high variability into more interpretable trends  
315 (Figure 5). By examining various time frames from previous data we concluded that by  
316 using a five year running average, no single good or bad year out of the five skewed the  
317 results into the red or green classification. A single good or bad year in either the two,

318 three or four year running averages could bias the mean, thus resulting in an inaccurate  
319 stoplight color.

320

321           There are natural background conditions that can result in nest failure that are  
322 unrelated to CERP or water management practices. Therefore, we need to control for  
323 natural background variation in foraging conditions. We dealt with this problem by using  
324 the northwestern region's success rate as control for natural background conditions.

325 While the northeastern region's production declined post SDCS, the northwestern regions  
326 production remained relatively high (1.24c/n) even though there was still a great deal of  
327 interannual variability. Lorenz and Frezza (2007) concluded that the interannual variation  
328 in productivity of the northwestern colonies reflects the natural variation while the  
329 variation in the northeast is affected by both this background and by water management  
330 practices. Therefore, we propose that the metric used to gage success in the northeastern  
331 region be tied to that of the northwestern, i.e., the metric should be calculated by dividing  
332 annual northeastern production by that of the northwest thereby resulting in a percentage  
333 (Figure 6). The thresholds for stoplight colors are presented in Table 1.

334

335           Although this metric solves the problem of natural interannual variation in nesting  
336 success, it is also dependant on the continued high rates of success of the northwestern  
337 colony. What happens if CERP or other issues begin to negatively affect the success of  
338 the northwestern colonies? This would result in the metric receiving higher scores even  
339 though there was actually a degradation of the bay for spoonbills. Therefore, stoplight  
340 metrics were developed to examine the northwestern regions (explained below in section

341 2.3.5). If all three of the metrics are yellow or red then the metric for northeastern  
342 success should be based on the long term mean production rate of 1.5 c/n for northeastern  
343 Florida Bay (Lorenz et al., 2002, Figure 5).

344

345 *2.3.2. Number of spoonbill nests in Florida Bay.* Spoonbill nest counts for  
346 Florida Bay have been performed intermittently since 1935 (Powell et al., 1989). Over  
347 that period, spoonbills have been recorded nesting on thirty-eight keys throughout the  
348 Bay (Figure 2; Lorenz et al., 2002). Spoonbills typically establish nests in Florida Bay in  
349 November or December of each year, however, nest initiation has started as early as  
350 October and as late as March (Powell et al., 1989, Alvear-Rodriguez, 2001). All known  
351 nesting keys are visited every twenty-one days during the nesting season. Our data show  
352 that prior to the establishment of the SDCS, the peak number of nests was 1258 in 1978  
353 (Figure 1, Lorenz et al., 2002). For this stoplight, annual nest counts are divided by 1258  
354 to get the annual percentage of the historic peak number of nests (Figure 7) and assigned  
355 the stoplight color as per Table 1.

356

357 *2.3.3 Spoonbill nesting location.* This stoplight indicator consists of two metrics:  
358 a return to pre-SDCS nest numbers in the northeastern region and return of spoonbills to  
359 nesting colonies along the southwest coast of the Everglades in the Shark River Slough  
360 and Lostman's Slough estuaries. Powell et al., (1989) reported that in the peak year of  
361 1978 more than half of the 1258 nests were located in the northeast region (688 nests).  
362 Following the completion of the SDCS, this number dropped to approximately 100 nests  
363 from 2000 to 2007. In 2008 there were a total of 47 nests in the region. For restoration

364 to be considered successful, we should expect a return to nesting numbers to pre-SDCS  
365 numbers. This metric is the percentage of 650 nests that occur annually (Figure 8).  
366 Similar to nest success and total nests for Florida Bay, the interannual variation can bias  
367 individual years and a five year mean was used for this metric (Table 1).

368

369           According to Scott (1889), spoonbills “nested in the thousands” along the  
370 southwest coast of the Everglades in the Shark River and Lostman’s slough estuaries.  
371 Restoration of more historic hydrological conditions should promote greater prey  
372 abundance and availability in this region, potentially leading to a return of spoonbill  
373 nesting in large numbers. In recent years, Everglades National Park has performed aerial  
374 wading bird surveys of this area and has documented spoonbill nesting (Pers. Comm,  
375 Sonny Bass, Supervisory Wildlife Biologist, Everglades National Park), however  
376 accurate surveys of spoonbills nest number can not be performed from aircraft because  
377 they tend to nest low in the canopy. Although it is imperative to get a baseline for pre-  
378 CERP nesting in this critical region, no funds have been identified to pay for this effort.  
379 As a result, no stoplight metrics can be established at the time of this publication.

380

381           *2.3.4 Prey Community Structure.* Spoonbills primarily feed on small demersal  
382 fishes found throughout the Everglades system (Allen, 1942, Dumas, 2000). Lorenz et  
383 al., (1997) developed a methodology that uniquely sampled fishes in the dwarf mangrove  
384 foraging grounds that are the preferred feeding locations for spoonbills nesting in Florida  
385 Bay. The sampling design uses a 9m<sup>2</sup> drop trap at fixed locations at known spoonbill

386 feeding sites. Data collection began in 1990 at four sites. Currently, there are 14  
387 sampling sites associated with Florida Bay's nesting spoonbill population (Figure 2)

388

389 Lorenz (1999) documented that these fish respond markedly to changes in water  
390 level and salinity and these factors can be altered by water management practices.  
391 Lorenz and Serafy (2006) performed a fish community analysis of eight years of these  
392 data from six sites. During the eight-year span reported by this study, there were three  
393 consecutive years of unusually high rainfall and freshwater flows to the estuary which  
394 resulted in low salinity similar those believed to have occurred in the region prior to  
395 water management influences. As part of their analysis, Lorenz and Serafy (2006),  
396 placed individual species in one of four salinity categories (freshwater, oligohaline,  
397 mesohaline or polyhaline) based on the Venice System of Estuarine Classification  
398 (Bulger et al., 1993). To accomplish this, the authors used the mean salinity for the thirty  
399 days prior to a given collection (based on the findings of Lorenz, 1999) to identify the  
400 range of salinities in which each species was found. The median score of each species  
401 salinity range was then used to classify the species into one of the four categories.  
402 During the period of low salinity and high fish abundance, Lorenz and Serafy (2006)  
403 found that more than 40% of the total fish community were freshwater affiliates (Figure  
404 3). Furthermore, they demonstrated that it took two to three years of low salinity for the  
405 freshwater populations to respond. Finally, they demonstrated these low salinity  
406 communities were much more productive based on both number and biomass of the  
407 standing stock (Figure 3). The stoplight for prey abundance will use the percentage of  
408 the fish community that was classified by Lorenz and Serafy (2006) as freshwater species

409 as per Table 1. Although the stoplight will be reported on an annual basis, it is  
410 integrative for the previous two years as well, i.e., this stoplight measures conditions on a  
411 three year time scale.

412

413 *2.3.5 Monitoring nesting success in northwestern Florida Bay as a control.* As  
414 stated above, comparing nesting success in the northeastern bay to that of the  
415 northwestern bay accounts for background fluctuations on an interannual basis. For this  
416 metric to work, however, there needs to be a control for any anthropogenically induced  
417 reduction in nesting and productivity in the northwestern bay. We propose three stoplight  
418 metrics to act as a control for the proposed comparison of the two regions. Lorenz et al.,  
419 (2002) indicated that the mean production rate for spoonbill nests in the northeastern  
420 region was 1.24 c/n. Based on this we expect the five year mean production rate to  
421 remain above 1.25 c/n and the control stoplight will remain green so long as this criterion  
422 is met (Figure 9, Table 1). Since the completion of the SDCS, the northwestern region of  
423 Florida Bay has produced a mean of 218 nests annually. Based on this metric, we set the  
424 control metric for nest number at 200 and use a five year running mean of the percentage  
425 of 200 as the stoplight indicator (Figure 9, Table 1). Finally, spoonbills have averaged  
426 success in more than six of every ten years in the northwest region. The percentage of  
427 successful years (mean production of >1.0 c/n) will also be used as a control with any  
428 metric above six of ten years receiving a green stoplight score (Figure 9, Table 1). If all  
429 three of the control metrics are yellow and/or red, than the metric for the northeastern bay  
430 should be re-evaluated based on the historic trends of the northeastern region (Figure 5).

431

432 **3. Longer-Term Science Needs**

433           Population dynamics of spoonbills in the Everglades and methods to monitor their  
434 responses to hydrologic management effectively, are relatively well understood. The  
435 techniques used to survey spoonbills is relatively well worked out, however, there are  
436 components of their basic biology that are unknown. For example, life expectancy and  
437 age at maturity are not known. Furthermore, migratory patterns are not well understood  
438 and need to be assessed to determine if spoonbills nest in multiple locations annually or if  
439 the nesting population in Florida Bay is distinct from other nesting locations around the  
440 state. Also, our knowledge of the dispersal of fledglings from the nesting colony is  
441 extremely limited. A banding program is underway to determine movements within the  
442 state, however, further funding for this effort has not been identified and the program will  
443 be eliminated without identification of funds. Furthermore, a satellite tagging program  
444 would provide a great deal of information on international movements (e.g. Cuba,  
445 Yucatan). This would also allow definitive data on local foraging flights. Currently, we  
446 use inferences (such as flight line counts) to track where birds are feeding.

447

448           Currently there are no efforts to survey wading bird nesting colonies in the  
449 estuaries of the southwestern coast of the Everglades even though this has been  
450 documented as an important nesting area prior to the plume hunting era. A return to  
451 nesting in this area has been identified as an important indicator for the restoration of  
452 flows through Shark River and Lostman's sloughs. Funding for such surveys may be  
453 expensive as they will require the use of a helicopter for access, however, it is imperative  
454 that such funds be identified so as to maximize the use of this versatile indicator species  
455 in the larger restoration plan.

456

457           Of the seventeen existing prey fish sampling sites, three critical sites in  
458 northeastern Florida Bay are not funded through any restoration effort. Secure funding  
459 for these sites needs to be identified to preserve the statistical integrity of this effort.

460

#### 461 **4. Discussion and Conclusions**

462

##### 463 *4.1 Effectiveness of spoonbills as an Indicator of Ecological Restoration*

464           Spoonbills provide information to assess restoration of the Everglades that are  
465 unique from other wading bird indicators and require different methods of assessing their  
466 population trends. Therefore, spoonbills were identified as a separate indicator from the  
467 other wading bird species for two reasons. First, spoonbills nest cryptically within the  
468 canopy of mangroves and are not conspicuous from the air requiring nesting surveys to  
469 be performed on the ground rather than aurally. As a result, different parameters have  
470 been used to monitor spoonbills. Since we have to enter the nesting colonies to monitor  
471 nesting effort, we are able to get more accurate counts of total nests, what region of  
472 Florida Bay the nests were located, and the success of individual nests is documented  
473 through mark and revisitation of the nests.

474

475           In southern Florida, spoonbills show a distinct fidelity to estuarine habitats with  
476 approximately 90% of all nests found within Florida Bay, Tampa Bay and Indian River  
477 Lagoon (although in recent years spoonbills have begun nesting at such inland freshwater  
478 habitats such as the Corkscrew Swamp, Water Conservations Areas and mainland



479 Everglades National Park). In contrast, other wading birds are much more plastic in there  
480 selection of breeding sites with a well documented switch from coastal mangrove habitats  
481 to the Water Conservation Areas in response to water management practices. Given  
482 these differences, spoonbills are an indicator for Florida Bay, the southwest coastal  
483 estuaries and, perhaps Biscayne Bay while other wading birds are indicators for central  
484 Everglades habitats.

485

486         The RECOVER Conceptual Ecological Models identify three major stressors to  
487 wetlands that are affecting the spoonbill nesting activities in Florida Bay: reduced  
488 freshwater flow volume and duration (affecting hydrology and salinity, fish abundance  
489 and availability); invasive exotic species (affecting primary producers and the prey base  
490 fish community); and sea level rise (affecting habitat loss, wetland function and  
491 geomorphology, preliminary and secondary production in the prey base) (Davis et al.,  
492 2005; CERP Monitoring and Assessment Plan; U. S. Army Corps of Engineers, 2004).  
493 Only the first of these stressors will be ameliorated by CERP and, therefore, the spoonbill  
494 assessment tool only addresses issues for water flow, volume and duration.

495

496         Changes in salinity patterns reduces primary production (through stresses caused  
497 by rapid and frequent fluctuations in salinity; Montague and Ley, 1993, Ross et al., 2000,  
498 Frezza and Lorenz, 2003) and alter the prey base fish community to a state of lower  
499 secondary production (Lorenz, 1999, Lorenz and Serafy, 2006). As a result, the overall  
500 abundance of spoonbill prey items is reduced. The spoonbill assessment tool includes a

501 parameter that examines fish community structure which has been shown to have a direct  
502 link to prey fish productivity thereby addressing this issue.

503

504 Changes in the timing and distribution of fresh-water deliveries, result in  
505 increased water levels on the primary foraging grounds of spoonbills nesting in  
506 northeastern Florida Bay (Lorenz, 2000). Studies performed in the mangrove foraging  
507 grounds indicate that the prey base fishes begin concentrating into deeper creeks and  
508 pools when water level on the wetlands drops to a certain depth threshold (Lorenz, 2000).  
509 Spoonbills time nesting with falling water levels on these wetlands such that prey will be  
510 concentrated at the time of egg hatching (Bjork and Powell, 1994). This provides a  
511 highly available and consistent prey resource at a time when the energetic demands of  
512 their rapidly growing young are highest. Out-of-season pulse releases resulting from  
513 upstream water management activities rapidly raise water levels above the concentration  
514 threshold and fish disperse across the surface of the wetland. This eliminates the needed  
515 abundant and easily captured food resources for the spoonbills. Even brief reversal  
516 events (3-5 days) can result in total failure of the spoonbill colonies. CERP and related  
517 projects will alleviate this situation leading to higher nesting success and a return to  
518 higher nest numbers in northeastern Florida Bay. The spoonbill metrics of nesting  
519 success, location and number assess these components of the impacts of water  
520 management practices.

521

522 The performance measure metrics chosen for spoonbills reflect current and  
523 historic ecosystem conditions. The metrics used to evaluate spoonbills have been well

524 documented in the literature and are based on the best understanding of how the Florida  
525 Bay estuary functioned historically, currently and how we expect it to function under  
526 restored conditions. The metrics used provide both spatial and temporal metrics to assess  
527 the state of recovery efforts. We conclude that the spoonbill assessment tool will provide  
528 a powerful and integrative means to evaluate CERP activities.

529

#### 530 *4.2. Communicating the Spoonbill Indicator*

531

532 Roseate spoonbills, being a species that Everglades visitors seek out and  
533 appreciate provide a valuable social as well as natural indicator. They are also well  
534 accepted by managers and policy makers as a species that is important to our  
535 understanding of estuarine systems. This is an important feature for system-wide  
536 integrative indicators and we can capitalize on these points with the spoonbill indicator.

537

538 Making environmental decisions requires both effective communication of  
539 environmental information to decision makers and consideration of what members of the  
540 public value about ecosystems (Schiller et al., 2001). As described above, spoonbills are  
541 good indicators (well-established relationships with environmental parameters under  
542 management control) and the metrics (nest number and location, nesting success, prey  
543 species composition) are remarkably easy to understand and communicate. The first  
544 MAP Annual Assessment Report for spoonbills and their prey summarizes the most  
545 recent advancements for spoonbills (System Status Report, 2006). The concepts of low  
546 nest numbers, nesting in less desirable habitats, declines in nest success and prey

547 abundance are all real concepts, with meaning to managers. Tracking improving or  
548 declining conditions due to restoration activities with these metrics is easily  
549 communicated and understood.

550

551 **Literature Cited**

552

553 Allen, R. P., 1942. The Roseate Spoonbill. Dover Publications Inc., New York.

554

555 Alvear-Rodriguez, E.A., 2001. The use of nesting initiation dates of Roseate Spoonbills  
556 in northeastern Florida Bay as an ecosystem indicator for water management  
557 practices. MS Thesis, Florida Atlantic University, Boca Raton, Florida.

558

559 Bartell, S.M., Lorenz, J.J., Nuttle, W.K., 2004. Ecological models for ENP evaluation of  
560 CERP activities; Roseate Spoonbill Habitat Suitability Index model. Report to  
561 South Florida Research Center, Everglades National Park, Homestead FL.

562

563 Bjork, R. D. and Powell, G.V.N., 1994. Relationships between hydrologic conditions  
564 and quality and quantity of foraging habitat for Roseate Spoonbills and other  
565 wading birds in the C-111 basin. Final report to the South Florida Research  
566 Center, Everglades National Park, Homestead, Florida.

567

568 Bulger A.J., Hayden, B.P., Monaco, M.E., Nelson, D.M. McCormick-Ray, M.G., 1993.  
569 Biologically based estuarine salinity zones derived from multivariate analysis.  
570 Estuaries 16, 311-322.  
571

572 Davis, S.M., Childers, D., Lorenz, J.J., Hopkins, T.E., 2005. A conceptual model of  
573 ecological interactions in the mangrove estuaries of the Florida Everglades.  
574 Wetlands 25, 832-842.  
575

576 DeAngelis, D. L., Loftus, W.F., Texler, J.C., Ulanowicz, R.E., 1997. Modeling fish  
577 dynamics and effects of stress in a hydrologically pulsed ecosystem. J. Aqua.  
578 Ecosys. Stress Rec. 6: 1-13.  
579

580 Doren, R.F. 2006. Indicators for Restoration: South Florida Ecosystem Restoration.  
581 Report to the South Florida Ecosystem Restoration Task Force.  
582

583 Dumas, J., 2000. Roseate Spoonbill (*Ajaia ajaja*). *In*: A. Poole and F. Gill (Eds.) The  
584 birds of North America. The Academy of Natural Science, Philadelphia.  
585

586 Frederick, P., Spalding, M.G., 1994. Factors affecting reproductive success of wading  
587 birds (Ciconiiformes) in the Everglades ecosystem. *In*: S. M. Davis and J. C.  
588 Ogden, (Eds.) Everglades. The Ecosystem and Its Restoration. St. Lucie Press,  
589 Boca Raton.  
590

591 Frezza, P.E., Lorenz, J.J., 2003. Distribution and abundance patterns of submerged  
592 aquatic vegetation in response to changing salinity in the mangrove ecotone of  
593 northeastern Florida Bay. Florida Bay Program and Abstracts from the Joint  
594 Conference on the Science and Restoration of the Greater Everglades and Florida  
595 Bay Ecosystem. University of Florida, Gainesville FL.  
596

597 Gawlik, D. E., 2002. The effects of prey availability on the numerical response of wading  
598 birds. Ecol. Monogr. 72, 329-346  
599

600 Kahl, M.P., 1964. Food ecology of the wood stork (Mycteria americana). Ecol. Monogr.  
601 34, 97-117.  
602

603 Light, S.S., Dineen, J.W., 1994. Water control in the Everglades: a historical perspective.  
604 In Davis S.M. & J.C. Ogden (Eds), Everglades: the Ecosystem and Its  
605 Restoration, St. Lucie press, Delray Beach, FL.  
606

607 Loftus, W. F., Kushlan, J.A., 1987. Freshwater fishes of southern Florida. Bull. Fla.  
608 State Mus. Biol. Sci. 31, 137-344.  
609

610 Lorenz, J.J., 1999. The response of fishes to physicochemical changes in the mangroves  
611 of northeast Florida Bay. Estuaries 22, 500-517.  
612

613 Lorenz, J.J., 2000. Impacts of water management on Roseate Spoonbills and their piscine  
614 prey in the coastal wetlands of Florida Bay. Ph.D. Dissertation, University of  
615 Miami, Coral Gables FL.  
616

617 Lorenz, J.J., Serafy J.E., 2006. Subtropical wetland fish assemblages and changing  
618 Salinity regimes: implications of Everglades Restoration. *Hydrobiologia* 569,  
619 401-422.  
620

621 Lorenz, J.J., Ogden, J.C., Bjork, R.D., Powell, G.V.N., 2002. Nesting patterns of Roseate  
622 Spoonbills in Florida Bay 1935 -1999: implications of landscape scale  
623 anthropogenic impacts. In: Porter, J.W. & K. G. Porter (Eds), *The Everglades,  
624 Florida Bay, and Coral Reefs of the Florida Keys: An Ecosystem Sourcebook*,  
625 CRC Press, Boca Raton, FL.  
626

627 Lovett, G.M., Burns, D.A., Driscoll, C.T., Jenkins, J.C., Mitchell, M.J., Rustad, L.,  
628 Shanley, J.B., Likens, G.E., Haeuber, R., 2007. Who needs environmental  
629 monitoring? *Front. Ecol. Environ.* 5(5), 253-260.  
630

631 McIvor, C.C., J.A. Ley & R.D. Bjork, 1994. Changes in freshwater inflow from the  
632 Everglades to Florida Bay including effects on the biota and biotic processes: a  
633 review. In Davis S.M. & J.C. Ogden (eds), *Everglades: the Ecosystem and Its  
634 Restoration*, St. Lucie Press, Delray Beach, FL: 117-146.  
635

636 Montague, C.L., Ley, J.A., 1993. A possible effect of salinity fluctuation on abundance  
637 of benthic vegetation and associated fauna in northeastern Florida Bay. *Estuaries*  
638 16, 703-717.  
639

640 Powell, G. V. N., Bjork, R.D., Ogden, J.C., Paul, R.T., Powell, A.H., Robertson, W.B.,  
641 1989. Population trends in some Florida Bay wading birds. *Wils. Bull.* 101, 436-  
642 457  
643

644 Ross, M. S., Meeder, J.F., Sah, J.P., Ruiz, P.L., Telesnicki, G., 2000. The Southeast  
645 Saline Everglades revisited: a half-century of coastal vegetation change. *J. Veg.*  
646 *Sci.* 11, 101-112.  
647

648 Scott W.E.D., 1889. A summary of observations on the birds of the gulf coast of Florida.  
649 *Auk* 6, 13-18.  
650

651 Schiller, A., Hunsaker, C.t., Kane, M.A., Wolfe, A.K., Dale, V.H., Suter, G.W., Russell,  
652 C.S., Pion, G., Jensen, M.H., Konar, V.C., 2001. Communicating Ecological  
653 Indicators to Decision Makers and the Public. *Cons. Ecol.* 5(1), 19.  
654

655 U. S. Army Corps of Engineers. 1999. CERP Central and Southern Florida  
656 Comprehensive Review Study. Final Integrated Feasibility Report and  
657 Programmatic Environmental Impact Statement. Jacksonville District, United  
658 States Army Corps of Engineers, Jacksonville, FL.



659

660 U. S. Army Corps of Engineers. 2004. CERP Comprehensive Monitoring and

661 Assessment Plan

662 [http://www.evergladesplan.org/pm/recover/recover\\_map\\_2004.cfm](http://www.evergladesplan.org/pm/recover/recover_map_2004.cfm)

663

664 Williams, B.K., Szaro, R.C., Shapiro, C.D., 2007. Adaptive Management: The U.S.

665 Department of the Interior technical guide. Adaptive Management Working

666 Group, U.S. Department of the Interior, Washington, DC.

667

668

669 Acknowledgements

670

671 We would like to thank Greg May, the Executive Director of the South Florida

672 Ecosystem Restoration Task Force, and Rock Salt, Co-chair of the Science Coordination

673 Group, for their support in making the publication of the special issue of Ecological

674 Indicators possible. We would also like to thank G. Ronnie Best, U.S. Geological

675 Survey, for additional financial support in the publication of this special issue. This

676 description of crocodilians as an ecological indicator for restoration of Greater

677 Everglades ecosystems borrowed both inspiration and words directly from the template

678 for Fish and Macroinvertebrates prepared by Joel Trexler and Bob Doren.

679 Table 1. Decision rule targets and scores for forming performance measure/suitability  
680 relationships for the Roseate Spoonbill indicator communication tool.

681

682 1. Northeastern Nesting Success: number of successful nesting attempts (average of >1 chick  
683 fledged per nest attempt) out of the previous 10 years in northeastern Florida Bay. Target is 7  
684 out of 10 successful years based on the pre-SDCS average (Lorenz et al., 2002)

685 a. 0 – 3 Red

686 b. 3 - 6 Yellow

687 c. 7 - 10 Green

688

689 2 Northeastern Nest Production:

690 A. Five year mean of northeastern Florida Bay nest production expressed as a percentage of  
691 northwestern Florida Bay nest production. This metric will be used if any of the control  
692 metrics for northwestern Florida Bay (number 7 below) are green. In the case of none of the  
693 controls being scored green than 2B will be used.

694 a. 0 - 33 Red

695 b. 33 - 66 Yellow

696 c. > 66 Green

697

698 B. Five year mean of the percentage of mean pre-SDCS nest production. Target is 1.5 chicks  
699 per nest attempt is based on the mean nest production from 1962 to 1982 (Lorenz et al.,  
700 2002). This metric will only be used when all of the northwestern Florida Bay control  
701 metrics (number 7 below) are scored as yellow and/or red. In the case of any of the controls  
702 being scored a green than 2A will be used.

703 a. 0 - 50 Red

704        b   50 - 100     Yellow

705        c. > 100        Green

706

707        3. Nest Number: five year mean of the percentage of pre-SDCS peak nest numbers found

708            throughout Florida Bay. Target is 1250 based on the peak number of nests found in 1978

709            (Powell et al., 1989).

710        a.  0 - 50         Red

711        b.  50 - 100      Yellow

712        c.  > 100        Green

713

714        4. Florida Bay Spoonbill Nesting Location: five year mean of the percentage of pre-SDCS peak

715            nest numbers found in northeastern Florida Bay. Target number is 625 based on the peak

716            number of nests found in 1978 (Powell et al., 1989).

717        a.  0 - 33         Red

718        b.  33 - 66      Yellow

719        c.  > 66         Green

720

721        5. Nesting in Southwestern Everglades Estuaries: No targets or stoplight scores can be set at this

722            time

723

724 6. Prey Community Structure: Annual percentage of prey base fish sampling that are classified as  
725 freshwater species according to Lorenz and Serafy (2007). Target is that 40% of the total  
726 annual catch collected at six sampling sites within the foraging grounds of spoonbills nesting  
727 in northeastern Florida Bay (Figure 2: TR, EC, WJ, JB, SB, and HC) are freshwater species  
728 using data. Note that this metric is integrative of three years.

729 a. 0 - 20 Red

730 b. 20 - 40 Yellow

731 c. > 40 Green

732

733 7. Northwestern Florida Bay Control Metrics:

734 A: Five year mean of the percentage of mean post-SDCS nest production in northwestern  
735 Florida Bay. Target is 1.24 chicks per nest attempt is based on the mean nest production  
736 from 1982-2002 (Lorenz et al., 2002).

737 a. 0 - 50 Red

738 b. 50 - 100 Yellow

739 c. > 100 Green

740

741 B: Five year mean of the percentage of post-SDCS mean nest numbers found in northwestern  
742 Florida Bay. Target number is 200 based on the number of nests from 1982-2002 (Lorenz et  
743 al., 2002).

744 a. 0 - 50 Red

745 b. 50 - 100 Yellow

746 c. > 100 Green

747

- 748 C. Number of successful nesting attempts (average of >1 chick fledged per nest attempt) out of  
749 the previous 10 years in northwestern Florida Bay. Target is 6 out of 10 successful years  
750 based on the post-SDCS average (Lorenz et al., 2002)
- 751 a. 0 – 2            Red
  - 752 b. 3 - 5            Yellow
  - 753 c. 6 - 10           Green
- 754
- 755 8. Cumulative Spoonbill Stoplight Metric: the mean of the 6 (or 7 if nesting location on the  
756 southwest coast of Florida can be calculated from future efforts) non-baseline  
757 stoplights where red is scored 1, yellow is scored 0.5 and red is zero.
- 758 a. 0 - 33            Red
  - 759 b. 33 - 66           Yellow
  - 760 c. > 66            Green
- 761




Figure 1. Annual number of roseate spoonbill nests for all of Florida Bay (Total) and for just the northeastern region of the bay from 1935 to 2008.





Figure 2. Top : Map of southern Florida indicating the major features discussed. Bottom: Map of Florida Bay indicating all the nesting locations for spoonbills since 1935, the primary foraging areas for five regions of Florida Bay and the fish sampling sites used to evaluate the spoonbill's forage base.




Figure 3. Top: Left Axis : Percent of total species collected annually at the three estuarine fish sampling sites (Figure : TR, JB, HC) by each salinity category as defined by Lorenz and Serafy 2006. Right Axis: Mean daily salinity from the three sites for the period of record. Note that years following a high salinity dry season have lower representation of freshwater species and higher representation of mesohaline and polyhaline species. The figure also indicates that it takes 2 to 3 consecutive years of low salinity for the freshwater species to become the dominate fish category. (Copyright: Hydrobiologia). Bottom: Differences in fish biomass between salinity categories as defined by Lorenz and Serafy (2006) using Non-Metric Multidimensional Scaling from eight years of fish collections at 6 sites. Their results show that samples dominated by lower salinity species have significantly higher biomass than those dominated by higher salinity species. (Copyright: Hydrobiologia).






Figure 4. Decadal metric for percent of years nesting was successful. The percentage years out of the previous ten in which spoonbills nesting in northeastern Florida Bay were successful (>1 chick per nest fledged). These data demonstrate the declining number of successful years in spoonbill nesting since 1998. Note that due to data limitations we used the five year average in the figure, however, the ten year mean will be used for the actual stoplight metric.




Figure 5. Five year metric used for nest production in northeastern Florida Bay. Left: Percentage of the target production rate of 1.5 chicks per nest fledged in northeastern Florida Bay since the completion of the South Dade Conveyance System (SDCS). The target is based on pre-SDCS nest production data presented by Lorenz et al (2002). Right: The five year running mean of data presented in the figure on the left. Note that due to data limitations the first 3 data points are four year averages, however, the five year mean will be used for the actual stoplight metric. This metric will only be used if the three control metrics for northwestern Florida Bay (Figure ) are scored yellow and/or red.




Figure 6. Five year metric used for nest production in northeastern Florida Bay. Left: Northeastern Florida Bay nest production (in chicks fledged per nest attempt) as a percentage of northwestern Florida Bay production since the completion of the South Dade Conveyance System. Right: The five year running mean of data presented in the figure on the right. This metric will be used as the stoplight metric for nest productivity unless the three control metrics for northwestern Florida Bay (Figure ) are scored yellow and/or red




Figure 7. Bay wide nest number metric. Left: Number of nests bay-wide as a percentage of a target of 1250 nests. The target was set based on the maximum number of nests in Florida Bay prior to the completion of the South Dade Conveyance System (SDCS) as reported by Powell et al (1989). Right: Five year running mean of the data presented to the right. Note that due to data limitations the earliest data point was a mean of only 3 years, however, the five year mean will be used for the actual stoplight metric.




Figure 8. Nest location metric for northeastern Florida Bay. Left: Number of nests in northeastern Florida Bay as a percentage of a target of 625 nests. The target was set based on the maximum number of nests in northeastern Florida Bay prior to the completion of the SDCS as reported by Powell et al (1989). Right: Five year running mean of the data presented to the right. Note that due to data limitations the earliest data point was a mean of only 3 years, however, the five year mean will be used for the actual stoplight metric.




Figure 9 Control metric for using northwestern Florida Bay production as the standard for calculating the stoplight metric in northeastern Florida Bay (Figure ^). Top Right: Percentage of the target production rate of 1.25 chicks per nest fledged in northwestern Florida Bay since the completion of the SDCS. The target is based on the post-SDCS nest production data presented by Lorenz et al (2002). Top Left: Five year mean of the number of nests in northwestern Florida Bay as a percentage of a target of 200 nests. The target was set based on the average number of nests in northwestern Florida Bay since the completion of the SDCS as reported by Lorenz et al (2002). Bottom: The percentage years out of the previous ten in which spoonbills nesting in northeastern Florida Bay were successful (>1 chick per nest fledged). Note that due to data limitations we used the five year average in the figure, however, the ten year mean will be used for the actual stoplight metric.