

5-15-2014

Vorinostat positively regulates synaptic plasticity genes expression and spine density in HIV infected neurons: role of nicotine in progression of HIV-associated neurocognitive disorder

Venkata Atluri

Herbert Wertheim College of Medicine, Florida International University, valturi@fiu.edu

Sudheesh Pilakka-Kanthikeel

Herbert Wertheim College of Medicine, Florida International University, spilakka@fiu.edu

Thangavel Samikkannu

Herbert Wertheim College of Medicine, Florida International University, Samikkannu.Thangavel@fiu.edu

Vidya Sagar

Herbert Wertheim College of Medicine, Florida International University, vsagar@fiu.edu

Kesava Rao Venkata Kurapati

Herbert Wertheim College of Medicine, Florida International University, krao@fiu.edu

See next page for additional authors

Follow this and additional works at: http://digitalcommons.fiu.edu/com_facpub

 Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Atluri, Venkata; Pilakka-Kanthikeel, Sudheesh; Samikkannu, Thangavel; Sagar, Vidya; Rao Venkata Kurapati, Kesava; Saxena, Shailendra K.; Yndart, Adriana; Raymond, Andrea; Ding, Hong; Hernandez, Oscar; and Nair, Madhavan, "Vorinostat positively regulates synaptic plasticity genes expression and spine density in HIV infected neurons: role of nicotine in progression of HIV-associated neurocognitive disorder" (2014). *HWCOC Faculty Publications*. Paper 65.
http://digitalcommons.fiu.edu/com_facpub/65

Authors

Venkata Atluri, Sudheesh Pilakka-Kanthikeel, Thangavel Samikkannu, Vidya Sagar, Kesava Rao Venkata Kurapati, Shailendra K. Saxena, Adriana Yndart, Andrea Raymond, Hong Ding, Oscar Hernandez, and Madhavan Nair



RESEARCH

Open Access

Vorinostat positively regulates synaptic plasticity genes expression and spine density in HIV infected neurons: role of nicotine in progression of HIV-associated neurocognitive disorder

Venkata Subba Rao Atluri¹, Sudheesh Pilakka-Kanthikeel¹, Thangavel Samikkannu¹, Vidya Sagar¹, Kesava Rao Venkata Kurapati¹, Shailendra K Saxena², Adriana Yndart¹, Andrea Raymond¹, Hong Ding¹, Oscar Hernandez¹ and Madhavan PN Nair^{1*}

Abstract

Background: HIV-associated neurocognitive disorder (HAND) is characterized by development of cognitive, behavioral and motor abnormalities, and occurs in approximately 50% of HIV infected individuals. In the United States, the prevalence of cigarette smoking ranges from 35-70% in HIV-infected individuals compared to 20% in general population. Cognitive impairment in heavy cigarette smokers has been well reported. However, the synergistic effects of nicotine and HIV infection and the underlying mechanisms in the development of HAND are unknown.

Results: In this study, we explored the role of nicotine in the progression of HAND using SK-N-MC, a neuronal cell line. SK-N-MC cells were infected with HIV-1 in the presence or absence of nicotine for 7 days. We observed significant increase in HIV infectivity in SK-N-MC treated with nicotine compared to untreated HIV-infected neuronal cells. HIV and nicotine synergize to significantly dysregulate the expression of synaptic plasticity genes and spine density; with a concomitant increase of HDAC2 levels in SK-N-MC cells. In addition, inhibition of HDAC2 up-regulation with the use of vorinostat resulted in HIV latency breakdown and recovery of synaptic plasticity genes expression and spine density in nicotine/HIV alone and in co-treated SK-N-MC cells. Furthermore, increased eIF2 alpha phosphorylation, which negatively regulates eukaryotic translational process, was observed in HIV alone and in co-treatment with nicotine compared to untreated control and nicotine alone treated SK-N-MC cells.

Conclusions: These results suggest that nicotine and HIV synergize to negatively regulate the synaptic plasticity gene expression and spine density and this may contribute to the increased risk of HAND in HIV infected smokers. Apart from disrupting latency, vorinostat may be a useful therapeutic to inhibit the negative regulatory effects on synaptic plasticity in HIV infected nicotine abusers.

Keywords: Human synaptic plasticity, PCR array, Spine density, HIV, Nicotine, HDAC2, eIF2 α , Vorinostat, SK-N-MC neuronal cells, Neurocognitive disorder

* Correspondence: nairm@fiu.edu

¹Department of Immunology, Institute of NeuroImmune Pharmacology, Herbert Wertheim College of Medicine, Florida International University, 11200 SW 8th Street, Miami, FL -33199, USA

Full list of author information is available at the end of the article

Background

HIV-1 infection, cigarette smoking and drug abuse are global public health concerns. Several studies report that nicotine, the psychoactive drug in tobacco products that induces dependence among smokers, is one of the major immunomodulatory agents [1-5]. Nicotine dependence is the most common form of chemical dependence in the United States. Research suggests that nicotine may be as addictive as heroin, cocaine and alcohol [3,5]. In the United States, the prevalence of cigarette smoking ranges from 35-70% in HIV-infected individuals compared to approximately 20% in general population [6-9].

Infection with HIV is often associated with a wide range of neurological and neuropathological abnormalities, which are collectively known as HIV-associated neurocognitive disorder (HAND). HAND is characterized by development of cognitive, behavioral and motor abnormalities, and occurs in approximately 50% of HIV infected individuals [10,11]. Previous studies indicate that HIV-1 clade B (predominant subtype found in the United States and in the western world) being more neuropathogenic than clade C (predominant in sub-Saharan Africa and Asia) [12-14]. Experimental evidence and clinical observations suggest a correlation between smoking and increased risk for neurological disorders [15]. A great deal of research has focused on the pathophysiological and clinical effects of nicotine. In human studies, nicotine and other agonists of the nicotinic acetylcholine receptor (nAChRs) have been reported to improve cognitive performance in healthy adults and in different psychiatric populations [16,17]. In studies with rodents, the results have been mixed with observations of improvement, impairment, or no effect on various measures of learning and memory following acute or chronic systemic treatment [16]. However, the effect of nicotine on HIV-1 pathogenesis has received limited research attention. Few studies have reported that nicotine exacerbates HIV-1 replication in *in vitro*-infected alveolar macrophages and primary human microglia [18,19]. Further, limited examination of the neuropathogenic effects of nicotine on HIV-1 infection of CNS cells has shown that galantamine and nicotine exert synergistic effect on the inhibition of microglial cell activation induced by HIV-1 gp120 [20]. A recent study reported that exposure to tobacco smoke extract increases the blood-brain barrier permeability and plays a critical role in pathophysiology of ischemia [21].

Although few studies have reported deficits in learning, memory, and global cognitive performance among cigarette smokers living with HIV [22,23], the underlying mechanisms associated with the use of nicotine and the development of HAND in these patients are unknown. New memory formation involves gene expression-dependent changes in synaptic structure and plasticity in

the hippocampus [24]. Recently, we have reported dysregulated synaptic plasticity genes expression and decreased spine density in HIV infected CNS cells [13]. Although epigenetic modifications are known to regulate gene expression during HIV infection [25], molecular and biochemical evidence on epigenetic mechanisms in CNS cells in response to HIV and nicotine are lacking. Acetylation and deacetylation of histone proteins associated with chromatin plays a pivotal role in the epigenetic regulation of transcription and other functions in cells, including neurons.

HDACs often function as a component of the transcriptional repressor complex to silence gene expression and induce chromatin compaction through histone protein deacetylation. HDAC2 functions in modulating synaptic plasticity and long-lasting changes of neural circuits, which in turn negatively regulates learning and memory [26]. HDAC inhibitors are emerging as an exciting new class of potential therapeutic agents for the treatment of solid and hematological malignancies. Vorinostat also known as suberanilohydroxamic acid (SAHA) was the first histone deacetylase inhibitor approved by the U.S. Food and Drug Administration for the treatment of Cutaneous T-Cell Lymphoma. Vorinostat binds to the catalytic domain of the histone deacetylases (HDACs). This allows the hydroxamic moiety to chelate zinc ion located in the catalytic pockets of HDAC, thereby inhibiting deacetylation and leading to an accumulation of both hyperacetylated histones and transcription factors. Vorinostat was also shown to cross the blood-brain barrier in a mouse model of Huntington's disease [27]. Therefore, use of vorinostat as a therapeutic agent in the CNS related diseases may be useful. In the down-stream, translation of eukaryotic mRNAs is regulated primarily at the level of initiation. Two main mechanisms by which translation is controlled are the formation of the ternary complex via eIF2 and the recruitment of the ribosome to the mRNA via 4E-BPs. Phosphorylation of an α subunit of eIF2 (at Ser51) converts the protein from a substrate to a competitive inhibitor of the GDP/GTP-exchange reaction by decreasing the rate of dissociation of eIF2 from eIF2B [28,29]. This causes a decrease in general translational initiation [29,30]. We hypothesize that by inducing the transcriptional repression, HDAC2 plays a major role in the neuropathogenesis caused by HIV and nicotine (Figure 1). In the down-stream, we hypothesize that translational process in HIV infected cells will be affected by eIF2 α phosphorylation (Figure 2).

In the current study, we have analyzed the effect of nicotine on the differential expression of synaptic plasticity genes, spine density in an *in vitro* HIV infection model using neuronal cells. Along with the dysregulated synaptic plasticity genes and altered spine density, we have found increased levels of HDAC2 expression indicating





Figure 1 (See legend on next page.)

(See figure on previous page.)

Figure 1 A hypothetical model for the role of HDAC2 in transcriptional repression of synaptic plasticity genes in neuronal cells infected with HIV and/or in the presence of nicotine. During the gene transcription, the DNA to be transcribed is associated with histone proteins (blue) that are modified by the addition of acetyl groups (green). This modification results in a relaxed chromatin configuration that allows the transcriptional machinery access to the DNA. Up-regulation of HDAC2 during HIV infection and/or nicotine treatment leads to deletion of acetyl groups from histone proteins, resulting in a condensed chromatin that limits the binding of the transcriptional machinery, thereby decreasing gene transcription. Thus, inhibition of HDAC2 by using vorinostat may block these enzymatic processes and return the chromatin to a relaxed state, allowing gene transcription.



the transcriptional repression of synaptic plasticity genes in these cells. We also found increased eIF2α subunit phosphorylation (at Ser51) which may associate with the translational repression of synaptic plasticity genes in HIV alone infected SK-N-MC cells and in the presence of nicotine. These negative regulatory effects of HIV infection and nicotine on the expression of synaptic plasticity genes and spine density were reversed by the HDAC inhibitor vorinostat. This study suggests that nicotine synergistically acts with HIV to down-regulate synaptic plasticity genes and alter spine density, potentially via upregulation of HDAC2. The use of vorinostat may help to reverse these negative regulatory effects (induced by HDAC2) on synaptic plasticity genes expression and spine density.


Results

Increased HIV-infectivity rate of neuronal cells in the presence of nicotine

We observed significant increase in HIV-infection in nicotine-treated HIV-infected SK-N-MC compared to the HIV- infected alone cells by using the LTR real-time PCR. We also observed significantly increased HIV infection (latency disruption) in HIV infected neuronal cells in the presence of vorinostat by both LTR real-time PCR (Figure 3) and p24 ELISA (data not shown). We did not see significant levels of vorinostat induced latency disruption in HIV infected neuronal cells in the presence of nicotine and vorinostat.

Differential expression of human synaptic plasticity genes in SK-N-MC cells infected with HIV and/or in the presence of nicotine

Out of 84 human synaptic plasticity genes expression analyzed by human synaptic plasticity PCR array, 23 synaptic plasticity genes were significantly down-regulated whereas 16 synaptic plasticity genes were significantly up-regulated in nicotine treated SK-N-MC cells. In the HIV infected SK-N-MC cells, while 28 synaptic plasticity genes were down-regulated, 8 synaptic plasticity genes were significantly up-regulated. However, in the presence of nicotine, in HIV infected SK-N-MC cells, a total of 47 genes were significantly down-regulated (3–22 fold) and 4 genes were significantly up-regulated (Table 1). Out of the total 84 genes, 21 genes were commonly down regulated in all the three groups (nicotine treated, HIV



infected and HIV along with nicotine). Figure 4 (Venn diagram) shows the number of down-regulated synaptic plasticity genes in SK-N-MC neuronal cells infected with HIV and/or nicotine combination. These results indicate that in the presence of nicotine, dysregulation of synaptic plasticity genes will be further increased in HIV infected neuronal cells.

Vorinostat regulates HDAC2 expression in neuronal cells infected with HIV and/or treated with nicotine

Using qPCR (Figure 5) and western blot analysis (Figure 6), we observed significantly higher HDAC2 levels in nicotine treated cells, HIV infected cells and HIV infected cells treated with nicotine compared to control cells, indicating possible epigenetic changes which are responsible for the transcriptional repression of synaptic plasticity genes. But we did not see additive/synergistic effect on HDAC2 expression in HIV infected cells in the presence of nicotine. Vorinostat is a potent inhibitor of Classes I and II histone deacetylases. To inhibit the up-regulation of HDAC2 expression in SK-N-MC cells infected with HIV and/or treated with nicotine, these test groups were treated with vorinostat on the 5th day of infection and/or nicotine treatment. Using vorinostat (at 1 μM conc.), we were able to inhibit the HDAC2 expression levels in the nicotine treated cells and HIV infected neuronal cells with/without

nicotine to the expression levels approximately equal to the control cells (Figures 5 and 6).

Vorinostat positively regulates synaptic plasticity genes expression in HIV infected and/or nicotine treated neuronal cells

In vorinostat alone treated SK-N-MC cells, total 19 synaptic plasticity genes were significantly up-regulated while only 4 genes were significantly down-regulated. 24 synaptic plasticity genes out of 28 genes, which were down-regulated with HIV infection, were recovered and 16 more genes were significantly up-regulated in the presence of vorinostat. Further, in the case of HIV infected cells treated with nicotine, out of 47 down-regulated genes, expression of 40 genes were recovered upon vorinostat treatment and 15 more genes were up-regulated. Surprisingly, four genes down-regulated (CREM, NFKB1, RHEB, TIMP1) in the presence of vorinostat alone were consistently down-regulated in all the groups containing vorinostat. Excluding these four genes, vorinostat failed to recover the expression of EGR4, HOMER1 and TNF in HIV infected cells in the presence of nicotine (Table 1 and Figure 7). These results indicate that vorinostat can inhibit the negative regulatory effects on synaptic plasticity genes expression (mediated by increased HDAC2 expression) in HIV infected cells in the presence of nicotine.

Synergistic decrease of spine density in HIV infected neuronal cells in the presence of Nicotine

Using the established protocol, we have observed for change in spine density in HIV infected SK-N-MC cells and/or nicotine/vorinostat combination in comparison to the control cells [13]. In nicotine alone treated cells, spine density was significantly less than the control cells. In HIV only infected cells, spine density was significantly less than the nicotine treated cells ($p < 0.02$). Interestingly, HIV infected cells in the presence of nicotine exhibit decreased spine density compared to nicotine/HIV alone infected cells. Spine density was recovered when these nicotine/HIV/HIV + nicotine infected cells were treated with the vorinostat (Figure 8).

eIF2 α phosphorylation in SK-N-MC cells infected with HIV and/or treated with nicotine

Phosphorylation of the α -subunit of eukaryotic initiation factor 2 (eIF2 α) inhibits the initiation of translational process. To investigate whether the translational process is affected or not in the HIV infected and/or nicotine treated neuronal cells, eIF2 α phosphorylation (at Ser51) was analyzed. In the SK-N-MC cells, significant eIF2 α phosphorylation (at Ser51) was observed in HIV infected cells alone and in combined treatment with nicotine. We did not observe significant eIF2 α phosphorylation in

Table 1 Human synaptic plasticity genes expression in HIV infected neuronal cells and in combination with nicotine and/or vorinostat (fold change)

Genes	Vorinostat	HIV	HIV + Vorinostat	Nicotine	Nicotine + Vorinostat	HIV + Nicotine	HIV + Nicotine + Vorinostat
<i>ADAM10</i> ^{5,6,9}	1.47	1.89	1.24	4.12	1.45	-1.03	1.11
<i>ADCY1</i> ³	1.12	-3.39	1.12	-3.15	2.26	-3.99	1.17
<i>ADCY8</i> ³	1.12	-3.39	1.12	-3.15	2.26	-3.99	-1.96
<i>AKT1</i> ⁷	3.10	2.96	3.84	3.42	4.59	-1.32	4.23
<i>ARC</i> ^{1,9}	-1.09	1.30	1.54	1.07	2.79	-4.81	1.55
<i>BDNF</i> ^{1,3}	1.12	-3.39	1.12	-3.15	2.26	-3.99	-1.96
<i>CAMK2A</i> ³	1.12	-3.39	1.12	-3.15	2.26	-3.99	-1.96
<i>CAMK2G</i> ^{3,7}	1.48	1.11	1.46	1.16	1.62	-3.38	1.33
<i>CDH2</i> ^{3,5}	1.12	-3.06	1.12	-3.15	2.26	-3.99	-1.37
<i>CEBPB</i> ¹	1.37	2.50	1.64	1.98	2.07	-3.07	2.04
<i>CEBPD</i> ¹	1.39	1.89	1.96	1.35	3.03	-3.07	1.96
<i>CNR1</i> ³	1.12	-3.39	1.12	-3.15	2.26	-3.99	-1.96
<i>CREM</i> ¹	-35.90	-9.78	-15.78	-8.26	-16.02	-25.41	-16.77
<i>EGR1</i> ¹	2.37	1.13	1.34	-5.68	1.15	-9.24	1.20
<i>EGR2</i> ¹	6.89	1.94	4.29	9.22	3.34	1.43	2.43
<i>EGR3</i> ¹	1.86	1.01	3.89	-1.76	2.94	-3.99	2.43
<i>EGR4</i> ¹	-1.95	-6.69	-1.96	-3.20	1.03	-8.74	-4.28
<i>EPHB2</i> ⁸	1.71	-2.97	1.74	-3.15	2.36	-3.99	1.43
<i>FOS</i> ¹	1.15	-1.24	1.07	1.07	1.43	-3.83	-1.12
<i>GABRA5</i> ^{3,8}	1.91	-3.39	1.12	-1.62	2.26	-3.99	-1.96
<i>GRIA1</i> ^{3,4,8,9}	1.12	-3.39	1.12	-3.15	2.26	-3.99	-1.96
<i>GRIA2</i> ^{3,4,8}	4.04	2.38	2.55	2.09	4.40	1.10	3.49
<i>GRIA4</i> ^{4,8,9}	12.85	9.24	4.35	2.39	5.73	-1.50	3.12
<i>GRIN1</i> ^{3,7,8,9}	31.65	5.89	28.25	4.70	39.89	-1.02	35.80
<i>GRIN2A</i> ^{3,5,7,8,9}	2.26	-3.06	1.12	-2.14	2.26	-3.99	-1.07
<i>GRIN2B</i> ^{3,5,7,8,9}	2.26	-3.39	1.12	-3.15	2.26	-3.99	-1.07
<i>GRIN2C</i> ^{3,7,8,9}	2.26	-1.73	1.80	-3.15	2.26	-3.99	-1.07
<i>GRIN2D</i> ^{3,7,8}	4.25	1.42	2.17	4.97	4.25	-1.34	2.80
<i>GRIP1</i> ⁴	3.07	-1.09	2.36	2.72	3.07	-1.90	1.96
<i>GRM1</i> ^{4,8,9}	1.12	-3.39	1.12	-3.15	2.26	-3.99	-1.96
<i>GRM2</i> ^{4,8}	1.85	1.93	1.12	3.79	2.26	-1.47	1.01
<i>GRM3</i> ^{8,9}	1.12	-3.39	1.12	-2.11	2.26	-3.99	-1.96
<i>GRM4</i> ⁸	2.26	-3.06	1.12	1.34	2.26	-3.99	-1.21
<i>GRM5</i> ⁸	1.12	3.38	1.12	-1.15	2.26	-2.75	-1.96
<i>GRM7</i> ⁸	1.12	-3.06	1.12	2.45	2.26	-2.33	-1.10
<i>HOMER1</i> ^{1,9}	-1.64	-1.13	-2.95	1.30	-2.68	-1.93	-3.13
<i>IGF1</i> ⁴	1.96	9.13	1.32	11.75	1.19	3.37	1.17
<i>INHBA</i> ²	4.15	-2.11	2.03	-1.43	2.22	-5.89	2.25
<i>JUN</i> ¹	8.84	-1.39	6.73	-1.65	8.04	-11.61	6.33
<i>JUNB</i> ¹	3.93	2.73	4.06	1.42	10.84	-3.97	3.79
<i>KIF17</i> ¹⁰	2.30	-2.99	3.20	-1.89	4.19	-9.30	3.34
<i>MMP9</i> ^{1,3,6}	3.06	-2.68	2.00	-2.94	2.99	-10.32	1.62
<i>NCAM1</i> ⁵	1.17	-4.59	1.41	-6.80	1.40	-39.88	1.25

Table 1 Human synaptic plasticity genes expression in HIV infected neuronal cells and in combination with nicotine and/or vorinostat (fold change) (Continued)

<i>NFKB1</i> ¹	-20.48	-6.41	-35.26	-5.45	-26.76	-25.59	-48.77
<i>NFKB1B</i> ¹	9.61	12.91	13.09	14.56	14.20	13.54	13.66
<i>NGF</i> ^{1,4}	5.79	1.45	4.92	1.30	4.72	-3.01	6.55
<i>NGFR</i> ⁴	1.30	2.50	1.13	2.65	1.37	-2.18	1.04
<i>NOS1</i> ⁴	1.12	-3.39	1.12	1.22	2.26	-3.99	-1.96
<i>NPTX2</i> ¹	1.12	2.22	1.12	1.28	2.26	-3.29	3.19
<i>NR4A1</i> ¹	5.11	1.47	6.73	2.54	4.82	-1.79	4.67
<i>NTF3</i> ¹	1.12	-3.06	1.12	1.92	2.26	-1.53	-1.96
<i>NTF4</i> ³	1.98	1.14	4.82	-1.30	5.85	-6.14	4.63
<i>NTRK2</i> ³	1.12	-3.40	1.12	-3.15	2.26	-3.99	-1.96
<i>PCDH8</i> ^{1,5}	2.37	2.53	1.29	3.37	2.04	-1.68	1.35
<i>PICK1</i> ^{4,9}	1.58	1.54	2.64	1.18	2.65	-4.21	2.30
<i>PIM1</i> ¹	11.19	4.86	25.46	10.59	33.08	2.55	32.95
<i>PLAT</i> ^{1,4,6}	-1.03	-1.82	-1.03	-2.34	1.97	-4.59	-2.25
<i>PLCG1</i> ³	2.19	1.23	2.35	1.39	3.78	-3.62	2.91
<i>PPP1CA</i> ^{3,4,7}	-2.04	1.54	1.37	-1.69	1.16	-13.25	1.48
<i>PPP1R14A</i> ⁴	1.12	-3.06	1.12	-3.15	2.26	-3.99	-1.96
<i>PPP2CA</i> ⁴	6.52	10.06	6.87	13.31	7.20	3.28	6.64
<i>PRKCA</i> ^{3,4}	5.40	2.80	3.29	7.23	4.43	2.24	2.93
<i>PRKCG</i> ³	2.30	-5.07	2.16	-4.84	3.22	-6.62	2.12
<i>PRKG1</i> ⁴	1.12	-3.06	1.12	-3.15	2.26	-3.99	-1.47
<i>RAB3A</i> ³	86.46	56.49	106.15	75.27	91.65	15.81	103.39
<i>RELA</i> ¹	1.30	2.00	1.34	3.26	1.45	-1.12	1.25
<i>RELN</i> ^{5,6}	1.12	-3.06	1.12	-3.15	2.26	-3.99	-1.96
<i>RGS2</i> ¹	5.29	-1.40	2.62	-1.12	4.46	-2.90	2.38
<i>RHEB</i> ¹	-49.73	-32.90	-57.68	-35.16	-46.92	-84.30	-69.94
<i>SIRT1</i> ¹⁰	2.29	2.58	1.91	4.70	1.89	1.52	1.28
<i>SRF</i> ¹	2.70	2.83	1.82	3.19	2.33	-1.44	1.60
<i>TIMP1</i> ⁶	-18.20	-38.00	-20.11	-24.02	-15.69	-79.20	-17.73
<i>TNF</i> ^{1,5}	-2.65	-5.47	-2.28	-2.63	-1.42	-5.97	-9.30

Out of 84 genes analyzed, only genes significantly ($\pm \geq 3$ fold) dysregulated were shown in this table.

¹Immediate Early Response Genes; ²Late Response Genes; ³Long Term Potentiation; ⁴Long Term Depression; ⁵Cell Adhesion; ⁶Extracellular Matrix & Proteolytic Processing; ⁷CREB Cofactors; ⁸Neuronal Receptors; ⁹Postsynaptic Density; ¹⁰Others.



If the synaptic plasticity genes expression change is $\pm \geq 3$ fold, the fold change was represented in bold letters.

nicotine alone treated SK-N-MC cells compared to the control cells (Figure 9).

Discussion

Nicotine is the single tobacco constituent demonstrated to have the potential to stimulate the production of HIV-1, as has been shown using in vitro-infected alveolar macrophages [13]. A growing body of evidence suggests chronic cigarette smoking alone is associated with abnormalities in brain morphology and neurophysiology [31,32]. Reports on effect of nicotine on HIV associated neurocognitive disorders are scanty. We have used SK-N-MC cells to study the HIV infectivity and for the

analysis of synaptic plasticity genes expression and spine density changes as these cells have been reported to be susceptible for the productive HIV-1 infection [33]. In this study, with the use of neuronal cells, we have observed that the presence of nicotine have significantly increased HIV infection which are in agreement with the previous reports observed in other types of cells [18,19,34]. Rock et al., reported that TGF- β 1 may be involved in the effects of nicotine on enhanced HIV-1 expression in microglial cells [19]. Out of 84 synaptic plasticity genes analyzed, in this study, we have observed down-regulation of 28 synaptic plasticity genes (3–38 folds) and up-regulation (3–56 folds) of 8 genes in HIV




infected cells. Our results indicate that HIV infection alters the expression of synaptic plasticity genes in neuronal cells which may contribute to the disrupted synaptic plasticity.

Nicotine treated SK-N-MC cells exhibit down-regulation in 23 synaptic plasticity genes (3–35 fold) and up-regulation in 16 genes (3–75 folds). Among these genes, 20 were commonly down-regulated in both nicotine-treated (alone) and HIV infected cells (alone). Recent studies have shown that nicotinic mechanisms influence forms of synaptic plasticity that are thought to underlie learning and memory [35,36]. Although nicotine is known to have neuroprotective effects, reports indicate that nicotine do not provide any evidence of the amelioration of the learning deficit (and slightly worsened performance during acquisition) observed in the behavioral HIV-1 transgenic (HIV-1Tg) rat model [37]. In few studies with nicotine exposure, improved attention, learning and memory processes were reported in both experimental animals and humans [16,17,38]. But the positive effect of nicotine on memory and attention is still a matter of debate in case of chronic nicotine exposure. In this study, we observed both positive and negative regulatory effects of nicotine on synaptic plasticity genes expression in neuronal cells after 7 days of chronic nicotine exposure.

This is the first report to study the impact of nicotine on synaptic plasticity genes expression in HIV infected neuronal cells. HIV infected SK-N-MC cells in the presence of nicotine display a significant reduction in total 47 synaptic plasticity genes expression (3–84 folds) and significant up-regulation in only 4 genes expression (IGF1, NFKB1B, PPP2CA, RAB3A) (3–15 folds). All the genes down-regulated in nicotine/HIV infection alone are also down-regulated (few genes synergistically) in HIV infected cells co-treated with nicotine. The fold down-regulation ranges from 3–84 folds indicating increased neuropathogenicity in HIV infected cells treated with nicotine. Activation of immediate early transcription factors [immediate early genes (IEGs)] leads to the transcriptional activation of synaptic plasticity genes that presumably play an important role in the structural and functional neuronal changes involved in mammalian LTM [39,40]. Out of 30 immediate early response genes we have analyzed, 18 genes were significantly down-regulated in HIV infected SK-N-MC neuronal cells in the presence of nicotine, indicating transcriptional repression of other synaptic plasticity genes, therefore decreased LTM. Synaptic plasticity gene-gene interactions/relationships were built using the GNCPro Gene Network Central for genes significantly down-regulated in SK-N-MC neuronal cells Infected with HIV (Figure 10a); Nicotine treated (Figure 10b) and HIV infected cells in the presence of nicotine (Figure 10c).


Transcription factor NF- κ B (NF-kB) regulates spatial memory formation, synaptic transmission, and



plasticity. Inhibition of NF-κB leads to decrease in forskolin-induced CREB phosphorylation [41]. In this study, along with down-regulation of NF-κB1, we have also observed up-regulation of NFKBIB that belongs to the NF-kappa-B inhibitor family, which inhibit NF-κB1 by complexing with, and trapping it in the cytoplasm

that may further disrupt the synaptic plasticity in HIV infected neurons in the presence of nicotine.

NCAM1 is another highly down-regulated synaptic plasticity gene in the HIV infected neuronal cells in the presence of nicotine. NCAM1 is a multifunction trans-membrane protein expressed in both neurons and glial






Figure 8 Synergistic decrease of spine density in HIV infected neuronal cells in the presence of nicotine. SK-N-MC cells were infected with HIV and/or treated with nicotine for 7 days. On the 5th day of HIV infection and/or nicotine treatment, vorinostat was added to the culture. **a)** Control SK-N-MC cells: High number of long spines on the dendritic length, high dendrite diameter, total dendrite area and spine area. **b)** Vorinostat treated Cells: Cells with the higher number of spines on a dendrite length similar to the control cells. **c)** Nicotine treated cells: Loss of spine on a dendrite length when compared to the control cells. **d)** Nicotine along with vorinostat: Restoration of spines on the dendrite length. **e)** HIV infected cells: Loss of spines on the dendrite length, decreased dendrite diameter, dendrite and spine area was observed than the control neuronal cells. **f)** HIV infected cells in the presence of vorinostat: Recovery of spines on the dendrite length was observed. **g)** HIV infected cells in the presence of nicotine: Synergistic loss of number of spine on a dendrite length was observed when compare to the control or nicotine treated or HIV infected cells. **h)** HIV infected cells co-treated with nicotine in the presence of vorinostat: In the presence of vorinostat, spine density was restored when compare to the HIV + nicotine treated cells. (*, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.001$; ****, $p \leq 0.0001$; NS-Not Significant).

cells with cell recognition properties involved in cellular migration, synaptic plasticity [42,43] and central nervous system development [42]. In this study, expression of NACAM1 was recovered with the use of vorinostat treatment.

Two other highly down-regulated synaptic plasticity genes in HIV infected neuronal cells and in combination with nicotine are the RHEB and TIMP1. While RHEB plays an important role in long term activity-dependent neuronal responses [44], TIMP1 is a candidate plasticity gene induced by seizures and by stimuli leading to long

term potentiation. Down-regulation of these two genes may lead to the altered learning and memory.

We have also analyzed the dendritic spine density changes in HIV infected SK-N-MC cells co-treated with nicotine. We have got supporting results for the gene expression studies in SK-N-MC showing the significantly reduced spine density in the HIV infected cells co-treated with nicotine than the control, nicotine/HIV infected cells. We have also observed significantly decreased spine density in nicotine/HIV infected neuronal cells than the control cells indicating neurotoxicity in




Figure 9 Pattern of eIF2 α phosphorylation (Ser51) in SK-N-MC neuronal cells infected with HIV alone and together with nicotine. SK-N-MC cells were treated with HIV/nicotine alone and in combination. After 7 days of infection and/or nicotine treatment, eIF2 α phosphorylation (at Ser51) was analyzed by western blot assay. We have observed significant eIF2 α phosphorylation in HIV and HIV + nicotine treated groups when compared with the control cells. (*, $p \leq 0.05$; NS-Not Significant).

nicotine/HIV alone infected cells as well. Reduced cortical synaptic density has been reported in mild to moderate HAND patient's autopsy samples [45]. Our study shows that nicotine severely affects spine density during HIV infection. This report provides evidence of increased neuropathogenicity in HIV-infected neuronal cells exposed to nicotine and shows the induced synergistic decrease in synaptic plasticity genes expression and spine density than HIV/Nicotine alone infected cells. Among different nicotinic mechanisms influencing synaptic plasticity, role of overexpressed $\alpha 7$ -nAChRs, which play a role in neurological disorders [46] was studied with nicotine/HIV. Chronic exposure to nicotine was reported to cause up-regulation of $\alpha 7$ -nAChRs [47] and up-regulation of $\alpha 7$ -nAChR has been reported as a

contributor to the neurotoxicity associated with HIV infection as well [48]. Another important mechanism is the accumulation of excess glutamate in the extracellular space as a consequence of CNS trauma, neurodegenerative diseases, infection, or deregulation of glutamate clearance results in excitotoxicity. Most acute and chronic neuronal diseases, including HAND, have implicated this type of bystander pathology of excitotoxicity [49]. The presence of excess glutamate in the synaptic clefts activates glutamate gated ion channels and results in high levels of ion influx into neuronal cells allowing the over activation of downstream calcium ion-dependent effectors and signaling pathways, culminating in neuronal damage. Nicotine also reported to enhance the glutamate release [50,51]. Therefore, it may be possible that HIV in combination of




Figure 10 Gene-Gene interaction network for human synaptic plasticity genes. Human synaptic plasticity gene-gene interactions/relationships network was built using the GNCPro Gene Network Central for genes significantly down-regulated in SK-N-MC neuronal cells **a)** Infected with HIV; **b)** Nicotine treated; **c)** HIV infected cells in the presence of nicotine. The interactions are determined and color-coded by down-regulation, up-regulation, regulation, co-expression, chemical modification, physical interaction, predicted protein interaction, predicted TFactor regulation and other.

nicotine may increase the excitotoxicity mediated by glutamate release.

Further, we have investigated the mechanisms attributed for the transcriptional and translational repression of synaptic plasticity genes in HIV infected cells and in combination with nicotine. We analyzed one of the important mechanisms, the role of HDAC2 in the transcriptional repression of synaptic plasticity genes. Histone acetylation is an epigenetic mark in the chromatin that favors gene expression and is modulated by the balance between histone acetyltransferase (HAT) and histone deacetylase (HDAC) enzyme activities, which add and remove, respectively, acetyl groups from histones. With the use of SKNMC, we showed significant up-regulation of HDAC2 in all the three study groups (1. HIV infected; 2. Nicotine treated; 3. HIV infected and co-treated with nicotine) indicating the possible role of HDAC2 in the transcriptional repression of synaptic plasticity genes in neurons. The evidence for HDAC2 in the negative regulation of memory and synaptic plasticity has been well documented [26,52] and the HDAC inhibitors especially class-I were reported as cognitive enhancers [53,54]. Loss of HDAC2 has been reported to improve the associative working memory and accelerated extinction learning [52]. Recently, Hanson et al., reported enhanced excitatory and reduced inhibitory synaptic transmission in HDAC2 knockdown single pyramidal neurons, whereas reduced excitatory and enhanced inhibitory synaptic transmission in HDAC2 overexpressed neurons [55]. Earlier, from our lab, we have reported the up-regulation of HDAC2 in human neurons in response to the HIV-1 Tat protein induced neurotoxicity [56]. Janus kinase (JAK)/signal transducer and activator of transcription (STAT) pathway is known to regulate the transcription of many genes [57-59]. Recently, the role of JAK/STAT pathway in synaptic plasticity has been reported [60].

Further we have used vorinostat, a potent, reversible HDAC inhibitor (both Class I and Class II) to inhibit HDAC2 expression in SK-N-MC cells infected with HIV and/or treated with nicotine. Interestingly, we observed increased HIV infection (latency disruption) in neuronal cells when treated with vorinostat at 1 μ M concentration (no significant change in HIV infection in combined treatment with nicotine and vorinostat). Vorinostat reported to disrupt the HIV latency in resting CD4⁺ cells in vitro [61,62] and in vivo as well [63]. Our results suggest that vorinostat can disrupt the HIV latency in HIV infected CNS cells as well. Therefore, reactivation of latent HIV in the neuronal cells (by using vorinostat) along with antiretrovirals may provide a precise approach to treat and eradicate latent HIV infection in the brain. Vorinostat could also be able to inhibit the HDAC2 up-regulation in neuronal cells infected with HIV and/or treated with nicotine. This results in a more relaxed

chromatin configuration that allows the transcription of synaptic plasticity genes. Figure 1 shows the schematic representation of role of vorinostat in recovery of synaptic plasticity genes expression in neuronal cells infected with HIV and/or treated with nicotine. We have found that in all the groups (Nicotine, HIV infected, HIV + Nicotine), synaptic plasticity genes expression and spine density were recovered and many synaptic plasticity genes were further up-regulated when treated with vorinostat compared to the control cells. In vorinostat alone treated cells, 19 synaptic plasticity genes were significantly up-regulated and 4 genes (CREM, NFKB1, RHEB, TIMP1) were significantly down-regulated than control cells. Effect of down-regulation of these 4 genes in synaptic plasticity needs to be further evaluated. These results indicate that vorinostat positively regulate synaptic plasticity genes expression and spine density in HIV infected neurons and may be useful as a therapeutic agent in the treatment of HAND. Zolinda (Merck, Whitehouse Station, NJ), a prescription drug contains vorinostat is currently being used for the treatment of cutaneous T-cell lymphoma [64], multiple myeloma [65], mesothelioma [66]. Therefore, further studies to use vorinostat as a therapeutic agent against HAND and in nicotine users need to be studied.

In the down-stream, we have also analyzed one of the important mechanisms i.e. eukaryotic initiation factor 2 α phosphorylation (at Ser51), which is responsible for the translational repression of synaptic plasticity genes. Due to partial phosphorylation of eIF2 α , general translation is reduced, and ATF4 mRNA translation is augmented. As a consequence, expression of synaptic plasticity and memory-related genes is depressed (Figure 2). Decreased eIF2 α phosphorylation reduces ATF4 mRNA translation and enhances general mRNA translation, thus facilitating the induction of synaptic plasticity genes expression, which leads to L-LTP and long-term memory (LTM) consolidation [67]. In our study, we have found increased eIF2 α phosphorylation in two-study group of neurons (1. HIV infected; and 2. HIV infected and co-treated with nicotine). It is indicating that translation of synaptic plasticity genes will also be effected in the neuronal cells infected with HIV and in combined treatment with nicotine. To the best of our knowledge, till now, there are no reports of translational repression in HIV infected cells mediated by eIF2 α phosphorylation. Therefore, further regulation of eIF2 α phosphorylation in HIV infected neurons should be studied to control the translation of synaptic plasticity genes.

Conclusions

There are few reports showing the cognitive impairment in heavy smokers [68-71]. Effects of cigarette smoking on cognitive disorders in HIV infected patients are

scanty [23,72]. This is the first study in analyzing the molecular mechanisms behind the increased risk of HAND in HIV infected nicotine users. This study concludes that HIV infected nicotine users are more prone to develop HAND symptoms compared to HIV infected patients who are not nicotine addicts, due to the HDAC2 induced transcriptional repression. These negative regulatory effects on synaptic plasticity genes expression and spine density may be reversed by using the vorinostat (class I and II HDACs inhibitor). Vorinostat may be a useful therapeutic to inhibit the neurotoxic effects on synaptic plasticity in HIV infected nicotine abusers.

Materials and methods

Cell culture and virus

SK-N-MC neuronal cells were obtained from ATCC (Cat # HTB-10). HIV-1Ba-L (clade B, macrophage-tropic [R5] virus) was obtained through AIDS Research and Reference Reagent Program, Division of AIDS, NIAID, NIH (NIH AIDS Reagent Program Cat. # 510).

HIV-1 infection and/or nicotine treatment of SK-N-MC cells

SK-N-MC neuronal cells were infected with HIV-1 using the previously described protocol [13,73-75] with slight modifications. Briefly, SK-N-MC cells (1×10^6 cells) were cultured overnight in T-75 flasks using complete minimum essential medium. The cells were activated by treating with polybrene (10 μ g/ml) for 8 hrs before the infection. The cells were infected with TCID₅₀ of HIV-1 and/or treated with 1 μ M-1 mM nicotine (Sigma, Cat # N3876) for 7–10 days under same experimental conditions. In every second day, 2 ml of fresh medium (with/without nicotine) was added to the culture. Controls cells (without HIV-1/Nicotine) were included in the set-up of all experiments.

Inhibition of HDAC2 up-regulation

SK-N-MC cells were infected with HIV and/or treated with nicotine as explained above. On the 5th day of HIV infection and/or nicotine treatment, SK-N-MC cells were treated with 0.1-10 μ M vorinostat. After total 7 days of HIV infection and/or nicotine treatment, expression of HDAC2 was analyzed by qPCR and western blot.

mRNA and protein extraction; and first strand cDNA synthesis

After 7 days of HIV infection and/or co-treatment with optimum concentration of nicotine (100 μ M)/Vorinostat (1 μ M), SK-N-MC cells were harvested and the pellet was used for the mRNA isolation using illustra triple-Prep Kit (GE Healthcare Life Sciences, UK; Cat # 28-9425-44) and on-column DNase treatment step was also performed in the procedure. Purity and concentration of the RNA was measured by microspot RNA reader

(Synergy HT Multi-Mode Microplate Reader from Bio-Tek, US) and RNAs with an OD₂₆₀ nm/OD₂₈₀ nm absorbance ratio of at least 2.0 were used for PCR array. This mRNA was also used for the long terminal repeat (LTR) and HDAC2 gene expression using RT-qPCR. One microgram of RNA from all the control and test groups (1. Uninfected/untreated; 2. Nicotine; 3. HIV; 4. HIV + Nicotine; 5. Vorinostat; 6. Nicotine + Vorinostat; 7. HIV + Vorinostat and 8. HIV + Nicotine + Vorinostat) was used for the first strand cDNA synthesis using SABiosciences's RT² First Strand Kit (Cat. # 330401) as per supplier's protocol. Genomic DNA elimination step was performed prior to reverse transcription. With the use of illustra triplePrep Kit, from the same set of samples, protein was precipitated and preceded for SDS-PAGE after the estimation of protein concentration.

QPCR of viral transcripts in the presence of nicotine

The cDNA for m-RNA isolated from all control and test groups (as mentioned above) was synthesized using the high-capacity reverse transcriptase cDNA kit (Applied Biosystems, Cat # 4368814) to perform qRT-PCR by Taqman gene expression assay as described by manufacturer's protocols. A quantitative HIV-1 DNA protocol (LTR real-time PCR) was used to analyze the viral transcripts in HIV infected SK-N-MC cells and in combination with nicotine and vorinostat. The following previously published [76] primers and probes were used: LTR U5/R-sense-5'-GGCTAACTAGGGAACCCACTG-3' and antisense-5'-CTGCTAGAGATTTTCCACACTGAC-3', probe 5'-FAM-TGTGCCCCGTCTGTTGTGTG-TAMRA-3'. Values were normalized against GAPDH.

Expression of HDAC2 in SK-N-MC neuronal cells infected with HIV and/or treated with nicotine/vorinostat combination

In the SK-N-MC cells infected with HIV and/or treated with nicotine/vorinostat combination (as explained above), expression levels of HDAC2 was measured by qPCR with the use of Taqman assay (Assay ID # Hs00231032_m1) and by western blot using anti-HDAC2 antibody (Millipore cat # 07-222).

Human synaptic plasticity RT² profile PCR array

Synaptic plasticity gene expression profiling was done in all the neuronal cell control and test groups (all the groups as mentioned above) using 96 well format RT² Profile PCR Array human Synaptic Plasticity kit (SABiosciences, Cat. # PAHS-126A-2) using Stratagene Mx3000p qRT-PCR instrument. The human Synaptic Plasticity RT² Profiler PCR Array interrogates 84 genes related to the human synaptic plasticity. This kit was chosen because it includes diverse genes important in the human synaptic plasticity, including Immediate-Early Response (n = 30), Late Response (n = 2), Long Term Potentiation (n = 28),

Long Term Depression (n = 21), Cell Adhesion (n = 9), Extracellular Matrix & Proteolytic Processing (n = 5), CREB Cofactors (n = 10), Neuronal Receptors (n = 19), Postsynaptic Density (n = 15), as well as other genes involved in the synaptic plasticity (n = 2). Few genes have role in multiple functions listed above. Relative abundance of each mRNA species was assessed using RT² SYBR Green/ROX PCR Master mix (SABiosciences, Cat # 330520) and aliquoted in equal volumes (25 μ l) to each well of the real-time PCR arrays. The real-time PCR cycling program (as indicated by the manufacturer) was run on a Stratagene Mx3000p qRT-PCR thermal cycler. The threshold cycle (Ct) of each gene was determined by using the Stratagene MaxPro software. CT data were uploaded into the data analysis template on the manufacturer's website (<http://pcrdataanalysis.sabiosciences.com/pcr/arrayanalysis.php>). The relative expression of each gene in each test group was compared with the expression in control cells and it was calculated on the website using the $\Delta\Delta$ CT method with five housekeeping genes as controls. Controls are also included on each array for genomic DNA contamination, RNA quality, and general PCR performance.

Measurement of spine density

Dil staining

Established protocol to stain the neuronal cells and measurement of the spine density (in all the control and test group neuronal cells) was used with few modifications [13,74,77-81]. In brief, SK-N-MC cells were grown in Eagle's minimal essential medium (MEM) containing 10% fetal bovine serum (FBS), 5 mM sodium pyruvate, 100 units/ml penicillin, 100 mg/ml streptomycin and retinoic acid at 37°C with 5% CO₂. SK-N-MC cells were grown onto 22 mm \times 50 mm glass coverslips placed in a petri-dish. Cells were treated with polybrene for 8 hrs followed by addition of HIV and/or nicotine. On the 5th day of HIV infection and/or nicotine treatment, vorinostat (1 μ M) was added to the respective test groups. After 7 days of infection and/or treatment with nicotine, cells were fixed in 4% Formaldehyde in PBS for 30 min at RT. The fluorescent membrane tracer 1, 1'-Diiododecyl-3, 3,3',3'-tetramethylindocarbocyanine perchlorate (DiI) at 5 μ g/ml in PBS concentration was directly added onto the fixed cultures and allowed to incubate for 90 min at RT. The stained coverslips were placed at 4°C in small petri dishes containing PBS and allowed for overnight for the transport of dye. Cover-slips were mounted on the slides with Prolong[®]Gold antifade reagent (Invitrogen) before proceeding for confocal microscopy.

Confocal microscopy

Confocal images were obtained using TCS SP2 Confocal Laser Scanning Microscope (Leica Microsystems, Germany) at 488 nm (100%) illumination of an argon-ion laser using 60X

oil immersion objectives with high numeric aperture and 2.5X confocal electronic zoom settings to visualize individual cells and dendrites. Twenty Optical serial sections of 0.14 μ m/section (~2.8 μ m total) through the cells were reconstructed to yield complete "two dimensional" images of individual cells in focus.

Western blot assay

For SDS-PAGE, similar amounts of control and test group SK-N-MC cellular protein, typically 40 μ g per lane were used. All the protein samples were separated by using Any KD Mini-Protean TGX precast Gels (Bio-Rad, Cat # 456-9034). Proteins were transferred to nitrocellulose membranes, and the quality of protein measurement, electrophoresis, and transfer was checked by staining with Ponceau S. Membranes were blocked with 5% skimmed milk in TBS-T (20 mM Tris buffer, pH 7.5, 0.5 M NaCl, 0.1% Tween 20) for 1 hr at room temperature and incubated at 4°C overnight in the primary antibody diluted in 2% skimmed milk in TBS-T. The primary antibodies used were as follows: anti-HDAC2 antibody (Millipore cat # 07-222, 1/5000); rabbit anti-phospho-eIF2 α (Ser51) antibody (Cat # 9721, 1/1000), rabbit anti-eIF2 α antibody (Cat # 9722, 1/1000) from Cell Signaling (Beverly, MA). Subsequently, blots were washed in TBS-T (4 times, 10 min each) and incubated for 1 hr at room temperature in horse radish peroxidase-goat anti-rabbit antibody (Promega, Cat # W401B) diluted 1/2500 in 2% skim milk in TBS-T. After additional washings, protein bands were detected by chemiluminescence using SuperSignal West Pico Luminol/Enhancer (Thermo Scientific, Cat # 1856136) and SuperSignal West Pico substrate (Thermo Scientific, Cat # 1856135). Band density was measured by using ImageJ software.

Data analysis

In the expression studies, a gene was considered differentially regulated if the difference was ≥ 3 fold in comparison with the control. Experiments were performed at least three times and the values obtained were averaged. All the results are expressed as mean \pm standard error of the mean. Statistical analysis of two groups was performed by Student's t test, while more than two groups were analyzed using one way ANOVA followed by Bonferroni's multiple comparison test. Differences were considered significant at $p \leq 0.05$. If the combined effect observed is significantly greater than the expected (additive) effect, it was considered synergism [82]. Data analysis was performed with the Statistical Program, GraphPad Prism software (La Jolla, CA).

ImageJ software program was used to quantify DiI-labeled cells. Dendritic segments were chosen randomly from the apical and basal regions and at least one soma's length away from the cell soma.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

VSRA, MN, and SKS conceived and designed the experiments. VSRA performed the experiments. VSRA, SPK, VS, KRVK, ST, HD, AR and OH analyzed the data. MN contributed reagents/materials/analysis tools. VSRA, AR, SPK and KRVK prepared the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by National Institute of Health Grants: 1R03DA025576, 5R01DA021537, 1R01MH085259 and R01DA034547-01.

Author details

¹Department of Immunology, Institute of NeuroImmune Pharmacology, Herbert Wertheim College of Medicine, Florida International University, 11200 SW 8th Street, Miami, FL -33199, USA. ²CSIR- Centre for Cellular and Molecular Biology, Uppal Road, Hyderabad, India.

Received: 28 January 2014 Accepted: 28 April 2014

Published: 15 May 2014

References

- Sopori ML, Kozak W: Immunomodulatory effects of cigarette smoke. *J Neuroimmunol* 1998, **83**:148–156.
- Keller PA, Christiansen B, Kim SY, Piper ME, Redmond L, Adsit R, Fiore MC: Increasing consumer demand among Medicaid enrollees for tobacco dependence treatment: the Wisconsin "Medicaid covers it" campaign. *Am J Health Promot* 2011, **25**:392–395.
- U.S. Department of Health and Human Services: *How Tobacco Smoke Causes Disease: The Biology and Behavioral Basis for Smoking-Attributable Disease: A Report of the Surgeon General*. Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2010. [accessed 2013 Feb 26] In 2011/04/01 edition.
- U.S. Department of Health and Human Services: *Reducing Tobacco Use: A Report of the Surgeon General*. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health: Atlanta: U.S.; 2000. accessed 2013 Feb 26.
- National Institute on Drug Abuse: *Research Report Series: Tobacco Addiction*. Bethesda (MD): National Institutes of Health, National Institute on Drug Abuse; 2009. accessed 2013 Feb 26.
- Marshall MM, McCormack MC, Kirk GD: Effect of cigarette smoking on HIV acquisition, progression, and mortality. *AIDS Educ Prev* 2009, **21**:28–39.
- Marshall MM, Kirk GD, Caporaso NE, McCormack MC, Merlo CA, Hague JC, Mehta SH, Engels EA: Tobacco use and nicotine dependence among HIV-infected and uninfected injection drug users. *Addict Behav* 2011, **36**:61–67.
- Burkhalter JE, Springer CM, Chhabra R, Ostroff JS, Rapkin BD: Tobacco use and readiness to quit smoking in low-income HIV-infected persons. *Nicotine Tob Res* 2005, **7**:511–522.
- Gritz ER, Vidrine DJ, Lazev AB, Amick BC 3rd, Arduino RC: Smoking behavior in a low-income multiethnic HIV/AIDS population. *Nicotine Tob Res* 2004, **6**:71–77.
- Ellis R, Langford D, Masliah E: HIV and antiretroviral therapy in the brain: neuronal injury and repair. *Nat Rev Neurosci* 2007, **8**:33–44.
- McArthur JC, Brew BJ: HIV-associated neurocognitive disorders: is there a hidden epidemic? *Aids* 2010, **24**:1367–1370.
- Samikkannu T, Rao KVK, Pilakka-Kanthikeel S, Atluri VSR, Agudelo M, Roy U, Nair MP: Immunoneuropathogenesis of HIV-1 clades B and C: Role of redox expression and thiol modification. *Free Radic Biol Med* 2014, **69**:136–144.
- Atluri VS, Kanthikeel SP, Reddy PV, Yndart A, Nair MP: Human synaptic plasticity gene expression profile and dendritic spine density changes in HIV-infected human CNS cells: role in HIV-associated neurocognitive disorders (HAND). *PLoS One* 2013, **8**:e61399.
- Reddy P, Agudelo M, Atluri VR, Nair M: Inhibition of nuclear factor erythroid 2-related factor 2 exacerbates HIV-1 gp120-induced oxidative and inflammatory response: role in HIV associated neurocognitive disorder. *Neurochem Res* 2012, **37**:1697–1706.
- Hawkins BT, Brown RC, Davis TP: Smoking and ischemic stroke: a role for nicotine? *Trends Pharmacol Sci* 2002, **23**:78–82.
- Deiana S, Platt B, Riedel G: The cholinergic system and spatial learning. *Behav Brain Res* 2011, **221**:389–411.
- Levin E, McClernon FJ, Rezvani A: Nicotinic effects on cognitive function: behavioral characterization, pharmacological specification, and anatomic localization. *Psychopharmacology* 2006, **184**:523–539.
- Abbud RA, Finegan CK, Guay LA, Rich EA: Enhanced production of human immunodeficiency virus type 1 by in vitro-infected alveolar macrophages from otherwise healthy cigarette smokers. *J Infect Dis* 1995, **172**:859–863.
- Rock RB, Gekker G, Aravalli RN, Hu S, Sheng WS, Peterson PK: Potentiation of HIV-1 expression in microglial cells by nicotine: involvement of transforming growth factor-beta 1. *J Neuroimmune Pharmacol* 2008, **3**:143–149.
- Giunta B, Ehrhart J, Townsend K, Sun N, Vendrame M, Shytle D, Tan J, Fernandez F: Galantamine and nicotine have a synergistic effect on inhibition of microglial activation induced by HIV-1 gp120. *Brain Res Bull* 2004, **64**:165–170.
- Hossain MST, Fazio V, Mazzone P, Weksler B, Janigro D, Rapp E, Cucullo L: Tobacco smoke: a critical etiological factor for vascular impairment at the blood-brain barrier. *Brain Res* 2009, **1287**:192–205.
- Bryant VE, Kahler CW, Devlin KN, Monti PM, Cohen RA: The effects of cigarette smoking on learning and memory performance among people living with HIV/AIDS. *AIDS Care* 2013, **25**:1308–1316.
- Durazzo TC, Rothlind JC, Cardenas VA, Studholme C, Weiner MW, Meyerhoff DJ: Chronic cigarette smoking and heavy drinking in human immunodeficiency virus: consequences for neurocognition and brain morphology. *Alcohol* 2007, **41**:489–501.
- Kandel ER: The molecular biology of memory storage: a dialogue between genes and synapses. *Science* 2001, **294**:1030–1038.
- Imai K, Ochiai K: Role of histone modification on transcriptional regulation and HIV-1 gene expression: possible mechanisms of perioral diseases in AIDS progression. *J Oral Sci* 2011, **53**:1–13.
- Guan JS, Haggarty SJ, Giacometti E, Dannenberg JH, Joseph N, Gao J, Nieland TJ, Zhou Y, Wang X, Mazitschek R, Bradner JE, DePinho RA, Jaenisch R, Tsai L-H: HDAC2 negatively regulates memory formation and synaptic plasticity. *Nature* 2009, **459**:55–60.
- Hockly E, Richon VM, Woodman B, Smith DL, Zhou X, Rosa E, Sathasivam K, Ghazi-Noori S, Mahal A, Lowden PAS, Steffan JS, Marsh JL, Thompson LM, Lewis CM, Marks PA, Bates GP: Suberoylanilide hydroxamic acid, a histone deacetylase inhibitor, ameliorates motor deficits in a mouse model of Huntington's disease. *Proc Natl Acad Sci U S A* 2003, **100**:2041–2046.
- Hinnebusch AG: Translational regulation of GCN4 and the general amino acid control of yeast. *Annu Rev Microbiol* 2005, **59**:407–450.
- Hinnebusch AG, Dever TE, Asano K: *Mechanism of Translation Initiation in the Yeast Saccharomyces cerevisiae*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press; 2007.
- Dever TE: Gene-specific regulation by general translation factors. *Cell* 2002, **108**:545–556.
- Brody AL, Mandelkern MA, Jarvik ME, Lee GS, Smith EC, Huang JC, Bota RG, Bartzokis G, London ED: Differences between smokers and nonsmokers in regional gray matter volumes and densities. *Biol Psychiatry* 2004, **55**:77–84.
- Gallinat J, Meisenzahl E, Jacobsen LK, Kalus P, Bierbrauer J, Kienast T, Witthaus H, Leopold K, Seifert F, Schubert F, Staedtgen M: Smoking and structural brain deficits: a volumetric MR investigation. *Eur J Neurosci* 2006, **24**:1744–1750.
- Alvarez Losada S, Canto-Nogues C, Munoz-Fernandez MA: A new possible mechanism of human immunodeficiency virus type 1 infection of neural cells. *Neurobiol Dis* 2002, **11**:469–478.
- Zhao L, Li F, Zhang Y, Elbourkadi N, Wang Z, Yu C, Taylor EW: Mechanisms and genes involved in enhancement of HIV infectivity by tobacco smoke. *Toxicology* 2010, **278**:242–248.
- Ji D, Lape R, Dani JA: Timing and location of nicotinic activity enhances or depresses hippocampal synaptic plasticity. *Neuron* 2001, **31**:131–141.
- Mansvelder HD, McGehee DS: Long-term potentiation of excitatory inputs to brain reward areas by nicotine. *Neuron* 2000, **27**:349–357.
- Vigorito M, Cao J, Li M, Chang S: Acquisition and long-term retention of spatial learning in the human immunodeficiency virus-1 transgenic rat: effects of repeated nicotine treatment. *J NeuroViral* 2013, **19**:157–165.

38. Rezvani AH, Levin ED: **Cognitive effects of nicotine.** *Biol Psychiatry* 2001, **49**:258–267.
39. Silva AJ, Kogan JH, Frankland PW, Kida S: **CREB and memory.** *Annu Rev Neurosci* 1998, **21**:127–148.
40. Josselyn SA, Shi C, Carlezon WA Jr, Neve RL, Nestler EJ, Davis M: **Long-term memory is facilitated by cAMP response element-binding protein overexpression in the amygdala.** *J Neurosci* 2001, **21**:2404–2412.
41. Kaltschmidt B, Ndiaye D, Korte M, Pothion S, Arbibe L, Prullage M, Pfeiffer J, Lindecke A, Staiger V, Israel A, Kaltschmidt C, Memet S: **NF-kappaB regulates spatial memory formation and synaptic plasticity through protein kinase A/CREB signaling.** *Mol Cell Biol* 2006, **26**:2936–2946.
42. Ronn LC, Hartz BP, Bock E: **The neural cell adhesion molecule (NCAM) in development and plasticity of the nervous system.** *Exp Gerontol* 1998, **33**:853–864.
43. Kiss JZ, Muller D: **Contribution of the neural cell adhesion molecule to neuronal and synaptic plasticity.** *Rev Neurosci* 2001, **12**:297–310.
44. Yamagata K, Sanders LK, Kaufmann WE, Yee W, Barnes CA, Nathans D, Worley PF: **rheb, a growth factor- and synaptic activity-regulated gene, encodes a novel Ras-related protein.** *J Biol Chem* 1994, **269**:16333–16339.
45. Everall IP, Heaton RK, Marcotte TD, Ellis RJ, McCutchan JA, Atkinson JH, Grant I, Mallory M, Masliah E: **Cortical synaptic density is reduced in mild to moderate human immunodeficiency virus neurocognitive disorder.** HNRC Group. HIV Neurobehavioral Research Center. *Brain Pathol* 1999, **9**:209–217.
46. Counts SE, He B, Che S, Ikonovic MD, DeKosky ST, Ginsberg SD, Mufson EJ: **alpha7 nicotinic receptor up-regulation in cholinergic basal forebrain neurons in alzheimer disease.** *Arch Neurol* 2007, **64**:1771–1776.
47. Brown KC, Perry HE, Lau JK, Jones DV, Pulliam JF, Thornhill BA, Crabtree CM, Luo H, Chen YC, Dasgupta P: **Nicotine induces the up-regulation of the alpha7-nicotinic receptor (alpha7-nAChR) in human squamous cell lung cancer cells via the Sp1/GATA protein pathway.** *J Biol Chem* 2013, **288**:33049–33059.
48. Ballester LY, Capo-Velez CM, Garcia-Beltran WF, Ramos FM, Vazquez-Rosa E, Rios R, Mercado JR, Melendez RI, Lasalde-Dominicci JA: **Up-regulation of the neuronal nicotinic receptor alpha7 by HIV glycoprotein 120: potential implications for HIV-associated neurocognitive disorder.** *J Biol Chem* 2012, **287**:3079–3086.
49. Potter M, Figuera-Losada M, Rojas C, Slusher B: **Targeting the glutamatergic system for the treatment of HIV-associated neurocognitive disorders.** *J Neuroimmune Pharmacol* 2013, **8**:594–607.
50. Lambe EK, Picciotto MR, Aghajanian GK: **Nicotine induces glutamate release from thalamocortical terminals in prefrontal cortex.** *Neuropsychopharmacology* 2003, **28**:216–225.
51. Reno LA, Zago W, Markus RP: **Release of [(3)H]-L-glutamate by stimulation of nicotinic acetylcholine receptors in rat cerebellar slices.** *Neuroscience* 2004, **124**:647–653.
52. Morris MJ, Mahgoub M, Na ES, Pranav H, Monteggia LM: **Loss of histone deacetylase 2 improves working memory and accelerates extinction learning.** *J Neurosci* 2013, **33**:6401–6411.
53. Graff J, Rei D, Guan JS, Wang WY, Seo J, Hennig KM, Nieland TJ, Fass DM, Kao PF, Kahn M, Su SC, Samiei A, Joseph N, Haggarty SJ, Delalle I, Tsai L-H: **An epigenetic blockade of cognitive functions in the neurodegenerating brain.** *Nature* 2012, **483**:222–226.
54. Hawk JD, Florian C, Abel T: **Post-training intrahippocampal inhibition of class I histone deacetylases enhances long-term object-location memory.** *Learn Mem* 2011, **18**:367–370.
55. Hanson JE, Deng L, Hackos DH, Lo S-C, Lauffer BE, Steiner P, Zhou Q: **Histone deacetylase 2 cell autonomously suppresses excitatory and enhances inhibitory synaptic function in CA1 pyramidal neurons.** *J Neurosci* 2013, **33**:5924–5929.
56. Saiyed ZM, Gandhi N, Agudelo M, Napuri J, Samikkannu T, Reddy PV, Khatavkar P, Yndart A, Saxena SK, Nair MP: **HIV-1 Tat upregulates expression of histone deacetylase-2 (HDAC2) in human neurons: implication for HIV-associated neurocognitive disorder (HAND).** *Neurochem Int* 2011, **58**:656–664.
57. Aaronson DS, Horvath CM: **A road map for those who don't know JAK-STAT.** *Science* 2002, **296**:1653–1655.
58. Levy DE, Darnell JE Jr: **Stats: transcriptional control and biological impact.** *Nat Rev Mol Cell Biol* 2002, **3**:651–662.
59. Li WX: **Canonical and non-canonical JAK-STAT signaling.** *Trends Cell Biol* 2008, **18**:545–551.
60. Nicolas CS, Peineau S, Amici M, Csaba Z, Fafouri A, Javalet C, Collett VJ, Hildebrandt L, Seaton G, Choi SL, Sim SE, Bradley C, Lee K, Zhuo M, Kaang BK, Gressens P, Dournaud P, Fitzjohn SM, Bortolotto ZA, Cho K, Collingridge GL: **The Jak/STAT pathway is involved in synaptic plasticity.** *Neuron* 2012, **73**:374–390.
61. Archin NM, Espeseth A, Parker D, Cheema M, Hazuda D, Margolis DM: **Expression of latent HIV induced by the potent HDAC inhibitor suberoylanilide hydroxamic acid.** *AIDS Res Hum Retroviruses* 2009, **25**:207–212.
62. Contreras X, Schweneker M, Chen C-S, McCune JM, Deeks SG, Martin J, Peterlin BM: **Suberoylanilide hydroxamic acid reactivates HIV from latently infected cells.** *J Biol Chem* 2009, **284**:6782–6789.
63. Archin NM, Liberty AL, Kashuba AD, Choudhary SK, Kuruc JD, Crooks AM, Parker DC, Anderson EM, Kearney MF, Strain MC, Richman DD, Hudgens MG, Bosch RJ, Coffin JM, Eron JJ, Hazuda DJ, Margolis DM: **Administration of vorinostat disrupts HIV-1 latency in patients on antiretroviral therapy.** *Nature* 2012, **487**:482–485.
64. Kavanaugh SM, White LA, Kolesar JM: **Vorinostat: A novel therapy for the treatment of cutaneous T-cell lymphoma.** *Am J Health Syst Pharm* 2010, **67**:793–797.
65. Mitsiades CS, Mitsiades NS, McMullan CJ, Poulaki V, Shringarpure R, Hideshima T, Akiyama M, Chauhan D, Munshi N, Gu X, Bailey C, Joseph M, Libermann TA, Richon VM, Marks PA, Anderson KC: **Transcriptional signature of histone deacetylase inhibition in multiple myeloma: Biological and clinical implications.** *Proc Natl Acad Sci U S A* 2004, **101**:540–545.
66. Hurwitz JL, Stasik I, Kerr EM, Holohan C, Redmond KM, McLaughlin KM, Busacca S, Barbone D, Broaddus VC, Gray SG, O'Byrne KJ, Johnston PG, Fennell DA, Longley DB: **Vorinostat/SAHA-induced apoptosis in malignant mesothelioma is FLIP/caspase 8-dependent and HR23B-independent.** *Eur J Cancer* 2012, **48**:1096–1107.
67. Costa-Mattioli M, Gobert D, Stern E, Gamache K, Colina R, Cuello C, Sossin W, Kaufman R, Pelletier J, Rosenblum K, Krnjevic K, Lacaille JC, Nader K, Sonenberg N: **elF2alpha phosphorylation bidirectionally regulates the switch from short- to long-term synaptic plasticity and memory.** *Cell* 2007, **129**:195–206.
68. Richards M, Jarvis MJ, Thompson N, Wadsworth ME: **Cigarette smoking and cognitive decline in midlife: evidence from a prospective birth cohort study.** *Am J Public Health* 2003, **93**:994–998.
69. Sabia S, Elbaz A, Dugravot A, Head J, Shipley M, Hagger-Johnson G, Kivimaki M, Singh-Manoux A: **Impact of smoking on cognitive decline in early old age: the Whitehall II cohort study.** *Arch Gen Psychiatry* 2012, **69**:627–635.
70. Jacobsen LK, Krystal JH, Mendl WE, Westerveld M, Frost SJ, Pugh KR: **Effects of smoking and smoking abstinence on cognition in adolescent tobacco smokers.** *Biol Psychiatry* 2005, **57**:56–66.
71. Anstey KJ, von Sanden C, Salim A, O'Kearney R: **Smoking as a risk factor for dementia and cognitive decline: a meta-analysis of prospective studies.** *Am J Epidemiol* 2007, **166**:367–378.
72. Turner J, Page-Shafer K, Chin DP, Osmond D, Mossar M, Markstein L, Huitsing J, Barnes S, Clemente V, Chesney M: **Adverse impact of cigarette smoking on dimensions of health-related quality of life in persons with HIV infection.** *AIDS Patient Care STDS* 2001, **15**:615–624.
73. Ohagen A, Ghosh S, He J, Huang K, Chen Y, Yuan M, Osathanondh R, Gartner S, Shi B, Shaw G, Gabuzda D: **Apoptosis induced by infection of primary brain cultures with diverse human immunodeficiency virus type 1 isolates: evidence for a role of the envelope.** *J Virol* 1999, **73**:897–906.
74. Kurapati KR, Atluri VS, Samikkannu T, Nair MP: **Ashwagandha (Withania somnifera) reverses beta-amyloid1-42 induced toxicity in human neuronal cells: implications in HIV-associated neurocognitive disorders (HAND).** *PLoS One* 2013, **8**:e77624.
75. Eugenin EA, Berman JW: **Gap junctions mediate human immunodeficiency virus-bystander killing in astrocytes.** *J Neurosci* 2007, **27**:12844–12850.
76. Schmidtmayerova H, Alfano M, Nuovo G, Bukrinsky M: **Human immunodeficiency virus type 1 T-lymphotropic strains enter macrophages via a CD4- and CXCR4-mediated pathway: replication is restricted at a postentry level.** *J Virol* 1998, **72**:4633–4642.
77. Park JS, Bateman MC, Goldberg MP: **Rapid alterations in dendrite morphology during sublethal hypoxia or glutamate receptor activation.** *Neurobiol Dis* 1996, **3**:215–227.
78. Papa M, Bundman MC, Greenberger V, Segal M: **Morphological analysis of dendritic spine development in primary cultures of hippocampal neurons.** *J Neurosci* 1995, **15**:1–11.

79. Smith DL, Pozueta J, Gong B, Arancio O, Shelanski M: **Reversal of long-term dendritic spine alterations in Alzheimer disease models.** *Proc Natl Acad Sci U S A* 2009, **106**:16877–16882.
80. Pilakka-Kanthikeel S, Atluri VS, Sagar V, Saxena SK, Nair M: **Targeted brain derived neurotropic factors (BDNF) delivery across the blood-brain barrier for neuro-protection using magnetic nano carriers: an in-vitro study.** *PLoS One* 2013, **8**:e62241.
81. Ding H, Sagar V, Agudelo M, Pilakka-Kanthikeel S, Atluri VSR, Raymond A, Samikkannu T, Nair MP: **Enhanced blood–brain barrier transmigration using a novel transferrin embedded fluorescent magneto-liposome nanoformulation.** *Nanotechnology* 2014, **25**:5, 055101. doi:10.1088/0957-4484/25/5/055101.
82. Tallarida RJ: **Quantitative methods for assessing drug synergism.** *Genes Cancer* 2011, **2**:1003–1008.

doi:10.1186/1756-6606-7-37

Cite this article as: Atluri et al.: Vorinostat positively regulates synaptic plasticity genes expression and spine density in HIV infected neurons: role of nicotine in progression of HIV-associated neurocognitive disorder. *Molecular Brain* 2014 **7**:37.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

