

Secondhand smoke exposure, parental depressive symptoms and preschool behavioral outcomes

Short Title: Secondhand smoke, depression and preschool behavior
Short Title Word Count with Spaces: 51

Authors:

Nerissa S. Bauer, MD, MPH^{1,2}

Vibha Anand, PhD³

Aaron E. Carroll, MD, MS^{1,2}

Stephen M. Downs, MD, MS^{1,2}

¹Department of General & Community Pediatrics, Section of Children's Health Services Research, Indiana University School of Medicine, Indianapolis, Indiana, USA

²Regenstrief Institute for Healthcare, Indianapolis, Indiana, USA

³Pediatric Institute, Cleveland Clinic, Cleveland, OH, USA

Corresponding Author:

Nerissa S. Bauer, MD, MPH

410 W. 10th Street, Suite 1000, Indianapolis, IN 46202

Email: nsbauer@iu.edu; Office: 317-278-0552; Fax: 317-278-0456

Key Words: child, preschool; mental health; primary health care; screening; secondhand smoke

Acknowledgements: The authors would like to thank the Child Health Informatics Research and Development Lab (CHIRDL) with the Child Health Improvement through Computer Automation (CHICA) system and to Marc Rosenman, MD with IU School of Medicine Department of Pediatrics, Section of Children's Health Services Research; Jane Wang, PhD; and Roberta Ambuehl, BA of the Regenstrief Institute for their assistance with the creation of the database used in the study. The project was supported by grants 1R01HS018453 and R01 HS017939 from the Agency for Healthcare Research and Quality and R01 LM010031 from the National Library of Medicine.

Financial Disclosure for all authors: none

This is the author's manuscript of the article published in final edited form as:

Bauer, N. S., Anand, V., Carroll, A. E., & Downs, S. M. (2015). Secondhand smoke exposure, parental depressive symptoms and preschool behavioral outcomes. *Journal of Pediatric Nursing*, 30(1), 227–235. <http://doi.org/10.1016/j.pedn.2014.06.004>

Conflict of Interest for all authors: none

Abbreviations: Secondhand smoke (SHS); Child Health Improvement through Computer Automation (CHICA); Pre-screener form (PSF); Adjusted Odds Ratio (AOR); Confidence Interval (CI), Attention deficit-hyperactivity disorder (ADHD)

ABSTRACT

Little is known about the association of secondhand smoke (SHS) exposure and behavioral conditions among preschoolers. A cross-sectional analysis was used to examine billing and pharmacy claims from November 2004 to June 2012 linked to medical encounter-level data for 2,441 children from four pediatric community health clinics. Exposure to SHS was associated with attention deficit-hyperactivity disorder/ADHD and disruptive behavior disorder/DBD after adjusting for potential confounding factors. Assessment of exposure to SHS and parental depressive symptoms in early childhood may increase providers' ability to identify children at higher risk of behavioral issues and provide intervention at the earliest stages.

INTRODUCTION

The prevalence of behavioral conditions among preschool aged children has been increasing steadily (Egger & Angold, 2006). There also is a nationwide shortage of child psychiatrists and other community mental health providers with expertise in working with young children. Therefore, there is a critical need to examine efficient strategies to support pediatric health care providers who are asked to identify at-risk children as early as possible (Thomas & Holzer, 2006). Shifting the focus towards screening for psychosocial risk factors associated with pediatric behavioral issues may help identify at-risk children such that behavioral issues can be prevented or identified as they are emerging. This is important as pediatric providers can then feasibly manage these issues in the primary care setting. This understanding may aid in prioritizing screening and intervention efforts by pediatric healthcare providers and public health officials for at-risk children. SHS exposure can be screened for effectively in primary care settings and is amendable to brief counseling (Anand, Carroll, & Downs, 2012; Bandiera, Richardson, Lee, He, & Merikangas, 2011; Downs, Zhu, Anand, Biondich, & Carroll, 2008;

Hamer, Ford, Stamatakis, Dockray, & Batty, 2011; Rosen, Noach, Winickoff, & Hovell, 2012; Twardella, Bolte, Fromme, Wildner, & von Kries, 2010). In a 2007 national survey, 5.5 million children resided in households with a smoker (Singh, Siahpush, & Kogan, 2010). Despite successful US public health efforts aimed at reducing SHS exposure among non-smokers, children of non-Hispanic whites and blacks continue to be at highest risk (Pirkle, Bernert, Caudill, Sosnoff, & Pechacek, 2006; Sexton et al., 2004; Wilson, Kahn, Khoury, & Lanphear, 2005). Additionally, SHS exposure in the prenatal and postnatal periods has been associated with increased rates of childhood behavior problems (Herrmann, King, & Weitzman, 2008).

Despite the growing attention to the science on early brain and child development (Garner & Shonkoff, 2012), relatively little research has focused on secondhand smoke and its association with preschool behavioral outcomes. The majority of studies examining the negative effects of secondhand smoke and behavioral outcomes have relied upon samples of school-age children. One example of this is attention-deficit hyperactivity disorder (ADHD) (Banerjee, Middleton, & Faraone, 2007; Faraone, Sergeant, Gillberg, & Biederman, 2003; Froehlich et al., 2007; Langley, Rice, van den Bree, & Thapar, 2005; Yoshimasu et al., 2009). However, ADHD is increasingly being diagnosed among preschoolers such that clinical care guidelines have been revised to encourage providers to consider this diagnosis among this age group (Kollins et al., 2006; Wolraich et al., 2011). Moreover, identification rates of other common psychiatric disorders among preschoolers have been increasing as well (Carter et al., 2010; Egger & Angold, 2006; Wilens et al., 2002b). Pediatric providers routinely screen for various psychosocial risk factors during frequent well-child visits in the first five years of life. Yet, no studies were found in our review of the literature that have utilized data collected in community clinics to understand what risk factors are associated with earlier behavioral outcomes.

Prior work also has not accounted for important risk factors such as parental mental health conditions. For instance, parental depression has been correlated with poor childhood behavioral outcomes and is being increasingly screened for by pediatric healthcare providers (Earls, 2010; Field, 2011; Murray et al., 2011). Yet, past studies have not always controlled for this important confounder, i.e. depressive symptoms among parents of young children (Bandiera et al., 2011; Hamer et al., 2011; Twardella et al., 2010). Our findings add to existing literature by showing that secondhand smoke is associated with behavioral outcomes as early as preschool age. It also focuses on a community-based sample using data collected in primary care and adjusts for exposure to parental depressive symptoms.

METHODS

Study Design

This is a cross-sectional study of children between the ages of 0 and 6 years seen at 4 community pediatric clinics between November 2004 and June 2012. Data for this study was derived from a computerized decision support system used in conjunction with our electronic health record. Children were included in the study sample if complete visit level data for exposure to SHS and parental depressive symptoms were available.

Data Sources

We used data collected through the pre-screener form generated by the Child Health Improvement through Computer Automation (CHICA) system, which has been described elsewhere (Anand, Biondich, Liu, Rosenman, & Downs, 2004; Anand et al., 2012). Briefly, CHICA provides clinical decision support for both pediatric preventive care and disease management services (Anand et al., 2004; Anand et al., 2012; Downs et al., 2008). When a parent brings a child to a clinic that uses CHICA, a pre-screener form is generated for the parent

to complete in the waiting room. The pre-screener form contains 20 questions based on information contained in the child's health record and the age of the child at the visit as displayed in Figure 1 (Downs, Biondich, Anand, Zore, & Carroll, 2006). Once completed by the parent, the pre-screener form is scanned back into the system and the data is incorporated immediately into the child's health record. CHICA then generates a separate physician worksheet with 6 selected prompts, based on information the parent provided on the pre-screener form and information in the health record. These prompts are designed to call the pediatric provider's attention to specific health risks during the face-to-face encounter as displayed on a sample physician worksheet in Figure 2. At the end of the visit, the physician worksheet is scanned back into CHICA. Data for this study was extracted from the electronic medical record. We linked data from the parent-completed pre-screener form containing information about individual child and family health risks to administrative data of mental health diagnoses and prescriptions data from our statewide clinical network, the Indiana Network for Patient Care, which houses billing data, physician orders, filled prescriptions, and laboratory reports (McDonald et al., 2005).

Measures

SHS exposure. SHS exposure was defined by any positive response to CHICA's pre-screener question asking families whether any household member smoked. Parents of children 0 – 11 years of age were asked about SHS exposure if no previous information regarding the child's exposure to SHS existed in CHICA during the last 18 months or if the question had not been asked before. A positive response during any visit resulted in the child being classified as having had SHS exposure. Likewise, if no affirmative responses were captured for any visits, the child was categorized as not having had SHS exposure.

Socio-demographic characteristics. Child gender, race/ethnicity and insurance category were obtained from CHICA's database. Insurance category served as a proxy for socio-economic status.

Parental depressive symptoms. CHICA aids pediatric providers' monitoring of parental depressive symptoms using questions adapted from validated screening instruments as part of routine surveillance of pertinent health risks of the child and family. From 2004 to 2010, the pre-screener form included items from the Patient Health Questionnaire-2 (PHQ-2) measuring parental depressive mood and loss of interest or pleasure in doing things that previously was perceived as pleasurable (Kroenke, Spitzer, & Williams, 2003). In 2010, the Edinburgh Postnatal Depression Scale (EPDS-3) anxiety subscale items were substituted for the PHQ-2 to improve detection of postpartum depression. The EPDS-3 has been shown to have a high (95%) sensitivity and negative predictive value (98%) and measures parental symptoms of panic, self-blame and anxiety for postpartum depression (K. Kabir, Sheeder, & Kelly, 2008). Even though the use of the PHQ-2 is a valid way to detect parental depression in primary care settings, the items on the pre-screener form were changed to the EPDS-3 items as these were validated specifically for identification of postpartum depression. CHICA performs surveillance of parental mood every 90 days in the first 15 months of life or if there is no previous data on paternal depression in child's record in the electronic health record. If a parent responded affirmatively to any of the three surveillance items, the child was classified as being exposed to parental depressive symptoms; likewise children of parents with no affirmative responses to any of the symptoms at any visit were classified as not being exposed to parental depressive symptoms.

Behavioral Conditions. As we sought to examine the association of preschool children's exposure to SHS with children's behavior, we linked data captured from the pre-screener form to behavioral diagnoses that existed in the child's electronic health record. These behavioral diagnoses were identified using International Classification of Diseases-ninth revision (ICD-9) administrative billing codes for the following conditions: disruptive behavior disorder (DBD) (312.*), attention-deficit hyperactivity disorder (ADHD) (314.*), anxiety (300.*), depression (311), sleep disturbance (307.4), or adjustment disorder (309*).

Psychotropic Treatment. Psychotropic medications of interest included the following classes: stimulant medications, non-stimulants, alpha-2-agonists, sleep agents, atypical antipsychotic drugs, and selective serotonin reuptake inhibitors.

Statistical Analysis

Bivariate analyses of SHS exposure and socio-demographic variables were performed using chi-square test. Multivariable logistic regression models were used to assess the association between SHS exposure and each mental health diagnosis, adjusting for child gender, race/ethnicity, insurance status and parental depressive symptoms. An additional logistic regression model tested the association between SHS exposure and ever having been prescribed a psychotropic drug. All four covariates were selected *a priori*. Adjusted odds ratios (AOR) and 95% confidence intervals (CI) were calculated for each model. All analyses were performed using Stata11 (StataCorp, College Station, TX, 2010). This study was approved by our institution's Office of Research Administration-Human Subjects.

RESULTS

A total of 2,441 children were included in the study and their socio-demographics are reported in Table 1. The sample was 52% male, and a large proportion was Black (40.7%) or

Hispanic/Latino (45.3%), followed by White (10.5%). The majority of the families in the sample had public insurance (90.2%), followed by self-pay and uninsured (5.9%) and a minority with private insurance (3.9%). Approximately 27% of children (n=669) were exposed to SHS and 31.5% of children were exposed to parental depressive symptoms.

Race/ethnicity was significantly associated with SHS exposure. A majority of the children whose parents reported SHS exposure were Black (57%), followed by Hispanic/Latino (21%) and White (20%) parents reported the lowest rate of SHS exposure. A majority (89.4%) of families reporting SHS exposure were publically insured. A similar proportion of male and female children lived in households with reported SHS exposure.

The prevalence of behavioral conditions among preschoolers aged 3 to 6 years varied by diagnosis, with ADHD (3.3%) and disruptive behavior disorder (8.6%) being the most common as is shown in Table 1. Among our sample, 1.9% of preschoolers had ever been prescribed psychotropic medication.

Multivariable logistic regression analysis revealed significant associations between SHS and ADHD (AOR 1.9; 95% CI: 1.2-3.1) and DBD (AOR 1.6; 95% CI: 1.1-2.1) after adjusting for child's gender, race/ethnicity, insurance category and parental depressive symptoms. Parental depression was associated with a higher likelihood of a child ever being prescribed a psychotropic medication as shown in Table 2 (AOR 1.9; 95% CI 1.1-3.3). There were no other significant associations between SHS exposure and all other behavioral outcomes examined.

DISCUSSION

In this cross-sectional study of 2,441 children, 27% were exposed to SHS. SHS exposure was significantly associated with a diagnosis of ADHD and DBD among preschoolers, even after

adjusting for all other factors. Exposure to parental depressive symptoms was associated with having ever been prescribed a psychotropic medication.

Our study adds to the growing literature showing an association between SHS exposure and behavioral outcomes in early childhood (DiFranza, Aligne, & Weitzman, 2004; Lam, Leung, & Ho, 2001; Mannino, Moorman, Kingsley, Rose, & Repace, 2001; Stoddard & Gray, 1997; Yolton, Dietrich, Auinger, Lanphear, & Hornung, 2005). To our knowledge, ours is the first study to examine the association of SHS exposure and a range of preschool behavioral outcomes in a pediatric primary care setting (Bandiera et al., 2011; Hamer et al., 2011; Twardella et al., 2010). Our study differs from previous work in several significant ways and strengthens previously reported findings. First, while our study focused on preschool-aged children, similar to the cross-sectional study conducted by Twardella et al, (Twardella et al., 2010) the measurement of behavioral outcomes was assessed using different data sources. In that study, behavior problems were assessed using the “Strengths and Difficulties” questionnaire, a parent-reported behavioral screening tool (Twardella et al., 2010) while we used available administrative billing data to classify preschool behavioral outcomes. Given that parents with mood disorders are more likely to have concerns about children’s behavior, reliance on a parent-reported screening measure may overestimate the prevalence (Fergusson, Lynskey, & Horwood, 1993) due to reporting bias. We believe our data sources remove this source of bias.

Furthermore, we were able to adjust for the presence of parental depressive symptoms, a potential confounder. This is important not only because mental health conditions can be heritable, but also because individuals with mood disorders are more likely to smoke (Glazier, 2010; Lawrence, Mitrou, & Zubrick, 2009). In addition, caregivers affected by mental illness may experience sub-optimal parent-child interactions and diminished parenting capacity, both of

which place their children at higher risk of behavioral conditions (Crawford, Schrock, & Woodruff-Borden, 2011; Keown, 2011). For these reasons, controlling for parental depressive symptoms as a confounder is an important advantage of our study.

There is no known “safe” or acceptable level of SHS exposure. Given the mounting evidence for the negative effects of SHS and the rising prevalence of behavioral conditions among children, our study underscores the continued need for health promotion and intervention at the earliest stages. In 2006, mental illness was the highest healthcare cost among children, surpassing asthma, trauma and infectious disease (Soni, April 2009). Thus far, ADHD among school-aged children is the mental health condition with the greatest evidence of its association with SHS exposure (Braun, Kahn, Froehlich, Auinger, & Lanphear, 2006; Froehlich et al., 2009; Z. Kabir, Connolly, & Alpert, 2011). We found a similar significant association with ADHD and disruptive behaviors among preschool age children.

Our findings underscore the importance of continued smoking cessation counseling efforts. However, overall rates of SHS screening and counseling in both pediatrics and family practice clinics have remained low (Tanski, Klein, Winickoff, Auinger, & Weitzman, 2003; Winickoff et al., 2003; Zapka et al., 1999). Smoking cessation efforts may be strengthened if pediatric providers are able to emphasize to parents the health risks of SHS to children, especially to parents who are interested in quitting (Alwan, Siddiqi, Thomson, & Cameron, 2010; Katz, Muehlenbruch, Brown, Fiore, & Baker, 2002). Brief interventions lasting 3 minutes in primary care can lead to decreases in smoking cessation rates ("A clinical practice guideline for treating tobacco use and dependence: A US Public Health Service report. The Tobacco Use and Dependence Clinical Practice Guideline Panel, Staff, and Consortium Representatives," 2000). The growing evidence base of the association of SHS exposure and behavioral outcomes

may be a motivator for a subset of parents with a strong family history of mental illness. Pediatric providers can discuss these risks with parents and other caregivers and provide smoking cessation advice, perform targeted screening and ongoing surveillance of child's behavior and functioning for early-onset manifestations of behavioral conditions (Chen, Stanton, Hopper, & Khankari, 2011). Several randomized controlled trials of smoking cessation counseling with parents to protect their children from potential negative health effects have been found to be effective and doable within the scope of primary care practice (Abdullah, Mak, Loke, & Lam, 2005; Curry et al., 2003; Emmons et al., 2001; Rosen et al., 2012).

Some limitations to our study must be considered when interpreting the results. These include the retrospective design. We are not able to account for confounders that may contribute both to parental smoking and preschool behavioral conditions (McDonnell & Glod, 2003; Wilens et al., 2002a). However, we have tried to control for these by adjusting our analyses for socio-demographic characteristics and parental depressive symptoms. Over the course of the study, the method of capturing parental depressive symptoms changed from the PHQ-2 to the EPDS-3. Scores from the EPDS or the PHQ-9, from which the EPDS-3 and PHQ-2 are derived, were often concordant when using either instrument to screen for parental depression in the clinical care setting (Yawn et al., 2009). In this study, we classified SHS exposure by parent report rather than using a biomarker, such as cotinine. However, parental report has been shown to correlate to biomarkers and is a reliable indicator of exposure (Puig et al., 2008; Yeager & Krosnick, 2010). Moreover, the CHICA pre-screener question is similar to that used in the National Health and Nutrition Examination Survey (NHANES) that used the question, "Does anyone who lives here smoke cigarettes, cigars or pipes anywhere inside this home?" (Braun et al., 2008). Parental response to the SHS surveillance question is subject to social desirability

bias; however, the prevalence in our sample is similar to national estimates from the National Health Interview Survey (NHIS) examining parent-reported SHS exposure (Soliman, Pollack, & Warner, 2004). We were unable to control for maternal prenatal smoking history, nor whether any parent had a history of ADHD. While in utero exposure to SHS is a known risk factor for childhood ADHD, the postnatal presence of other smokers in the home, has been shown to be associated with ADHD symptoms in the absence of maternal smoking (Langley, Heron, Smith, & Thapar, 2012). Lastly, we acknowledge that there are inherent challenges to the reliance on ICD-9 codes of behavioral and mental health conditions in children less than 6 years of age (Egger & Emde, 2011); however, published studies have shown these classification systems can be used reliably for mental health disorders among children 2-5 years of age (Briggs-Gowan, Carter, Skuban, & Horwitz, 2001; Lavigne et al., 1998a, 1998b).

As pediatric healthcare providers build therapeutic relationships with families over time, they appreciate the risk and protective factors that may put a particular child at higher or lower need for behavioral concerns. Our study supports the recommendation by the American Academy of Pediatrics that primary care providers continue to pre-screen families for associated psychosocial risk factors, such as exposure to SHS and parental depression, as these risk factors are associated with behavioral issues and psychotropic drug treatment among preschoolers. Moreover, our findings support health promotion efforts that routinely occur during primary care visits that can more accurately aid in identifying families that may benefit from educational efforts to promote optimal parent-child interactions (Regalado & Halfon, 2001). Our study takes a critical first step towards understanding how pre-screening families for modifiable risk factors associated with negative childhood behavioral outcomes may increase providers' ability to identify children at higher risk of behavioral issues at the earliest stages.

CONCLUSION

Understanding what modifiable risk factors are associated with pediatric behavioral outcomes can help prioritize screening by pediatric healthcare providers so that efforts can be aimed at prevention or as these conditions are emerging. Given the need for primary care providers to perform active surveillance for a host of conditions that can affect childhood behavioral health, multi-faceted efforts to understand how to identify and feasibly intervene in primary care are needed. Preschoolers' exposure to SHS and parental depressive symptoms are potentially modifiable risk factors that can be screened for in primary care practice.

REFERENCES

- Abdullah, A. S., Mak, Y. W., Loke, A. Y., & Lam, T. H. (2005). Smoking cessation intervention in parents of young children: a randomised controlled trial. [Randomized Controlled Trial Research Support, Non-U.S. Gov't]. *Addiction*, *100*(11), 1731-1740. doi: 10.1111/j.1360-0443.2005.01231.x
- Alwan, N., Siddiqi, K., Thomson, H., & Cameron, I. (2010). Children's exposure to second-hand smoke in the home: a household survey in the North of England. [Research Support, Non-U.S. Gov't]. *Health & social care in the community*, *18*(3), 257-263. doi: 10.1111/j.1365-2524.2009.00890.x
- Anand, V., Biondich, P. G., Liu, G., Rosenman, M., & Downs, S. M. (2004). Child Health Improvement through Computer Automation: the CHICA system. [Research Support, N.I.H., Extramural Research Support, Non-U.S. Gov't Research Support, U.S. Gov't, P.H.S.]. *Studies in health technology and informatics*, *107*(Pt 1), 187-191.
- Anand, V., Carroll, A. E., & Downs, S. M. (2012). Automated primary care screening in pediatric waiting rooms. [Research Support, N.I.H., Extramural Research Support, Non-U.S. Gov't Research Support, U.S. Gov't, P.H.S.]. *Pediatrics*, *129*(5), e1275-1281. doi: 10.1542/peds.2011-2875
- Bandiera, F. C., Richardson, A. K., Lee, D. J., He, J. P., & Merikangas, K. R. (2011). Secondhand smoke exposure and mental health among children and adolescents. [Research Support, N.I.H., Extramural Research Support, N.I.H., Intramural Research Support, Non-U.S. Gov't]. *Archives of pediatrics & adolescent medicine*, *165*(4), 332-338. doi: 10.1001/archpediatrics.2011.30
- Banerjee, T. D., Middleton, F., & Faraone, S. V. (2007). Environmental risk factors for attention-deficit hyperactivity disorder. [Research Support, N.I.H., Extramural Review]. *Acta paediatrica*, *96*(9), 1269-1274. doi: 10.1111/j.1651-2227.2007.00430.x
- Braun, J. M., Froehlich, T. E., Daniels, J. L., Dietrich, K. N., Hornung, R., Auinger, P., & Lanphear, B. P. (2008). Association of environmental toxicants and conduct disorder in U.S. children: NHANES 2001-2004. [Research Support, N.I.H., Extramural Research Support, U.S. Gov't, Non-P.H.S.]. *Environ Health Perspect*, *116*(7), 956-962. doi: 10.1289/ehp.11177
- Braun, J. M., Kahn, R. S., Froehlich, T., Auinger, P., & Lanphear, B. P. (2006). Exposures to environmental toxicants and attention deficit hyperactivity disorder in U.S. children. [Research Support, N.I.H., Extramural Research Support, U.S. Gov't, Non-P.H.S.]. *Environmental health perspectives*, *114*(12), 1904-1909.
- Briggs-Gowan, M. J., Carter, A. S., Skuban, E. M., & Horwitz, S. M. (2001). Prevalence of social-emotional and behavioral problems in a community sample of 1- and 2-year-old children. [Research Support, U.S. Gov't, P.H.S.]. *J Am Acad Child Adolesc Psychiatry*, *40*(7), 811-819. doi: 10.1097/00004583-200107000-00016
- Carter, A. S., Wagmiller, R. J., Gray, S. A., McCarthy, K. J., Horwitz, S. M., & Briggs-Gowan, M. J. (2010). Prevalence of DSM-IV disorder in a representative, healthy birth cohort at school entry: sociodemographic risks and social adaptation. [Research Support, N.I.H., Extramural Research Support, U.S. Gov't, P.H.S.]. *Journal of the American Academy of Child and Adolescent Psychiatry*, *49*(7), 686-698. doi: 10.1016/j.jaac.2010.03.018

- Chen, X., Stanton, B., Hopper, J., & Khankari, N. (2011). Sources, locations, and predictors of environmental tobacco smoke exposure among young children from inner-city families. [Research Support, Non-U.S. Gov't]. *Journal of pediatric health care : official publication of National Association of Pediatric Nurse Associates & Practitioners*, 25(6), 365-372. doi: 10.1016/j.pedhc.2010.04.014
- A clinical practice guideline for treating tobacco use and dependence: A US Public Health Service report. The Tobacco Use and Dependence Clinical Practice Guideline Panel, Staff, and Consortium Representatives. (2000). [Consensus Development Conference Guideline Practice Guideline Research Support, Non-U.S. Gov't Research Support, U.S. Gov't, P.H.S. Review]. *JAMA*, 283(24), 3244-3254.
- Crawford, N. A., Schrock, M., & Woodruff-Borden, J. (2011). Child internalizing symptoms: contributions of child temperament, maternal negative affect, and family functioning. *Child Psychiatry Hum Dev*, 42(1), 53-64. doi: 10.1007/s10578-010-0202-5
- Curry, S. J., Ludman, E. J., Graham, E., Stout, J., Grothaus, L., & Lozano, P. (2003). Pediatric-based smoking cessation intervention for low-income women: a randomized trial. [Clinical Trial Randomized Controlled Trial Research Support, U.S. Gov't, P.H.S.]. *Archives of pediatrics & adolescent medicine*, 157(3), 295-302.
- DiFranza, J. R., Aligne, C. A., & Weitzman, M. (2004). Prenatal and postnatal environmental tobacco smoke exposure and children's health. [Review]. *Pediatrics*, 113(4 Suppl), 1007-1015.
- Downs, S. M., Biondich, P. G., Anand, V., Zore, M., & Carroll, A. E. (2006). Using Arden Syntax and adaptive turnaround documents to evaluate clinical guidelines. *AMIA Annu Symp Proc*, 214-218. doi: 86547 [pii]
- Downs, S. M., Zhu, V., Anand, V., Biondich, P. G., & Carroll, A. E. (2008). The CHICA smoking cessation system. [Research Support, N.I.H., Extramural Research Support, Non-U.S. Gov't]. *AMIA ... Annual Symposium proceedings / AMIA Symposium. AMIA Symposium*, 166-170.
- Earls, M. F. (2010). Incorporating recognition and management of perinatal and postpartum depression into pediatric practice. *Pediatrics*, 126(5), 1032-1039. doi: 10.1542/peds.2010-2348
- Egger, H. L., & Angold, A. (2006). Common emotional and behavioral disorders in preschool children: presentation, nosology, and epidemiology. [Review]. *Journal of child psychology and psychiatry, and allied disciplines*, 47(3-4), 313-337. doi: 10.1111/j.1469-7610.2006.01618.x
- Egger, H. L., & Emde, R. N. (2011). Developmentally sensitive diagnostic criteria for mental health disorders in early childhood: the diagnostic and statistical manual of mental disorders-IV, the research diagnostic criteria-preschool age, and the diagnostic classification of mental health and developmental disorders of infancy and early childhood-revised. [Research Support, Non-U.S. Gov't]. *Am Psychol*, 66(2), 95-106. doi: 10.1037/a0021026
- Emmons, K. M., Hammond, S. K., Fava, J. L., Velicer, W. F., Evans, J. L., & Monroe, A. D. (2001). A randomized trial to reduce passive smoke exposure in low-income households with young children. [Clinical Trial Randomized Controlled Trial Research Support, Non-U.S. Gov't Research Support, U.S. Gov't, P.H.S.]. *Pediatrics*, 108(1), 18-24.
- Faraone, S. V., Sergeant, J., Gillberg, C., & Biederman, J. (2003). The worldwide prevalence of ADHD: is it an American condition? *World Psychiatry*, 2(2), 104-113.

- Fergusson, D. M., Lynskey, M. T., & Horwood, L. J. (1993). The effect of maternal depression on maternal ratings of child behavior. [Research Support, Non-U.S. Gov't]. *Journal of abnormal child psychology*, *21*(3), 245-269.
- Field, T. (2011). Prenatal depression effects on early development: a review. [Research Support, N.I.H., Extramural Research Support, Non-U.S. Gov't]. *Infant behavior & development*, *34*(1), 1-14. doi: 10.1016/j.infbeh.2010.09.008
- Froehlich, T. E., Lanphear, B. P., Auinger, P., Hornung, R., Epstein, J. N., Braun, J., & Kahn, R. S. (2009). Association of tobacco and lead exposures with attention-deficit/hyperactivity disorder. [Research Support, N.I.H., Extramural Research Support, Non-U.S. Gov't]. *Pediatrics*, *124*(6), e1054-1063. doi: 10.1542/peds.2009-0738
- Froehlich, T. E., Lanphear, B. P., Epstein, J. N., Barbaresi, W. J., Katusic, S. K., & Kahn, R. S. (2007). Prevalence, recognition, and treatment of attention-deficit/hyperactivity disorder in a national sample of US children. [Research Support, N.I.H., Extramural Research Support, Non-U.S. Gov't Research Support, U.S. Gov't, P.H.S.]. *Archives of pediatrics & adolescent medicine*, *161*(9), 857-864. doi: 10.1001/archpedi.161.9.857
- Garner, A. S., & Shonkoff, J. P. (2012). Early childhood adversity, toxic stress, and the role of the pediatrician: translating developmental science into lifelong health. *Pediatrics*, *129*(1), e224-231. doi: 10.1542/peds.2011-2662
- Glazier, K. (2010). Tobacco-Induced Diseases: An Emphasis on Mental Health Conditions that May Result from Cigarette Use. *The New School Psychology Bulletin*, *8*(1), 56-61.
- Hamer, M., Ford, T., Stamatakis, E., Dockray, S., & Batty, G. D. (2011). Objectively measured secondhand smoke exposure and mental health in children: evidence from the Scottish health survey. *Archives of pediatrics & adolescent medicine*, *165*(4), 326-331. doi: 10.1001/archpediatrics.2010.243
- Herrmann, M., King, K., & Weitzman, M. (2008). Prenatal tobacco smoke and postnatal secondhand smoke exposure and child neurodevelopment. [Review]. *Curr Opin Pediatr*, *20*(2), 184-190. doi: 10.1097/MOP.0b013e3282f56165
- Kabir, K., Sheeder, J., & Kelly, L. S. (2008). Identifying postpartum depression: are 3 questions as good as 10? [Comparative Study]. *Pediatrics*, *122*(3), e696-702. doi: 10.1542/peds.2007-1759
- Kabir, Z., Connolly, G. N., & Alpert, H. R. (2011). Secondhand smoke exposure and neurobehavioral disorders among children in the United States. [Comparative Study Research Support, Non-U.S. Gov't]. *Pediatrics*, *128*(2), 263-270. doi: 10.1542/peds.2011-0023
- Katz, D. A., Muehlenbruch, D. R., Brown, R. B., Fiore, M. C., & Baker, T. B. (2002). Effectiveness of a clinic-based strategy for implementing the AHRQ Smoking Cessation Guideline in primary care. [Clinical Trial Research Support, Non-U.S. Gov't Research Support, U.S. Gov't, P.H.S.]. *Prev Med*, *35*(3), 293-301.
- Keown, L. J. (2011). Predictors of Boys' ADHD Symptoms from Early to Middle Childhood: The Role of Father-Child and Mother-Child Interactions. *J Abnorm Child Psychol*. doi: 10.1007/s10802-011-9586-3
- Kollins, S., Greenhill, L., Swanson, J., Wigal, S., Abikoff, H., McCracken, J., . . . Bauzo, A. (2006). Rationale, design, and methods of the Preschool ADHD Treatment Study (PATs). [Comment Multicenter Study Randomized Controlled Trial Research Support, N.I.H.,

- Extramural]. *Journal of the American Academy of Child and Adolescent Psychiatry*, 45(11), 1275-1283. doi: 10.1097/01.chi.0000235074.86919.dc
- Kroenke, K., Spitzer, R. L., & Williams, J. B. (2003). The Patient Health Questionnaire-2: validity of a two-item depression screener. *Med Care*, 41(11), 1284-1292. doi: 10.1097/01.MLR.0000093487.78664.3C
- Lam, T. H., Leung, G. M., & Ho, L. M. (2001). The effects of environmental tobacco smoke on health services utilization in the first eighteen months of life. [Research Support, Non-U.S. Gov't]. *Pediatrics*, 107(6), E91.
- Langley, K., Heron, J., Smith, G. D., & Thapar, A. (2012). Maternal and paternal smoking during pregnancy and risk of ADHD symptoms in offspring: testing for intrauterine effects. *Am J Epidemiol*, 176(3), 261-268. doi: 10.1093/aje/kwr510
- Langley, K., Rice, F., van den Bree, M. B., & Thapar, A. (2005). Maternal smoking during pregnancy as an environmental risk factor for attention deficit hyperactivity disorder behaviour. A review. [Research Support, Non-U.S. Gov't Review]. *Minerva Pediatr*, 57(6), 359-371.
- Lavigne, J. V., Arend, R., Rosenbaum, D., Binns, H. J., Christoffel, K. K., & Gibbons, R. D. (1998a). Psychiatric disorders with onset in the preschool years: I. Stability of diagnoses. *J Am Acad Child Adolesc Psychiatry*, 37(12), 1246-1254.
- Lavigne, J. V., Arend, R., Rosenbaum, D., Binns, H. J., Christoffel, K. K., & Gibbons, R. D. (1998b). Psychiatric disorders with onset in the preschool years: II. Correlates and predictors of stable case status. [Research Support, U.S. Gov't, P.H.S.]. *J Am Acad Child Adolesc Psychiatry*, 37(12), 1255-1261. doi: 10.1097/00004583-199812000-00008
- Lawrence, D., Mitrou, F., & Zubrick, S. R. (2009). Smoking and mental illness: results from population surveys in Australia and the United States. [Research Support, N.I.H., Extramural Research Support, Non-U.S. Gov't Research Support, U.S. Gov't, Non-P.H.S.]. *BMC public health*, 9, 285. doi: 10.1186/1471-2458-9-285
- Mannino, D. M., Moorman, J. E., Kingsley, B., Rose, D., & Repace, J. (2001). Health effects related to environmental tobacco smoke exposure in children in the United States: data from the Third National Health and Nutrition Examination Survey. *Archives of pediatrics & adolescent medicine*, 155(1), 36-41.
- McDonald, C. J., Overhage, J. M., Barnes, M., Schadow, G., Blevins, L., Dexter, P. R., . . . Comm, I. M. (2005). The Indiana network for patient care: A working local health information infrastructure. *Health affairs*, 24(5), 1214-1220. doi: Doi 10.1377/Hlthaff.24.5.1214
- McDonnell, M. A., & Glod, C. (2003). Prevalence of psychopathology in preschool-age children. [Review]. *J Child Adolesc Psychiatr Nurs*, 16(4), 141-152.
- Murray, L., Arteche, A., Fearon, P., Halligan, S., Goodyer, I., & Cooper, P. (2011). Maternal postnatal depression and the development of depression in offspring up to 16 years of age. [Research Support, Non-U.S. Gov't]. *Journal of the American Academy of Child and Adolescent Psychiatry*, 50(5), 460-470. doi: 10.1016/j.jaac.2011.02.001
- Pirkle, J. L., Bernert, J. T., Caudill, S. P., Sosnoff, C. S., & Pechacek, T. F. (2006). Trends in the exposure of nonsmokers in the U.S. population to secondhand smoke: 1988-2002. *Environ Health Perspect*, 114(6), 853-858.
- Puig, C., Garcia-Algar, O., Monleon, T., Pacifici, R., Zuccaro, P., Sunyer, J., . . . Vall, O. (2008). A longitudinal study of environmental tobacco smoke exposure in children: parental

- self reports versus age dependent biomarkers. [Research Support, Non-U.S. Gov't]. *BMC public health*, 8, 47. doi: 10.1186/1471-2458-8-47
- Regalado, M., & Halfon, N. (2001). Primary care services promoting optimal child development from birth to age 3 years: review of the literature. [Research Support, Non-U.S. Gov't Review]. *Archives of pediatrics & adolescent medicine*, 155(12), 1311-1322.
- Rosen, L. J., Noach, M. B., Winickoff, J. P., & Hovell, M. F. (2012). Parental smoking cessation to protect young children: a systematic review and meta-analysis. [Meta-Analysis Research Support, Non-U.S. Gov't Review]. *Pediatrics*, 129(1), 141-152. doi: 10.1542/peds.2010-3209
- Sexton, K., Adgate, J. L., Church, T. R., Hecht, S. S., Ramachandran, G., Greaves, I. A., . . . Geisser, M. S. (2004). Children's exposure to environmental tobacco smoke: using diverse exposure metrics to document ethnic/racial differences. [Research Support, Non-U.S. Gov't Research Support, U.S. Gov't, Non-P.H.S. Research Support, U.S. Gov't, P.H.S.]. *Environ Health Perspect*, 112(3), 392-397.
- Singh, G. K., Siahpush, M., & Kogan, M. D. (2010). Disparities in children's exposure to environmental tobacco smoke in the United States, 2007. *Pediatrics*, 126(1), 4-13. doi: 10.1542/peds.2009-2744
- Soliman, S., Pollack, H. A., & Warner, K. E. (2004). Decrease in the prevalence of environmental tobacco smoke exposure in the home during the 1990s in families with children. [Research Support, Non-U.S. Gov't]. *American journal of public health*, 94(2), 314-320.
- Soni, A. (April 2009). *The Five Most Costly Conditions Children's Conditions, 2006: Estimates for the U.S. Civilian Noninstitutionalized Children, Ages 0-17. Statistical Brief #242*. Rockville, MD: Agency for Healthcare Research and Quality.
- Stoddard, J. J., & Gray, B. (1997). Maternal smoking and medical expenditures for childhood respiratory illness. *American journal of public health*, 87(2), 205-209.
- Tanski, S. E., Klein, J. D., Winickoff, J. P., Auinger, P., & Weitzman, M. (2003). Tobacco counseling at well-child and tobacco-influenced illness visits: opportunities for improvement. [Research Support, U.S. Gov't, P.H.S.]. *Pediatrics*, 111(2), E162-167.
- Thomas, C. R., & Holzer, C. E., 3rd. (2006). The continuing shortage of child and adolescent psychiatrists. *Journal of the American Academy of Child and Adolescent Psychiatry*, 45(9), 1023-1031. doi: 10.1097/01.chi.0000225353.16831.5d
- Twardella, D., Bolte, G., Fromme, H., Wildner, M., & von Kries, R. (2010). Exposure to secondhand tobacco smoke and child behaviour - results from a cross-sectional study among preschool children in Bavaria. [Research Support, Non-U.S. Gov't]. *Acta paediatrica*, 99(1), 106-111. doi: 10.1111/j.1651-2227.2009.01522.x
- Wilens, T. E., Biederman, J., Brown, S., Monuteaux, M., Prince, J., & Spencer, T. J. (2002a). Patterns of psychopathology and dysfunction in clinically referred preschoolers. *J Dev Behav Pediatr*, 23(1 Suppl), S31-36.
- Wilens, T. E., Biederman, J., Brown, S., Monuteaux, M., Prince, J., & Spencer, T. J. (2002b). Patterns of psychopathology and dysfunction in clinically referred preschoolers. *Journal of developmental and behavioral pediatrics : JDBP*, 23(1 Suppl), S31-36.
- Wilson, S. E., Kahn, R. S., Khoury, J., & Lanphear, B. P. (2005). Racial differences in exposure to environmental tobacco smoke among children. [Clinical Trial Comparative Study

- Randomized Controlled Trial Research Support, N.I.H., Extramural Research Support, U.S. Gov't, P.H.S.]. *Environ Health Perspect*, 113(3), 362-367.
- Winickoff, J. P., McMillen, R. C., Carroll, B. C., Klein, J. D., Rigotti, N. A., Tanski, S. E., & Weitzman, M. (2003). Addressing parental smoking in pediatrics and family practice: a national survey of parents. [Research Support, Non-U.S. Gov't Research Support, U.S. Gov't, P.H.S.]. *Pediatrics*, 112(5), 1146-1151.
- Wolraich, M., Brown, L., Brown, R. T., DuPaul, G., Earls, M., Feldman, H. M., . . . Visser, S. (2011). ADHD: clinical practice guideline for the diagnosis, evaluation, and treatment of attention-deficit/hyperactivity disorder in children and adolescents. [Research Support, Non-U.S. Gov't Review]. *Pediatrics*, 128(5), 1007-1022. doi: 10.1542/peds.2011-2654
- Yawn, B. P., Pace, W., Wollan, P. C., Bertram, S., Kurland, M., Graham, D., & Dietrich, A. (2009). Concordance of Edinburgh Postnatal Depression Scale (EPDS) and Patient Health Questionnaire (PHQ-9) to assess increased risk of depression among postpartum women. [Research Support, U.S. Gov't, P.H.S.]. *Journal of the American Board of Family Medicine : JABFM*, 22(5), 483-491. doi: 10.3122/jabfm.2009.05.080155
- Yeager, D. S., & Krosnick, J. A. (2010). The validity of self-reported nicotine product use in the 2001-2008 National Health and Nutrition Examination Survey. [Validation Studies]. *Medical care*, 48(12), 1128-1132. doi: 10.1097/MLR.0b013e3181ef9948
- Yolton, K., Dietrich, K., Auinger, P., Lanphear, B. P., & Hornung, R. (2005). Exposure to environmental tobacco smoke and cognitive abilities among U.S. children and adolescents. *Environmental health perspectives*, 113(1), 98-103.
- Yoshimasu, K., Kiyohara, C., Minami, T., Yoshikawa, N., Kihira, S., Toyonaga, K., . . . Miyashita, K. (2009). Maternal smoking during pregnancy and offspring attention-deficit/hyperactivity disorder: a case-control study in Japan. [Research Support, Non-U.S. Gov't]. *Atten Defic Hyperact Disord*, 1(2), 223-231. doi: 10.1007/s12402-009-0015-1
- Zapka, J. G., Fletcher, K., Pbert, L., Druker, S. K., Ockene, J. K., & Chen, L. (1999). The perceptions and practices of pediatricians: tobacco intervention. [Research Support, Non-U.S. Gov't Research Support, U.S. Gov't, P.H.S.]. *Pediatrics*, 103(5), e65.

Table 1. Sample characteristics (n=2,441)

Variable	N	%
Socio-demographics		
Male Gender	1270	52
Race/Ethnicity		
White	256	10.5
Black	994	40.7
Hispanic/Latino	1106	45.3
Other	85	3.5
Insurance Type		
Commercial/Private	95	3.9
Medicaid/Public	2186	90.2
Self-Pay/Special Payer	142	5.9
Report of any secondhand smoke exposure	669	27.4
Report of any exposure to parental depressive symptoms	769	31.5
ICD-9 diagnoses between 3 and 6 years of age		
Attention-deficit hyperactivity disorder	80	3.3
Disruptive behavior disorder	209	8.6
Anxiety	17	0.7
Depression	9	0.4

Sleep Problems	7	0.3
Adjustment Disorder	41	1.7
Psychotropic Treatment		
Any Psychotropic Treatment	47	1.9

ICD-9= International Classification of Diseases, Ninth Edition

* Totals vary due to missing data

Table 2. Association between SHS exposure and behavioral conditions among preschoolers*

Characteristic	AOR	95% CI
Attention deficit-hyperactivity disorder	1.9	1.2-3.1
Disruptive behavior disorder	1.6	1.1-2.1
Anxiety	1.5	0.6-4.2
Depression	2.0	0.6-6.8
Sleep Problems	3.4	0.8-14.8
Adjustment Disorder	1.5	0.8-2.8

*Multivariable logistic regression with robust estimates, adjusted for gender, race/ethnicity, insurance type and parental depressive symptoms.

SHS: secondhand smoke; AOR: adjusted odds ratio; CI: confidence interval.

Variables that achieved statistical significance are bolded

5494266983 **CHICA Pre-Screening** MRN: #99-2
Name: Patient, Ima Great
Age: 13 mo DOB: Jan 18 2013
Date: Mar 7 2014

Uncooperative / Unable to Screen:
 Vision Hearing BP

Height: [] [] [] cm. Vision Left: 20/[] []
Weight: [] [] [] kg. Vision Right: 20/[] []
HC: [] [] [] cm. Vision Corrected?
BP: [] [] [] / [] [] [] Pulse Ox: [] [] [] %
Temp: [] [] [] deg. F Oral Axillary Rectal P F
Pulse: [] [] [] /min RR: [] [] []
 Sick Visit
 Patient left without treatment
 Patient refused to complete form
 Two IDs checked

Box For Nursing Use Only - Box For Nursing Use Only - Box For Nursing Use Only - Box For Nursing Use Only

Parents: Thank you for answering these questions about your child. The answers will help your doctor provide better quality of care. If your child is age 11 or older, he/she should answer the questions privately. Answers are confidential, but if you prefer not to answer that is allowed. You may want to talk about these questions with your doctor.
Please fill in the circles completely with a pencil or pen.

Y	N		Y	N	
<input type="radio"/>	<input type="radio"/>	Can Ima be allergic to any medicines?	<input type="radio"/>	<input type="radio"/>	Is Ima having pain today?
<input type="radio"/>	<input type="radio"/>	Can Ima let you know what she wants without crying or whining? Examples of this are pointing or pulling on you.	<input type="radio"/>	<input type="radio"/>	Does Ima say 'Dada' when she wants or sees her father? Does she say 'Mama' when she wants or sees her mother? Fill in YES if she says either one.
<input type="radio"/>	<input type="radio"/>	In the past 7 days, have you felt scared or panicky for not a very good reason?	<input type="radio"/>	<input type="radio"/>	In the past 7 days, have you blamed yourself unnecessarily when things went wrong?
<input type="radio"/>	<input type="radio"/>	In the past 7 days, have you been anxious or worried for no good reason?	<input type="radio"/>	<input type="radio"/>	Does anyone in Ima's home smoke?
<input type="radio"/>	<input type="radio"/>	Has your partner your children?	<input type="radio"/>	<input type="radio"/>	In the past 7 days, have you felt scared or panicky for not a very good reason?
<input type="radio"/>	<input type="radio"/>	Has Ima seen a doctor?	<input type="radio"/>	<input type="radio"/>	In the past 7 days, have you blamed yourself unnecessarily when things went wrong?
<input type="radio"/>	<input type="radio"/>	Is Ima on the WIC?	<input type="radio"/>	<input type="radio"/>	Does anyone in Ima's home smoke?
<input type="radio"/>	<input type="radio"/>	Does Ima spend time with anyone who has been in jail or a shelter, uses illegal drugs, or has HIV?	<input type="radio"/>	<input type="radio"/>	Has Ima been exposed to anyone with tuberculosis (TB) disease?
<input type="radio"/>	<input type="radio"/>	Is there any lye drain cleaner (Drano) in Ima's home?	<input type="radio"/>	<input type="radio"/>	Does Ima's home have a pool?
<input type="radio"/>	<input type="radio"/>	Was Ima born premature (more than one month early)?	<input type="radio"/>	<input type="radio"/>	Is Ima's outdoor play area fenced in?

Preguntas





Figure 1. Sample pre-screener form with secondhand smoke and parental depressive symptom exposure surveillance items

1225519072 **CHICA Physician Encounter Form** #99-2

Patient: Patient, Ima Great (F) **MRN:** #99-2 **A Vital Signs:**
DOB: 01/18/13 **Age:** 13 mo **Date:** Mar 7 2014
Doctor: Other **Time:** 4:01PM

Physical Exam: Nil Abnl **History / Exam Comments:** Additional notes on back...
 Two ID's Checked
 Informant: _____

General:
Head:
Skin:
Eyes:
Ears:
Nose / Throat:
Teeth / Gums:
Nodes:
Chest / Lungs:
Heart / Pulses:
Abdomen:
Ext. Genitalia:
Back:
Neuro:
Extremities:
Legend: * = Previous Examination

Vital Signs:
Height: cm. (%)
Weight: kg. (%)
BMI: (%)
Head Circ: cm. (%)
Temp: F ()
Pulse:
RR:
BP: ()
Pulse Ox: %
Hear (L): Hear (R):
Vision (L):
Vision (R):

**** ATTENTION **** Ima's parent identified symptoms of anxiety on 03/07/14. Ima has a smoker in the home. This may pose health risk to her.
Symptoms of sadness, anhedonia, sleep disturbance or appetite disturbance may indicate depression.

May have depression ----> Referred Discuss Smoking risks
 Counseled & gave handout
 Not depressed Mom not present

INSTRUCTIONS: Check all applicable boxes. COMPLETELY fill space to right of each box to "uncheck" misfilled boxes.

**** ATTENTION **** Ima's parent identified symptoms of anxiety on 03/07/14. Ima has a smoker in the home. This may pose health risk to her.
Symptoms of sadness, anhedonia, sleep disturbance or appetite disturbance may indicate depression.

May have depression ----> Referred Discuss Smoking risks
 Counseled & gave handout
 Not depressed Mom not present

Pb level indicated if: Ima 1) exposed to sib/playmate with hi Pb, 2) someone who works with lead, 3) lives in home built before 1950, 4) home built before 1978 recently renovated.

Pre-1950 house Pre-1978 renov. house Parent was interested in info-> Provided contact information
 Lead + contact Contact that works with lead Advise rear facing Provided brochure
 Ordered / Done elsewhere No risk factors Parent not interested in info

The AAP has age-specific recommendations for the prevention of childhood burns and fires. Please review the following, and check those you review:

Working smoke detector Store matches/lighters safely NEVER carry child & hot liquids NEVER play near hot appliances
 Test batteries monthly Adjust water max temp 120F/49C
 Rec no smoking in house-> Smoking = #1 cause of fires Safe place for play while cooking

Assessments and Plan: Medication Education Performed and/or Counseled on Vaccines: Y N N/A
 The medical student acted as a scribe for this note.

Staff: _____ Signature: _____

Figure 2. Sample physician worksheet with reminders to assess exposure to parental depressive symptoms and secondhand smoke