

1
2
3
4 **Title**

5
6 The Effect of Diluted Triple and Double Antibiotic Pastes on Dental Pulp Stem Cells and Established
7 *Enterococcus faecalis* Biofilm.
8
9

10
11
12 **AUTHORS**

13
14 Alaa HA. Sabrah,¹ Ghaeth H. Yassen,² Wai-Ching. Liu,² W. Scott. Goebel, ,³ Richard L. Gregory,⁴
15 Jeffrey A. Platt.²
16

17
18 ¹Department of Conservative Dentistry, The University of Jordan, Amman, Jordan.

19
20 ²Department of Restorative Dentistry, Indiana University School of Dentistry, Indianapolis, IN, USA.

21
22 ³Department of Pediatrics, Indiana University School of Medicine, Indianapolis, IN, USA.

23
24 ⁴Department of oral biology, Indiana University School of Dentistry, Indianapolis, IN, USA.
25
26
27

28
29 **CORRESPONDING AUTHOR**

30
31 Alaa H.A. Sabrah

32
33 The University of Jordan

34
35 Department of Conservative Dentistry

36
37 Queen Rania Al-Abdullah street

38
39 Amman 11942 Jordan

40
41 Tel: +962-077-9862670; fax: +966-06-5355522.
42

43
44 *E-mail address:* sabralolo@yahoo.com
45
46
47
48
49
50
51
52
53
54
55
56
57
58

59 This is the author's manuscript of the article published in final edited form as:

60 Sabrah, A. H. A., Yassen, G. H., Liu, W.-C., Goebel, W. S., Gregory, R. L., & Platt, J. A. (2015). The effect of
61 diluted triple and double antibiotic pastes on dental pulp stem cells and established *Enterococcus faecalis*
62 biofilm. *Clinical Oral Investigations*, 19(8), 2059–2066. <http://doi.org/10.1007/s00784-015-1423-6>
63
64
65

1
2
3
4 **ABSTRACT**
5

6 **Objectives:** To investigate the effect of various dilutions of antibiotic medicaments used in
7 endodontic regeneration on the survival of human dental pulp stem cells (DPSC) and to determine their
8 antibacterial effect against established *Enterococcus faecalis* biofilm.
9

10 **Materials and Methods:** The cytotoxic and antibacterial effects of different triple (TAP) and
11 double antibiotic (DAP) paste dilutions (0.125, 0.25, 0.5, 1, and 10 mg/ml) were tested against
12 *Enterococcus faecalis* established biofilm and DPSC. Established bacterial biofilm were exposed to
13 antibiotic dilutions for 3 days. Then, biofilms were collected, spiral plated and the numbers of bacterial
14 colony forming units (CFU/mL) were determined. For the cytotoxic effect, lactate dehydrogenase activity
15 assays (LDH) and cell viability assays (WST-1) were used to measure the percentage of DPSC
16 cytotoxicity after 3 day treatment with the same antibiotic dilutions. A general linear mixed model was
17 used for statistical analyses ($\alpha=0.05$).
18
19
20
21
22
23
24
25
26
27
28
29
30

31 **Results:** All antibiotic dilutions significantly decreased the bacterial CFU/mL. For WST-1
32 assays, all antibiotic dilutions except 0.125 mg/ml significantly reduced the viability of DPSC. For LDH
33 assays, the three lowest tested concentrations of DAP (0.5, 0.25, 0.125 mg/ml) and the two lowest
34 concentrations of TAP (0.25 and 0.125 mg/ml) were non-toxic to DPSC.
35
36
37
38
39

40 **Conclusions:** All tested dilutions had an antibacterial effect against *Enterococcus faecalis*.
41 However, 0.125 mg/ml of DAP and TAP showed a significant antibacterial effect with no cytotoxic
42 effects on DPSC.
43
44

45 **Clinical relevance:** Using appropriate antibiotic concentrations of intracanal medicament during
46 endodontic regeneration procedures is critical to disinfect root canal and decrease the adverse effects on
47 stem cells.
48
49
50
51
52

53 **KEYWORDS**
54

55 Triple antibiotic paste, double antibiotic paste, *Enterococcus faecalis*, established biofilm, dental pulp
56 stem cells.
57
58
59
60
61
62
63
64
65

1
2
3
4 **Introduction**
5

6 Endodontic regeneration techniques have received great attention in recent years. These
7 techniques may offer a continuation of root development and improve the prognosis of necrotic immature
8 teeth treatment [1-3]. Regeneration treatment protocol involves the use of an intracanal dressing with
9 antibacterial properties followed by the induction of bleeding to create a matrix for the growth of new
10 dento-pulpal tissue in the root canal space [4, 5]. Stem cells in the remaining vital pulp or apical papilla
11 have been hypothesized to mediate tissue regeneration [1, 5].
12
13
14
15
16
17
18
19

20 Since its first use by Hoshino and his research group [6, 7], triple antibiotic paste (TAP), a
21 combination of metronidazole, ciprofloxacin, and minocycline, is the most widely used intracanal
22 medicament in endodontic regeneration [5, 8-10]. TAP use has been reported in 51% of published cases
23 of endodontic regeneration [10]. To overcome the discoloration problem associated with the presence of
24 minocycline in TAP, a combination of metronidazole and ciprofloxacin (DAP) was proposed [11] and
25 used successfully in endodontic regeneration [12].
26
27
28
29
30
31
32

33 Due to the critical role of stem cells in endodontic regeneration, their viability in the presence of
34 antibiotic pastes is essential for success. Recently, studies suggested that intracanal medicaments (TAP,
35 DAP) are directly toxic to stem cells of apical papilla (SCAP), human periodontal ligament fibroblasts,
36 human dental pulp cells, and apical papilla cells [13-16] and indirectly toxic to SCAP [17] in the
37 concentrations currently used in endodontic regeneration. Most of the studies proposed that 0.1-2 mg/mL
38 of TAP or DAP did not have any cytotoxic effect on dental pulp cells [13, 15, 16]. TAP and DAP were
39 effective against newly formed bacterial biofilm at 0.03 and 0.01 mg/mL, respectively [18]. However,
40 these relatively low concentrations of medicaments may not be effective against established bacterial
41 biofilm. It is essential to explore the antibacterial effectiveness against established bacterial biofilm
42 because of the critical involvement of biofilm in the etiology of periapical periodontitis [19]. It is also
43 crucial to explore the optimum dilutions of antibiotic medicaments that have antibacterial effect against
44 established bacterial biofilm without compromising the survival of human dental pulp stem cells.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60 Therefore, the aim of this study was to explore the effect of various dilutions of antibiotic medicaments
61
62
63
64
65

1
2
3
4 used in endodontic regeneration on human dental pulp stem cells (DPSC) survival and to determine the
5
6 antibiofilm effect of these dilutions against established biofilm of *Enterococcus faecalis* (*E. faecalis*).
7
8

9 **MATERIALS AND METHODS**

10 **Antibacterial experiment**

11 **Bacterial Strain and Media**

12
13 *E. faecalis* (ATCC 29212 strain) was initially grown on anaerobic blood agar plates (CDC,
14
15 BioMerieux, Durham, NC USA). Then, bacteria cultures were grown in brain heart infusion broth
16
17 supplemented with 5 g/l yeast extract (BHI-YE) at 37°C in an anaerobic environment using gas
18
19 generating sachets (GaPak EZ, Becton, Dickinson and Company, Franklin Lakes, NJ, USA).
20
21
22

23 **Antibiotic medicament preparation**

24
25 For TAP, 100 mg of USP grade antibiotic powder compounded of equal portions of
26
27 metronidazole, ciprofloxacin, and minocycline (CHAMPS Medical, San Antonio, TX, USA) was
28
29 dissolved in 1 mL of sterile water (33.3 mg of each antibiotic/mL). For DAP (CHAMPS Medical), 100
30
31 mg of USP grade antibiotic powder compounded of equal portions of metronidazole and ciprofloxacin
32
33 was dissolved in 1 mL of sterile water (50 mg of each antibiotic/mL). The stock solution of each
34
35 antibiotic mixture was used to prepare the five experimental concentrations (10, 1, 0.5, and 0.25 and
36
37 0.125 mg/mL) by mixing the appropriate amount of the antibiotic solution with BHI-YE culture media for
38
39 *E. faecalis* biofilm assay. The ranges of dilutions used in this study were selected according to the most
40
41 recent available literature regarding both antibacterial and cytotoxic effect of these medicaments [7, 13,
42
43 15, 18]. The different dilutions were filter sterilized using a sterile 20 µm filter (Fisher Scientific, Newark,
44
45 DE, USA). A control group of BHI-YE culture media with no antibiotics was also used.
46
47
48
49
50

51 **Antibiotic effects on established bacterial biofilm**

52
53 To determine the effectiveness of the different dilutions of TAP and DAP antibiotic medicaments on the
54
55 established bacterial biofilm, 100 µl of an overnight *E. faecalis* culture (10⁶ CFU/mL) in BHI-YE and 900
56
57 µl fresh BHI-YE were incubated for 3 days in independent wells of 12-well plates (Becton, Dickinson and
58
59 Company). **The culture media were not changed during the 3 days of biofilm cultivation in an attempt to**
60
61
62
63
64
65

1
2
3
4 grow biofilm that is closer to the biofilm grows inside an infected root canal. A recent study found that
5
6 frequent replacement of fresh media may lead to overgrowth of facultative cocci and formation of a
7
8 biofilm that is untypical to root canal environment [20]. After the incubation time, planktonic bacterial in
9
10 culture supernatants were removed from each well, and the remaining established biofilm was washed
11
12 with saline solution (5mL over 60 minutes) and treated, in triplicate, with antibiotic dilutions in BHI-YE
13
14 for 3 days. At the end of the incubation time, antibiotic dilutions were removed from each well, and the
15
16 biofilm cells were gently washed with saline. The biofilm cells were collected by cell scrapers (Fisher
17
18 Scientific), vortexed, diluted in sterile saline and spirally plated on tryptic soy agar plates (TSA). The
19
20 plates were incubated for 48 h in a 5% CO₂ incubator at 37°C. Then, the number of colony forming units
21
22 per volume (CFUs/mL) of each treated well was determined using an automated colony counter
23
24 (Synbiosis, Inc., Frederick, MD, USA) and compared to the values from the control culture.
25
26
27
28

29 **Cytotoxicity experiment**

30 **Human dental pulp stem cells (DPSC)**

31
32 DPSC (Cook General BioTechnology LLC, Indianapolis, IN, USA) obtained from immature third
33
34 molars were used in this study. The stem-like nature of these cells has been previously established
35
36 utilizing flow cytometric analysis [21, 22]. The cells were cultured in DMEM media supplemented with
37
38 10% fetal bovine serum (Atlanta Biologicals Inc. Flowery Branch, GA, USA), 5% penicillin-streptomycin
39
40 (Life Technologies Corporation, Grand Island, NY, USA) and 2% Amphotericin B (Life Technologies
41
42 Corporation). Cells at passages between 3 and 5 were utilized in this experiment. Sub-confluent cells
43
44 were detached from the culture plate with 0.05% trypsin and EDTA (Life Technologies Corporation). The
45
46 evaluation of cytotoxic effects of the various TAP and DAP dilutions were determined using lactate
47
48 dehydrogenase (LDH) activity and cell viability colorimetric assays utilizing water soluble tetrazolium
49
50 salts (WST-1).
51
52
53
54

55 **LDH assays**

56
57 Sub-confluent DPSC were seeded into a 96-well plate (10,000 cells/well) and incubated for 24
58
59 hours in DMEM supplemented with 10% fetal bovine serum. After 24 hours, the old media was removed,
60
61
62
63
64
65

1
2
3
4 and the cells were incubated with serum free DMEM in the presence of different concentrations of TAP or
5
6 DAP mixed with DMEM culture media. Additionally, DPSC incubated without any medicament in serum
7
8 free DMEM were used as a negative control. After 3 days, the media were collected and utilized for the
9
10 LDH assays. The levels of LDH released by the cells into the media due to membrane damage were
11
12 determined using the Pierce™ LDH cytotoxicity assay kit (Thermo Scientific, Pierce Biotechnology,
13
14 Rockford, IL, USA). Briefly, 50 µl of DPSCs culture media were transferred into another 96-well plate.
15
16 Then, 50 µl of LDH assay mix was prepared according to the manufacturer's protocol, added and mixed
17
18 in each well, followed by incubation for 30 minutes at room temperature. A spectrophotometer
19
20 (SpectraMax 190; Molecular Devices, Sunnyvale, CA, USA) was used to quantitatively measure the
21
22 colorimetric change at a wavelength of 490 nm. Positive control wells (Maximum LDH release) were also
23
24 used by incubating DPSCs with the lysis solution provided by the manufacturer for 45 minutes (100 µl of
25
26 serum-free DMEM and 10 µl lysis solution). The following equation was used to calculate the percentage
27
28 of cytotoxicity due to LDH release from the treated cells relative to the untreated control:
29
30

$$31 \text{ Cytotoxicity (\%)} = (\text{experimental value} - \text{negative control}) / (\text{positive control} - \text{negative control}) \times 100.$$

32
33 The positive control cytotoxicity percentage was considered to be 100%, while the percentage
34
35 cytotoxicity of the negative control was 0%.
36
37

38 **WST-1 assays**

39
40 After collecting the media for LDH assays and washing the cells with phosphate buffer solution,
41
42 the DPSCs were incubated with 10 µl WST-1 and 90 µl serum-free DMEM for 4 h in a humidified
43
44 atmosphere at 37°C and 5% CO₂. After that, a spectrophotometer was used to measure the optical
45
46 absorbance of each well at 450 nm. Negative controls consisting of 10 µl WST-1 incubated with 90 µl
47
48 serum-free DMEM without the presence of cells were used as a blank. The following equation was used
49
50 to calculate the percentage of the absorbance values of the antibiotic-treated cells at each concentration in
51
52 comparison to the absorbance values of the untreated cells.
53
54

55
56 The percentage of cell viability (%) = (absorbance value of the treated cells / absorbance value of
57
58 untreated cells) × 100. The percentage cell viability of the untreated cells was considered 100%.
59
60
61
62
63
64
65

1
2
3
4
5
6
7 **Statistical Analysis**

8
9 Each experimental treatment was conducted in triplicate and repeated individually three times. **Data**
10 **were analyzed using a linear mixed model (SAS software 9.4, SAS Institute, NC, USA) to test the effect**
11 **of treatment on both bacteria and DPSC.** The untreated control groups in both experiments were used as
12
13 reference groups and they were removed from the model. The significance level was set at 0.05.
14
15

16
17
18 **Results**

19
20 A significant decrease in the log CFU/mL of *E. faecalis* bacteria was noticed with all dilutions of
21
22 TAP and DAP compared with the untreated control ($P < 0.0001$). For TAP, a gradual and significant
23
24 reduction in antibacterial activity was observed between different dilutions ($P = 0.003-0.0001$) (Figure 1).
25
26 However, 0.125 and 0.25 mg/mL concentrations of TAP were not significantly different from each other.
27
28 For DAP, a gradual and significant reduction in antibacterial activity was observed between different
29
30 dilutions ($P = 0.03-0.0001$) (Figure 1). However, 0.125, 0.25, and 0.5 mg/mL concentrations of DAP were
31
32 not significantly different from each other. **The log reductions in bacterial CFU/mL were 8.4, 6.5, 3.3,**
33 **1.7, and 2.3 for 10, 1, 0.5, 0.25, and 0.125 mg/ml TAP, respectively. The log reductions in bacterial**
34 **CFU/mL were 8.4, 6.8, 3.4, 3.9, and 2.9 for 10, 1, 0.5, 0.25, and 0.125 mg/ml DAP, respectively.**
35
36
37
38
39

40 For the LDH cytotoxic assays, the two highest tested dilutions of DAP (10 and 1 mg/mL)
41
42 produced a significant cytotoxic effect on DPSC compared to the untreated control ($P < 0.001$) (Figure 2).
43
44 For TAP, the three highest tested dilutions (10, 1, and 0.5 mg/mL) produced a significant cytotoxic effect
45
46 on DPSC compared to the untreated control ($P = 0.006- 0.0001$). For WST-1 assays, all dilutions of TAP
47
48 and DAP significantly impaired the viability of DPSC ($P < 0.0001$) except 0.125 mg/mL (Figure 3). On
49
50 the other hand, 0.125 mg/mL of TAP produced a significant enhancement in DPSC viability compared to
51
52 untreated control ($P < 0.0001$). **Images of DPSC in the presence of different dilutions of TAP and DAP**
53 **(0.25-10 mg/ml) are presented in Figure 4.**
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7 **Discussion**
8

9 One of the distinguishing characteristics of endodontic microorganisms is their ability to form
10 biofilms. Indeed, apical periodontitis has been classified as a biofilm-induced disease [19]. The use of
11 minimum or no instrumentation during endodontic regeneration requires an effective chemical challenge
12 to eradicate the endodontic pathogens. However, a root canal in an infected immature tooth has a large,
13 blunder buss apical foramen, which may cause the disinfecting agent to pass through the open apex and
14 induce a cytotoxic effect to the stem cells present in the periapical area. Furthermore, remaining vital pulp
15 in immature teeth may be in direct contact with the disinfecting agent. Therefore, it is essential to achieve
16 a balance between the antibacterial effect and the cytotoxic effect of any suggested intracanal
17 medicament.
18
19

20 In this study, both TAP and DAP at a concentration of 10 mg/mL were able to eradicate all the
21 established biofilm of *E. faecalis* after 3 days. All other concentrations of both antibiotic pastes
22 significantly reduced the established *E. faecalis* biofilm but did not completely eradicate it. The antibiotic
23 concentration required to completely eradicate the biofilm in this study (10 mg/mL) was higher than the
24 concentrations reported in previous studies [6, 7], which ranged from 0.075 to 1.5 mg/mL of TAP.
25 However, an agar diffusion test rather than bacterial biofilm was used in one study [6], and early biofilm
26 formation was measured in the other study [7]. A recent study reported that TAP and DAP were able to
27 significantly reduce new biofilm formation of *E. faecalis* and *Porphyromonas gingivalis* at
28 concentrations of 0.01 mg/mL for DAP and 0.03 mg/mL for TAP using a microtiter plate method [18].
29 However, the previous study tested the antibacterial effects against early biofilm formation rather than
30 established biofilm. To the best of our knowledge, this is the first report that investigates the antibacterial
31 effect of diluted antibiotic medicaments on established bacterial biofilm which is more representative of
32 the actual clinical situation. The established biofilm in this study was grown for only 3 days. However,
33 bacterial established biofilm in the root canal might be formed in a longer period of time before patient
34 seeks treatment. Future researches should investigate the antibacterial concentrations needed to eradicate
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 older established bacterial biofilm. *E. faecalis* was used as *in-vitro* model because it is a commonly
5
6 isolated microorganism from persistent endodontic infections [23], it poses various survivals and
7
8 virulence factors that enables it to resist nutritional deprivation [24] and hence resist treatment.
9

10
11 Additionally, *E. faecalis* is described as a high biofilm producer [25, 26]. **Future studies are needed to**
12
13 **identify antibiotic concentrations that are effective against bacterial biofilm taken from immature teeth**
14
15 **with infected root canals.**

16
17 **A concentration of 1 mg/ml of TAP or DAP was also effective in eradicating more than**
18
19 **99.9999 % of *E. faecalis* established biofilm (more than 6 unites log reduction in bacterial CFU/mL). A**
20
21 **material can be considered effective in eradicating the bacterial biofilm when it can produce a reduction**
22
23 **of more than 5 logarithmic units in the log CFU/mL (99.999 % reduction) compared to untreated control**
24
25 **[27-29]. The lowest three concentrations of both TAP and DAP (0.125-0.5 mg/mL) did not produce a**
26
27 **reduction of 5 logarithmic units in the log CFU/mL compared to untreated control. However, the**
28
29 **antibacterial effect measured in this study was accomplished after only 3 days of antibiotic application.**
30
31 **The use of these concentrations in clinical situations might be more efficient since the application time of**
32
33 **intracanal medicaments usually range between 1 to 4 weeks [10]. A material can be considered**
34
35 **bacteriocidal when it can produce a reduction of more than 3 logarithmic units in the log CFU/mL (99.9**
36
37 **% reduction in bacterial biofilm). A recent study has reported that the minimum bactericidal**
38
39 **% reduction in bacterial biofilm). A recent study has reported that the minimum bactericidal**
40
41 **concentration of TAP and DAP against *E. faecalis* were 0.3, and 0.14 mg/ml, respectively [18]. This is in**
42
43 **agreement with our results which show that TAP concentrations lower than 0.3 mg/ml (0.25, and 0.125**
44
45 **mg/ml) did not produce 3 log reductions in CFU/mL. Similarly, 0.125 mg/ml of DAP in our study did not**
46
47 **produce 3 log reductions in CFU/mL because it falls under the minimum bactericidal concentration**
48
49 **identified previously [18].**
50
51

52
53 **Different substrates have been used to develop bacterial biofilms in endodontics such as dentin,**
54
55 **polystyrene culture plates and hydroxyapatite disks [30]. Polystyrene culture plates, a popular substrate in**
56
57 ***in-vitro* endodontic experiments [18, 25, 31], was used in the current study. This substrate was suggested**
58
59 **to offer rapid retrieval and quantification of bacterial biofilm [30, 31]. On the other hand, the use of**
60
61
62
63
64
65

1
2
3
4 dentin substrate in antibiofilm studies is closer to the clinical situation. However, a relatively recent study
5
6 suggested similar pattern of antibiofilm effect against *E. faecalis* developed in both polystyrene plates and
7
8 root canal dentine [32].
9

10
11 For LDH cytotoxicity, the three lowest tested concentrations of DAP (0.125, 0.25 and 0.5
12 mg/mL) and the two lowest concentrations of TAP (0.125 and 0.25 mg/mL) were not toxic to DPSCs. A
13
14 recent study reported that 0.3-2.5 mg/mL of TAP and 0.5-1 mg/mL of DAP were not toxic to dental pulp
15
16 fibroblasts using LDH assays [15]. The difference in detected cytotoxic concentrations using LDH assays
17
18 between the current and previous study could be explained by the different type of cells used (pulp stem
19
20 cells versus pulp fibroblasts).
21
22
23
24

25 For WST-1 assays, only 0.125 mg/mL of both antibiotic mixtures was not toxic to DPSCs. This is
26
27 in agreement with previously reported concentrations of both DAP and TAP (0.1 mg/mL) that did not
28
29 have any toxic effects against stem cells of dental papilla (SCAP) [13]. In this study, DPSC were in direct
30
31 contact with the antibiotic medicament. The concentration used in our study (0.125-10 mg/mL) fall under
32
33 the concentrations reported previously for saturated solution of TAP and DAP (100, and 50 mg/mL,
34
35 respectively) [18]. Therefore, direct cytotoxicity from undissolved medicament is not expected. Our
36
37 results are in agreement with the results reported by Ruperal et al., [13] which demonstrate that the safest
38
39 concentration of DAP or TAP on stem cells is 0.1 mg/mL. Regardless of the different methods to detect
40
41 cytotoxicity (colorimetric assays vs cell counting), different mesenchymal stem cells (DPSC and SCAP)
42
43 seems to behave similarly in the presence of antibiotic medicaments compared to differentiated cells (i.e.,
44
45 dental pulp cells) which have been reported to withstand higher concentrations of TAP and DAP (2 and
46
47 0.3 mg/mL, respectively) [15].
48
49
50
51

52 In the current study, cytotoxicity was detected earlier using WST-1 assays compared to LDH
53
54 assays. This could be explained by the capability of the WST-1 to detect change in mitochondrial
55
56 metabolic activities inside the cells that may precede the cell membrane destruction detected by LDH
57
58 assays. Previous studies that used pulp fibroblasts or other types of cells have also reported earlier
59
60
61
62
63
64
65

1
2
3
4 detection of cytotoxicity using WST-1 assays compared to LDH assays [15, 33, 34]. Various cytotoxic
5
6 assays may give different cytotoxic results depending on the measured physiological endpoint. Therefore,
7
8 any clinical extrapolation of a detected cytotoxic concentration should be based on more than a single
9
10 cytotoxic assay.

11
12
13 According to both antibacterial and cytotoxicity assays, a concentration of 0.125 mg/mL of both
14
15 TAP and DAP was able to significantly reduce the bacterial biofilm without causing any cytotoxicity to
16
17 DPSC. The American Association of Endodontists (AAE) has recently recommended the use of antibiotic
18
19 medicaments in endodontic regeneration in concentrations ranging from 0.01 to 0.1 mg/mL [10, 35].
20
21 However, this concentration might not be effective against established bacterial biofilm. Additionally, no
22
23 clinical studies have reported the use of these recommended diluted concentrations of antibiotics.
24
25 Furthermore, a full concentration of TAP (1000 mg/mL) is still used in recently published clinical studies
26
27 [36, 37]. One of the challenges associated with the use of diluted antibiotic concentrations in immature
28
29 infected teeth is that they are in liquid form and do not have a pasty consistency. Incorporation of the
30
31 diluted antibiotics within a vehicle may be a good option to efficiently deliver the intended concentration
32
33 of antibiotics into the root canal. Another approach to deliver these low antibiotic concentrations into the
34
35 root canal could be through the use of a three-dimensional scaffold system. Recent studies have proposed
36
37 the use of a bio-resorbable antibiotic-loaded scaffold to disinfect the root canal space in endodontic
38
39 regeneration [38, 39]

40 41 42 43 44 **Conclusions**

45
46 Within the limitation of this *in vitro* study, only 0.125 mg/ml of TAP and DAP did not negatively
47
48 affect the viability of DPSC. However, this concentration was not enough to completely eradicated
49
50 established biofilm of *E. faecalis*. Antibiotic intracanal medicament with a concentration that can offer
51
52 antibiotic effect against endodontic pathogens with minimum adverse effect against pulp stem cells
53
54 should be considered during endodontic regeneration procedures.
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Conflict of interest

WSG is consulting medical director of Cook General BioTechnology, LLC. The other authors declare no potential conflicts of interest.

1
2
3
4 **References:**
5

- 6 1. Nosrat A, Seifi A, Asgary S (2011) Regenerative endodontic treatment (revascularization) for
7 necrotic immature permanent molars: a review and report of two cases with a new biomaterial.
8 J Endod 37: 562-567.
9
- 10 2. Thibodeau B, Trope M (2007) Pulp revascularization of a necrotic infected immature permanent
11 tooth: case report and review of the literature. Pediat Dent 29: 47-50.
- 12 3. Lenzi R, Trope M (2012) Revitalization procedures in two traumatized incisors with different
13 biological outcomes. J Endod 38: 411-414.
- 14 4. Murray PE, Garcia-Godoy F, Hargreaves KM (2007) Regenerative endodontics: a review of
15 current status and a call for action. J Endod 33: 377-390.
- 16 5. Lovelace TW, Henry MA, Hargreaves KM, Diogenes A (2011) Evaluation of the delivery of
17 mesenchymal stem cells into the root canal space of necrotic immature teeth after clinical
18 regenerative endodontic procedure. J Endod 37: 133-138.
- 19 6. Hoshino E, Kurihara-Ando N, Sato I, Uematsu H, Sato M, Kota K, et al. (1996) In-vitro
20 antibacterial susceptibility of bacteria taken from infected root dentine to a mixture of
21 ciprofloxacin, metronidazole and minocycline. Int Endod J 29: 125-130.
- 22 7. Sato I, Ando-Kurihara N, Kota K, Iwaku M, Hoshino E (1996) Sterilization of infected root-canal
23 dentine by topical application of a mixture of ciprofloxacin, metronidazole and minocycline in
24 situ. Int Endod J 29: 118-124.
- 25 8. Miller EK, Lee JY, Tawil PZ, Teixeira FB, Vann WF (2012) Emerging therapies for the management
26 of traumatized immature permanent incisors. Pedia Dent 34: 66-69.
- 27 9. Garcia-Godoy F, Murray PE (2012) Recommendations for using regenerative endodontic
28 procedures in permanent immature traumatized teeth. Dent Traumatol 28: 33-41.
- 29 10. Diogenes A, Henry MA, Teixeira FB, Hargreaves KM (2013) An update on clinical regenerative
30 endodontics. Endodontic Topics 28: 2-23.
- 31 11. Trope M (2010) Treatment of the immature tooth with a non-vital pulp and apical periodontitis.
32 Dent Clin North Am 54: 313-324.
- 33 12. Iwaya SI, Ikawa M, Kubota M (2001) Revascularization of an immature permanent tooth with
34 apical periodontitis and sinus tract. Dent Traumatol 17: 185-187.
- 35 13. Ruparel NB, Teixeira FB, Ferraz CC, Diogenes A (2012) Direct effect of intracanal medicaments
36 on survival of stem cells of the apical papilla. J Endod 38: 1372-1375.
- 37 14. Yadlapati M, Souza LC, Dorn S, Garlet GP, Letra A, Silva RM (2014) Deleterious effect of triple
38 antibiotic paste on human periodontal ligament fibroblasts. Int Endod J 47: 769-775.
- 39 15. Labban N, Yassen GH, Windsor LJ, Platt JA (2014) The direct cytotoxic effects of medicaments
40 used in endodontic regeneration on human dental pulp cells. Dent Traumatol 30: 429-434.
- 41 16. Phumpatrakom P, Srisuwan T (2014) Regenerative capacity of human dental pulp and apical
42 papilla cells after treatment with a 3-antibiotic mixture. J Endod 40: 399-405.
- 43 17. Althumairy RI, Teixeira FB, Diogenes A (2014) Effect of Dentin Conditioning with Intracanal
44 Medicaments on Survival of Stem Cells of Apical Papilla. J Endod 40: 521-525.
- 45 18. Sabrah AH, Yassen GH, Gregory RL (2013) Effectiveness of antibiotic medicaments against
46 biofilm formation of *Enterococcus faecalis* and *Porphyromonas gingivalis*. J Endod 39: 1385-
47 9138.
- 48 19. Chavez de Paz LE (2007) Redefining the persistent infection in root canals: possible role of
49 biofilm communities. J Endod 33: 652-662.
- 50 20. Stojcic S, Shen Y, Haapasalo M (2013) Effect of the source of biofilm bacteria, level of biofilm
51 maturation, and type of disinfecting agent on the susceptibility of biofilm bacteria to
52 antibacterial agents. J Endod 39: 473-477.
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
- 2
- 3
- 4 21. Perry BC, Zhou D, Wu X, Yang FC, Byers MA, Chu TM, et al. (2008) Collection, cryopreservation, and characterization of human dental pulp-derived mesenchymal stem cells for banking and clinical use. *Tissue Eng Part C: Methods* 14: 149-156.
- 5
- 6
- 7
- 8 22. Woods EJ, Perry BC, Hockema JJ, Larson L, Zhou D, Goebel WS (2009) Optimized cryopreservation method for human dental pulp-derived stem cells and their tissues of origin for banking and clinical use. *Cryobiology* 59: 150-157.
- 9
- 10
- 11 23. Stuart CH, Schwartz SA, Beeson TJ, Owatz CB (2006) *Enterococcus faecalis*: its role in root canal treatment failure and current concepts in retreatment. *J Endod* 32: 93-98.
- 12
- 13
- 14 24. Kayaoglu G, Orstavik D (2004) Virulence factors of *Enterococcus faecalis*: relationship to endodontic disease. *Crit Rev Oral Biol M* 15: 308-320.
- 15
- 16 25. Al-Ahmad A, Ameen H, Pelz K, Karygianni L, Wittmer A, Anderson AC, et al. (2014) Antibiotic resistance and capacity for biofilm formation of different bacteria isolated from endodontic infections associated with root-filled teeth. *J Endod* 40: 223-230.
- 17
- 18
- 19 26. Al-Ahmad A, Muller N, Wiedmann-Al-Ahmad M, Sava I, Hubner J, Follo M, et al. (2009) Endodontic and salivary isolates of *Enterococcus faecalis* integrate into biofilm from human salivary bacteria cultivated in vitro. *J Endod* 35: 986-991.
- 20
- 21
- 22 27. British Standards Institution(2006), Quantitative suspension test for the evaluation of basic bactericidal activity of chemical disinfectants and antiseptics: in test methods and requirements (phase 1). London, UK. EN 1040:2005.
- 23
- 24
- 25
- 26 28. Arias-Moliz MT, Ferrer-Luque CM, Espigares-Garcia M, Baca P (2009) *Enterococcus faecalis* biofilms eradication by root canal irrigants. *J Endod* 35: 711-714.
- 27
- 28
- 29 29. Athanassiadis B, Abbott PV, George N, Walsh LJ (2010) An in vitro study of the antimicrobial activity of some endodontic medicaments against *Enterococcus faecalis* biofilms. *Aust Dent J* 55: 150-155.
- 30
- 31
- 32 30. Kishen A, Haapasalo M (2010) Biofilm models and methods of biofilm assessment. *Endodontic Topics* 22: 58-78.
- 33
- 34
- 35 31. Barros J, Silva MG, Rocas IN, Goncalves LS, Alves FF, Lopes MA, Pina-Vaz I, Siqueira JF, Jr. (2014) Antibiofilm effects of endodontic sealers containing quaternary ammonium polyethylenimine nanoparticles. *J Endod* 40: 1167-1171.
- 36
- 37
- 38 32. George S, Kishen A (2007) Photophysical, photochemical, and photobiological characterization of methylene blue formulations for light-activated root canal disinfection. *J Biomed Opt* 12: 034029.
- 39
- 40
- 41 33. Kikkawa R, Yamamoto T, Fukushima T, Yamada H, Horii I (2005) Investigation of a hepatotoxicity screening system in primary cell cultures --"what biomarkers would need to be addressed to estimate toxicity in conventional and new approaches?". *J Toxicol Sci* 30: 61-72.
- 42
- 43
- 44 34. Huhtala A, Mannerstrom M, Alajuuma P, Nurmi S, Toimela T, Tahti H, et al. (2002) Comparison of an immortalized human corneal epithelial cell line and rabbit corneal epithelial cell culture in cytotoxicity testing. *J Ocul Pharmacol Ther* 18: 163-175.
- 45
- 46
- 47 35. American Association of Endodontists (2014) *AAE Clinical Considerations for a Regenerative Procedure*. USA.
- 48
- 49
- 50 https://www.aae.org/uploadedfiles/publications_and_research/research/currentregenerative_endodonticconsiderations.pdf. Accessed 5 August 2014.
- 51
- 52
- 53 36. Nagata JY, Gomes BP, Rocha Lima TF, Murakami LS, de Faria DE, Campos GR, et al. (2014) Traumatized immature teeth treated with 2 protocols of pulp revascularization. *J Endod* 40: 606-612.
- 54
- 55
- 56 37. Becerra P, Ricucci D, Loghin S, Gibbs JL, Lin LM (2014) Histologic study of a human immature permanent premolar with chronic apical abscess after revascularization/revitalization. *J Endod* 40: 133-139.
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

38. Bottino MC, Kamocki K, Yassen GH, Platt JA, Vail MM, Ehrlich Y, et al. (2013) Bioactive nanofibrous scaffolds for regenerative endodontics. *J Dent Res* 92: 963-969.

39. Bottino MC, Yassen GH, Platt JA, Labban N, Windsor LJ, Spolnik KJ, et al. (2013) A novel three-dimensional scaffold for regenerative endodontics: materials and biological characterizations. *J Tissue Eng Regen Med*, doi: 10.1002/term.1712.

1
2
3
4 Captions:
5

6 Figure 1: The mean log CFU/mL of *E. faecalis* after incubation with different TAP and DAP antibiotic
7 medicament dilutions for 72 h. The different groups are color coded in ascending order representing the
8 increase in the mean log CFU/mL. Different letters represent statistically significant differences.
9

10 Figure 2: The percent cytotoxicity to DPSC in the presence of different TAP and DAP antibiotic dilutions
11 for 72 h using LDH assay compared to the negative control set at 0% and the positive control set at 100%
12 cytotoxicity. * Significantly higher than the control. ** Significantly lower than the control.
13

14 Figure 3: The percent viability of DPSC in the presence of different TAP and DAP antibiotic dilutions for
15 72 h using WST-1 assay compared to a negative control set at 100% and a positive control set at 0%
16 viability. * Significantly higher than the control. ** Significantly lower than the control.
17
18

19 Figure 4: Images of DPSC in the presence of different TAP and DAP antibiotic dilutions for 72 h. (A)
20 control group. (B) The number of DPSC decreased in the presence of 0.25 mg/ml DAP. (C, D) The
21 decrease in number and change in shape of DPSC in the presence of 0.5 and 1 mg/ml DAP, respectively.
22 (E) Complete loss of viability and cell membrane integrity of DPSC in the presence of 10 mg/ml DAP.
23 (F) The number of DPSC decreased in the presence of 0.25 mg/ml TAP. (G, H) The decrease in number
24 and change in shape of DPSC in the presence of 0.5 and 1 mg/ml TAP, respectively. (I) Complete loss of
25 viability and cell membrane integrity of DPSC in the presence of 10 mg/ml TAP.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 1
[Click here to download high resolution image](#)




Figure 2

[Click here to download high resolution image](#)




Figure 3

[Click here to download high resolution image](#)




Figure 4A
[Click here to download high resolution image](#)




Figure 4B

[Click here to download high resolution image](#)




Figure 4C

[Click here to download high resolution image](#)




Figure 4D

[Click here to download high resolution image](#)




Figure 4E

[Click here to download high resolution image](#)




Figure 4F

[Click here to download high resolution image](#)




Figure 4G

[Click here to download high resolution image](#)




Figure 4H

[Click here to download high resolution image](#)




Figure 41
[Click here to download high resolution image](#)

