



Published in final edited form as:

*Clin Pharmacol Ther.* 2014 November ; 96(5): 542–548. doi:10.1038/clpt.2014.159.

## Clinical Pharmacogenetics Implementation Consortium (CPIC) Guidelines for *CYP2C9* and *HLA-B* Genotype and Phenytoin Dosing

Kelly E. Caudle<sup>1</sup>, Allan E. Rettie<sup>2</sup>, Michelle Whirl-Carrillo<sup>3</sup>, Lisa H. Smith<sup>4</sup>, Scott E. Mintzer<sup>5</sup>,  
Ming Ta Michael Lee<sup>6,7,8</sup>, Teri E. Klein<sup>3</sup>, and J. Thomas Callaghan<sup>9,10</sup>

<sup>1</sup>Department of Pharmaceutical Sciences, St. Jude Children's Research Hospital, Memphis, TN

<sup>2</sup>Department of Medicinal Chemistry, University of Washington School of Pharmacy, Seattle, Washington, USA

<sup>3</sup>Department of Genetics, Stanford University, Palo Alto, California

<sup>4</sup>Indiana University Department of Neurology, Division of Pediatric Neurology, Indianapolis, Indiana

<sup>5</sup>Thomas Jefferson University, Department of Neurology, Philadelphia, Pennsylvania

<sup>6</sup>Laboratory for International Alliance on Genomic Research, RIKEN Center for Integrative Medical Sciences, Yokohama, Japan

<sup>7</sup>National Center for Genome Medicine, Institute of Biomedical Sciences, Academia Sinica, Taipei, Taiwan

<sup>8</sup>School of Chinese Medicine, China Medical University, Taichung, Taiwan

<sup>9</sup>Division of Clinical Pharmacology, Department of Medicine, Indiana University School of Medicine, Indianapolis, IN

<sup>10</sup>Department of Veterans Affairs, RLR VA Medical Center, Indianapolis, IN

### Abstract

---

Corresponding Author: Kelly E. Caudle, Pharm.D., Ph.D., BCPS, CPIC Coordinator, Pharmaceutical Sciences Department, St. Jude Children's Research Hospital, 262 Danny Thomas Place MS: 313, Memphis, TN 38105, Office: 901-595-3994, Cell: 901-289-7392, Kelly.caudle@stjude.org, cpic@pharmgkb.org.

#### Conflicts of Interest

T.E.K. is a consultant for Personalis Inc. The other authors declared no conflict of interest.

#### Disclaimer

Clinical Pharmacogenetics Implementation Consortium (CPIC) guidelines reflect expert consensus based on clinical evidence and peer-reviewed literature available at the time they are written, and are intended only to assist clinicians in decision-making, as well as to identify questions for further research. New evidence may have emerged since the time a guideline was submitted for publication. Guidelines are limited in scope and are not applicable to interventions or diseases not specifically identified. Guidelines do not account for all individual variation among patients and cannot be considered inclusive of all proper methods of care or exclusive of other treatments. It remains the responsibility of the health care provider to determine the best course of treatment for the patient. Adherence to any guideline is voluntary, with the ultimate determination regarding its application to be solely made by the clinician and the patient. CPIC assumes no responsibility for any injury to persons or damage to property related to any use of CPIC's guidelines, or for any errors or omissions.

Phenytoin is a widely used antiepileptic drug with a narrow therapeutic index and large inter-patient variability partly due to genetic variations in *CYP2C9*. Furthermore, the variant allele *HLA-B\*15:02* is associated with an increased risk of Stevens-Johnson syndrome and toxic epidermal necrolysis in response to phenytoin treatment. We summarize evidence from the published literature supporting these associations and provide recommendations for the use of phenytoin based on *CYP2C9* and/or *HLA-B* genotype (also available on PharmGKB: [www.pharmgkb.org](http://www.pharmgkb.org)).

### Keywords

Phenytoin; HLA-B; CPIC; CYP2C9; pharmacogenetics; fosphenytoin; Stevens-Johnson syndrome; toxic epidermal necrolysis

---

### Introduction

The purpose of this guideline is to provide information for the interpretation of HLA-B and/or *CYP2C9* genotype tests so that the results can guide dosing and/or use of phenytoin. Detailed guidelines for use of phenytoin as well as analyses of cost effectiveness are out of scope. CPIC guidelines are periodically updated at <http://www.pharmgkb.org>.

### Focused Literature Review

A literature review focused on *CYP2C9* and *HLA-B\*15:02* genotype and phenytoin use (see Supplemental Material online) was conducted. Reviews were included to summarize the available literature.

### Genes: *HLA-B* and *CYP2C9*

#### Background

In this guideline, human leukocyte antigen B (*HLA-B*) will be discussed as it relates to phenytoin-induced cutaneous adverse drug reactions (ADRs) of Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN) and hepatic *CYP2C9* and its alleles are discussed as they relate to phenytoin metabolism and dosing.

**HLA-B**—*HLA-B* is part of a gene cluster designated as the human major histocompatibility complex (MHC) located on the short arm of chromosome 6. The cluster contains three classes (I, II and III). MHC class I contains three genes: *HLA-B* plus *HLA-A* and *HLA-C*. *HLA-B* encodes a cell surface protein that binds peptides generated by proteolysis and extruded from proteasomes. The presentation of these peptides on the cell surface enables the immune system to distinguish self-proteins from foreign proteins typically introduced by infectious organisms (*e.g.*, viruses and bacteria) (see supplemental material for further discussion).

*HLA* genes, and specifically *HLA-B*, are among the most highly polymorphic genes in the human genome. *HLA* polymorphisms were previously ascertained serologically, but genotyping and DNA sequencing methods reveal much greater genetic complexity. More

than 2,000 *HLA-B* alleles, many of which differ by more than one nucleotide from each other, were deposited to the World Health Organization (WHO) Nomenclature Committee for Factors of the HLA System (<http://hla.alleles.org>). Each allele is designated by the gene name followed by an asterisk and up to an eight-digit (four pairs) identifier giving information about the allele type (designated by the first two digits) and specific protein subtypes (second set of digits). For more information and a diagram outlining the description of the current HLA allele nomenclature see <http://hla.alleles.org/nomenclature/naming.html>. The details of HLA nomenclature are also described in a previous CPIC guideline (1). This guideline specifically discusses only the *HLA-B\*15:02* allele as it relates to the phenytoin-induced cutaneous adverse drug reactions of SJS and TEN.

**CYP2C9**—Hepatic CYP2C9 enzyme contributes to the metabolism of many clinically relevant drugs, including phenytoin (<http://www.pharmgkb.org/pathway/PA145011115>). The *CYP2C9* gene is highly polymorphic, having more than 50 known variant alleles (<http://www.cypalleles.ki.se/cyp2c9.htm>, Supplemental Table S1 and S2). Individuals homozygous for the reference *CYP2C9* allele (*CYP2C9\*1*) have the “normal metabolizer” phenotype. Each named *CYP2C9* star (\*) allele is defined by a genotype at one or more specific single-nucleotide polymorphisms (SNPs) with variable enzyme activity. The two most common variants with reduced enzyme activity in Europeans are *CYP2C9\*2* (rs1799853) and *CYP2C9\*3* (rs1057910) (2).

### Genetic Test Interpretation

**HLA-B**—Clinical genotyping test results for *HLA-B\*15:02* are interpreted as “positive” if one or two copies of *HLA-B\*15:02* are present or “negative” if no copies of *HLA-B\*15:02* are present. Phenotype assignments for *HLA-B\*15:02* genotypes are summarized in Table 1. The allele frequencies of *HLA-B* vary greatly among populations. Specifically, *HLA-B\*15:02* is most prevalent in Oceania, East Asian and South/Central Asian populations ranging from 1% to over 10%. It is less frequent in European populations (0–1%) and apparently absent in several African populations (Supplemental Table S3 and S4). The global average derived from over 46,000 individuals is 1.37%.

**CYP2C9**—Most clinical laboratories reporting *CYP2C9* genotype use the star (\*) allele nomenclature and may interpret the patient’s predicted metabolizer phenotype (Table 1, Supplemental Table S1). The combination of alleles is used to determine a patient’s diplotype. Not all *CYP2C9* allelic variants may be tested, influencing the accuracy of the genotype-based dose prediction, primarily in individuals of Asian or African ancestry who carry other common functionally decreased activity *CYP2C9* variant alleles (Supplemental Table S5). The frequencies of the *CYP2C9\*2* and *\*3* alleles and diplotypes derived from these and other alleles differ between racial/ethnic groups (Supplementary Table S5, S6 and S7) (2). *CYP2C9* alleles are typically characterized as wild-type (normal function) or decreased function depending on the reported activity of the enzyme for which they encode.

### Available Genetic Test Options

Several methods of *CYP2C9* and *HLA-B* genotyping are commercially available. The Supplemental Material online and at [www.pharmgkb.org](http://www.pharmgkb.org) contains more information on available clinical testing options.

### Incidental findings

*HLA-B* alleles are associated with hypersensitivity reactions to other drugs. CPIC guidelines are available for *HLA-B\*57:01* and abacavir-induced hypersensitivity reactions, *HLA-B\*58:01* and allopurinol-induced severe cutaneous adverse reactions, and *HLA-B\*15:02* and carbamazepine-induced SJS and TEN (1, 3, 4).

No diseases have been linked to genetic variations in *CYP2C9*, except for a small study that linked *CYP2C9\*2* and *\*3* variants and phenytoin to a higher frequency of cerebellar atrophy (5). *CYP2C9* poor metabolizers may be predisposed to serious bleeding during warfarin therapy (6).

### Other Considerations

Not applicable

### Drugs: Phenytoin and Fosphenytoin

Phenytoin and its prodrug fosphenytoin are one of the mainstays of treatment for both focal and generalized convulsive status epilepticus. Dosing is complex owing to its highly unusual pharmacokinetics and requiring adjustments be made in line with patient weight, sex, and age. Outpatient therapy is generally initiated at 5–7 mg/kg/day in adults (slightly higher in children) and may be given once daily (or twice daily in children). Starting dose must be lower in the setting of hepatic impairment. Careful dose adjustments must then be made -- generally 30 – 40 mg at a time in 2-week intervals in adults -- to stabilize the level within the typical therapeutic range (10 – 20 ug/dL). In urgent situations such as status epilepticus, intravenous loading doses of 15–20 mg/kg are given, followed by maintenance doses, IV or oral, as above. Acute dose-related side effects include sedation, ataxia, dizziness, nystagmus, nausea, and cognitive impairment. The drug is highly allergenic, and rashes ranging from mild eruptions to life-threatening hypersensitivity reactions may be seen. *HLA-B\*15:02* is associated with the phenytoin-induced SJS and TEN. SJS is characterized by epidermal detachment involving up to 10% of body surface area (BSA) while TEN usually affects more than 30% of the BSA. Subacutely, hematologic and hepatic toxicity can occur; the latter is likely a hypersensitivity reaction itself, as it is usually accompanied by rash (7), while the former may consist of leukopenia or pancytopenia.

Because of the acute dose-related side effects, initial maintenance dose selection is important. Higher plasma concentrations increase the probability of these toxicities. However, non-linear saturable pharmacokinetics, auto-inductive effects with maintenance dosing, and *CYP2C9* pharmacogenetic polymorphisms complicate dose-selection. The *CYP2C9* poor metabolizer phenotype and *CYP2C9* drug interactions, such as produced by voriconazole, can significantly augment phenytoin exposure (8). Variability in protein

binding, primarily related to changes in albumin concentrations, can confound the relationship of therapeutic drug monitoring and pharmacodynamic expectations. Further discussion of the metabolism of phenytoin and fosphenytoin can be found in the supplemental material.

### Linking genetic variability to variability in drug-related phenotypes

Substantial evidence links *CYP2C9* and *HLA-B\*15:02* genotype with phenotypic variability (see Supplemental Tables S8 and S9). Application of a grading system to evidence linking genotypic to phenotypic variability indicates a high quality of evidence in the majority of cases (Supplemental Tables S8 and S9). The evidence presented here and in Supplemental Tables S8 and S9 provides the basis for the dosing recommendations in Table 2.

**HLA-B**—An increased risk of SJS/TEN has been associated with the *HLA-B\*15:02* allele in Han Chinese and other Asian groups (see Supplemental Material; Supplemental Table S8). Cheung et al conducted a meta-analysis of two studies in Taiwan (9) and Hong Kong (10), comprising in total 41 cases and 188 controls and showed a positive association of *HLA-B\*15:02* with phenytoin-induced SJS/TEN ( $p < 3 \times 10^{-4}$ ; OR 4.26; 95% CI 1.93–9.39) under a fixed-effect model with statistically insignificant heterogeneity. By pooling data directly, the association had a sensitivity of 36.6% (95% CI 23.6–51.9) and specificity of 87.2% (95% CI 81.7–91.3). Therefore, the absence of these variants does not rule out the possibility of a patient developing phenytoin-induced SJS/TEN. The strength of the association between phenytoin use and SJS/TEN is weaker than the association between carbamazepine use and SJS/TEN due to the limited number of studies and observations with phenytoin or fosphenytoin in the literature. However, taken together with the known association of carbamazepine and SJS/TEN in carriers of *HLA-B\*15:02* the association supports the FDA recommendations to avoid these agents as substitutes for carbamazepine in individuals who test positive for *HLA-B\*15:02* (4).

**CYP2C9**—Available model estimates predict that variant *CYP2C9* alleles lower phenytoin intrinsic clearance based on the allele and number of variant. Several studies indicate that individuals with *CYP2C9\*1/\*3* and *CYP2C9\*1/\*2* genotypes have mild-to-moderately reduced clearance values (Supplemental Table S9) and are classified as Intermediate Metabolizers. Individuals with two decreased activity alleles (*CYP2C9\*2/\*2*, *CYP2C9\*3/\*3*) have reduced clearance of several drugs and are classified as CYP2C9 poor metabolizers. Phenytoin maintenance doses were reported to be reduced 23–38% in heterozygous individuals with one decreased activity allele (11–13) and 31–52% for carriers with two decreased activity *CYP2C9* alleles versus *CYP2C9\*1/\*1* (12, 13). Furthermore, case reports indicate that poor metabolizers appear to be at higher risk for exposure-related toxicities than patients homozygous for the wild-type alleles (14–17).

### Therapeutic Recommendations

**HLA-B\*15:02 recommendations**—The FDA warning for phenytoin states, “Consideration should be given to avoiding phenytoin as an alternative for carbamazepine in patients positive for *HLA-B\*15:02*’ due to the increased risk of SJS/TEN in patients of Asian ancestry.” The evidence linking *HLA-B\*15:02* to phenytoin-induced SJS/TEN was

generated in individuals of Asian ancestry as the frequency of *HLA-B\*15:02* is very low in other populations (see Supplemental Table S3 and S4 for frequency information) that have been studied. However, it may also occur in other populations throughout the world yet to be studied and patients may be unaware of or fail to disclose more distant Asian ancestry in their families. Furthermore, much of the evidence (summarized in Supplemental Table S8) linking *HLA-B\*15:02* to phenytoin-induced SJS/TEN was generated in both children and adults. Therefore, regardless of the *CYP2C9* genotype and individual's ancestry or age, if the *HLA-B\*15:02* test result is "positive", the recommendation is to consider using an anticonvulsant other than carbamazepine and phenytoin unless the benefits of treating the underlying disease clearly outweigh the risks (see Table 2). Alternative medications such as oxcarbazepine, eslicarbazepine acetate, and lamotrigine have some evidence linking SJS/TEN with the *HLA-B\*15:02* allele and thus caution should be used in choosing alternatives to phenytoin (see Supplement for details).

**CYP2C9 recommendations**—Phenytoin and fosphenytoin dose should first be adjusted according to a patient's clinical characteristics. Table 2 summarizes the gene-based dosing recommendations for phenytoin based on *CYP2C9* phenotype. The recommended phenytoin maintenance dose does not need adjustment based on genotype for *CYP2C9* extensive metabolizers. Available evidence does not clearly indicate the amount of dose reduction needed to prevent phenytoin-related toxicities in *CYP2C9* intermediate and poor metabolizers; thus, our recommendation should be considered conservative estimates given the variability surrounding phenytoin dosing to an individual. Based on the doses reported in the pharmacokinetic and pharmacogenetic studies mentioned above (11–13) and in Supplemental Table S9, at least a 25% reduction of the recommended starting maintenance dose may be considered for *CYP2C9* intermediate metabolizers with subsequent maintenance doses adjusted based on therapeutic drug monitoring and response. For *CYP2C9* poor metabolizers, consider at least a 50% reduction of starting maintenance dose with subsequent maintenance doses adjusted based on therapeutic drug monitoring or response.

Furthermore, while *in vitro* data suggest that the degree of reduction of catalytic activity is greater for the *CYP2C9\*3* variant than for the *CYP2C9\*2* variant (18), clinical pharmacokinetic studies indicate similar dose reductions and pharmacokinetic parameters (e.g., trough levels, serum p-HPPH/P ratio) for these variants as compared to wild-type (12, 19, 20). Thus, our recommendation is to start with at least the above recommended reduction of the maintenance dose followed by an adjustment of dose based on therapeutic drug monitoring.

**Pediatrics**—Special consideration should be taken with the pediatric population. Phenytoin is used in the treatment of neonatal seizures and subsequently after discharge from the neonatal ICU. Maintaining therapeutic levels can be particularly problematic in this population. This may be due to the developmental expression of hepatic *CYP2C*. P450 expression and functional activities have been shown to develop at different rates within subfamilies (21). It has been found that activity levels of *CYP2C9* are at 1 to 2% of adult values in the fetus during the first trimester. These levels gradually increase to 30% of adult



values at term. There is a high variability in these levels during the first five months of life, eventually approach adult values somewhere between five months to 2 years of age (22). Other considerations include the clearance of phenytoin being twice that of adult values in children under 6 years of age. This is attributed to the finding that the maximal rate of phenytoin metabolism is inversely related to age. However, this varied significantly within age subgroups (23). For these reasons, phenytoin therapeutic recommendations based on *CYP2C9* genotype in this population are difficult. There is only one published report describing a two-year old patient (*CYP2C9*\*2/\*2 and *CYP2C19*\*1/\*4) presenting with phenytoin toxicity two hours after a 15mg/kg phenytoin loading dose with symptoms lasting 122 hours (24). The half-life was much higher than expected (112 hours versus 46.7) which could be explained by the influence of *CYP2C9* and *CYP2C19* genetic polymorphisms (other predisposing factors such as malnourishment, renal failure, hepatic dysfunction, and inhibition of phenytoin metabolism by other drugs were ruled out). Therefore, for pediatric patients who are *CYP2C9* intermediate or poor metabolizers dose adjustment is recommended with close therapeutic drug monitoring.

**HLA-B\*15:02 and CYP2C9 dosing recommendation**—If both *HLA-B\*15:02* and *CYP2C9* genotype are known, consider the *HLA-B\*15:02* genotype first, then the *CYP2C9* genotype (Figure 1; Table 2).

#### Other considerations

**HLA-B**—*HLA-B\*15:02* is linked to SJS and TEN but not to a predisposition for other phenytoin-induced cutaneous adverse reactions such as mild maculopapular eruptions (MPE) or drug hypersensitivity syndrome (HSS) (25).

**CYP2C9**—Because of its potent CYP450-inducing properties, phenytoin is involved in a very large number of drug interactions, especially involving increased metabolism of other agents with subsequent decrease in their levels (26). A full discussion of these is beyond the scope of this guideline, but agents prominently and significantly affected include antineoplastic and immunosuppressive agents, lipid-lowering agents, psychotropics, oral contraceptives, and warfarin, just to name a few. Furthermore, inhibitors of *CYP2C9* can generate phenytoin overexposure and toxicity. Although fluconazole and amiodarone are recognized as potent *CYP2C9* enzyme inhibitors, other less potent drugs can produce significant elevations in phenytoin plasma concentrations. Therefore, it is important to interpret the results of genetic testing in the context of other co-administered drugs.

*CYP2C9* genetic variation does not account for all of the pharmacogenetic variability in phenytoin metabolism. Some studies have implicated variants in other genes associated with phenytoin metabolism (e.g., *CPY2C19*, *CYP1A1*, and *EPHX1*; see (27) for a review) and combined genetic analysis might improve the predictability of *CYP2C9* alone (11, 13). However, this has not been consistently replicated and there are limited studies evaluating the effect of multiple gene variation and phenytoin dose adjustment requirement. Consequently, this guideline on genotype-directed phenytoin dosing is limited to *CYP2C9* variant alleles.

## Recommendations for Incidental Findings

Several drugs structurally and therapeutically similar to phenytoin, such as oxcarbazepine and carbamazepine, have also been associated with SJS/TEN and *HLA-B\*15:02* in Asian populations (28–34) (see Supplemental Material). The drug-specific evidence linking *HLA-B\*15:02* and SJS/TEN is discussed in the CPIC Guideline for HLA-B genotype and carbamazepine dosing (4) and may have implications for choosing alternatives to phenytoin in those who carry the *HLA-B\*15:02* allele. Case reports have identified cross-reactions to lamotrigine and other anti-epileptic drugs in the presence of *HLA-B\*15:02* (see Supplemental Material for further discussion). However, larger studies appear to be needed for confirmation.

CYP2C9 metabolism includes substrates from several drug classes, including NSAIDs, oral hypoglycemics/sulfonylureas, as well as a miscellaneous group of drugs. Reports support that patients with enhanced sensitivity to warfarin are likely to have a decreased capacity to metabolize phenytoin (35).

## Potential Benefits and Risks for the Patient

The potential benefit for patients with existing *CYP2C9* and/or *HLA-B\*15:02* genotyping information is in avoiding adverse effects in those patients who are *CYP2C9* poor metabolizers by making significant reductions in their starting maintenance dose or by selecting alternative agents for those who are *HLA-B\*15:02* carriers. For *HLA-B\*15:02* carriers, a potential risk is that phenytoin therapy may have been needlessly avoided in patients who may not have developed SJS/TEN; however, this risk is mitigated because alternatives to phenytoin with comparable effectiveness exist. Another potential risk would be an error in genotyping. Also, many commercially available genotyping tests do not detect alleles that are rare or *de novo* variants. Other alleles are not well characterized, resulting in uncertainty when predicting the phenotype for some genetic test results. Due to the high rate of *HLA-B\*15:02* false-negative results (36) or in the event of a rare variant not detected on the genetic test, a high-risk patient could be prescribed phenytoin or prescribed a higher dose than needed. Moreover, because not all phenytoin-induced adverse events are attributed to *HLA-B\*15:02* or *CYP2C9* metabolizer status, clinicians should carefully monitor all patients according to standard practices.

## Caveats: Appropriate Use and/or Potential Misuse of Genetic Tests

The application of genotype-based dosing is most appropriate when initiating phenytoin therapy. Obtaining genetic information after months of drug therapy is less helpful, given that the drug dose may have already been adjusted based on plasma concentrations, response, or side effects. As with all diagnostic tests, genetic tests are only one of several pieces of clinical information that should be considered before initiating drug therapy.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.




## Acknowledgments

We acknowledge the critical input of members of CPIC of the Pharmacogenomics Research Network (PGRN) particularly Dr. Mary V. Relling (St Jude Children's Research Hospital) funded by the National Institutes of Health (NIH). This work is funded by NIH grants R24 GM61374, U01 GM092666, GM32165, UO1 GM092676, and U01 HL0105918.

## References

1. Martin MA, Klein TE, Dong BJ, Pirmohamed M, Haas DW, Kroetz DL. Clinical pharmacogenetics implementation consortium guidelines for HLA-B genotype and abacavir dosing. *Clinical pharmacology and therapeutics*. 2012; 91:734–8. [PubMed: 22378157]
2. Lee CR, Goldstein JA, Pieper JA. Cytochrome P450 2C9 polymorphisms: a comprehensive review of the in-vitro and human data. *Pharmacogenetics*. 2002; 12:251–63. [PubMed: 11927841]
3. Hershfield MS, et al. Clinical Pharmacogenetics Implementation Consortium Guidelines for Human Leukocyte Antigen-B Genotype and Allopurinol Dosing. *Clinical pharmacology and therapeutics*. 2012
4. Leckband SG, et al. Clinical Pharmacogenetics Implementation Consortium guidelines for HLA-B genotype and carbamazepine dosing. *Clinical pharmacology and therapeutics*. 2013; 94:324–8. [PubMed: 23695185]
5. Twardowschy CA, Werneck LC, Scola RH, Borgio JG, De Paola L, Silvado C. The role of CYP2C9 polymorphisms in phenytoin-related cerebellar atrophy. *Seizure*. 2013; 22:194–7. [PubMed: 23298603]
6. Johnson JA, et al. Clinical Pharmacogenetics Implementation Consortium Guidelines for CYP2C9 and VKORC1 genotypes and warfarin dosing. *Clinical pharmacology and therapeutics*. 2011; 90:625–9. [PubMed: 21900891]
7. Parker WA, Shearer CA. Phenytoin hepatotoxicity: a case report and review. *Neurology*. 1979; 29:175–8. [PubMed: 571061]
8. Purkins L, Wood N, Ghahramani P, Love ER, Eve MD, Fielding A. Coadministration of voriconazole and phenytoin: pharmacokinetic interaction, safety, and toleration. *British journal of clinical pharmacology*. 2003; 56 (Suppl 1):37–44. [PubMed: 14616412]
9. Hung SI, et al. Common risk allele in aromatic antiepileptic-drug induced Stevens-Johnson syndrome and toxic epidermal necrolysis in Han Chinese. *Pharmacogenomics*. 2010; 11:349–56. [PubMed: 20235791]
10. Cheung YK, Cheng SH, Chan EJ, Lo SV, Ng MH, Kwan P. HLA-B alleles associated with severe cutaneous reactions to antiepileptic drugs in Han Chinese. *Epilepsia*. 2013
11. Hung CC, Lin CJ, Chen CC, Chang CJ, Liou HH. Dosage recommendation of phenytoin for patients with epilepsy with different CYP2C9/CYP2C19 polymorphisms. *Therapeutic drug monitoring*. 2004; 26:534–40. [PubMed: 15385837]
12. van der Weide J, Steijns LS, van Weelden MJ, de Haan K. The effect of genetic polymorphism of cytochrome P450 CYP2C9 on phenytoin dose requirement. *Pharmacogenetics*. 2001; 11:287–91. [PubMed: 11434505]
13. Hung CC, et al. Effects of polymorphisms in six candidate genes on phenytoin maintenance therapy in Han Chinese patients. *Pharmacogenomics*. 2012; 13:1339–49. [PubMed: 22966884]
14. Brandolese R, Scordo MG, Spina E, Gusella M, Padrini R. Severe phenytoin intoxication in a subject homozygous for CYP2C9\*3. *Clinical pharmacology and therapeutics*. 2001; 70:391–4. [PubMed: 11673755]
15. Dorado P, Lopez-Torres E, Penas-Lledo EM, Martinez-Anton J, Llerena A. Neurological toxicity after phenytoin infusion in a pediatric patient with epilepsy: influence of CYP2C9, CYP2C19 and ABCB1 genetic polymorphisms. *The pharmacogenomics journal*. 2012
16. Hennessy S, et al. CYP2C9, CYP2C19, and ABCB1 genotype and hospitalization for phenytoin toxicity. *Journal of clinical pharmacology*. 2009; 49:1483–7. [PubMed: 19617466]

17. Ramasamy K, Narayan SK, Chanolean S, Chandrasekaran A. Severe phenytoin toxicity in a CYP2C9\*3\*3 homozygous mutant from India. *Neurology India*. 2007; 55:408–9. [PubMed: 18040121]
18. Rettie AE, Haining RL, Bajpai M, Levy RH. A common genetic basis for idiosyncratic toxicity of warfarin and phenytoin. *Epilepsy research*. 1999; 35:253–5. [PubMed: 10413320]
19. Aynacioglu AS, et al. Frequency of cytochrome P450 CYP2C9 variants in a Turkish population and functional relevance for phenytoin. *British journal of clinical pharmacology*. 1999; 48:409–15. [PubMed: 10510154]
20. Kerb R, et al. The predictive value of MDR1, CYP2C9, and CYP2C19 polymorphisms for phenytoin plasma levels. *The pharmacogenomics journal*. 2001; 1:204–10. [PubMed: 11908757]
21. Cresteil T, Beaune P, Kremers P, Celier C, Guengerich FP, Leroux JP. Immunoquantification of epoxide hydrolase and cytochrome P-450 isozymes in fetal and adult human liver microsomes. *European journal of biochemistry / FEBS*. 1985; 151:345–50. [PubMed: 2411555]
22. Koukouritaki SB, et al. Developmental expression of human hepatic CYP2C9 and CYP2C19. *The Journal of pharmacology and experimental therapeutics*. 2004; 308:965–74. [PubMed: 14634042]
23. Suzuki Y, Mimaki T, Cox S, Koepke J, Hayes J, Walson PD. Phenytoin age-dose-concentration relationship in children. *Therapeutic drug monitoring*. 1994; 16:145–50. [PubMed: 8009561]
24. Dorado P, Lopez-Torres E, Penas-Lledo EM, Martinez-Anton J, Llerena A. Neurological toxicity after phenytoin infusion in a pediatric patient with epilepsy: influence of CYP2C9, CYP2C19 and ABCB1 genetic polymorphisms. *The pharmacogenomics journal*. 2013; 13:359–61. [PubMed: 22641027]
25. Yip VL, Marson AG, Jorgensen AL, Pirmohamed M, Alfirevic A. HLA Genotype and Carbamazepine-Induced Cutaneous Adverse Drug Reactions: A Systematic Review. *Clinical pharmacology and therapeutics*. 2012; 92:757–65. [PubMed: 23132554]
26. Mintzer S, et al. Effects of antiepileptic drugs on lipids, homocysteine, and C-reactive protein. *Annals of neurology*. 2009; 65:448–56. [PubMed: 19296463]
27. Thorn CF, Whirl-Carrillo M, Leeder JS, Klein TE, Altman RB. PharmGKB summary: phenytoin pathway. *Pharmacogenetics and genomics*. 2012; 22:466–70. [PubMed: 22569204]
28. Shankarkumar U, Shah KN, Ghosh K. Letter: HLA B\*1502 allele association with oxcarbazepine-induced skin reactions in epilepsy patient from India. *Epilepsia*. 2009; 50:1837–8. [PubMed: 20831525]
29. Lin LC, Lai PC, Yang SF, Yang RC. Oxcarbazepine-induced Stevens-Johnson syndrome: a case report. *Kaohsiung J Med Sci*. 2009; 25:82–6. [PubMed: 19321411]
30. Chen YC, Chu CY, Hsiao CH. Oxcarbazepine-induced Stevens-Johnson syndrome in a patient with HLA-B\*1502 genotype. *J Eur Acad Dermatol Venereol*. 2009; 23:702–3. [PubMed: 18785891]
31. Lochareonkul C, et al. Carbamazepine and phenytoin induced Stevens-Johnson syndrome is associated with HLA-B\*1502 allele in Thai population. *Epilepsia*. 2008; 49:2087–91. [PubMed: 18637831]
32. Whirl-Carrillo M, et al. Pharmacogenomics knowledge for personalized medicine. *Clinical pharmacology and therapeutics*. 2012; 92:414–7. [PubMed: 22992668]
33. Hu FY, Wu XT, An DM, Yan B, Stefan H, Zhou D. Pilot association study of oxcarbazepine-induced mild cutaneous adverse reactions with HLA-B\*1502 allele in Chinese Han population. *Seizure*. 2011; 20:160–2. [PubMed: 21169036]
34. Hu FY, Wu XT, An DM, Yan B, Stefan H, Zhou D. Phenytoin-induced Stevens-Johnson syndrome with negative HLA-B\*1502 allele in mainland China: two cases. *Seizure*. 2011; 20:431–2. [PubMed: 21334226]
35. Bochner F, Hooper WD, Eadie MJ, Tyrer JH. Decreased capacity to metabolize diphenylhydantoin in a patient with hypersensitivity to warfarin. *Australian and New Zealand journal of medicine*. 1975; 5:462–6. [PubMed: 1061551]
36. Dong D, Sung C, Finkelstein EA. Cost-effectiveness of HLA-B\*1502 genotyping in adult patients with newly diagnosed epilepsy in Singapore. *Neurology*. 2012; 79:1259–67. [PubMed: 22955130]



**Figure 1. Algorithm for suggested clinical actions based *HLA-B\*15:02* and *CYP2C9* genotype**  
EM, extensive metabolizer; IM, intermediate metabolizer; PM, poor metabolizer

<sup>a</sup>If patient has previously used phenytoin for longer than 3 months without incidence of cutaneous adverse reactions, reinitiate phenytoin with caution. Adjust dose based on *CYP2C9* genotype if known.

<sup>b</sup>Carbamazepine should not be used as an alternative (4). Alternative medications such as oxcarbazepine, eslicarbazepine acetate, and lamotrigine have some evidence linking SJS/TEN with the *HLA-B\*15:02* allele and thus caution should be used in choosing alternatives to phenytoin (see Supplement for details).

<sup>c</sup>Recommended maintenance dose based on patient's clinical characteristics

Table 1

## Assignment of likely phenotype based on genotypes

Assignment of likely CYP2C9 phenotypes based on genotypes		
Likely Phenotype <sup>a</sup>	Genotypes	Examples of diplotypes
Extensive metabolizer (normal activity) (constitutes ~ 91% of patients)	An individual carrying two normal activity alleles	*1/*1
Intermediate metabolizer <sup>c</sup> (heterozygote or intermediate activity) (constitutes ~ 8% of patients)	An individual carrying one normal activity allele plus one decreased function allele	*1/*3, *1/*2
Poor metabolizer (homozygous variant, mutant, low, or deficient activity) (constitutes ~ 1% of patients)	An individual carrying two decreased function alleles	*2/*2, *3/*3, *2/*3

Assignment of likely HLA-B phenotypes based on genotypes		
Likely Phenotype <sup>b</sup>	Genotype	Examples of diplotypes
Homozygous for an allele other than *15:02; at "normal" or reduced risk of phenytoin-associated cutaneous adverse reactions (constitutes ~ 98.6% of patients)	<b>HLA-B*15:02 non-carrier.</b> No *15:02 alleles reported, often reported as "negative" on a genotyping test.	*X*X <sup>d</sup>
Heterozygote or homozygous variant; at significantly increased risk of phenytoin-associated cutaneous adverse reactions (constitutes ~ 1.4% of patients)	<b>HLA-B*15:02 Carrier.</b> One or two *15:02 alleles, often reported as "positive" on a genotyping test.	*15:02*X <sup>d</sup> , *15:02/*15:02

<sup>a,b</sup> Global frequencies presented in the parenthesis. Haplotype frequencies vary among populations, please see details for individual population frequencies in <sup>a</sup>Supplemental Table S5, S6 and S7 for *CYP2C9*\*2 and \*3, and <sup>b</sup>Supplemental Table S3 and S4 for *HLA-B*\*15:02.

<sup>c</sup>The enzyme activity in this grouping varies widely. See Supplemental Table S2 for activity ranges.

<sup>d</sup>Where \*X = any genotype other than \*15:02

Table 2

Recommended dosing of phenytoin/fosphenytoin based on *HLA-B\*15:02* and *CYP2C9* phenotype/genotype.

Phenotype/ Genotype	<i>HLA-B*15:02</i> carrier			<i>HLA-B*15:02</i> non-carrier		
	Implication	Therapeutic Recommendation	Classification of Recommendation <sup>d</sup>	Implication	Therapeutic Recommendation	Classification of Recommendation <sup>d</sup>
<b>CYP2C9 Extensive metabolizer</b>	Increased risk of phenytoin-induced SJS/TEN	If patient is phenytoin-naïve <sup>b</sup> , do not use phenytoin/fosphenytoin <sup>c</sup>	Strong	Normal phenytoin metabolism	Initiate therapy with recommended maintenance dose. <sup>d</sup>	Strong
<b>CYP2C9 Intermediate metabolizer</b>	Increased risk of phenytoin-induced SJS/TEN	If patient is phenytoin-naïve <sup>b</sup> , do not use phenytoin/fosphenytoin <sup>c</sup>	Strong	Reduced phenytoin metabolism Higher plasma concentrations will increase probability of toxicities	Consider 25% reduction of recommended starting maintenance dose. <sup>d</sup> . Subsequent maintenance doses should be adjusted according to therapeutic drug monitoring and response.	Moderate
<b>CYP2C9 Poor metabolizer</b>	Increased risk of phenytoin-induced SJS/TEN	If patient is phenytoin-naïve <sup>b</sup> , do not use phenytoin/fosphenytoin <sup>c</sup>	Strong	Reduced phenytoin metabolism Higher plasma concentrations will increase probability of toxicities.	Consider 50% reduction of recommended starting maintenance dose. <sup>d</sup> . Subsequent maintenance doses should be adjusted according to therapeutic drug monitoring and response.	Strong

<sup>a</sup>Rating scheme described in the Supplemental Material

<sup>b</sup>If patient has previously used phenytoin for longer than 3 months without incidence of cutaneous adverse reactions, reinitiate phenytoin with caution. Adjust dose based on *CYP2C9* genotype if known.

<sup>c</sup>Carbamazepine should not be used as an alternative (4). Alternative medications such as oxcarbazepine, eslicarbazepine acetate, and lamotrigine have some evidence linking SJS/TEN with the *HLA-B\*15:02* allele and thus caution should be used in choosing alternatives to phenytoin (see Supplement for details).

<sup>d</sup>Recommended maintenance dose based on patient's clinical characteristics