

## An Investigation of Whether Vitamin E Preferentially Interacts with Polyunsaturated Lipids

**Andres Cavazos**<sup>1</sup>, Jacob J. Kinnun<sup>1</sup>, Justin A. Williams<sup>1</sup>, Morris Bank<sup>1</sup>, Bruce D. Ray<sup>1</sup>, Paul E. Harper<sup>2</sup>, Jeffrey Atkinson<sup>3</sup>, Horia I. Petrache<sup>1</sup>, and Stephen R. Wassall<sup>1</sup>.

<sup>1</sup>Department of Physics, Indiana University-Purdue University-Indianapolis, Indianapolis, IN, 46202,

<sup>2</sup>Department of Physics and Astronomy, Calvin College, Grand Rapids, MI 49548, <sup>3</sup>Department of Chemistry, Brock University, St. Catharines, Ontario L2S 3A1, Canada

Vitamin E ( $\alpha$ -tocopherol) is a lipid-soluble antioxidant that has the role of protecting phospholipids from oxidation in membranes. A question that remains is how the low concentration of  $\alpha$ -tocopherol found in whole cells can protect the relatively large concentration of polyunsaturated phospholipids found in membranes that are particularly vulnerable to oxidative attack. We hypothesize that  $\alpha$ -tocopherol co-localizes with polyunsaturated phospholipids to optimize its role as an antioxidant. This project attempts to test this hypothesis by comparing the effect of  $\alpha$ -tocopherol on the molecular organization of 1-palmitoyl-2-docosahexaenoyl-sn-glycerophosphatidylethanolamine (16:0-22:6PE, PDPE) and, as a monounsaturated control, 1-palmitoyl-2-oleoyl-sn-glycerophosphatidylethanolamine (16:0-18:1PE, POPE) in mixtures with sphingomyelin (SM). By solid-state  $^2\text{H}$  NMR spectroscopy, we directly observe order and phase behavior of POPE- $\text{d}_{31}$  and PDPE- $\text{d}_{31}$  (analogs of POPE and PDPE with a perdeuterated sn-1 chain) in the mixed membranes. In complementary X-ray diffraction and differential scanning calorimetry experiments we further probe phase behavior. The spectra observed for POPE- $\text{d}_{31}$  in POPE/SM/ $\alpha$ -tocopherol (2:2:1 mol) reveal that a transition from gel to liquid crystalline phase is no longer apparent. At higher temperatures there is a superposition of two spectral components that we ascribe to  $\alpha$ -tocopherol promoting a transition from lamellar to inverted hexagonal ( $\text{H}_{\text{II}}$ ) phase. Analysis of depaked spectra shows that order is increased by about 8 % and that the amount of  $\text{H}_{\text{II}}$  phase increases with temperature, ranging from 7 (31 °C) to 41 % (65 °C). In mixed membranes where POPE- $\text{d}_{31}$  is replaced by PDPE- $\text{d}_{31}$ , we shall investigate whether there is a greater tendency for  $\alpha$ -tocopherol to increase order and destabilize bilayer structure for the polyunsaturated phospholipid.

Mentor: Stephen R. Wassall, Department of Physics, IUPUI