

Antenatal and Intrapartum Risk Factors for Use of Emergency and Restorative Medicaid Dental Services for Children

Juan F. Yepes DDS, MD, MPH, MS, DrPH
Associate Professor of Pediatric Dentistry
Indiana University School of Dentistry

Heather M. Bush PhD
Associate Professor
Department of Biostatistics
University of Kentucky College of Dentistry, Lexington, KY

Hsin-Fang Li MPH
Doctoral Student
University of Kentucky College of Public Health, Lexington, KY

Jeffrey Talbert PhD
Associate Professor
University of Kentucky College of Pharmacy, Lexington, KY

David A. Nash DMD, MS, EdD
William R. Willard Professor of Dental Education
Division of Pediatric Dentistry
University of Kentucky College of Dentistry, Lexington, KY

This is the author's manuscript of the article published in final edited form as:

Yepes, J. F., Bush, H. M., Li, H.-F., Talbert, J., & Nash, D. A. (2014). Antenatal and Intrapartum Risk Factors for Use of Emergency and Restorative Medicaid Dental Services for Children. *Pediatric Dentistry*, 36(5), 405–410.

**Antenatal and Intrapartum Risk Factors for Use of Emergency and Restorative Medicaid
Dental Services for Children**

ABSTRACT

Purpose: To investigate the relationship between antenatal/intrapartum factors affecting the utilization of Medicaid. **Methods:** A retrospective study was conducted. Three databases were used: 1) birth records 2) Medicaid files, and 3) Medicaid dental claims. Mode of delivery, mother's race, weight, and age-at-first dental visit were variables included. The outcome variables were dental claims.

Results: The analysis revealed children of Caucasian mothers were 34 percent more likely to have more-than one restorative claim than children of African-American mothers (OR=1.34, CI 1.10-1.65, $P=.0046$). Children born with low-birth-weight were 37 percent more likely to have emergency claims (OR= 1.37 CI 1.02-1.83, $P=.03$). The adjusted analysis found Caucasian mothers had higher odds ratio of having a dental claim than did African-American mothers ($P<.001$): 33 percent for a restorative claim, and 56 percent for an emergency claim. When race was analyzed, in African-American mothers, the odds of a restorative claim were 2.5 times higher in children delivered by C-section compared with vaginally delivered (OR= 2.52, CI 1.02-6.2, $P=.0001$). **Conclusions:** This study found an association between children of Caucasian mothers and the likelihood of experiencing claims. A relationship was found between children born with low-birth-weight and C-section and the likelihood of use of Medicaid services.

Key Words: *Medicaid, Early childhood Caries, Antenatal, C-section, Intrapartum, Children, vaginal birth*

INTRODUCTION

The importance of oral health to the general well-being of individuals is well established. Studies have focused on the impact association and correlation of the mother's oral health, habits, attitudes, and knowledge on the child's dental health, because mothers are traditionally regarded as the main caregivers of their children. Maternal factors may play an important role in shaping the oral health, attitudes, and behavior of their children.

The role of mother to child transmission of *Streptococcus mutans* has been a topic of several studies in the past decade. During and shortly after delivery, the epithelial surfaces in the oral cavity of microbiologically naïve infants become colonized by different bacterial species¹. As members of the indigenous biota, *Streptococcus oralis*, *S. mitis*, and *S. salivarius* are the predominant pioneer streptococci that colonize the oral cavity of infants during the first few days of life². Factors influencing the acquisition of *S. mutans* in infants include high maternal *S. mutans* levels and caries status, low infant birth weight, early tooth emergence, and low salivary IgA antibody levels³. Li et al¹ in a four-year follow-up study, observed a positive association between Caesarean delivery and initial acquisition and transmission of *Streptococcus mutans* in a mother-infant prospective cohort. Additionally, their analysis revealed that an infant delivered by C-section from a mother with a low socio-economic status acquired *S. mutans* earlier than did a vaginally delivered infant. Köhler⁴ et al. found that an early colonization of *S. mutans* was associated with higher incidence and more severe dental caries in children.

To date, there have been no reported studies that have investigated the relationship between antenatal and intrapartum maternal factors and dental claims that indirectly reflect early childhood caries. Therefore, the main purpose of this study was to determine the influence of antenatal and intrapartum risk factors on subsequent use of Medicaid dental services.

METHODS

Data and Sample Selection

A retrospective cohort study was completed subsequent to approval by University of Kentucky Institutional Review Board. Medicaid-enrolled children born in the Commonwealth of Kentucky in 2000, who had been were monitored longitudinally until their fifth birthday in 2005, were included in the analysis. The study was based on the following Kentucky databases: 1) Composite birth records (from Vital Health Statistics) from the 2000 calendar year; 2) Individual Medicaid eligibility files for all children born in 2000 and enrolled continuously in the Medicaid program from 2000 to 2005; and 3) Medicaid dental claims data covering 2000-2005. Children were excluded if there was more than one Medicaid identification number or the children were not enrolled continuously from 2000 to 2005.

Variable Definition

The study focused on claims filed through Medicaid for oral health services. Outcome variables were categorized according to type of dental visit (restorative or emergency care). Dental terminology codes from the Medicaid dental claims allowed determination of the type of visit. The dental codes used to reflect emergency visits and restorative treatment are presented in Table 1. Antenatal and intrapartum risk factors were obtained from the composite birth records. Mode of delivery (vaginal vs. C-section), mother's race (Caucasian vs. African-American), birth weight status (less than 2,500 grams vs. equal to or greater than 2,500 grams), and age at the first dental claim were the variables included in the analysis. The major outcome variables were the dental claims (restorative or emergency) filed with Medicaid from 2000 to 2005. Dental claims were assumed as an indirect or surrogate assessment of early childhood caries (**ECC**).

Statistical Analysis

The study's outcome measures were the number of claims filed through Medicaid for oral health services. The treatment codes were divided into restorative or emergency dental claims (Table 1). Each variable dichotomized into "no claim" or "equal to or greater than than one claim". All the risk factors were treated as categorical variables and were summarized using frequencies and percentages. The relationships between risk factors and dental claims were assessed using both adjusted and unadjusted odds ratios. Unadjusted odds ratios were computed for type of delivery, mother's race, and whether the child had low birth weight against the outcome variable in each type of dental visit. Chi-square tests of independence were used to compare the categorical variables. Adjusted odds ratios were computed using multivariable logistic regression for the same set of risk factors while adjusting for the child's age at first visit. Data were analyzed using the Statistical Analysis Software, SAS version 9.2, and a significance level of $p \leq 0.05$ was used for all statistical tests.

RESULTS

There were 55,978 live births in the Commonwealth of Kentucky in 2000, with 26,456 of those children being enrolled in Medicaid. Of the 26,456 Medicaid-enrolled children born in 2000, 5,342 were continuously enrolled in Medicaid for five years and generated a dental claim during the study period (2000-2005). At birth, the mothers' (n=5,342) mean age was 22.7 (SD=5.2). Eighty-three percent of the mothers were Caucasians, and 96 percent of the children were born with normal weight. A total of 5,051 infants were delivered vaginally and 291 were delivered by C-section. The number of dental claims generated during the study period was 21,780.

Bivariate analysis

Restorative claims by antenatal / intrapartum factors

Counts, percentages, and crude odds ratios were calculated using more than one claim for restorative treatment as the outcome variable (Table 2). While no difference was found between the type of delivery and whether the child had low birth weight, mother's race was found as an important antenatal factor. The odds of having more than one restorative claims for children whose mother was Caucasian was 1.34 times the odds of those whose mother was African-American (CI 1.10-1.65, $P<0.05$).

Emergency dental claims by antenatal / intrapartum factors

Children born with low birth weight had higher odds of more than one emergency dental claim during the five-year period (OR= 1.37 CI 1.02-1.83, $P<0.05$). Mother's race was also related to increased odds of having more than one emergency dental claim. Children of Caucasian mothers

were 58 percent more likely to have more than one emergency dental claim compared with children from African-American mothers (OR=1.58 CI 1.28-1.96, $P<0.05$). (Table 3)

Logistic Regression Analysis

Restorative / emergency claims and antenatal / intrapartum risk factors

Binary logistic regression was utilized to investigate the relationship between multiple variables with the restorative and emergency claim outcomes. After adjusting for variables, the odds of experiencing a restorative claim was 33 percent higher in children of Caucasian mothers compared with children of African-American mothers (OR=1.33, CI 1.07-1.63, $P<0.05$). (Table 4) Similarly, the adjusted odds of experiencing an emergency dental claim was 56 percent higher for children of Caucasian mothers compared with children of African-American mothers (OR=1.56, CI 1.26-1.96, $P<0.05$). When adjusted for the child's age of first dental visit, the association between whether the child had a low birth weight was not associated with having more than one emergency dental claim. In addition, it was found that the odds of having a restorative dental claim and an emergency dental claim are reduced by 29 percent and 34 percent respectively for every unit increase in age. In other words, an early dental visit was associated with an increased number of subsequent dental claims.

Whether age at first visit and race were effect modifiers was investigated. There was an interaction between mother's race and type of delivery for caries. Specifically, among African-American mothers, the odds of having a restorative claim experience for children born with C-section was 2.52 times (OR=2.52, CI 1.02 - 6.21, $P<0.05$) that of children with vaginal delivery, while adjusting for low birth weight and age at first dental visit. (Table 5) However, the estimated odds ratio for the Caucasian mothers was not significant (OR=1.06, CI 0.74, 1.52). The delivery type

did not demonstrate a strong effect on restorative claims as a main effect due to difference between the numbers of Caucasian mothers versus African-American ones (n=2177 vs. n=451). Race was not an effect modifier for emergency dental claims.

DISCUSSION

The main purpose of this study was to investigate the relationship between antenatal/intrapartum factors on subsequent use of Medicaid dental services.

Race and dental claims

Children born from African-American mothers were less likely to use Medicaid dental services. The odds of having more than one dental claim for restorative dentistry and emergency care was higher for children born to Caucasian mothers compared with children born to African-American mothers. The lower utilization of services by minority children on Medicaid was studied previously by Dasanayake et al ⁵, who found that among Alabama children on Medicaid, fewer African-American than Caucasian received dental care. Several factors might have influenced the low utilization rates among children born from African-American mothers. Lack of transportation and limited appointment availabilities were reported by caregivers as two reasons they felt discourage from seeking dental care for their children covered by Medicaid ⁶. In addition, it has been reported that dental patients feel more comfortable with a provider of their own race⁷ This could have influenced minority utilization in this study, because less than five percent of the practicing dentists in Kentucky represent minorities ⁸. A different approach to analyzing the results is to assume that children born from African-American mothers are interacting less with Medicaid dental services, because they have a better oral health status. However, this assumption is unlikely

in Kentucky where previous studies found a poor oral health in children from minority ethnic groups compared with Caucasians^{9,10}.

Low birth weight and dental claims

Preterm birth (delivery at less than 37 weeks of gestation) and low birth weight (less than 2,500 grams) are the leading causes of neonatal morbidity and mortality in the United States^{11,12}. Several factors are associated with preterm birth and low birth weight, such as low maternal body mass index, maternal smoking, and maternal infection¹³. In this study, children born with low birth weight were more likely to have emergency (37 percent) and restorative (28 percent) dental claims compared with children born with normal weight. The association was significant in the bivariate analysis but not significant in the logistic regression model. However, the analysis showed a potential trend that merits further investigation. The relationship between low birth weight and dental caries has been studied extensively in the last 10 years¹⁴⁻¹⁶. In a recent study, Rajshekar¹⁷ and co-researchers studied the relationship between low birth weight with caries experience in 250 children. They found a significant relationship between caries and children born with low birth weight. Nelson and co-researchers¹⁸ examined developmental enamel defects and dental caries in low birth weight adolescents compared with full term adolescent. They found that low birth weight was a significant risk factor for increased enamel defects in the permanent incisors and first molars. These enamel defects are associated with an increase in caries.

Albert and Nelson¹⁹ examined the effect of low birth weight (exposure) in dental caries (outcome) and assessed if whether the exposure was a direct or an indirect effect on the outcomes. They concluded that low birth weight had a direct negative effect on the prevalence of dental caries. Despite the previous studies, it is important to recognize that other studies have failed to

demonstrate any association between low birth weight and dental caries. Berg ²⁰ and Shulman ²¹ in 2006 and 2005 respectively, published two systematic reviews. Neither of these reviews supported the association between low birth weight and dental caries. Using a unique design with Medicaid data in the Commonwealth of Kentucky this study found that children born with low birth weight were more likely to have restorative or emergency dental claims.

Mode of delivery and dental claims

The first exposure to microorganisms in vaginally delivered infants occurs during the passage through the birth canal; on the other hand, the first exposure to bacteria in infants born by C-section is from the skin of the parents and health providers. The concept of differences in the microbiology and health status between children born by C-section and vaginal delivery has been extensively reported in the literature. Glass and co-researchers ²² in a population-based study found that infants delivered by C-section carried the highest risk for developing seizures later in life when compared with infants delivered vaginally. Furthermore, vaginally delivered children have been reported to have a more diverse gut microbiology environment, whereas children delivered by C-section had higher numbers of *Clostridium difficile* with delayed acquisition of bifidobacteria and *Escherichia coli* ²³⁻²⁵.

Gibbons and colleagues ²⁶ demonstrated that the initial pioneer bacteria entering the oral cavity influence the pattern of microbial succession, and that succession is associated with the availability of colonization sites. Bacteria that subsequently attempt colonization must compete with other microorganisms for colonization sites and essential nutrients. Additionally, they must survive in the presence of adverse metabolic end-products that may be produced by other members of the environment. Once established, early colonization bacteria tend to persist in the mouth^{27, 28}. It is

reasonable hypothesize that C-section delivered infants may have experienced less exposure to maternal environmental microbial challenges at birth, and an atypical microbial environment may prevail, providing more potential biological binding for *S. mutans*. A recent study by Holgerson and co-researchers ²⁹ in Sweden compared the oral microbiology in infants delivered by C-section and vaginal delivery. The oral biofilm was assayed by the Human Oral Microbe Identification Microarray (**HOMIM**) in healthy three months old infants (38 delivered by C-section and 25 delivered vaginally). More bacteria were detected in the infants delivered vaginally compared with infants delivered by C-section.

A previous study by Li and colleagues ¹ explored the mode of delivery and other maternal factors in the acquisition of *Streptococcus mutans* in infants. The study found that among infants, those delivered by C-section acquired *S. mutans* 12 months early than did vaginally delivery infants. Children delivered by C-section had less exposure to the maternal microbiota at birth and maybe a different environment for the colonization by *S. Mutans*. This current study design was significantly different from previous studies. This study explored the association between mode of delivery (C-section vs. vaginal delivery) and the subsequent use of Medicaid dental services. Using a retrospective cohort, this study found an interesting trend. The likelihood of experiencing restorative and emergency claims were 25 percent higher for children born by C-section compared with vaginal delivery. These results add more evidence to the literature regarding the different oral environments based on the mode of delivery. They suggest potential additional risk factor for caries to be included by the dentist in the risk assessment for caries.

This study found an interesting interaction between mother's race and type of delivery for restorative claims that merits future investigations. Specifically, among African-American mothers, the odds of having a restorative claim experience for children delivered with C-section

was 2.52 times (OR=2.52, CI 1.02 - 6.21, $P<0.05$) that of children with vaginal delivery, while adjusting for low birth weight and age at first dental visit. While the missing data on race (over 50 percent) is a major limitation, this preliminary finding, may generate new hypothesis of differences between vaginal and C-section delivery in relation with dental caries.

Age of the first dental visit and dental claims

Children who had an early claim for restorative or emergency dental treatment were more likely to have subsequent dental claims. Savage and colleagues³⁰ found that children who had their first preventive dental visit by age one were more likely to have subsequent preventive visits but were not more likely to have subsequent restorative or emergency visits. In the same direction of Savage's research, the present study confirms, the importance of early anticipatory guidance and prevention. Also, prevention programs must prioritize specific designs and tools for the sub-population of patients who had an early visit for restorative or emergency care.

LIMITATIONS

Though this was a large population-based study to examine antenatal and intrapartum risk factors for caries, the data were limited in several ways. Treatment codes for the diagnosis of caries were relied upon rather than direct examination and detection of caries. On the basis of previous research among low-income children, (31, 32) it is possible to anticipate that this population would be at high risk for caries. However, specific disease rates for the population of interest were unknown. Second, as in any investigation that relies solely on administrative data, the study was susceptible to coding errors. The accuracy of coding dental claims in the Commonwealth of Kentucky is unknown. Third, because this was a retrospective cohort study, there was the potential for selection

bias that affected the interpretation of the results. Fourth, the amount of missing data in relation with race potentially could have affected the final results.

CONCLUSIONS

This study retrospectively explored antenatal and intrapartum risk factors associated with dental claims in the Commonwealth of Kentucky. This study concluded:

- 1) The odds of experiencing a restorative dental claim was 33 percent higher for children of Caucasian mothers compared with children of African-American mothers.
- 2) The odds of experiencing an emergency dental claim was 56 percent higher for children of Caucasian mothers compared with children of African-American mothers. These two conclusions demonstrated a lower utilization of Medicaid dental services by minority children, assuming that the burden of disease was higher in the minority population.
- 3) Children born with low birth weight were more likely to use Medicaid dental services suggesting a trend of more severe dental disease associated with low birth weight.
- 4) Children delivered by C-section were 25 percent more likely to have a restorative or emergency dental claim compared with vaginally delivered children. This result is based on extrapolated data and is affected by several confounders. However, this finding maybe suggests different times of colonization by cariogenic bacteria in the oral cavity according to the mode of delivery.

TABLES

Table 1. Emergency and Restorative Dental Codes

Emergency Dental Codes		Restorative Dental Codes	
D0140	Limited oral evaluation	D2140	Amalgam –one surface
D9110	Emergency palliative	D2330	Composite –one surface anterior
D7140	Extraction –Erupted tooth-	D2391	Composite –one surface posterior
		D2150	Amalgam –two surface
		D2331	Composite –two surface anterior
		D2392	Composite –two surface posterior
		D2160	Amalgam –three surface
		D2332	Composite –three surface anterior
		D2393	Composite –three surface posterior
		D2161	Amalgam –four or more surface
		D2335	Composite incisal angle
		D2390	Composite crown anterior
		D2394	Composite –four or more posterior
		D2933	Crown prefabricated with resin
		D2930	Stainless Steel Crown primary
		D3220	Pulpotomy
		D3240	Pulp therapy
		D3120	Indirect pulp cap

Table 2 - Restorative claims

	Total Participants (N=5342)	No Claim (N=2804)	>1 Claim (N=2538)	Odds Ratio (95% CI)	P-value
Type of Delivery					0.8331
Vaginal delivery	5051 (94.6%)	2653 (94.6%)	2398 (94.5%)		
C-section	291 (5.4%)	151 (5.4%)	140 (5.5%)	1.03 (0.81, 1.30)	
Mother race*					0.0046
African-American	451 (17.2%)	264 (19.1%)	187 (15.0%)		
Caucasian	2177 (82.8%)	1115 (80.9%)	1062 (85.0%)	1.34 (1.10, 1.65)	
Birth weight					0.0965
≥2,500g	5153 (96.5%)	2716 (96.9%)	2437 (96.0%)		
<2,500g	189 (3.5%)	88 (3.1%)	101 (4.0%)	1.28 (0.96, 1.71)	

Note: 50% of the information on mother's race is missing (n=2628) where 2177 (83%) are Caucasian and 451(17%) are African-American.

Table 3 - Emergency Claims

	Total Participants (N=5342)	No Emergency Claim (N=2948)	≥1 Emergency Claim (N=2394)	Odds Ratio (95% CI)	P-value
Type of Delivery					0.2978
Vaginal delivery	5051 (94.6%)	2796 (94.8%)	2255 (94.2%)		
C-section	291 (5.4%)	152 (5.2%)	139 (5.8%)	1.13 (0.90, 1.44)	
Mother race*					<.0001
African- American	451 (17.2%)	295 (19.9%)	156 (13.6%)		
Caucasian	2177 (82.8%)	1185 (80.1%)	992 (86.4%)	1.58 (1.28, 1.96)	
Birth weight					0.0332
≥2,500g	5153 (96.5%)	2858 (96.9%)	2295 (95.9%)		
<2,500g	189 (3.5%)	90 (3.1%)	99 (4.1%)	1.37 (1.02, 1.83)	

Note: 50% of the information on mother's race is missing (n=2628) where 2177 (83%) are Caucasian and 451(17%) are African-American.

Table 4 – LRM determining factors associate with restorative and emergency claims

	Restorative claims (N=2,538) OR (95% CI)	Emergency claims (N=2,255) OR (95% CI)
Mother Race		
African-American	Reference	Reference
Caucasian	1.33 (1.07-1.63) *	1.56 (1.26-1.96) *
Baby weight		
Normal birth weight	Reference	Reference
low birth weight	1.42 (0.8-2.27)	1.42 (0.86-2.3)
Mode of delivery		
Vaginal	Reference	Reference
C-section	1.25 (0.86-1.66)	1.25 (0.9-1.66)
Age of first visit	0.71 (0.63-0.8) *	0.66 (0.63-0.7) *

* Statistically significant ($P < 0.05$)

LRM: Linear Regression Model

Table 5– Logistic regression model determining whether age at first visit and race were effect modifiers for African American mothers

	Restorative claims OR (95% CI)
Baby weight	
Normal birth weight	Reference
low birth weight	0.96 (0.3-2.7)
Mode of delivery	
Vaginal	Reference
C-section	2.52 (1.02-6.2)
Age of first visit	0.69 (0.57-0.84) *

* Statistically significant ($P < 0.05$)

REFERENCES

1. Li Y, Caufield PW, Dasanayake AP, Wiener HW, Vermund SH. Mode of delivery and other maternal factors influence the acquisition of *Streptococcus mutans* in infants. *J Dent Res*. 2005;84(9):806-11.
2. Smith DJ, Anderson JM, King WF, van Houte J, Taubman MA. Oral streptococcal colonization of infants. *Oral Microbiol Immunol*. 1993;8(1):1-4.
3. Milgrom P, Riedy CA, Weinstein P, Tanner AC, Manibusan L, Bruss J. Dental caries and its relationship to bacterial infection, hypoplasia, diet, and oral hygiene in 6- to 36-month-old children. *Community Dent Oral Epidemiol*. 2000;28(4):295-306.
4. Kohler B, Andreen I, Jonsson B. The earlier the colonization by mutans streptococci, the higher the caries prevalence at 4 years of age. *Oral Microbiol Immunol*. 1988;3(1):14-7.
5. Dasanayake AP, Li Y, Wadhawan S, Kirk K, Bronstein J, Childers NK. Disparities in dental service utilization among Alabama Medicaid children. *Community Dent Oral Epidemiol*. 2002;30(5):369-76.
6. Mofidi M, Rozier RG, King RS. Problems with access to dental care for Medicaid-insured children: what caregivers think. *Am J Public Health*. 2002;92(1):53-8.
7. Mofidi M, Konrad TR, Porterfield DS, Niska R, Wells B. Provision of care to the underserved populations by National Health Service Corps alumni dentists. *J Public Health Dent*. 2002;62(2):102-8.
8. Saman DM, Arevalo O, Johnson AO. The dental workforce in Kentucky: current status and future needs. *J Public Health Dent*. 2010;70(3):188-96.
9. Chattopadhyay A, Arevalo O, Cecil JC, 3rd. Kentucky's oral health indicators and progress towards Healthy People 2010 objectives. *J Ky Med Assoc*. 2008;106(4):165-74.
10. Chattopadhyay A. Oral health disparities in the United States. *Dent Clin North Am*. 2008;52(2):297-318, vi.
11. Martin JA, Hamilton BE, Ventura SJ, Menacker F, Park MM. Births: final data for 2000. *Natl Vital Stat Rep*. 2002;50(5):1-101.
12. Martin JA, Hamilton BE, Ventura SJ, Menacker F, Park MM, Sutton PD. Births: final data for 2001. *Natl Vital Stat Rep*. 2002;51(2):1-102.
13. Kramer MS. The epidemiology of adverse pregnancy outcomes: an overview. *J Nutr*. 2003;133(5 Suppl 2):1592S-6S.
14. O'Connell AC, O'Connell SM, O'Mullane E, Hoey HM. Oral health of children born small for gestational age. *Ir Med J*. 2010;103(9):275-8.
15. Ghasempour M, Ahmadpour-Kacho M, Sheikhi S. Dental caries in pre-term and low birth-weight children and related factors. *J Contemp Dent Pract*. 2009;10(4):51-8.
16. Saraiva MC, Bettiol H, Barbieri MA, Silva AA. Are intrauterine growth restriction and preterm birth associated with dental caries? *Community Dent Oral Epidemiol*. 2007;35(5):364-76.
17. Rajshekar SA, Laxminarayan N. Comparison of primary dentition caries experience in pre-term low birth-weight and full-term normal birth-weight children aged one to six years. *J Indian Soc Pedod Prev Dent*. 2011;29(2):128-34.
18. Nelson S, Albert JM, Lombardi G, Wishnek S, Asaad G, Kirchner HL, et al. Dental caries and enamel defects in very low birth weight adolescents. *Caries Res*. 2010;44(6):509-18.
19. Albert JM, Nelson S. Generalized causal mediation analysis. *Biometrics*. 2011;67(3):1028-38.
20. Berg J, Cunha-Cruz J. Low birth weight babies are not at increased risk of primary tooth caries. *J Evid Based Dent Pract*. 2006;6(2):195-6.

21. Shulman JD. Is there an association between low birth weight and caries in the primary dentition? *Caries Res.* 2005;39(3):161-7.
22. Glass HC, Pham TN, Danielsen B, Towner D, Glidden D, Wu YW. Antenatal and intrapartum risk factors for seizures in term newborns: a population-based study, California 1998-2002. *J Pediatr.* 2009;154(1):24-8 e1.
23. Wang M, Ahrne S, Jeppsson B, Molin G. Comparison of bacterial diversity along the human intestinal tract by direct cloning and sequencing of 16S rRNA genes. *FEMS Microbiol Ecol.* 2005;54(2):219-31.
24. Penders J, Thijs C, van den Brandt PA, Kummeling I, Snijders B, Stelma F, et al. Gut microbiota composition and development of atopic manifestations in infancy: the KOALA Birth Cohort Study. *Gut.* 2007;56(5):661-7.
25. Penders J, Thijs C, Vink C, Stelma FF, Snijders B, Kummeling I, et al. Factors influencing the composition of the intestinal microbiota in early infancy. *Pediatrics.* 2006;118(2):511-21.
26. Gibbons RJ. Bacterial adhesion to oral tissues: a model for infectious diseases. *J Dent Res.* 1989;68(5):750-60.
27. Cole MF, Bryan S, Evans MK, Pearce CL, Sheridan MJ, Sura PA, et al. Humoral immunity to commensal oral bacteria in human infants: salivary antibodies reactive with *Actinomyces naeslundii* genospecies 1 and 2 during colonization. *Infect Immun.* 1998;66(9):4283-9.
28. Kononen E, Kanervo A, Takala A, Asikainen S, Jousimies-Somer H. Establishment of oral anaerobes during the first year of life. *J Dent Res.* 1999;78(10):1634-9.
29. Lif Holgerson P, Harnevik L, Hernell O, Tanner AC, Johansson I. Mode of birth delivery affects oral microbiota in infants. *J Dent Res.* 2011;90(10):1183-8.
30. Savage MF, Lee JY, Kotch JB, Vann WF, Jr. Early preventive dental visits: effects on subsequent utilization and costs. *Pediatrics.* 2004;114(4):e418-23.
31. Vargas CM, Crall JJ, Schneider DA. Sociodemographic distribution of pediatric dental caries: NHANES III, 1988-1994. *J Am Dent Assoc.* 1998;129(9):1229-38.
32. Newacheck PW, Hughes DC, Hung YY, Wong S, Stoddard JJ. The unmet health needs of America's children. *Pediatrics.* 2000;105(4 Pt 2):989-97.