

## POSTER PRESENTATION


## Open Access

# Qualitatively different scenarios for co-activation of NMDA, AMPA and GABA receptor currents on dopaminergic neuron

Denis Zakharov<sup>1,2\*</sup>, Alexey Kuznetsov<sup>3</sup>From 24th Annual Computational Neuroscience Meeting: CNS\*2015  
Prague, Czech Republic. 18-23 July 2015

We introduced a reduced functional model that qualitatively describes fundamental properties of the dopaminergic (DA) neuron. The neuron is a low-frequency pacemaker, and requires special conditions to increase the frequency of its firing [1]. In more detail, only N-methyl-D-aspartate receptor (NMDAR) current can significantly elevate the frequency (more than 5 fold in comparison with pacemaking).  $\alpha$ -Amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor (AMPA) current is not able to evoke such a high frequency. Instead, it suppresses firing at a conductance much lower than NMDAR.  $\gamma$ -Aminobutyric acid receptor (GABAR) activation first decreases the firing frequency, and only then suppresses firing. Thus, the latter property determines type 1 excitability for the DA neuron, whereas the difference in responses to AMPA and NMDA receptor activation make the neuron different from other

well-studied types. We focus this report on the co-activation of these receptors. Co-activation of AMPA together with NMDA receptor may obscure the high frequency firing and lead directly to its suppression. There are contradictory experimental data in this case [2,3]. We show that AMPAR can do both: further contribute to the frequency increase produced by NMDAR, or impede the high-frequency firing. We show that the distinction between these two cases is the level of the background NMDAR activation. Co-activation of GABA together with NMDA receptor may also lead directly to the suppression of high-frequency oscillations (e.g. by shunting inhibition), or reduce the frequency first. Experiments clearly show the latter [4]: the frequency can be reduced back to that in control conditions when GABA and NMDA receptor currents are balanced. This property allows us to recalibrate the model so that it retains type

\* Correspondence: [zakharov@neuron.appl.sci-nnov.ru](mailto:zakharov@neuron.appl.sci-nnov.ru)<sup>1</sup>Department of Nonlinear Dynamics, Institute of Applied Physics, RAS, Nizhny Novgorod, 603950, Russia

Full list of author information is available at the end of the article

1 excitability not only in control, but also with activated synaptic currents. In general, activation of synaptic currents changes neural excitability, and we investigate the dynamical mechanisms responsible for the transition. As a result, we were able to construct a model of the DA neuron that collects all the properties described above. The model is essential for further research of behaviorally-relevant modulations of the DA neuron activity, which involves simultaneous excitatory and inhibitory inputs to the neuron.

#### Acknowledgements

This work was supported by the RFBR (project 14-02-00916-a) and by the grant NIH-NIAAA R01AA022821.

#### Authors' details

<sup>1</sup>Department of Nonlinear Dynamics, Institute of Applied Physics, RAS, Nizhny Novgorod, 603950, Russia. <sup>2</sup>Laboratory of Nonlinear Physics, Nizhny Novgorod State University, Nizhny Novgorod, 603022, Russia. <sup>3</sup>Department of Mathematical Sciences and Center for Mathematical Modeling and Computational Sciences, IUPUI, Indianapolis, IN 46202, USA.

Published: 18 December 2015

#### References

1. Zakharov DG, Kuznetsov AS: **A minimal model for a slow pacemaking neuron.** *Chaos, Solitons & Fractals* 2012, **45**:640-644.
2. Deister CA, Teagarden MA, Wilson CJ, Paladini CA: **An intrinsic neuronal oscillator underlies dopaminergic neuron bursting.** *J Neurosci* 2009, **29**:15888-97.
3. Blythe SN, Wokosin D, Atherton JF, Bevan MD: **Cellular Mechanisms Underlying Burst Firing in Substantia Nigra Dopamine Neurons.** *J Neurosci* 2009, **29**:15531-41.
4. Lobb CJ, Wilson CJ, Paladini CA: **A Dynamic Role for GABA Receptors on the Firing Pattern of Midbrain Dopaminergic Neurons.** *J Neurophysiology* 2010, **104**:403-413.

doi:10.1186/1471-2202-16-S1-P138

**Cite this article as:** Zakharov and Kuznetsov: **Qualitatively different scenarios for co-activation of NMDA, AMPA and GABA receptor currents on dopaminergic neuron.** *BMC Neuroscience* 2015 **16**(Suppl 1):P138.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

