

Targeted neurogenesis pathway-based gene analysis identifies *ADORA2A* associated with hippocampal volume in mild cognitive impairment and Alzheimer's disease

Emrin Horgusluoglu, M.S.^{1,2}, Kwangsik Nho, Ph.D.^{2,3*}, Shannon L. Risacher, Ph.D.², Andrew J. Saykin, Psy.D.^{1,2,3,4*}

¹Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis, IN, USA;

²Center for Neuroimaging, Department of Radiology and Imaging Sciences, Indiana University School of Medicine, Indianapolis, IN, USA;

³Center for Computational Biology and Bioinformatics, Indiana University School of Medicine, Indianapolis, IN, USA;

⁴Indiana Alzheimer's Disease Center, Indiana University School of Medicine, Indianapolis, IN, USA

Background: New neurons are generated throughout adulthood in the olfactory bulb and dentate gyrus of the hippocampus, and are incorporated into hippocampal networks during maintenance of neural circuits and in turn contribute to learning and memory. Numerous intrinsic and extrinsic factors such as growth factors, transcription factors, and cell cycle regulators control neural stem cells proliferation, differentiation, and maintenance into mature neurons. However, the genetic mechanisms controlling adult hippocampal neurogenesis remain unclear. We performed a gene-based association analysis of neurogenesis pathway-related candidate genes using data from the Alzheimer's Disease Neuroimaging Initiative (ADNI).

Methods: Neurogenesis-related genes were curated from existing databases (Qiagen RT² Profiler PCR Arrays, GoGene and MANGO). The gene list was filtered by AD susceptibility genes from the Alzgene database (<http://www.alzgene.org/>) and large-scale GWAS (Lambert, et al. 2013, *Nature*). Caucasian non-Hispanic individuals (N=1,525) with AD or mild cognitive impairment (MCI) and cognitively normal older adults from the ADNI cohort with MRI and genotyping data were included. Gene-based association analysis of neurogenesis pathway-related candidate genes was performed. Baseline bilateral hippocampus and hippocampal subfield (CA regions and dentate gyrus) volumes were extracted from MRI and served as phenotypes. Gender, age, intracranial volume, MRI field strength, and diagnosis at scanning were entered as covariates. The empirical *p* value from permutation testing for each gene was adjusted for the number of significant SNPs in each gene.

Results: *ADORA2A* was significantly associated with total hippocampal volume and hippocampal subfield volumes ($p < 0.001$). For the most significant SNP (rs9608282) in *ADORA2A*, dosage of the minor allele (T) increased hippocampal volume. rs9608282 was also associated with composite memory score ($p = 0.0076$).

Conclusion: *ADORA2A*-mediated control of neuroinflammation modulates adult neurogenesis and the inhibition of *ADORA2A* prevents A β -induced neurotoxicity. Targeted pathway-based genetic analysis combined with brain imaging endophenotypes appears promising to help elucidate disease pathophysiology and identify potential therapeutic targets.

**Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

*Please address correspondences to:

Kwangsik Nho, PhD
Center for Neuroimaging,
Department of Radiology and Imaging Sciences,
Indiana University School of Medicine, Indianapolis, IN, USA
Phone: 317-963-7503; Fax: 317-963-7547; E-mail: knho@iupui.edu

Andrew J. Saykin, PsyD
Center for Neuroimaging,
Department of Radiology and Imaging Sciences,
Indiana University School of Medicine, Indianapolis, IN, USA
Phone: 317-963-7501; Fax: 317-963-7547; E-mail: asaykin@iu.edu