



















in patients with asymptomatic and symptomatic left ventricular systolic dysfunction: a retrospective analysis of the SOLVD trials. *Studies of Left Ventricular Dysfunction. J Am Coll Cardiol.* 1998; 32: 695-703.

9. Hunt SA, Abraham WT, Chin MH, Feldman AM, Francis GS, Ganiats TG, Jessup M, Konstam MA, Mancini DM, Michl K, Oates JA, Rahko PS, Silver MA, Stevenson LW, Yancy CW, Antman EM, Smith SC, Jr, Adams CD, Anderson JL, Faxon DP, Fuster V, Halperin JL, Hiratzka LF, Jacobs AK, Nishimura R, Ornato JP, Page RL, Riegel B, American College of Cardiology, American Heart Association Task Force on Practice Guidelines, American College of Chest Physicians, International Society for Heart and Lung Transplantation, Heart Rhythm Society. ACC/AHA 2005 Guideline Update for the Diagnosis and Management of Chronic Heart Failure in the Adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure): developed in collaboration with the American College of Chest Physicians and the International Society for Heart and Lung Transplantation: endorsed by the Heart Rhythm Society. *Circulation.* 2005; 112: e154-235.

10. Maisel WH, Stevenson LW. Atrial fibrillation in heart failure: epidemiology, pathophysiology, and rationale for therapy. *Am J Cardiol.* 2003; 91: 2D-8D.

11. Mathew J, Hunsberger S, Fleg J, Mc Sherry F, Williford W, Yusuf S. Incidence, predictive factors, and prognostic significance of supraventricular tachyarrhythmias in congestive heart failure. *Chest*. 2000; 118: 914-922.
12. Benza RL, Tallaj JA, Felker GM, Zabel KM, Kao W, Bourge RC, Pearce D, Leimberger JD, Borzak S, O'connor CM, Gheorghiade M, OPTIME-CHF Investigators. The impact of arrhythmias in acute heart failure. *J Card Fail*. 2004; 10: 279-284.
13. Olsson LG, Swedberg K, Ducharme A, Granger CB, Michelson EL, McMurray JJ, Puu M, Yusuf S, Pfeffer MA, CHARM Investigators. Atrial fibrillation and risk of clinical events in chronic heart failure with and without left ventricular systolic dysfunction: results from the Candesartan in Heart failure-Assessment of Reduction in Mortality and morbidity (CHARM) program. *J Am Coll Cardiol*. 2006; 47: 1997-2004.
14. Swedberg K, Olsson LG, Charlesworth A, Cleland J, Hanrath P, Komajda M, Metra M, Torp-Pedersen C, Poole-Wilson P. Prognostic relevance of atrial fibrillation in patients with chronic heart failure on long-term treatment with beta-blockers: results from COMET. *Eur Heart J*. 2005; 26: 1303-1308.
15. Savelieva I, Camm J. Anti-arrhythmic drug therapy for atrial fibrillation: current anti-arrhythmic drugs, investigational agents, and innovative approaches. *Europace*. 2008; 10: 647-665.

16. Tayal AH, Tian M, Kelly KM, Jones SC, Wright DG, Singh D, Jarouse J, Brillman J, Murali S, Gupta R. Atrial fibrillation detected by mobile cardiac outpatient telemetry in cryptogenic TIA or stroke. *Neurology*. 2008; .
17. Wolf P, Kannel W, McGee D, Meeks S, Bharucha N, McNamara P. Duration of atrial fibrillation and imminence of stroke: the Framingham study. *Stroke*. 1983; 14: 664-667.
18. Pollak WM, Simmons JD, Interian A, Jr, Atapattu SA, Castellanos A, Myerburg RJ, Mitrani RD. Clinical utility of intraatrial pacemaker stored electrograms to diagnose atrial fibrillation and flutter. *Pacing Clin Electrophysiol*. 2001; 24: 424-429.
19. Seidl K, Meisel E, VanAgt E, Ottenhoff F, Hess M, Hauer B, Zahn R, Senges J. Is the atrial high rate episode diagnostic feature reliable in detecting paroxysmal episodes of atrial tachyarrhythmias? *Pacing Clin Electrophysiol*. 1998; 21: 694-700.
20. Israel CW, Gronefeld G, Ehrlich JR, Li YG, Hohnloser SH. Long-term risk of recurrent atrial fibrillation as documented by an implantable monitoring device: implications for optimal patient care. *J Am Coll Cardiol*. 2004; 43: 47-52.
21. Gillis AM, Morck M. Atrial fibrillation after DDDR pacemaker implantation. *J Cardiovasc Electrophysiol*. 2002; 13: 542-547.
22. Cheung JW, Keating RJ, Stein KM, Markowitz SM, Iwai S, Shah BK, Lerman BB, Mittal S. Newly detected atrial fibrillation following dual chamber pacemaker implantation. *J Cardiovasc Electrophysiol*. 2006; 17: 1323-1328.

23. Glotzer TV, Hellkamp AS, Zimmerman J, Sweeney MO, Yee R, Marinchak R, Cook J, Paraschos A, Love J, Radoslovich G, Lee KL, Lamas GA, MOST Investigators. Atrial high rate episodes detected by pacemaker diagnostics predict death and stroke: report of the Atrial Diagnostics Ancillary Study of the MOde Selection Trial (MOST). *Circulation*. 2003; 107: 1614-1619.
24. Bunch TJ, Day JD, Olshansky B, Stolen KQ, Mullins CM. Newly Detected Atrial Fibrillation in Patients with an Implantable Cardioverter Defibrillator is a Strong Risk Marker of Increased Mortality. *Heart Rhythm*. 2008; .
25. Deneke T, Lawo T, Gerritse B, Lemke B, The European GEM DR Trade Mark Investigators. Mortality of patients with implanted cardioverter/defibrillators in relation to episodes of atrial fibrillation. *Europace*. 2004; 6: 151-158.
26. Ryan S, Siemon G, Drogemuller A, Rameken M, Vater M, Senges J, Seidl K. 2 Year Follow-Up of 321 Patients with an Implantable Cardioverter/defibrillator: Comparison of Patients with and without Atrial Fibrillation. *Z Kardiol*. 2001; 90: 906-915.
27. Capucci A, Santini M, Padeletti L, Gulizia M, Botto G, Boriani G, Ricci R, Favale S, Zolezzi F, Di Belardino N, Molon G, Drago F, Villani GQ, Mazzini E, Vimercati M, Grammatico A, Italian AT500 Registry Investigators. Monitored atrial fibrillation duration predicts arterial embolic events in patients suffering from bradycardia and atrial fibrillation implanted with antitachycardia pacemakers. *J Am Coll Cardiol*. 2005; 46: 1913-1920.

28. Poole JE, Johnson GW, Hellkamp AS, Anderson J, Callans DJ, Raitt MH, Reddy RK, Marchlinski FE, Yee R, Guarnieri T, Talajic M, Wilber DJ, Fishbein DP, Packer DL, Mark DB, Lee KL, Bardy GH. Prognostic importance of defibrillator shocks in patients with heart failure. *N Engl J Med.* 2008; 359: 1009-1017.