

5-21-2012

RhoA/ROCK Pathway is the Major Molecular Determinant of Basal Tone in Intact Human Internal Anal Sphincter

Satish Rattan

Thomas Jefferson University, Satish.Rattan@jefferson.edu

Jagmohan Singh

Thomas Jefferson University

Let us know how access to this document benefits you

Follow this and additional works at: http://jdc.jefferson.edu/gastro_hepfpPart of the [Gastroenterology Commons](#), and the [Hepatology Commons](#)

Recommended Citation

Rattan, Satish and Singh, Jagmohan, "RhoA/ROCK Pathway is the Major Molecular Determinant of Basal Tone in Intact Human Internal Anal Sphincter" (2012). *Division of Gastroenterology and Hepatology Faculty Papers*. Paper 9.

http://jdc.jefferson.edu/gastro_hepfp/9

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Division of Gastroenterology and Hepatology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Backgrounds & Aims

Knowledge of molecular control mechanisms underlying the basal tone in the intact human IAS is critical for the pathophysiology and rational therapy for debilitating rectoanal motility disorders.

Results




Figure 1. A,B. Phasic and tonic activities of RSM and IAS respectively. **C.** % maximal basal tone in IAS and RSM. **D.** EFS response in IAS and RSM.

Methods

We determined the effects of ROCK and PKC-selective inhibitors Y 27632 and Gö 6850 (10^{-8} to 10^{-4} M), respectively on the basal tone in the IAS vs. the RSM. We performed Western blot analysis, confocal microscopy and enzymatic activity assay to determine the levels, membrane distribution and enzyme activity of RhoA/ROCKII, PKC α , MYPT1, CPI-17, and MLC $_{20}$ before and after Y 27632 and Gö 6850.





Figure 3. A, B, C. Immunofluorescence analysis shows higher levels of PKC α , RhoA and ROCK II in the periphery vs. the cytosol of the IAS vs. RSM SMCs. **D.** Data show higher relax. of the IAS SMCs with Y 27632 as compared with calphostin C and Gö 6850.




Figure 4. WB analysis shows higher levels of RhoA, ROCKII, p-CPI17, p-MYPT1 and p-MLC $_{20}$ in IAS vs RSM. **Figure 5** WB analysis shows conc.-dependent decrease in p-MYPT1 by ROCK inhibitor Y 27632.




Figure 6. WB analysis shows conc.-dependent decrease in p-CPI-17 following Y 27632. **Figure 7.** WB analysis shows conc.-dependent decrease in p-MLC $_{20}$ following Y27632.




Figure 8. WB analysis shows significant decrease in p-CPI-17 following both Gö 6850 and Y 27632. **Figure 9.** Data show decrease in ROCK activity with Y 27632 closely correlate with the decreases in the IAS tone.





Figure 10. A, B, C. ROCK inhibitor (Y27632) nearly abolishes the basal tone in the IAS while PKC inhibitors Gö 6850 and calphostin C have limited effects.

MOLECULAR PATHWAY FOR BASAL IAS TONE



Conclusions

- RhoA/ROCK are constitutively active in the IAS, and this pathway (in contrast with PKC) is the critical determinant of the basal tone in the human IAS.
- Therefore, RhoA/ROCK are novel therapeutic targets for a number of rectoanal motility disorders in humans.