

PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Urine Chemical Content May Be a False Measure of Environmental Exposure

Robert L. Brent

Pediatrics 2013;132:e747; originally published online August 19, 2013;

DOI: 10.1542/peds.2013-2054

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/132/3/e747.full.html>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2013 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



Urine Chemical Content May Be a False Measure of Environmental Exposure

AUTHOR: Robert L. Brent, MD, PhD, DSc

Department of Pediatrics, Radiology, Pathology, Anatomy, and Cell Biology, Jefferson Medical College, Cornell University, Ithaca, New York; and Department of Obstetrics and Gynecology and Laboratory of Clinical and Environmental Teratology, Research Department, Alfred I. duPont Hospital for Children, Thomas Jefferson University, Wilmington, Delaware

KEY WORDS

bisphenol A, endocrine disruptors, urine chemical content as a false surrogate of exposure, DEHP

ABBREVIATIONS

BPA—bisphenol A

DEHP—di-2-ethylhexylphthalate

Opinions expressed in these commentaries are those of the authors and not necessarily those of the American Academy of Pediatrics or its Committees.

www.pediatrics.org/cgi/doi/10.1542/peds.2013-2054

doi:10.1542/peds.2013-2054

Accepted for publication Jul 2, 2013

Address correspondence to Robert L. Brent, MD, PhD, DSc, Thomas Jefferson University, Alfred I. DuPont Hospital for Children, Room 308 A/R, 1600 Rockland Rd, Wilmington, DE 19803. E-mail: rbrent@nemours.org

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2013 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The author has indicated he has no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The author has indicated he has no potential conflicts of interest to disclose.

COMPANION PAPERS: Companions to this article can be found on pages e637 and e646, online at www.pediatrics.org/cgi/doi/10.1542/peds.2013-0106 and www.pediatrics.org/cgi/doi/10.1542/peds.2012-4022.

FREE

This month *Pediatrics* has published 2 articles on environmental chemical exposures that warrant commentary, “Urinary Phthalates and Increased Insulin Resistance in Adolescents” (Trasande et al¹) and “Bisphenol A and Chronic Disease Risk Factors in US Children” (Eng et al²). Both of these articles use the urinary content of the chemical as a surrogate for the chemical exposure.

Dr Trasande and his group in the article on phthalates conclude that urine phthalate concentrations (represented by urine di-2-ethylhexylphthalate [DEHP] concentrations) were associated with increased insulin resistance in this cross-sectional study of adolescents. Dr Trasande refers to this as an association and indicated that this study cannot rule out the possibility that insulin-resistant children ingest food with higher phthalate content or that insulin-resistant children excrete more DEHP. Without knowing the tissue or serum concentration of DEHP or referring to primate studies, the authors’ hypotheses are appropriate.

The second article by Eng et al deals with bisphenol A (BPA). It is most important for authors to be familiar with the literature concerning the chemical that they are evaluating. Unfortunately, the BPA article fails to cite important publications dealing with the metabolism of BPA. For example, Daniel Doerge, a well-known authority on the metabolism of BPA, has co-written 5 publications that Eng et al have not cited.^{3–8} In 2011, in a report entitled, “Twenty-Four Hour Human Urine and Serum Profiles of Bisphenol A during High-Dietary Exposure”³ these investigators found that urinary BPA is not a surrogate of a serum or tissue exposure to BPA, but reflects the intake of BPA in the diet of the individual whose urine has been collected. This study, published in *Toxicological Sciences*, studied 20 adult volunteers who ingested 100% of 1 of 3 specified meals comprising of standard grocery store items for breakfast, lunch, and dinner. They studied the serum concentration of BPA, which was only measurable in the serum of the patients with exposure of 1.3 to 3.9 times higher than the 95th percentile of aggregate US exposure. “BPA serum concentrations were primarily less than or equal to the limit of detection...and were on average, 42 times lower than urine concentrations. During these high dietary exposures, BPA concentrations in serum were undetectable in 83% of the 320 samples collected and BPA concentrations were determined to be less than or equal to LOD [limit of detection] in all samples.”

Because there is little BPA in the serum even with large intakes of BPA and the metabolism of BPA in the serum destroys its endocrine receptor affinity, there is little justification to assume that BPA has toxic exposure in the human, even with very high exposure in the diet.

Most BPA articles in humans use the urine content of BPA as a surrogate of exposure, when in reality it is a measure of the intake of BPA

from the diet. The actual serum levels are extremely low and in many instances not detectable. Therefore, the studies that determine the effect of BPA by using urinary BPA levels are unlikely to be valid. Duty et al (2012)⁸ evaluated BPA in neonates in the nursery and found that newborns who were breastfed or formula bottle-fed had similar amounts of BPA in their urine. BPA in nursing bottles has been banned. Yet in reality, the BPA was not derived from the nursing bottle, but from the breast milk and formula milk in the bottle. We are fortunate that there is not an attempt to ban breastfeeding.

Another important article worth citing is the work of Dr Tyl,⁹ which is an excellent animal study evaluating oral effects of BPA entitled, "Three-Generation

Reproductive Toxicity Study of Dietary Bisphenol A in CD Sprague Dawley Rats." This is almost a perfect study for the following reasons^{10,11}: bisphenol was placed in the diet ad libitum among 7 exposure groups with a range of exposures that were over a 500 000-fold range. Only at the highest dosages were any potential toxic effects of BPA observed. It was not until the animals received 500 mg/kg that any adverse effects were observed, far higher than the highest reported experimental point previously used in animal studies.

Yamada et al¹² studied maternal serum and amniotic fluid BPA concentrations in the second trimester. The average concentration of BPA was 0.32 nM/mL that is far below the no-effect level that Dr Tyl determined in her animal stud-

ies. Although research dealing with the toxicity of BPA has indicated an increase of birth defects, neural behavioral defects, hypospadias, cryptorchidism, growth retardation, and prematurity, actually, it is unlikely that these levels in the serum of animals and humans have any possibility of producing these effects, and that using the urinary concentration of BPA as a surrogate of exposure can lead to serious misleading conclusions. Both of the studies published in this issue combined with the others that I have referenced in this commentary strongly suggest that using urine as a marker of an environmental toxin is not necessarily reflective of a significant exposure. Readers should use caution in interpreting more than the data actually indicate.

REFERENCES

1. Trasande L, Spanier AJ, Sathyanarayana S, Attina TM, Blustein J. Urinary phthalates and increased insulin resistance in adolescents. *Pediatrics*. 2013;132(3). Available at: www.pediatrics.org/cgi/content/full/132/3/e646
2. Eng DS, Gebremariam A, Meeker JD, Peterson K, Padmanabhan V, Lee JM. Bisphenol A and chronic disease risk factors in US children. *Pediatrics*. 2013;132(3). Available at: www.pediatrics.org/cgi/content/full/132/3/e637
3. Teeguarden JG, Calafat AM, Ye X, et al. Twenty-four hour human urine and serum profiles of bisphenol A during high-dietary exposure. *Toxicol Sci*. 2011;123(1):48–57
4. Doerge DR, Twaddle NC, Churchwell MI, Chang HC, Newbold RR, Delclos KB. Mass spectrometric determination of p-nonylphenol metabolism and disposition following oral administration to Sprague-Dawley rats. *Reprod Toxicol*. 2002;16(1):45–56
5. Doerge DR, Twaddle NC, Woodling KA, Fisher JW. Pharmacokinetics of bisphenol A in neonatal and adult rhesus monkeys. *Toxicol Appl Pharmacol*. 2010;248(1):1–11
6. Twaddle NC, Churchwell MI, Delclos KB, Newbold RR, Doerge DR. Determination using LC-ES/MS/MS of ethinylestradiol serum pharmacokinetics in adult Sprague-Dawley rats. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2003;793:309–315
7. Twaddle NC, Churchwell MI, Vanlandingham M, Doerge DR. Quantification of deuterated bisphenol A in serum, tissues, and excreta from adult Sprague-Dawley rats using liquid chromatography with tandem mass spectrometry. *Rapid Commun Mass Spectrom*. 2010;24(20):3011–3020
8. Duty S, Mendonca K, Hauser R, et al. Identification of potential sources of bisphenol A in the neonatal intensive care unit. *Pediatrics*. 2013;131(3):483–489
9. Tyl RW, Myers CB, Marr MC, et al. Three-generation reproductive toxicity study of dietary bisphenol A in CD Sprague-Dawley rats. *Toxicol Sci*. 2002;68(1):121–146
10. Brent RL. Utilization of animal studies to determine the effects and human risks of environmental toxicants (drugs, chemicals, and physical agents). *Pediatrics*. 2004;113(4 suppl):984–995
11. Brent RL, Weitzman M. The current state of knowledge about the effects, risks, and science of children's environmental exposures. *Pediatrics*. 2004;113(4 suppl):1158–1166
12. Yamada H, Furuta I, Kato EH, et al. Maternal serum and amniotic fluid bisphenol A concentrations in the early second trimester. *Reprod Toxicol*. 2002;16(6):735–739

Urine Chemical Content May Be a False Measure of Environmental Exposure

Robert L. Brent

Pediatrics 2013;132:e747; originally published online August 19, 2013;

DOI: 10.1542/peds.2013-2054

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/132/3/e747.full.html>

References

This article cites 10 articles, 3 of which can be accessed free at:
<http://pediatrics.aappublications.org/content/132/3/e747.full.html#ref-list-1>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://pediatrics.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://pediatrics.aappublications.org/site/misc/reprints.xhtml>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2013 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

