

POSTER PRESENTATION

Open Access

A Phase I study of AD5-GUCY2C-PADRE in stage I and II colon cancer patients

Adam Snook^{1*}, Trevor Baybutt¹, Michael Mastrangelo¹, Nancy Lewis¹, Scott Goldstein¹, Walter Kraft¹, Yaa Oppong¹, Terry Hyslop², Ronald Myers¹, Vitali Alexeev¹, Laurence Eisenlohr¹, Takami Sato¹, Scott Waldman¹

From 30th Annual Meeting and Associated Programs of the Society for Immunotherapy of Cancer (SITC 2015) National Harbor, MD, USA. 4-8 November 2015

Background

Ad5-GUCY2C-PADRE is a replication-deficient human type 5 recombinant adenovirus (Ad5) vaccine encoding guanylyl cyclase C (GUCY2C) fused to the PAn DR Epitope (PADRE). GUCY2C, a paracrine hormone receptor producing the second messenger cyclic GMP (cGMP), is selectively expressed by intestinal epithelial cells and a subset of hypothalamic neurons, but not other tissues. Importantly, GUCY2C is over-expressed in nearly all primary and metastatic human colorectal tumors. Preclinical studies in mice demonstrated selective tolerance of GUCY2C-specific CD4+ T cells, but not CD8+ T or B cells, necessitating inclusion of the exogenous CD4+ T helper cell epitope PADRE to maximize GUCY2C-specific CD8+ T cell and antibody responses and antitumor efficacy, without autoimmunity.

Patients and methods

This is an open-label, single arm “proof-of-concept” study evaluating a single dose level of Ad5-GUCY2C-PADRE as a vaccine for surgically-treated, node-negative colon cancer subjects (NCT01972737). Patients received a single intramuscular administration of 10¹¹ Ad5-GUCY2C-PADRE viral particles. Safety and immunomonitoring were examined at 30, 90 and 180 days following vaccination. Primary objectives were to determine the safety, tolerability and toxicity of Ad5-GUCY2C-PADRE and to determine whether Ad5-GUCY2C-PADRE induces GUCY2C-specific immune responses. The study employed a joint efficacy-toxicity design and included stopping rules for either efficacy or toxicity. Results here were obtained during the planned interim analysis following accrual of 10 subjects.

Results

The vaccine was well tolerated, producing only mild adverse events (AEs). Short-lived injection site pain/swelling, body aches and chills were the most commonly observed AEs and occurred in 30-40% of subjects. GUCY2C-specific antibody and T-cell responses were observed in a subset of subjects. Consistent with preclinical mouse data, T-cell responses were composed of CD8+, but not CD4+, T cells. Importantly, GUCY2C-specific responses occurred only in subjects with low Ad5 neutralizing antibody (NAb) titers at the time of vaccination, suggesting that pre-existing Ad5 immunity limits Ad5-GUCY2C-PADRE immunogenicity.

Conclusions

Interim analysis of 10 subjects receiving Ad5-GUCY2C-PADRE demonstrates proof-of-concept that GUCY2C is immunogenic in humans and that GUCY2C-directed vaccination is safe. Moreover, the presence of GUCY2C-specific antibody and CD8+ T-cell, but not CD4+ T-cell, responses is consistent with selective CD4+ T-cell tolerance observed in mouse models. These data establish GUCY2C as a safe and immunogenic target for immunotherapy in cancer patients.

Trial registration

ClinicalTrials.gov identifier NCT01972737.

Authors' details

¹Thomas Jefferson University, Philadelphia, PA, USA. ²Duke University, Durham, NC, USA.

¹Thomas Jefferson University, Philadelphia, PA, USA
Full list of author information is available at the end of the article

Published: 4 November 2015

doi:10.1186/2051-1426-3-S2-P450

Cite this article as: Snook *et al.*: A Phase I study of AD5-GUCY2C-PADRE in stage I and II colon cancer patients. *Journal for ImmunoTherapy of Cancer* 2015 **3**(Suppl 2):P450.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

