



2-12-2016

Effects of High Intensity Exercise on Central Neural Drive in Healthy Populations

Nicholas LeGrand, SPT

Department of Physical Therapy, Thomas Jefferson University, njl005@jefferson.edu

Kyra Robb, SPT

Department of Physical Therapy, Thomas Jefferson University, kxr096@jefferson.edu

Emily Slobodian, SPT

Department of Physical Therapy, Thomas Jefferson University, exs049@jefferson.edu

Nicole Weaknecht, SPT

Department of Physical Therapy, Thomas Jefferson University, new003@jefferson.edu

Megan Wood, SPT

Department of Physical Therapy, Thomas Jefferson University, mmw009@jefferson.edu

See next page for additional authors

Follow this and additional works at: <http://jdc.jefferson.edu/dptcapstones>

 Part of the [Physical Therapy Commons](http://jdc.jefferson.edu/dptcapstones)

Recommended Citation

LeGrand, SPT, Nicholas; Robb, SPT, Kyra; Slobodian, SPT, Emily; Weaknecht, SPT, Nicole; Wood, SPT, Megan; and Muth, PT, PhD, Stephanie, "Effects of High Intensity Exercise on Central Neural Drive in Healthy Populations" (2016). *Department of Physical Therapy Capstone Posters*. 9.

<http://jdc.jefferson.edu/dptcapstones/9>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](http://www.jefferson.edu/ctl). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Physical Therapy Capstone Posters by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Nicholas LeGrand, SPT; Kyra Robb, SPT; Emily Slobodian, SPT; Nicole Weaknecht, SPT; Megan Wood, SPT; and Stephanie Muth, PT, PhD

Background

Current research suggests that strength gains related to central neural adaptation occur more rapidly than peripheral mechanisms⁵. Central neural drive (CND) is the measure of cortical output that coordinates up-regulation of agonist contraction and inhibition of antagonist musculature in voluntary muscle contractions². Two common techniques for measuring CND are twitch interpolation and motor evoked potential from transcranial magnetic stimulation.

Methods

Literature search performed June 2015

Databases searched: PubMed, OVID, CINAHL, Cocharan, Scopus

Keywords: “motor drive and exercise”, “motor evoked potential and exercise”, “voluntary activation and exercise”, “motor drive and high intensity”, “voluntary activation and high intensity”, “motor evoked potential and high intensity”

Inclusion Criteria: high intensity strength training, protocols including multiple sessions over time

Total articles: n= 1692

Abstract review: n= 201

Brief article review:
n= 27

Critical appraisal: n= 10

Articles for Review: n= 5

Results

Citation	MacDermid score ⁸ / Sackett rating	Participants	Training Protocol	Instrumentation	Key findings
Carroll, 2009	28/48 1B	17 participants aged 19-35 without a significant history of neurologic disease.	Control group- non resistance movements only Training group- 4 weeks of traditional radial deviation and wrist extensor high -intensity strength training.	Electromyography and Transcranial Magnetic Stimulation	Training group showed a significant increase in TMS-induced twitches in both wrist extension and radial deviation at the end of 4 weeks
Ekblom, 2010	28/48 1B	20 healthy participants (12M 8W) with no history of resistance training or exercise >3days/week	Control group- no change in daily activity Training group- 3 times a week high intensity eccentric plantar flexor training for 5 weeks	Twitch interpolation	Training group demonstrated improved central neural drive and increased plantar flexor strength at the end of 5 weeks
Goodwill, 2012	34/48 1B	14 university participants (7 men, 7 women) age range 18-35	Control group-no training Training group- unilateral strength training sessions over 3 weeks for the right quadriceps	Transcranial Magnetic Stimulation	Training group showed significant increases in strength, decreases in SICI, and increases in cortico-motor excitability in both the trained and untrained limb at the end of 3 weeks
Kidgell, 2010	32/48 1B	23 healthy participants (10 men, 13 women, 26.8 +/- 7.3)	Control group-no training Training group-4 week progressive overload strength training of the right biceps brachii	Electromyography and Transcranial Magnetic Stimulation	Training group showed significant increase in 1RM strength in absence of muscle hypertrophy. Increase in MEP amplitude at and above AMT at the end of 4 weeks
Pucci, 2006	29/48 1B	20 male participants (25+/- 5.5) from the university population	Control group-no training Training group-3 week quadriceps isometric training program	Twitch interpolation	Training group demonstrated significant increase in knee extensor MVC and an increase in percentage maximal activation after 4-5 training days. Insignificant increase in twitch amplitude by day 9

Summary of Key Findings

These studies demonstrated strength increases in healthy individuals as early as 10 days and within 5 weeks of high intensity strength training without muscle hypertrophy. TMS and twitch interpolation techniques provide evidence that these strength gains can be attributed to central mechanisms.

Clinical Applications


For patients with an inability to train both sides of the body, there is potential for an overflow effect into the immobilized side when the non-affected side is trained.

High intensity training may be used to capitalize on rapid cortical adaptations in patients that need strength gains in a limited time frame (e.g. preoperatively).


Additional research is needed to further explore the effects of high intensity exercise on CND, including investigation of effects in neuro-compromised individuals.

Conclusions

Current understanding of neural adaptation suggests a central component to muscle activation separate from peripheral mechanisms. Moderate evidence suggests that high intensity exercise results in rapid increases in strength through central mechanisms in the absence of muscle hypertrophy.



Twitch interpolation:
A supramaximal electrical stimulus is applied during a voluntary contraction. Those motor units that have not already been recruited generate a twitch response demonstrating untapped potential for motor unit recruitment⁹.



Transcranial magnetic stimulation:
Magnetic stimulation is applied over the primary motor cortex eliciting a peripheral motor response termed motor evoked potential (MEP). This response is a measure of cortical excitability⁹.

References

1. Burns PB, Rohrich RJ, Chung KC. The levels of Evidence and their role in Evidence-Based Medicine. *Plastic and reconstructive surgery*. 2011; 128(1):305-310. doi:10.1097/PRS.0b013e318219c171.
2. Carroll, T.J., et al. "The Effect of Strength Training on the Force of Twitches by Corticospinal Stimulation in Humans." *Acta Physiologica* 197.2 (2009): 161-73. Web.
3. Ekblom, M. M. "Improvements in Dynamic Plantar Flexor Strength After Resistance Training are Associated with Increased Voluntary Activation and V-to-M Ratio." *Journal of applied physiology* 109.1 (2010): 1926.
4. Goodwill, Alicia M.B. Sc, H.O.N.S., Alan J. Pearce, and Dawson J, Kidgell. "Corticospinal Plasticity Following Unilateral Strength Training." *Muscle & nerve* 46.3 (2012): 384-93. Web
5. Kidgell, Dawson J., et al. "Neurophysiological Responses After Short-Term Strength Training of the Biceps Brachii Muscle." *Journal of Strength & Conditioning Research* 24.11 (2010): 3123-32. Web.
6. Pucci, A. R., Griffin, and E. Cafarelli. "Maximal Motor Unit Firing Rates during Isometric Resistance Training in Men." *Experimental physiology* 91.1 (2006): 171-8. Web.
7. Folland, Jonathan P, and Alun G Williams. "The Adaptations To Strength Training." *Sports Medicine* 37.2 (2007): 145-68. Web.
8. Macdermid, Joy. "An Introduction to Evidence-based Practice for Hand Therapists." *Journal of Hand Therapy* 17 (2004): 105-17
9. Shield, Anthony, and Shi Zhou. "Assessing Voluntary Muscle Activation With The Twitch Interpolation Technique." *Sports Medicine* 34.4 (2004): 253-67. Web.
10. Taylor, Janet L. "Point:Counterpoint: The Interpolated Twitch does/does not Provide a Valid Measure of the Voluntary Activation of Muscle." *Applied Physiology*. 107 (2009): 1354-355. Web

Abbreviations:

HIST - High Intensity Strength Training
MVC - Max Volitional Contraction
CND - Central Neural Drive
TMS - Transcranial Magnetic Stimulation
MEP - Motor Evoked Potential
SICI - Short-latency Intra-Cortical Inhibition
AMT - Active Muscle Threshold