

2-25-2016

Microcystic, Elongated, and Fragmented (MELF) Pattern Invasion in Ovarian Endometrioid Carcinoma: Immunohistochemical Profile and Prognostic Implications

Allison F. Goldberg, MD

Thomas Jefferson University, Allison.Goldberg@jefferson.edu

Lauren Cooper, MD

Thomas Jefferson University, lauren.hand@jefferson.edu


D. De Cotiis

University of Pennsylvania

Norman G. Rosenblum, MD, PhD

Thomas Jefferson University, Norman.Rosenblum@jefferson.edu

Joanna Chan, MD

*Department of Pathology, Jefferson Medical College of Thomas Jefferson University, Philadelphia, PA,
joanna.chan@jefferson.edu*Follow this and additional works at: <http://jdc.jefferson.edu/pacresidentposters> Part of the [Medical Anatomy Commons](#), [Medical Cell Biology Commons](#), and the [Medical Pathology Commons](#)

Recommended Citation

Goldberg, MD, Allison F.; Cooper, MD, Lauren; Cotiis, D. De; Rosenblum, MD, PhD, Norman G.; and Chan, MD, Joanna, "Microcystic, Elongated, and Fragmented (MELF) Pattern Invasion in Ovarian Endometrioid Carcinoma: Immunohistochemical Profile and Prognostic Implications" (2016). *Pathology, Anatomy and Cell Biology Resident's Posters*. Paper 8.
<http://jdc.jefferson.edu/pacresidentposters/8>

Microcystic, Elongated, and Fragmented (MELF) Pattern Invasion in Ovarian Endometrioid Carcinoma: Immunohistochemical Profile and Prognostic Implications

AF Goldberg¹, L Cooper², D De Cotiis³, N Rosenblum², JSY Chan¹

¹Department of Pathology, Anatomy, and Cell Biology, Thomas Jefferson University Hospital, Philadelphia, PA

²Department of Obstetrics and Gynecology, Thomas Jefferson University Hospital, Philadelphia, PA


³Department of Radiology, University of Pennsylvania, Philadelphia PA

BACKGROUND

- Microcystic, Elongated and Fragmented (MELF) is a well-recognized pattern of uterine endometrioid carcinoma (UEC) associated with lymphovascular space invasion and occult lymph node metastasis
- MELF in UEC may be seen with Lynch Syndrome
- MELF in UEC is hypothesized to be histologic evidence of an epithelial mesenchymal transition
- MELF pattern invasion in ovarian endometrioid carcinoma (OEC) was first described at USCAP 2015
- Current study evaluates MELF in OEC for
 - Prognostic implications
 - Immunohistochemical (IHC) profile related to
 - Lynch Syndrome
 - Epithelial mesenchymal transition

DESIGN

- 42 consecutive cases of OEC without concurrent UEC (1996-2014) evaluated by 2 pathologists
- MELF defined as at least three glands fulfilling histologic criteria
- 32 cases had blocks available for staining
 - MLH1, PMS2, MSH2 and MSH6 for mismatch repair (MMR) protein expression
 - Graded as “retained” or “lost”
 - β -catenin, e-cadherin, CK19 and cyclin D1 for evidence of epithelial mesenchymal transition
 - Graded as “rare” (<25% cells stain), “moderate” (25-75% cells stain), or “strong” (>75% cells stain)
- Retrospective chart review of clinical and demographic features and overall survival
- Data analyzed using Fisher exact test analysis
- Survival analyzed using Kaplan-Meier method



RESULTS

- MELF pattern invasion was identified in 45% of the cases reviewed
- Clear cell features were only seen in cases with MELF pattern invasion (p-value=0.044)
- Overall, 13% of cases demonstrate MMR protein loss
 - MELF: MSH2/MSH6 deficiency (n=2)
 - Non-MELF: PMS2 deficiency (n=2)
- No difference was identified in:
 - Overall survival
 - Cancer recurrence
 - IHC staining for β -catenin, e-cadherin, CK19 and cyclin D1
 - Serous features
 - Concurrent endometriosis
 - Lymphovascular space invasion
 - Lymph node metastasis
 - Bilaterality of disease
 - Extranodal metastasis

CONCLUSIONS

- MELF occurs in ovarian endometrioid carcinoma at a similar or higher frequency than in uterine endometrioid carcinoma.
- Clear cell features were identified exclusively in MELF pattern invasion cases.
- Different MMR proteins are lost in MELF and non-MELF pattern invasion carcinomas.
- As there is no current consensus on Lynch screening in patients with ovarian endometrioid carcinoma, perhaps the presence of MELF pattern invasion should prompt screening.
- MELF should be considered when assessing ovarian endometrioid carcinoma, as the pattern may be confused with endometriosis or endosalpingiosis.