



## Trends in urinary incontinence in women between 4 and 24months postpartum in the EDEN cohort.

Emeline Quiboef, Marie-Josèphe Saurel-Cubizolles, Xavier Fritel

► **To cite this version:**

Emeline Quiboef, Marie-Josèphe Saurel-Cubizolles, Xavier Fritel. Trends in urinary incontinence in women between 4 and 24months postpartum in the EDEN cohort.: Urinary incontinence between 4 and 24 months postpartum. *BJOG: An International Journal of Obstetrics and Gynaecology*, Wiley, 2015, <10.1111/1471-0528.13545>. <inserm-01185917>

**HAL Id: inserm-01185917**

**<http://www.hal.inserm.fr/inserm-01185917>**

Submitted on 24 Aug 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

- 1 Figure 2  
 2 Continence status 4-24 months postpartum (N= 1354):  
 3 A) Remain continent: women who were continent at baseline (4 months) and follow-up (24  
 4 months);  
 5 B) Remain incontinent: women incontinent at baseline and follow-up;  
 6 C) Become incontinent: women continent at baseline who became incontinent at follow-up;  
 7 D) Become continent: women incontinent at baseline who became continent at follow-up.  
 8

