



ON THE SYSTEM OF PARTIAL DIFFERENTIAL EQUATIONS ARISING IN MEAN FIELD TYPE CONTROL

Yves Achdou, Mathieu Laurière

► **To cite this version:**

Yves Achdou, Mathieu Laurière. ON THE SYSTEM OF PARTIAL DIFFERENTIAL EQUATIONS ARISING IN MEAN FIELD TYPE CONTROL. 2015. <hal-01132008>

HAL Id: hal-01132008

<https://hal.archives-ouvertes.fr/hal-01132008>

Submitted on 16 Mar 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON THE SYSTEM OF PARTIAL DIFFERENTIAL EQUATIONS ARISING IN MEAN FIELD TYPE CONTROL

ABSTRACT. We discuss the system of Fokker-Planck and Hamilton-Jacobi-Bellman equations arising from the finite horizon control of McKean-Vlasov dynamics. We give examples of existence and uniqueness results. Finally, we propose some simple models for the motion of pedestrians and report about numerical simulations in which we compare mean field games and mean field type control.

YVES ACHDOU

Université Paris Diderot
Laboratoire Jacques-Louis Lions, UMR 7598, UPMC, CNRS
Sorbonne Paris Cité F-75205 Paris, France

MATHIEU LAURIÈRE

Université Paris Diderot
Laboratoire Jacques-Louis Lions, UMR 7598, UPMC, CNRS
Sorbonne Paris Cité F-75205 Paris, France

(Communicated by the associate editor name)

1. Introduction. In the recent years, an important research activity has been devoted to the study of stochastic differential games with a large number of players. In their pioneering articles [11, 12, 13], J-M. Lasry and P-L. Lions have introduced the notion of mean field games, which describe the asymptotic behavior of stochastic differential games (Nash equilibria) as the number N of players tends to infinity. In these models, it is assumed that the agents are all identical and that an individual agent can hardly influence the outcome of the game. Moreover, each individual strategy is influenced by some averages of functions of the states of the other agents. In the limit when $N \rightarrow +\infty$, a given agent feels the presence of the other agents through the statistical distribution of the states of the other players. Since perturbations of a single agent's strategy does not influence the statistical distribution of the states, the latter acts as a parameter in the control problem to be solved by each agent.

Another kind of asymptotic regime is obtained by assuming that all the agents use the same distributed feedback strategy and by passing to the limit as $N \rightarrow \infty$ before optimizing the common feedback. Given a common feedback strategy, the asymptotics are given by the McKean-Vlasov theory, [16, 20] : the dynamics of a given agent is found by solving a stochastic differential equation with coefficients depending on a mean field, namely the statistical distribution of the states, which may also affect the objective function. Since the feedback strategy is common to all agents, perturbations of the latter affect the mean field. Then, having each

2010 *Mathematics Subject Classification.* Primary: 49J20; Secondary: 35K55.

Key words and phrases. mean field type control, existence and uniqueness.

player optimize its objective function amounts to solving a control problem driven by the McKean-Vlasov dynamics. The latter is named control of McKean-Vlasov dynamics by R. Carmona and F. Delarue [8, 7] and mean field type control by A. Bensoussan et al, [4, 5].

When the dynamics of the players are independent stochastic processes, both mean field games and control of McKean-Vlasov dynamics naturally lead to a coupled system of partial differential equations, a forward Fokker-Planck equation (which may be named FP equation in the sequel) and a backward Hamilton-Jacobi-Bellman equation (which may be named HJB equation). For mean field games, the coupled system of partial differential equations has been studied by Lasry and Lions in [11, 12, 13]. Besides, many important aspects of the mathematical theory developed by J-M. Lasry and P-L. Lions on MFG are not published in journals or books, but can be found in the videos of the lectures of P-L. Lions at Collège de France: see the web site of Collège de France, [15]. One can also see [9] for a brief survey.

In the present paper, we aim at studying the system of partial differential equations arising in mean field type control, when the horizon of the control problem is finite: we will discuss the existence and the uniqueness of classical solutions. In the last paragraph of the paper, we briefly discuss some numerical simulations in the context of motion of pedestrians, and we compare the results obtained with mean field games and with mean field type control.

1.1. Model and assumptions. For simplicity, we assume that all the functions used below (except in § 4) are periodic with respect to the state variables x_i , $i = 1, \dots, d$, of period 1 for example. This will save technical arguments on either problems in unbounded domains or boundary conditions. We denote by \mathbb{T}^d the d -dimensional unit torus: $\mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d$. Let \mathbb{P} be the set of probability measures on \mathbb{T}^d and $\mathbb{P} \cap L^1(\mathbb{T}^d)$ be the set of probability measures which are absolutely continuous with respect to the Lebesgue measure. For $m \in \mathbb{P} \cap L^1(\mathbb{T}^d)$, the density of m with respect to the Lebesgue measure will be still be noted m , i.e. $dm(x) = m(x)dx$. Let g be a map from \mathbb{P} to a subset of $\mathcal{C}^1(\mathbb{T}^d \times \mathbb{R}^n; \mathbb{R}^d)$ (the image of $m \in \mathbb{P}$ will be noted $g[m] \in \mathcal{C}^1(\mathbb{T}^d \times \mathbb{R}^n; \mathbb{R}^d)$) such that

- there exists a constant M such that for all $m \in \mathbb{P}$ and $x \in \mathbb{T}^d$, $|g[m](x, 0)| \leq M$
- there exists a constant L such that
 - for all $m \in \mathbb{P}$, $a \in \mathbb{R}^n$ and $x, y \in \mathbb{T}^d$, $|g[m](x, a) - g[m](y, a)| \leq Ld(x, y)$ where $d(x, y)$ is the distance between x and y in \mathbb{T}^d .
 - for all $m \in \mathbb{P}$, $a, b \in \mathbb{R}^n$ and $x \in \mathbb{T}^d$, $|g[m](x, a) - g[m](x, b)| \leq L|a - b|$
 - for all $m, m' \in \mathbb{P}$, $a \in \mathbb{R}^n$ and $x \in \mathbb{T}^d$, $|g[m](x, a) - g[m'](x, a)| \leq Ld_2(m, m')$ where d_2 is the Wasserstein distance:

$$d_2(m, m') \equiv \inf_{\gamma \in \Gamma(m, m')} \left(\int_{\mathbb{T}^d \times \mathbb{T}^d} d^2(x, y) d\gamma(x, y) \right)^{\frac{1}{2}},$$

$$\Gamma(m, m') \equiv \left\{ \gamma : \text{transport plan between } m \text{ and } m' \right\},$$

and a transport plan γ between m and m' is a Borel probability measure on $\mathbb{T}^d \times \mathbb{T}^d$ such that, for all Borel subset E of \mathbb{T}^d ,

$$\gamma(E \times \mathbb{T}^d) = m(E) \quad \text{and} \quad \gamma(\mathbb{T}^d \times E) = m'(E).$$

- there exists a map \tilde{g} from $L^1(\mathbb{T}^d)$ to $\mathcal{C}^1(\mathbb{T}^d \times \mathbb{R}^n; \mathbb{R}^d)$ such that $g|_{\mathbb{P} \cap L^1(\mathbb{T}^d)} = \tilde{g}|_{\mathbb{P} \cap L^1(\mathbb{T}^d)}$ and that for any $x \in \mathbb{T}^d$ and $a \in \mathbb{R}^n$, $m \rightarrow \tilde{g}[m](x, a)$ is Fréchet differentiable in $L^1(\mathbb{T}^d)$ and $(x, a) \mapsto \frac{\partial \tilde{g}}{\partial m}[m](x, a)$ belongs to

$\mathcal{C}^1(\mathbb{T}^d \times \mathbb{R}^n; L^\infty(\mathbb{T}^d; \mathbb{R}^d))$. Hereafter, we will not make the distinction between g and \tilde{g} .

Consider a probability space $(\Omega, \mathcal{A}, \mathcal{P})$ and a filtration \mathcal{F}^t generated by a d -dimensional standard Wiener process (W_t) and the stochastic process $(X_t)_{t \in [0, T]}$ in \mathbb{R}^d adapted to \mathcal{F}^t which solves the stochastic differential equation

$$dX_t = g[m_t](X_t, a_t) dt + \sqrt{2\nu} dW_t \quad \forall t \in [0, T], \quad (1.1)$$

given the initial state X_0 which is a random variable \mathcal{F}^0 -measurable whose probability density is noted m_0 . In (1.1), ν is a positive number, m_t is the probability distribution of X_t and a_t is the control which we take to be

$$a_t = v(t, X_t), \quad (1.2)$$

where $v(t, \cdot)$ is a continuous function on \mathbb{T}^d . To the pair (v, m) , we associate the objective

$$J(v, m) := \mathbb{E} \left[\int_0^T f[m_t](X_t, a_t) dt + h[m_T](X_T) \right] \quad (1.3)$$

where f (resp. h) is a map from \mathbb{P} to a subset of $\mathcal{C}^1(\mathbb{T}^d \times \mathbb{R}^n)$, resp. to a subset of $\mathcal{C}^1(\mathbb{T}^d)$. We assume that

- $\lim_{|a| \rightarrow \infty} \inf_{m \in \mathbb{P}, x \in \mathbb{T}^d} \frac{f[m](x, a)}{|a|} = +\infty$
- there exists a map \tilde{f} from $L^1(\mathbb{T}^d)$ to $\mathcal{C}^1(\mathbb{T}^d \times \mathbb{R}^n)$ such that $f|_{\mathbb{P} \cap L^1(\mathbb{T}^d)} = \tilde{f}|_{\mathbb{P} \cap L^1(\mathbb{T}^d)}$ and that for any $x \in \mathbb{T}^d$ and $a \in \mathbb{R}^n$, $m \rightarrow \tilde{f}[m](x, a)$ is Fréchet differentiable in $L^1(\mathbb{T}^d)$ and $(x, a) \mapsto \frac{\partial \tilde{f}}{\partial m}[m](x, a)$ belongs to $\mathcal{C}^1(\mathbb{T}^d \times \mathbb{R}^n; L^\infty(\mathbb{T}^d))$. Hereafter, we will not make the distinction between f and \tilde{f} .

We also assume that there exists a map \tilde{h} from $L^1(\mathbb{T}^d)$ to $\mathcal{C}^1(\mathbb{T}^d)$ such that $h|_{\mathbb{P} \cap L^1(\mathbb{T}^d)} = \tilde{h}|_{\mathbb{P} \cap L^1(\mathbb{T}^d)}$ and that for any $x \in \mathbb{T}^d$, $m \rightarrow \tilde{h}[m](x)$ is Fréchet differentiable in $L^1(\mathbb{T}^d)$ and $x \mapsto \frac{\partial \tilde{h}}{\partial m}[m](x)$ belongs to $\mathcal{C}^1(\mathbb{T}^d; L^\infty(\mathbb{T}^d))$. Hereafter, we will not make the distinction between h and \tilde{h} .

It will be useful to define the Lagrangian and Hamiltonian as follows: for any $x \in \mathbb{T}^d$, $a \in \mathbb{R}^n$ and $p \in \mathbb{R}^d$,

$$\begin{aligned} L[m](x, a, p) &:= f[m](x, a) + p \cdot g[m](x, a) \\ H[m](x, p) &:= \min_{a \in \mathbb{R}^n} L[m](x, a, p). \end{aligned}$$

where $p \cdot q$ denotes the scalar product in \mathbb{R}^d .

It is consistent with the previous assumptions to suppose that

- there exists a map \tilde{H} from $L^1(\mathbb{T}^d)$ to $\mathcal{C}(\mathbb{T}^d \times \mathbb{R}^d)$ such that $H|_{\mathbb{P} \cap L^1(\mathbb{T}^d)} = \tilde{H}|_{\mathbb{P} \cap L^1(\mathbb{T}^d)}$ and that for any $x \in \mathbb{T}^d$ and $p \in \mathbb{R}^d$, $m \rightarrow \tilde{H}[m](x, p)$ is Fréchet differentiable in $L^1(\mathbb{T}^d)$ and $(x, p) \mapsto \frac{\partial \tilde{H}}{\partial m}[m](x, p)$ belongs to $\mathcal{C}^1(\mathbb{T}^d \times \mathbb{R}^d; L^\infty(\mathbb{T}^d))$. We will not make the distinction between H and \tilde{H} .
- if $m \in \mathbb{P} \cap L^1(\mathbb{T}^d)$ and $a^* = \operatorname{argmin}_a f[m](x, a) + p \cdot g[m](x, a)$, then

$$\frac{\partial H}{\partial m}(x, p) = \frac{\partial f}{\partial m}(x, a^*) + p \cdot \frac{\partial g}{\partial m}(x, a^*).$$

As explained in [5], page 13, if the feedback function v is smooth enough and if $m_0 \in \mathbb{P} \cap L^1(\mathbb{T}^d)$, then the probability distribution $m_v(t, \cdot)$ has a density with

respect to the Lebesgue measure, $m_v(t, \cdot) \in \mathbb{P} \cap L^1(\mathbb{T}^d)$ for all t , and its density m_v is solution of the Fokker-Planck equation

$$\frac{\partial m_v}{\partial t}(t, x) - \nu \Delta m_v(t, x) + \operatorname{div} \left(m_v(t, \cdot) g[m_v(t, \cdot)](\cdot, v(t, \cdot)) \right)(x) = 0, \quad t \in (0, T], x \in \mathbb{T}^d, \quad (1.4)$$

with the initial condition

$$m_v(0, x) = m_0(x), \quad x \in \mathbb{T}^d. \quad (1.5)$$

Therefore, the control problem consists of minimizing

$$J(v, m_v) = \int_{[0, T] \times \mathbb{T}^d} f[m_v(t, \cdot)](x, v(t, x)) m_v(t, x) dx dt + \int_{\mathbb{T}^d} h[m_v(T, \cdot)](x) m_v(T, x) dx,$$

subject to (1.4)-(1.5). In [5], A. Bensoussan, J. Frehse and P. Yam have proved that a necessary condition for the existence of a smooth feedback function v^* achieving $J(v^*, m_{v^*}) = \min J(v, m_v)$ is that

$$v^*(t, x) = \operatorname{argmin}_v \left(f[m(t, \cdot)](x, v) + \nabla u(t, x) \cdot g[m(t, \cdot)](x, v) \right),$$

where (m, u) solve the following system of partial differential equations

$$0 = \frac{\partial u}{\partial t}(t, x) + \nu \Delta u(t, x) + H[m(t, \cdot)](x, \nabla u(t, x)) \quad (1.6)$$

$$+ \int_{\mathbb{T}^d} \frac{\partial H}{\partial m}[m(t, \cdot)](\xi, \nabla u(t, \xi))(x) m(t, \xi) d\xi,$$

$$0 = \frac{\partial m}{\partial t}(t, x) - \nu \Delta m(t, x) + \operatorname{div} \left(m(t, \cdot) \frac{\partial H}{\partial p}[m(t, \cdot)](\cdot, \nabla u(t, \cdot)) \right)(x), \quad (1.7)$$

with the initial and terminal conditions

$$m(0, x) = m_0(x) \quad \text{and} \quad u(T, x) = h[m(T, \cdot)](x) + \int_{\mathbb{T}^d} \frac{\partial h}{\partial m}[m(T, \cdot)](\xi)(x) m(T, \xi) d\xi. \quad (1.8)$$

It will be useful to write

$$G[m, q](x) := \int_{\mathbb{T}^d} m(\xi) \frac{\partial}{\partial m} H[m](\xi, q(\xi))(x) d\xi \quad (1.9)$$

for functions $m \in \mathbb{P} \cap L^1(\mathbb{T}^d)$ and $q \in \mathcal{C}(\mathbb{T}^d; \mathbb{R}^d)$, so that (1.6) can be written

$$0 = \frac{\partial u}{\partial t}(t, x) + \nu \Delta u(t, x) + H[m(t, \cdot)](x, \nabla u(t, x)) + G[m(t, \cdot), \nabla u(t, \cdot)](x).$$

Remark 1. Note the difference with the system of partial differential equations arising in mean field games, namely

$$0 = \frac{\partial u}{\partial t}(t, x) + \nu \Delta u(t, x) + H[m(t, \cdot)](x, \nabla u(t, x)), \quad (1.10)$$

$$0 = \frac{\partial m}{\partial t}(t, x) - \nu \Delta m(t, x) + \operatorname{div} \left(m(t, \cdot) \frac{\partial H}{\partial p}[m(t, \cdot)](\cdot, \nabla u(t, \cdot)) \right)(x), \quad (1.11)$$

with the initial and terminal conditions

$$m(0, x) = m_0(x) \quad \text{and} \quad u(T, x) = h[m(T, \cdot)](x). \quad (1.12)$$

Both the HJB equation (1.6) and the terminal condition on u in (1.8) involve additional nonlocal terms, which account for the variations of m_v caused by variations of the common feedback v .

Remark 2. At least formally, it is possible to consider situations when H and h depend locally on m , i.e. $H[m](x, p) = \tilde{H}(x, p, m(x))$ and $h[m](x) = \tilde{h}(x, m(x))$: in this case, (1.6)-(1.8) become

$$0 = \frac{\partial u}{\partial t}(t, x) + \nu \Delta u(t, x) + \tilde{H}(x, \nabla u(t, x), m(t, x)) + m(t, x) \frac{\partial \tilde{H}}{\partial m}(x, \nabla u(t, x), m(t, x)), \quad (1.13)$$

$$0 = \frac{\partial m}{\partial t}(t, x) - \nu \Delta m(t, x) + \operatorname{div} \left(m(t, \cdot) \frac{\partial \tilde{H}}{\partial p}(\cdot, \nabla u(t, \cdot), m(t, \cdot)) \right) (x), \quad (1.14)$$

with the initial and terminal conditions

$$m(0, x) = m_0(x) \quad \text{and} \quad u(T, x) = \tilde{h}(x, m(T, x)) + m(T, x) \frac{\partial \tilde{h}}{\partial m}(x, m(T, x)). \quad (1.15)$$

2. Existence results. We focus on the system (1.6)-(1.8). We are going to state existence results in some typical situations.

2.1. Notations. Let Q be the open set $Q := (0, T) \times \mathbb{T}^d$. We shall need to use spaces of Hölder functions in Q : For $\alpha \in (0, 1)$, the space of Hölder functions $\mathcal{C}^{\alpha/2, \alpha}(\bar{Q})$ is classically defined by

$$\mathcal{C}^{\alpha/2, \alpha}(\bar{Q}) := \left\{ w \in \mathcal{C}(\bar{Q}) : \exists C > 0 \text{ s.t. } \forall (t_1, x_1), (t_2, x_2) \in \bar{Q}, \right. \\ \left. |w(t_1, x_1) - w(t_2, x_2)| \leq C (d(x_1, x_2)^2 + |t_1 - t_2|)^{\alpha/2} \right\}$$

and we define

$$|w|_{\mathcal{C}^{\alpha/2, \alpha}(\bar{Q})} := \sup_{(t_1, x_1) \neq (t_2, x_2) \in \bar{Q}} \frac{|w(t_1, x_1) - w(t_2, x_2)|}{(d(x_1, x_2)^2 + |t_1 - t_2|)^{\alpha/2}}$$

and $\|w\|_{\mathcal{C}^{\alpha/2, \alpha}(\bar{Q})} := \|w\|_{\mathcal{C}(\bar{Q})} + |w|_{\mathcal{C}^{\alpha/2, \alpha}(\bar{Q})}$. Then the space $\mathcal{C}^{(1+\alpha)/2, 1+\alpha}(\bar{Q})$ is made of all the functions $w \in \mathcal{C}(\bar{Q})$ which have partial derivatives $\frac{\partial w}{\partial x_i} \in \mathcal{C}^{\alpha/2, \alpha}(\bar{Q})$ for all $i = 1, \dots, d$ and such that for all $(t_1, x) \neq (t_2, x) \in \bar{Q}$, $|w(t_1, x) - w(t_2, x)| \leq C |t_1 - t_2|^{(1+\alpha)/2}$ for a positive constant C . The space $\mathcal{C}^{(1+\alpha)/2, 1+\alpha}(\bar{Q})$, endowed with the semi-norm

$$|w|_{\mathcal{C}^{(1+\alpha)/2, 1+\alpha}(\bar{Q})} := \sum_{i=1}^d \left\| \frac{\partial w}{\partial x_i} \right\|_{\mathcal{C}^{\alpha/2, \alpha}(\bar{Q})} + \sup_{(t_1, x) \neq (t_2, x) \in \bar{Q}} \frac{|w(t_1, x) - w(t_2, x)|}{|t_1 - t_2|^{(1+\alpha)/2}}$$

and norm $\|w\|_{\mathcal{C}^{(1+\alpha)/2, 1+\alpha}(\bar{Q})} := \|w\|_{\mathcal{C}(\bar{Q})} + |w|_{\mathcal{C}^{(1+\alpha)/2, 1+\alpha}(\bar{Q})}$ is a Banach space.

Finally, the space $\mathcal{C}^{1+\alpha/2, 2+\alpha}$ is made of all the functions $w \in \mathcal{C}^1(\bar{Q})$ which are twice continuously differentiable w.r.t. x , with partial derivatives $\frac{\partial w}{\partial x_i} \in \mathcal{C}^{(1+\alpha)/2, 1+\alpha}(\bar{Q})$ for all $i = 1, \dots, d$, and $\frac{\partial w}{\partial t} \in \mathcal{C}^{\alpha/2, \alpha}(\bar{Q})$. It is a Banach space with the norm

$$\|w\|_{\mathcal{C}^{1+\alpha/2, 2+\alpha}(\bar{Q})} := \|w\|_{\mathcal{C}(\bar{Q})} + \sum_{i=1}^d \left\| \frac{\partial w}{\partial x_i} \right\|_{\mathcal{C}^{(1+\alpha)/2, 1+\alpha}(\bar{Q})} + \left\| \frac{\partial w}{\partial t} \right\|_{\mathcal{C}^{\alpha/2, \alpha}(\bar{Q})}.$$

2.2. The case when $\partial_p H$ is bounded. We make the following assumptions on h , m_0 , H and G , in addition to the regularity assumptions on H already made in § 1:

(H_0) For simplicity only, the map h is invariant w.r.t. m , i.e. $h[m](x) = u_T(x)$, where u_T is a smooth function defined on \mathbb{T}^d . Moreover, m_0 is a smooth positive function.

(H₁) There exists a constant $\gamma_0 > 0$ such that

$$|H[m](x, 0)| \leq \gamma_0 \quad \forall (m, x) \in (\mathbb{P} \cap L^1(\mathbb{T}^d)) \times \mathbb{T}^d$$

(H₂) There exists a constant $\gamma_1 > 0$ such that

$$\left\| \frac{\partial H}{\partial p}[m] \right\|_{\text{Lip}(\mathbb{T}^d \times \mathbb{R}^d)} \leq \gamma_1 \quad \forall m \in \mathbb{P} \cap L^1(\mathbb{T}^d)$$

(H₃) For all $(m, x, p) \in (\mathbb{P} \cap L^1(\mathbb{T}^d)) \times \mathbb{T}^d \times \mathbb{R}^d$, $\frac{\partial H}{\partial m}[m](x, p)$ is a \mathcal{C}^1 function on \mathbb{T}^d and there exists a constant $\gamma_2 > 0$ such that for all $(m, x, p) \in (\mathbb{P} \cap L^1(\mathbb{T}^d)) \times \mathbb{T}^d \times \mathbb{R}^d$,

$$\left\| \frac{\partial H}{\partial m}[m](x, p) \right\|_{\mathcal{C}^1(\mathbb{T}^d)} \leq \gamma_2(1 + |p|)$$

(H₄) There exists a constant $\gamma_3 > 0$ such that:

$$\left\| \frac{\partial H}{\partial p}[m_1](\cdot, 0) - \frac{\partial H}{\partial p}[m_2](\cdot, 0) \right\|_{\mathcal{C}(\mathbb{T}^d)} \leq \gamma_3 \|m_1 - m_2\|_{L^1(\mathbb{T}^d)} \quad \forall m_1, m_2 \in L^1(\mathbb{T}^d).$$

(H₅) There exists $\gamma_4 > 0$ such that for $m_1, m_2 \in \mathbb{P} \cap L^1(\mathbb{T}^d)$, $p_1, p_2 \in L^\infty(\mathbb{T}^d)$,

$$\|G[m_1, p_1] - G[m_2, p_2]\|_{L^\infty(\mathbb{T}^d)} \leq \gamma_4 (\|p_1 - p_2\|_{L^\infty(\mathbb{T}^d)} + \|m_1 - m_2\|_{L^1(\mathbb{T}^d)}).$$

Example. All the assumptions above are satisfied by the map H :

$$H[m](x, p) = -\frac{\Phi(p)}{(c + (\rho_1 * m)(x))^\alpha} + F(x, (\rho_2 * m)(x)),$$

where Φ is a \mathcal{C}^2 function from \mathbb{R}^d to \mathbb{R}_+ such that $D^2\Phi$ and $D\Phi$ are bounded, α and c are positive numbers, ρ_1 and ρ_2 are smoothing kernels in $\mathcal{C}^\infty(\mathbb{T}^d)$, ρ_1 is nonnegative, and F is a \mathcal{C}^2 function defined on $\mathbb{T}^d \times \mathbb{R}^d$. Here, $\rho * m(x) = \int_{\mathbb{T}^d} \rho(x - z)m(z)dz$. It is easy to check that

$$G[m, q](x) = \left(\alpha \tilde{\rho}_1 * \left(m \frac{\Phi(q)}{(c + \rho_1 * m)^{\alpha+1}} \right) \right) (x) + \tilde{\rho}_2 * (mF'(\cdot, \rho_2 * m))(x)$$

where $\tilde{\rho}_1(x) = \rho_1(-x)$ and $\tilde{\rho}_2(x) = \rho_2(-x)$.

Such a Hamiltonian models situations in which there are congestion effects, i.e. the cost of displacement increases in the regions where the density is large. The term $F(x, (\rho_2 * m)(x))$ may model aversion to crowded regions. The prototypical situation is $g[m](x, a) = a$ and $\Phi(q) = \min_{b \in \mathcal{K}} (q \cdot b + \Phi^*(b))$, where \mathcal{K} is a compact subset of \mathbb{R}^d . Setting $\Phi^*(b) = +\infty$ if $b \notin \mathcal{K}$, H corresponds to the cost $f[m](x, a) = \frac{1}{(c + (\rho_1 * m)(x))^\alpha} \Phi^*(a(c + (\rho_1 * m)(x))^\alpha) + F(x, (\rho_2 * m)(x))$.

2.2.1. *A priori estimates.* We first assume that (1.6)-(1.8) has a sufficiently smooth solution and we look for a priori estimates.

Step 1: uniform bounds on $\|m\|_{L^p(0, T; W^{1, p}(\mathbb{T}^d))} + \|m\|_{\mathcal{C}^{\alpha/2, \alpha}(\bar{Q})}$, $p \in [1, \infty)$, $\alpha \in [0, 1)$. First, standard arguments yield that $m(t, \cdot) \in \mathbb{P}$ for all $t \in [0, T]$.

From Assumption (H₂), the function $b : (t, x) \mapsto \partial_p H[m(t, \cdot)](x, \nabla u(t, x))$ is such that $\|b\|_{L^\infty(Q)} \leq \gamma_1$. The Cauchy problem satisfied by m can be written

$$\begin{aligned} \frac{\partial m}{\partial t}(t, x) - \nu \Delta m(t, x) + \text{div}(b(t, \cdot)m(t, \cdot))(x) &= 0, \\ m(0, x) &= m_0(x), \end{aligned} \tag{2.1}$$

and from the classical theory on weak solutions to parabolic equations, see e.g. Theorem 6.1 in [14], there exists a constant C_0 depending only on $\|m_0\|_{L^2(\mathbb{T}^d)}$ such that

$$\|m\|_{L^2(0,T;H^1(\mathbb{T}^d))} + \|m\|_{C([0,T];L^2(\mathbb{T}^d))} \leq C_0.$$

Moreover, since the operator in (2.1) is in divergence form, we have maximum estimates on m , see Corollary 9.10 in [14]: there exists a constant C_1 depending only on $\|m_0\|_\infty$ and γ_1 such that

$$m(t, x) \leq C_1 \quad \forall (t, x) \in [0, T] \times \mathbb{T}^d. \quad (2.2)$$

Therefore, the Fokker-Planck equation in (2.1) can be rewritten

$$\frac{\partial m}{\partial t}(t, x) - \nu \Delta m(t, x) + \operatorname{div}(B(t, \cdot))(x) = 0, \quad (2.3)$$

where $\|B\|_\infty \leq \gamma_1 C_1$. From standard results on the heat equation, see [10], this implies that for all $p \in [1, \infty)$ there exists a constant $C_2(p)$ which depends on $\|m_0\|_\infty$ and γ_1 , such that

$$\|m\|_{L^p(0,T;W^{1,p}(\mathbb{T}^d))} + \left\| \frac{\partial m}{\partial t} \right\|_{L^p(0,T;W^{-1,p}(\mathbb{T}^d))} \leq C_2(p). \quad (2.4)$$

Finally, Hölder estimates for the heat equation with a right hand side in divergence form, see for example Theorem 6.29 in [14], yield that for any $\alpha \in (0, 1)$, there exists a positive constant $C_3(\alpha) \geq C_1$ which only depends on γ_1 and on $\|m_0\|_{C^\alpha(\mathbb{T}^d)}$ such that

$$\|m\|_{C^{\alpha/2, \alpha}(\bar{Q})} \leq C_3(\alpha). \quad (2.5)$$

Step 2: uniform bounds on $\|u\|_{C^{(1+\theta)/2, 1+\theta}(\bar{Q})}$, $\theta \in (0, 1)$. Defining

$$a(t, x) := -H[m(t, \cdot)](x, 0) \quad \text{and} \quad A(t, x) := \int_0^1 \frac{\partial H}{\partial p}[m(t, \cdot)](x, \zeta \nabla u(t, x)) d\zeta,$$

the HJB equation (1.6) can be rewritten

$$\frac{\partial u}{\partial t}(t, x) + \nu \Delta u(t, x) + A(t, x) \cdot \nabla u(t, x) = a(t, x) - G[m(t, \cdot), \nabla u(t, \cdot)](x). \quad (2.6)$$

For some smooth function \hat{u} , let us consider

$$\frac{\partial \hat{u}}{\partial t}(t, x) + \nu \Delta \hat{u}(t, x) + A(t, x) \cdot \nabla \hat{u}(t, x) = a(t, x) - G[m(t, \cdot), \nabla \hat{u}(t, \cdot)](x) \quad (2.7)$$

instead of (2.6), with the same terminal condition as in (1.8). From Assumption (H_1) and (H_2) , $\|a\|_\infty \leq \gamma_0$ and $\|A\|_\infty \leq \gamma_1$. From Assumption (H_3) ,

$$\|G[m, \nabla \hat{u}]\|_{L^2(\mathbb{T}^d)} \leq c(1 + \|\nabla \hat{u}\|_{L^2(\mathbb{T}^d)}), \quad (2.8)$$

where $c > 0$ depends on C_1 in (2.2) and γ_2 . Multiplying (2.7) by $u(t, x)e^{-2\Lambda t}$ and integrating on \mathbb{T}^d , then using the bounds on $\|a\|_\infty$, $\|A\|_\infty \leq \gamma_1$ and (2.8), a standard argument yields that there exist constants Λ and \tilde{C}_4 which depend only on γ_0 , γ_1 , γ_2 , $\|m_0\|_\infty$ such that

$$\begin{aligned} & - \frac{d}{dt} \left(\|u(T-t, \cdot)\|_{L^2(\mathbb{T}^d)}^2 e^{-2\Lambda(T-t)} \right) + \nu \|\nabla u(T-t, \cdot)\|_{L^2(\mathbb{T}^d)}^2 e^{-2\Lambda(T-t)} \\ & \leq \tilde{C}_4 + \frac{\nu}{2} \|\nabla \hat{u}(T-t, \cdot)\|_{L^2(\mathbb{T}^d)}^2 e^{-2\Lambda(T-t)}. \end{aligned} \quad (2.9)$$

Hence, if

$$\nu \int_{t=0}^T \|\nabla \hat{u}(T-t, \cdot)\|_{L^2(\mathbb{T}^d)}^2 e^{-2\Lambda(T-t)} dt \leq C_4, \quad (2.10)$$

with

$$C_4 = 2\tilde{C}_4 T + 2 \int_{\mathbb{T}^d} u_T^2(x) dx, \quad (2.11)$$

then

$$\sup_t e^{-2\Lambda(T-t)} \int_{\mathbb{T}^d} u^2(T-t, x) dx + \nu \int_{t=0}^T \|\nabla u(T-t, \cdot)\|_{L^2(\mathbb{T}^d)}^2 e^{-2\Lambda(T-t)} dt \leq C_4. \quad (2.12)$$

Similarly, a solution of (1.6)-(1.8) satisfies (2.12) with the same constants Λ and C_4 . Note that Λ can be chosen large enough such that the function $(t, x) \mapsto u_T(x)$ satisfies (2.12).

For a solution of (1.6)-(1.8), this implies that $\partial_t u + \nu \Delta u$ is bounded in $L^2(Q)$, hence that u is bounded in $\mathcal{C}^0(0, T; H^1(\mathbb{T}^d))$ by a constant $\tilde{C}_4 > \|u_T\|_{H^1(\mathbb{T}^d)}$ which depends on Λ , C_4 , γ_1 and $\|u_T\|_{H^1(\mathbb{T}^d)}$, i.e.

$$\|\nabla u\|_{L^\infty(0, T; H^1(\mathbb{T}^d))} \leq \tilde{C}_4. \quad (2.13)$$

As a consequence, the left-hand side of (2.6) is bounded in $L^\infty(Q)$, and this yields Hölder estimates on u : by using Theorem 6.48 in [14], we see that for all $\theta \in (0, 1)$, there exists a constant $C_5(\theta)$ which depends on θ , $\|m_0\|_\infty$, $\|u_T\|_{\mathcal{C}^{1+\theta}(\mathbb{T}^d)}$, γ_0 , γ_1 , γ_2 such that

$$\|u\|_{\mathcal{C}^{(1+\theta)/2, 1+\theta}(\bar{Q})} \leq C_5(\theta), \quad (2.14)$$

which holds for a solution of (2.7) with the terminal condition (1.8), as soon as \hat{u} satisfies (2.12) and (2.13).

Step 3: uniform bound on $\|m\|_{\mathcal{C}^{(1+\theta)/2, 1+\theta}(\bar{Q})}$, $\theta \in (0, 1)$. Let us go back to (1.7). From Assumptions $(H_1) - (H_4)$, and from the previous two steps, we see that for any $\theta \in (0, 1)$, m and $\frac{\partial H}{\partial p}[m](\cdot, \nabla u)$ are both bounded in $\mathcal{C}^{\theta/2, \theta}(\bar{Q})$ by constants which depend on m_0 and u_T , and $\gamma_0, \dots, \gamma_3$. Thus, the function B in (2.3) is bounded in $\mathcal{C}^{\theta/2, \theta}(\bar{Q})$. Using Theorem 6.48 in [14] for the heat equation with a data in divergence form, we see that for all $\theta \in (0, 1)$, there exists a constant $C_6(\theta)$ which depends on θ , $\|m_0\|_{\mathcal{C}^{1+\theta}(\mathbb{T}^d)}$, $\|u_T\|_{\mathcal{C}^{1+\theta}(\mathbb{T}^d)}$, $\gamma_0, \dots, \gamma_3$ such that

$$\|m\|_{\mathcal{C}^{(1+\theta)/2, 1+\theta}(\bar{Q})} \leq C_6(\theta).$$

Step 4: uniform bounds on $\|u\|_{\mathcal{C}^{1+\theta/2, 2+\theta}(\bar{Q})}$, $\theta \in (0, 1)$. From the previous steps and Assumptions $(H_1) - (H_4)$, we see that there exists a constant c such that the functions in (2.6) satisfy $\|a\|_{\mathcal{C}^{\theta/2, \theta}(\bar{Q})} \leq c$ and $\|A\|_{\mathcal{C}^{\theta/2, \theta}(\bar{Q})} \leq c$. Similarly, from Assumptions (H_3) and (H_5) , $\|G[m, \nabla u]\|_{\mathcal{C}^{\theta/2, \theta}(\bar{Q})} \leq c$. Standard regularity results on parabolic equations, for instance Theorem 4.9 in [14] lead to the existence of $C_7(\theta)$ such that

$$\|u\|_{\mathcal{C}^{1+\theta/2, 2+\theta}(\bar{Q})} \leq C_7(\theta).$$

2.2.2. The existence theorem.

Theorem 1. *Under the Assumptions (H_0) - (H_5) , for $\alpha \in (0, 1)$ there exist functions $u \in \mathcal{C}^{1+\alpha/2, 2+\alpha}(\bar{Q})$ and $m \in \mathcal{C}^{(1+\alpha)/2, 1+\alpha}(\bar{Q})$ which satisfy (1.6)-(1.8), (note that (1.7) is satisfied in a weak sense).*

Proof. The argument is reminiscent of that used by J-M. Lasry and P-L. Lions for mean field games: it is done in two steps

Step A. For $R > 0$, let $\eta_R : \mathbb{R} \rightarrow \mathbb{R}$ be a smooth, nondecreasing and odd function such that

1. $\eta_R(y) = y$ if $|y| \leq R$, $\eta_R(y) = 2R$ if $y \geq 3R$
2. $\|\eta'_R\|_\infty \leq 1$

We consider the modified set of equations

$$0 = \frac{\partial u}{\partial t}(t, x) + \nu \Delta u(t, x) + H[m(t, \cdot)](x, \nabla u(t, x)) + \eta_R(G[m(t, \cdot), \nabla u(t, \cdot)](x)) \quad (2.15)$$

$$0 = \frac{\partial m}{\partial t}(t, x) - \nu \Delta m(t, x) + \operatorname{div} \left(m(t, \cdot) \frac{\partial H}{\partial p} [m(t, \cdot)](\cdot, \nabla u(t, \cdot)) \right) (x) \quad (2.16)$$

We are going to apply Leray-Schauder fixed point theorem to a map χ defined for example in $X = \{m \in \mathcal{C}^0([0, T]; L^2(\mathbb{T}^d) \cap \mathbb{P})\}$: consider first the map $\psi : X \rightarrow X \times L^2(0, T; H^1(\mathbb{T}^d))$, $m \mapsto (m, u)$ where u is a weak solution of (2.15) and $u|_{t=T} = u_T$. Existence and uniqueness for this problem are well known. Moreover, from the estimates above, for every $0 < \alpha < 1$, $\|u\|_{\mathcal{C}^{1/2+\alpha/2, 1+\alpha}(\bar{Q})}$ is bounded by a constant independent of m and $m \mapsto u$ is continuous from X to $\mathcal{C}^{1/2+\alpha/2, 1+\alpha}(\bar{Q})$.

Fix $\theta \in (0, 1)$, and consider the map $\zeta : X \times \mathcal{C}^{1/2+\theta/2, 1+\theta}(\bar{Q}) \rightarrow L^2(0, T; H^1(\mathbb{T}^d))$, $(m, u) \mapsto \tilde{m}$ where \tilde{m} is a weak solution of the Fokker-Planck equation

$$0 = \frac{\partial \tilde{m}}{\partial t}(t, x) - \nu \Delta \tilde{m}(t, x) + \operatorname{div} \left(\tilde{m}(t, \cdot) \frac{\partial H}{\partial p} [m(t, \cdot)](\cdot, \nabla u(t, \cdot)) \right) (x).$$

and $\tilde{m}|_{t=0} = m_0$. Existence and uniqueness are well known, and moreover, the estimates above tell us that for all $0 < \alpha < 1$, there exists $R_\alpha > 0$ such that $\|\tilde{m}\|_{\mathcal{C}^{\alpha/2, \alpha}(\bar{Q})} \leq R_\alpha$ uniformly with respect to m and u . Moreover from the assumptions, it can be seen that ζ maps continuously $X \times \mathcal{C}^{1/2+\theta/2, 1+\theta}(\bar{Q})$ to X .

Let K be the set $\{\|m\|_{\mathcal{C}^{\alpha/2, \alpha}(\bar{Q})} \leq R_\alpha; m|_{t=T} = m_T\} \cap X$: this set is a compact and convex subset of X and the map $\chi = \zeta \circ \psi : m \mapsto \tilde{m}$ is continuous in X and leaves K invariant. We can apply Leray-Schauder fixed point theorem the map χ , which yields the existence of a solution (u_R, m_R) to (2.15)-(2.16). Moreover the a priori estimates above tell us that $u_R \in \mathcal{C}^{1+\alpha/2, 2+\alpha}(\bar{Q})$ and $m_R \in \mathcal{C}^{(1+\alpha)/2, 1+\alpha}(\bar{Q})$.

Step B. Looking at all the a priori estimates above, it can be seen that m_R , (resp u_R) belongs to a bounded subset of $\mathcal{C}^{\alpha/2, \alpha}(\bar{Q})$ (resp. $\mathcal{C}^{1/2+\alpha/2, 1+\alpha}(\bar{Q})$) independent of R . Hence, for R large enough, $\eta_R(G[m_R, \nabla u_R]) = G[m_R, \nabla u_R]$, and (m_R, u_R) is a weak solution of (1.6)-(1.8), with $u_R \in \mathcal{C}^{1+\alpha/2, 2+\alpha}(\bar{Q})$ and $m_R \in \mathcal{C}^{(1+\alpha)/2, 1+\alpha}(\bar{Q})$. \square

Remark 3. It is possible to weaken some of the assumptions in Theorem 1: for example, we can assume the following weaker version of (H_2) , namely:

(H'_2) There exists a constant $\gamma_1 > 0$ and $\eta \in (0, 1)$ such that

- $\forall m \in \mathbb{P} \cap L^1(\mathbb{T}^d)$, $\|\frac{\partial H}{\partial p}[m]\|_{\mathcal{C}(\mathbb{T}^d \times \mathbb{R}^d)} \leq \gamma_1$
- $\forall m \in \mathbb{P} \cap L^1(\mathbb{T}^d)$, $x, y \in \mathbb{T}^d$, $p, q \in \mathbb{R}^d$,

$$\left| \frac{\partial H}{\partial p}[m](x, p) - \frac{\partial H}{\partial p}[m](y, q) \right| \leq \gamma_1 (d(x, y) + |p - q|^\eta)$$

Indeed, the regularity of $\frac{\partial H}{\partial p}$ with respect to p is only used in Steps 3 and 4 above: with this weaker assumptions, the conclusions of steps 3 and 4 hold with $0 < \theta < \eta$, and this is enough for proving the existence of $u \in \mathcal{C}^{1+\alpha/2, 2+\alpha}(\bar{Q})$ and $m \in \mathcal{C}^{(1+\alpha)/2, 1+\alpha}(\bar{Q})$ for some α , $0 < \alpha < \eta$ which satisfy (1.6)-(1.8).

2.3. Hamiltonian with a subquadratic growth in p : a specific case. For a smooth nonnegative periodic function ρ , two constants $\alpha > 0$ and β , $1 < \beta \leq 2$, let us focus on the following Hamiltonian:

$$H[m](x, p) := -\frac{|p|^\beta}{(1 + (\rho * m)(x))^\alpha}. \quad (2.17)$$

The map G defined in (1.9) is

$$G[m, q](x) = \alpha \left(\tilde{\rho} * \left(m \frac{|q|^\beta}{(1 + (\rho * m))^{\alpha+1}} \right) \right) (x),$$

where $\tilde{\rho}(x) := \rho(-x)$.

Assuming that m_0 is smooth, let us call $\bar{m}_0 = \|m_0\|_\infty$: for all $x \in \mathbb{T}^d$, $0 < m_0(x) \leq \bar{m}_0$. We assume that

$$\|\rho\|_{L^1(\mathbb{T}^d)} < \frac{\beta - 1}{\alpha \bar{m}_0}. \quad (2.18)$$

Remark 4. It would be interesting to make further investigations to see if the assumption on the regularizing kernel ρ in (2.18) is really necessary, since it is not necessary in the context of mean field games with congestion. Yet, in the a priori estimates proposed below, (2.18) is useful for getting a bound on $\|mH[m](\cdot, \nabla u)\|_{L^1(Q)}$, see (2.20).

2.3.1. A priori estimates. We first assume that (1.6)-(1.8) has a sufficiently smooth weak solution and we look for a priori estimates.

Step 1: a lower bound on u . Since G is non negative, by comparison, we see that

$$u(t, x) \geq \underline{u}_T := \min_{\xi \in \mathbb{T}^d} u(T, \xi) \quad \forall (t, x) \in [0, T] \times \mathbb{T}^d.$$

Step 2: an energy estimate and its consequences. Let us multiply (1.6) by $m - \bar{m}_0$ and (1.7) by u and integrate the two resulting equations on \mathbb{T}^d . Summing the resulting identities, we obtain:

$$\begin{aligned} & \int_Q \frac{\partial}{\partial t} (u(t, x)(m(t, x) - \bar{m}_0)) dx dt + \int_Q H[m(t, \cdot)](x, \nabla u(t, x))(m(t, x) - \bar{m}_0) dx dt \\ & + \int_Q G[m(t, \cdot), \nabla u(t, \cdot)](x)(m(t, x) - \bar{m}_0) dx dt \\ & + \int_Q \operatorname{div} \left(m(t, x) \frac{\partial}{\partial p} H[m(t, \cdot)](x, \nabla u(t, x)) \right) u(t, x) dx dt = 0 \end{aligned}$$

Hence

$$\begin{aligned} & \int_{\mathbb{T}^d} u(T, x)(m(T, x) - \bar{m}_0) dx + \int_{\mathbb{T}^d} u(0, x)(\bar{m}_0 - m(0, x)) dx \\ & = \int_Q H[m(t, \cdot)](x, \nabla u(t, x))(\bar{m}_0 - m(t, x)) dx dt \\ & + \int_Q G[m(t, \cdot), \nabla u(t, \cdot)](x)(\bar{m}_0 - m(t, x)) dx dt \\ & + \int_Q m(t, x) \frac{\partial}{\partial p} H[m(t, \cdot)](x, \nabla u(t, x)) \cdot \nabla u(t, x) dx dt \end{aligned} \quad (2.19)$$

In (2.19), the first term in the left hand side is bounded from below by $-\|u_T\|_\infty(1 + \bar{m}_0)$, The second term is larger than $\underline{u}_T \int_{\mathbb{T}^d} (\bar{m}_0 - m(0, x)) dx = (\bar{m}_0 - 1)\underline{u}_T$. Therefore, the left hand side of (2.19), is bounded from below by a constant c which only depends on \bar{m}_0 and $\|u_T\|_\infty$; we obtain that

$$\begin{aligned} c \leq & (\beta - 1) \int_Q m(t, x) H[m(t, \cdot)](x, \nabla u(t, x)) dx dt + \int_Q \bar{m}_0 H[m(t, \cdot)](x, \nabla u(t, x)) dx dt \\ & + \alpha \int_Q (\bar{m}_0 - m(t, x)) \tilde{\rho} * \left(m(t, \cdot) \frac{|\nabla u(t, \cdot)|^\beta}{(1 + \rho * m(t, \cdot))^{\alpha+1}} \right) (x) dx dt. \end{aligned}$$

We see that last term can be bounded as follows:

$$\begin{aligned} & \int_Q (\bar{m}_0 - m(t, x)) \tilde{\rho} * \left(m(t, \cdot) \frac{|\nabla u(t, \cdot)|^\beta}{(1 + \rho * m(t, \cdot))^{\alpha+1}} \right) (x) dx dt \\ \leq & \bar{m}_0 \int_Q \tilde{\rho} * \left(m(t, \cdot) \frac{|\nabla u(t, \cdot)|^\beta}{(1 + \rho * m(t, \cdot))^\alpha} \right) (x) dx dt \\ \leq & \bar{m}_0 \|\rho\|_{L^1(\mathbb{T}^d)} \int_Q m(t, x) \frac{|\nabla u(t, x)|^\beta}{(1 + \rho * m(t, x))^\alpha} dx dt \\ = & -\bar{m}_0 \|\rho\|_{L^1(\mathbb{T}^d)} \int_Q m(t, x) H[m(t, \cdot)](x, \nabla u(t, x)) dx dt. \end{aligned}$$

Therefore,

$$\begin{aligned} c \leq & (\beta - 1 - \alpha \bar{m}_0 \|\rho\|_{L^1(\mathbb{T}^d)}) \int_Q m(t, x) H[m(t, \cdot)](x, \nabla u(t, x)) dx dt \\ & + \int_Q \bar{m}_0 H[m(t, \cdot)](x, \nabla u(t, x)) dx dt. \end{aligned}$$

From (2.17) and (2.18), we see that there exists a constant C_1 which depends on \bar{m}_0 and $\|u_T\|_\infty$ such that

$$\|mH[m](\cdot, \nabla u)\|_{L^1(Q)} + \|H[m](\cdot, \nabla u)\|_{L^1(Q)} \leq C_1. \quad (2.20)$$

Using (2.20), we deduce from a comparison argument applied to the HJB equation that there exists a constant C_2 which depends on \bar{m}_0 and $\|u_T\|_\infty$ such that

$$\|u\|_{L^\infty(Q)} \leq C_2. \quad (2.21)$$

Since $1 < \beta \leq 2$, there exists a constant c such that $|\frac{\partial H[m]}{\partial p}(x, p)|^2 \leq c(1 - H[m](x, p))$. We deduce from (2.20) and the latter observation that there exists a constant $C_3 > 0$ such that

$$\int_Q (m(t, x) + 1) \left| \frac{\partial H[m(t, \cdot)]}{\partial p}(x, \nabla u(t, x)) \right|^2 dx dt \leq C_3. \quad (2.22)$$

Step 3: uniform estimates from the Fokker-Planck equation. The following estimates can be proved exactly as in [19], Lemma 2.3 and Corollary 2.4, (see also [6], Lemma 2.5 and Corollary 2):

Lemma 2. For $\gamma = \frac{d+2}{d}$ if $d > 2$ and all $\gamma < 2$ if $d = 2$, there exists an constant $c > 0$, (independent from m_0 and u_T) such that

$$\begin{aligned} & \sup_{t \in [0, T]} \|m(t, \cdot) \log(m(t, \cdot))\|_{L^1(\mathbb{T}^d)} + \|\sqrt{m}\|_{L^2(0, T; H^1(\mathbb{T}^d))}^2 + \|m\|_{L^\gamma(Q)}^\gamma \\ & \leq c \left(\int_Q m(t, x) \left| \frac{\partial H[m(t, \cdot)]}{\partial p}(x, \nabla u(t, x)) \right|^2 dx dt + \int_{\mathbb{T}^d} m_0(x) \log(m_0(x)) dx \right). \end{aligned} \quad (2.23)$$

Corollary 1. For $q = \frac{d+2}{d+1}$ if $d > 2$ and $q < 4/3$ if $d = 2$, there exists a constant $c > 0$ such that

$$\begin{aligned} & \|\nabla m\|_{L^q(Q)}^q + \left\| \frac{\partial m}{\partial t} \right\|_{L^q(0, T; W^{-1, q}(\mathbb{T}^d))}^q \\ & \leq c \left(\int_Q m(t, x) \left| \frac{\partial H[m(t, \cdot)]}{\partial p}(x, \nabla u(t, x)) \right|^2 dx dt + \int_{\mathbb{T}^d} m_0(x) \log(m_0(x)) dx \right). \end{aligned} \quad (2.24)$$

From (2.24) and (2.22), we have a uniform bound on $\left\| \frac{\partial m}{\partial t} \right\|_{L^q(0, T; W^{-1, q}(\mathbb{T}^d))}^q$ by a constant depending only on u_T and m_0 . We infer that (1.6) can be written

$$\frac{\partial u}{\partial t}(t, x) + \nu \Delta u(t, x) + a(t, x) |\nabla u|^\beta(t, x) = b(t, x), \quad (2.25)$$

where a is a function which belongs to $\mathcal{C}([0, T]; \mathcal{C}^p(\mathbb{T}^d))$ for all $p \in \mathbb{N}$, with corresponding norms bounded by constants depending only on u_T and m_0 . From (2.20), we deduce that for all $p \in \mathbb{N}$, $\|b\|_{L^1(0, T; W^{p, \infty}(\mathbb{T}^d))}$ is bounded by a constant depending only on u_T and m_0 , because

$$\begin{aligned} \|b\|_{L^1(0, T; W^{p, \infty}(\mathbb{T}^d))} &= \|G[m, \nabla u]\|_{L^1(0, T; W^{p, \infty}(\mathbb{T}^d))} \\ &\leq c \left\| \frac{m |\nabla u|^\beta}{(1 + (\rho * m))^{\alpha+1}} \right\|_{L^1(Q)} \\ &\leq c \|m H[m](\cdot, \nabla u)\|_{L^1(Q)}. \end{aligned}$$

Step 4: uniform estimates on $|\nabla u|$. Since $a \in \mathcal{C}([0, T]; \mathcal{C}^p(\mathbb{T}^d))$ and $b \in L^1(0, T; W^{p, \infty}(\mathbb{T}^d))$, we can apply Bernstein method to (2.25) and estimate $|\nabla u|$. By a slight modification of the proof of Theorem 11.1 in [14], (the only difference is that in [14], b is supposed to belong to $L^\infty(0, T; W^{p, \infty}(\mathbb{T}^d))$, but it can be checked that this assumption can be weakened), we prove that there exists a constant C_4 which depends on u_T and m_0 such that

$$\|\nabla u\|_{L^\infty(Q)} \leq C_4. \quad (2.26)$$

The proof adapted from [14] is rather long, so we do not reproduce it here.

Step 5: stronger a priori estimates. Since $|\nabla u|$ is bounded, we can recover all the a priori estimates in § 2.2.1, except that the estimates in Step 3 and 4 of § 2.2.1 only hold with $0 < \theta < \beta - 1$, in view of Remark 3. We obtain that for all $\gamma \in (0, 1)$, there exist two constants $C_5(\gamma)$ and $C_6(\gamma)$ such that $\|m\|_{\mathcal{C}^{\gamma/2, \gamma}(\bar{Q})} \leq C_5(\gamma)$ and $\|u\|_{\mathcal{C}^{(1+\gamma)/2, 1+\gamma}(\bar{Q})} \leq C_5(\gamma)$, and that for all $\theta \in (0, \beta - 1)$, there exist two constants $C_7(\theta)$ and $C_8(\theta)$ such that $\|m\|_{\mathcal{C}^{(1+\theta)/2, 1+\theta}(\bar{Q})} \leq C_7(\theta)$ and $\|u\|_{\mathcal{C}^{1+\theta/2, 2+\theta}(\bar{Q})} \leq C_8(\theta)$.

2.3.2. The existence theorem.

Theorem 3. *We assume (H_0) and (2.18). For γ , $0 < \gamma < \beta - 1$, there exists a function $u \in \mathcal{C}^{1+\gamma/2, 2+\gamma}(\bar{Q})$ and $m \in \mathcal{C}^{(1+\gamma)/2, 1+\gamma}(\bar{Q})$ which satisfy (1.6)-(1.8) with H given by (2.17).*

Proof. We start by suitably truncating the Hamiltonian H and the map G : for $R > 1$, define

$$H_R[m](x, p) = \begin{cases} -\frac{|p|^\beta}{(1 + (\rho * m)(x))^\alpha} & \text{if } |p| < R, \\ -\frac{\beta R^{\beta-1}|p| + (1 - \beta)R^\beta}{(1 + (\rho * m)(x))^\alpha} & \text{if } |p| \geq R, \end{cases} \quad (2.27)$$

and

$$G_R[m, q](x) = \alpha \left(\tilde{\rho} * \left(m \frac{\min(|q|^\beta, R^\beta)}{(1 + (\rho * m))^{\alpha+1}} \right) \right) (x). \quad (2.28)$$

Note that

$$-H_R[m](x, p) + \frac{\partial}{\partial p} H_R[m](x, p) \cdot p = \begin{cases} -(\beta - 1) \frac{|p|^\beta}{(1 + (\rho * m)(x))^\alpha} & \text{if } |p| \leq R, \\ -(\beta - 1) \frac{R^\beta}{(1 + (\rho * m)(x))^\alpha} & \text{if } |p| \geq R. \end{cases} \quad (2.29)$$

Thanks to Remark 3, we can use a slightly modified version of Theorem 1: for some γ , $0 < \gamma < \beta - 1$, there exists a solution (u_R, m_R) of

$$\begin{aligned} 0 &= \frac{\partial u_R}{\partial t}(t, x) + \nu \Delta u_R(t, x) + H_R[m_R(t, \cdot)](x, \nabla u_R(t, x)) + G_R[m_R(t, \cdot), \nabla u_R(t, \cdot)](x), \\ 0 &= \frac{\partial m_R}{\partial t}(t, x) - \nu \Delta m_R(t, x) + \operatorname{div} \left(m_R(t, \cdot) \frac{\partial H_R}{\partial p}[m_R(t, \cdot)](\cdot, \nabla u_R(t, \cdot)) \right) (x), \end{aligned}$$

with the initial and terminal conditions (1.8), such that $u_R \in \mathcal{C}^{1+\gamma/2, 2+\gamma}(\bar{Q})$ and $m_R \in \mathcal{C}^{(1+\gamma)/2, 1+\gamma}(\bar{Q})$.

Then it is possible to carry out the same program as in Step 1 and 2 in § 2.3.1: using (2.27)-(2.29), we obtain that there exists a constant c independent of R such that

$$\begin{aligned} c &\leq -(\beta - 1 - \alpha \bar{m}_0 \|\rho\|_{L^1(\mathbb{T}^d)}) \int_Q m_R(t, x) \frac{|\nabla u_R(t, x)|^\beta}{(1 + \rho * m_R(t, x))^\alpha} 1_{\{|\nabla u_R(t, x)| < R\}} dx dt \\ &\quad - (\beta - 1 - \alpha \bar{m}_0 \|\rho\|_{L^1(\mathbb{T}^d)}) \int_Q m_R(t, x) \frac{R^\beta}{(1 + \rho * m_R(t, x))^\alpha} 1_{\{|\nabla u_R(t, x)| \geq R\}} dx dt \\ &\quad + \int_Q \bar{m}_0 H_R[m_R(t, \cdot)](x, \nabla u_R(t, x)) dx dt, \end{aligned}$$

and this implies the counterpart of (2.20): there exists a constant C independent of R such that

$$\left\| (1 + m_R) \frac{\min(|\nabla u_R|^\beta, R^\beta)}{(1 + (\rho * m_R))^\alpha} \right\|_{L^1(Q)} \leq C. \quad (2.30)$$

From this, we obtain the counterpart of (2.22):

$$\int_Q (m_R(t, x) + 1) \left| \frac{\partial H_R}{\partial p}[m_R(t, \cdot)](x, \nabla u_R(t, x)) \right|^2 dx dt \leq C, \quad (2.31)$$

where C is a constant independent of R . This estimate allows one for carrying out Steps 3 and 4 in § 2.3.1 and obtaining estimates independent of R : in particular, the same Bernstein argument can be used, and we obtain that there exists a constant independent of R such that $\|\nabla u_R\|_{L^\infty(Q)} \leq C$. In turn, step 5 in § 2.3.1 can be used and leads to estimates independent of R .

From this, taking R large enough yields the desired existence result. \square

3. Uniqueness.

3.1. Uniqueness for (1.6)-(1.8): a sufficient condition. In what follows, we prove sufficient conditions leading to the uniqueness of a classical solution of (1.6)-(1.8). For simplicity, we still assume that the final cost does not depend on the density, i.e. that there exists a smooth function u_T such that $h[m](x) = u_T(x)$. In order to simplify the discussion, we assume that the operator H depends smoothly enough on its argument to give sense to the calculations that follow.

We consider two classical solutions (u, m) and (\tilde{u}, \tilde{m}) of

$$\begin{aligned} 0 &= \frac{\partial u}{\partial t}(t, x) + \nu \Delta u(t, x) + H[m(t, \cdot)](x, \nabla u(t, x)) \\ &\quad + \int_{\mathbb{T}^d} \frac{\partial H}{\partial m}[m(t, \cdot)](\xi, \nabla u(t, \xi))(x) m(t, \xi) d\xi, \end{aligned} \quad (3.1)$$

$$0 = \frac{\partial m}{\partial t}(t, x) - \nu \Delta m(t, x) + \operatorname{div} \left(m(t, \cdot) \frac{\partial H}{\partial p}[m(t, \cdot)](\cdot, \nabla u(t, \cdot)) \right) (x), \quad (3.2)$$

and

$$\begin{aligned} 0 &= \frac{\partial \tilde{u}}{\partial t}(t, x) + \nu \Delta \tilde{u}(t, x) + H[\tilde{m}(t, \cdot)](x, \nabla \tilde{u}(t, x)) \\ &\quad + \int_{\mathbb{T}^d} \frac{\partial H}{\partial \tilde{m}}[\tilde{m}(t, \cdot)](\xi, \nabla \tilde{u}(t, \xi))(x) \tilde{m}(t, \xi) d\xi, \end{aligned} \quad (3.3)$$

$$0 = \frac{\partial \tilde{m}}{\partial t}(t, x) - \nu \Delta \tilde{m}(t, x) + \operatorname{div} \left(\tilde{m}(t, \cdot) \frac{\partial H}{\partial p}[\tilde{m}(t, \cdot)](\cdot, \nabla \tilde{u}(t, \cdot)) \right) (x). \quad (3.4)$$

We subtract (3.3) from (3.1), multiply the resulting equation by $(m(t, x) - \tilde{m}(t, x))$, and integrate over Q . Similarly, we subtract (3.4) from (3.2), multiply the resulting equation by $(u(t, x) - \tilde{u}(t, x))$, and integrate over Q . We sum the two resulting identities: we obtain

$$\begin{aligned} 0 &= \int_{\mathbb{T}^d} (u(T, x) - \tilde{u}(T, x))(m(T, x) - \tilde{m}(T, x)) dx \\ &\quad - \int_{\mathbb{T}^d} (u(0, x) - \tilde{u}(0, x))(m(0, x) - \tilde{m}(0, x)) dx \\ &\quad + \int_{t=0}^T E[m(t, \cdot), \nabla u(t, \cdot), \tilde{m}(t, \cdot), \nabla \tilde{u}(t, \cdot)] dt. \end{aligned} \quad (3.5)$$

where

$$\begin{aligned} E[m_1, p_1, m_2, p_2] &= \int_{\mathbb{T}^d} (H[m_1](x, p_1(x)) - H[m_2](x, p_2(x)))(m_1(x) - m_2(x)) dx \\ &\quad + \int_{\mathbb{T}^d} (m_1(x) - m_2(x)) \int_{\mathbb{T}^d} \left(\frac{\partial H}{\partial m}[m_1](\xi, p_1(\xi))(x) m_1(\xi) - \frac{\partial H}{\partial m}[m_2](\xi, p_2(\xi))(x) m_2(\xi) \right) d\xi dx \\ &\quad - \int_{\mathbb{T}^d} \left(m_1(x) \frac{\partial}{\partial p} H[m_1](x, p_1(x)) - m_2(x) \frac{\partial}{\partial p} H[m_2](x, p_2(x)) \right) (p_1(x) - p_2(x)) dx. \end{aligned}$$

Call $\delta m = m_2 - m_1$ and $\delta p = p_2 - p_1$ and consider the function $e : [0, 1] \rightarrow \mathbb{R}$ defined by

$$\begin{aligned} e(\theta) &= \frac{1}{\theta} E[m_1, p_1, m_1 + \theta \delta m, p_1 + \theta \delta p], & \theta > 0, \\ e(0) &= 0. \end{aligned} \quad (3.6)$$

It can be checked that e is \mathcal{C}^1 on $[0, 1]$ and that its derivative is

$$\begin{aligned} e'(\theta) &= 2 \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \frac{\partial H}{\partial m}[m_1 + \theta \delta m](\xi, p_1 + \theta \delta p(\xi))(x) \delta m(\xi) \delta m(x) \\ &+ \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} \int_{\mathbb{T}^d} (m_1(\xi) + \theta \delta m(\xi)) \frac{\partial^2 H}{\partial m \partial m}[m_1 + \theta \delta m](\xi, p_1 + \theta \delta p(\xi))(x)(y) \delta m(x) \delta m(y) \\ &- \int_{\mathbb{T}^d} (m_1(x) + \theta \delta m(x)) \delta p(x) \cdot D_{p,p}^2 H[m_1 + \theta \delta m](x, p_1(x) + \theta \delta p(x)) \delta p(x). \end{aligned} \quad (3.7)$$

Let us introduce the functional defined on $\mathcal{C}(\mathbb{T}^d) \times \mathcal{C}(\mathbb{T}^d; \mathbb{R}^d)$ by

$$\mathcal{H}[m, p] := \int_{\mathbb{T}^d} m(x) H[m](x, p(x)) dx. \quad (3.8)$$

The second order Fréchet derivative of \mathcal{H} with respect to m (respectively p) at (m, p) is a bilinear form on $\mathcal{C}(\mathbb{T}^d)$, (resp. $\mathcal{C}(\mathbb{T}^d; \mathbb{R}^d)$), noted $D_{m,m}^2 \mathcal{H}[m, p]$, (resp. $D_{p,p}^2 \mathcal{H}[m, p]$). For all $m \in \mathcal{C}(\mathbb{T}^d) \cap \mathbb{P}$ and all $p \in \mathcal{C}(\mathbb{T}^d; \mathbb{R}^d)$, let us define the quadratic form $\mathcal{Q}[m, p]$ on $\mathcal{C}(\mathbb{T}^d) \times \mathcal{C}(\mathbb{T}^d; \mathbb{R}^d)$ by

$$\mathcal{Q}[m, p](\mu, \pi) = D_{m,m}^2 \mathcal{H}[m, p](\mu, \mu) - D_{p,p}^2 \mathcal{H}[m, p](\pi, \pi). \quad (3.9)$$

We see that (3.7) can be written as follows:

$$e'(\theta) = \mathcal{Q}[m_1 + \theta \delta m, p_1 + \theta \delta p](\delta m, \delta p). \quad (3.10)$$

Theorem 4. *We assume (H_0) and that $(m, x, p) \mapsto H[m](x, p)$ is \mathcal{C}^2 on $\mathcal{C}(\mathbb{T}^d) \times \mathbb{T}^d \times \mathbb{R}^d$. A sufficient condition for the uniqueness of a classical solution of (1.6)-(1.8) is that*

1. *for all $m \in \mathcal{C}(\mathbb{T}^d) \cap \mathbb{P}$ and all $p \in \mathcal{C}(\mathbb{T}^d; \mathbb{R}^d)$, the quadratic form $\mu \mapsto D_{m,m}^2 \mathcal{H}[m, p](\mu, \mu)$ is positive definite*
2. *for all $m \in \mathcal{C}(\mathbb{T}^d) \cap \mathbb{P}$ and $x \in \mathbb{T}^d$, the real valued function $p \in \mathbb{R}^d \mapsto H[m](x, p)$ is strictly concave.*

Proof. From the concavity of $p \mapsto H[m](x, p)$, $-D_{p,p}^2 \mathcal{H}[m, p]$ is positive semi-definite. Therefore, $\mathcal{Q}[m, p]$ is positive semi-definite, and $\mathcal{Q}[m, p](\mu, \pi) = 0$ implies that $D_{m,m}^2 \mathcal{H}[m, p](\mu, \mu) = 0$ and $-D_{p,p}^2 \mathcal{H}[m, p](\pi, \pi) = 0$, and therefore $\mu = 0$.

From (3.5), two solutions (u, m) and (\tilde{u}, \tilde{m}) of (1.6)-(1.8) satisfy

$$\int_{t=0}^T E[m(t, \cdot), \nabla u(t, \cdot), \tilde{m}(t, \cdot), \nabla \tilde{u}(t, \cdot)] dt = 0, \quad (3.11)$$

because $\tilde{m}(0, \cdot) = m(0, \cdot)$ and $\tilde{u}(T, \cdot) = u(T, \cdot)$.

But, from (3.6) and (3.10), the properties of the quadratic form $\mathcal{Q}[(1-\theta)m(t, \cdot) + \theta \tilde{m}(t, \cdot), (1-\theta)\nabla u(t, \cdot) + \theta \nabla \tilde{u}(t, \cdot)]$ imply that

$\int_{t=0}^T E[m(t, \cdot), \nabla u(t, \cdot), \tilde{m}(t, \cdot), \nabla \tilde{u}(t, \cdot)] dt > 0$ if $m \neq \tilde{m}$.

Therefore, (3.11) implies that $m = \tilde{m}$. Then,

0 =

$$\int_Q m(t, x) \left(\frac{\partial H}{\partial p}[m(t, \cdot)](x, \nabla u(t, x)) - \frac{\partial H}{\partial p}[m(t, \cdot)](x, \nabla \tilde{u}(t, x)) \right) \cdot (\nabla u - \nabla \tilde{u})(t, x). \quad (3.12)$$

If $\nu > 0$, then the maximum principle implies that $m(t, x) > 0$ for all $t > 0, x \in \mathbb{T}^d$. This observation, (3.12) and the strict concavity of H with respect to p imply that $\nabla u(t, x) = \nabla \tilde{u}(t, x) > 0$ for all t, x , which yields immediately that $u = \tilde{u}$ by using (1.6). \square

Remark 5. Let us give an alternative argument which does not require the knowledge that $m(t, x) > 0$ for all $t > 0, x \in \mathbb{T}^d$. Such an argument may be useful in situations when $\nu = 0$ or ν is replaced in (1.1) by a function of x which vanishes in some regions of \mathbb{T}^d . The strict concavity of H with respect to p and (3.12) yield the fact that $u = \tilde{u}$ in the region where $m > 0$. This implies that $G[m(t, \cdot)](x, \nabla u(t, x)) = G[m(t, \cdot)](x, \nabla \tilde{u}(t, x))$: hence, for all t and x ,

$$\begin{aligned} & \frac{\partial u}{\partial t}(t, x) + \nu \Delta u(t, x) + H[m(t, \cdot)](x, \nabla u(t, x)) \\ &= \frac{\partial \tilde{u}}{\partial t}(t, x) + \nu \Delta \tilde{u}(t, x) + H[m(t, \cdot)](x, \nabla \tilde{u}(t, x)). \end{aligned}$$

We can then apply standard results on the uniqueness of the Cauchy problem with the HJB equation $\frac{\partial u}{\partial t}(t, x) + \nu \Delta u(t, x) + H[m(t, \cdot)](x, \nabla u(t, x)) = g$ and obtain that $u = \tilde{u}$.

Corollary 2. *In the case when H depends locally on m , i.e.*

$$H[m](x, p) = \tilde{H}(x, p, m(x)),$$

the sufficient condition in Theorem 4 is implied by the strict concavity of $p \in \mathbb{R}^d \mapsto \tilde{H}(x, p, m)$ for all $m > 0$ and $x \in \mathbb{T}^d$ and the strict convexity of the real valued function $m \in \mathbb{R}_+ \mapsto m\tilde{H}(x, p, m)$, for all $p \in \mathbb{R}^d$.

Example. Consider for example the Hamiltonian

$$H[m](x, p) = \tilde{H}(x, p, m(x)) = -\frac{|p|^\beta}{(c + m(x))^\alpha} + F(m(x)), \quad (3.13)$$

with $c > 0, \alpha > 0, \beta > 1, F$ a smooth function defined on \mathbb{R}_+ . One can check that if $\alpha \leq 1$ and F is strictly convex, then uniqueness holds.

Such a Hamiltonian arise in a local model for congestion, see [15].

Remark 6. The same analysis can be carried out for mean field games, see [15]: for example, under Assumption (H_0) and in the case when H depends locally on m , i.e. $H[m](x, p) = \tilde{H}(x, p, m(x))$, a sufficient condition for the uniqueness of a classical solution of (1.10)-(1.12) is that

$$\begin{pmatrix} 2\frac{\partial \tilde{H}}{\partial m}(x, p, m) & -\frac{\partial}{\partial m}\nabla_p^T \tilde{H}(x, p, m) \\ -\frac{\partial}{\partial m}\nabla_p \tilde{H}(x, p, m) & -2D_{p,p}^2 \tilde{H}(x, p, m) \end{pmatrix}$$

be positive definite for all $x \in \mathbb{T}^d, m > 0$ and $p \in \mathbb{R}^d$. Here, we see that the sufficient condition involves the mixed partial derivatives of \tilde{H} with respect to m and p , which is not the case for mean field type control. If \tilde{H} depends separately on p and m as in [13], then $\frac{\partial}{\partial m}\nabla_p \tilde{H}(x, p, m) = 0$ and the condition becomes: \tilde{H} is

strictly concave with respect to p for $m > 0$ and non decreasing with respect to m , (or concave with respect to p and strictly increasing with respect to m).

Remark 7. The extension of the result on uniqueness to weak solutions is not trivial. In the context of mean filed games, one can find such results in [18] and [17]: roughly speaking they rely on some new uniqueness results for weak solutions of the Fokker-Planck equation and on crossed regularity lemmas, see Lemma 5 in [18]. In the context of mean field type control, the same kind of analysis has not been done yet.

In the case when $n = d$, $g[m](x, v) = v$ and $v \mapsto f[m](x, v)$ is strictly convex for all $m \in \mathbb{P}$ and $x \in \mathbb{T}^d$, it is well known that $f[m](x, v) = \sup_{q \in \mathbb{R}^d} (H[m](x, q) - q \cdot v)$. Furthermore if $p \mapsto H[m](x, p)$ is strictly concave for all $m \in \mathbb{P}$ and $x \in \mathbb{T}^d$, then

$$f[m](x, v) = \max_{q \in \mathbb{R}^d} (H[m](x, q) - q \cdot v) \quad (3.14)$$

and the maximum is achieved by a unique q . This observation leads to the following necessary condition for the assumption of Theorem 4 to be satisfied.

Proposition 1. *Assume that $n = d$, $g[m](x, v) = v$, that $v \mapsto f[m](x, v)$ is strictly convex for all $m \in \mathbb{P}$ and $x \in \mathbb{T}^d$, and that $p \mapsto H[m](x, p)$ is strictly concave for all $m \in \mathbb{P}$ and $x \in \mathbb{T}^d$. If for all $p \in \mathcal{C}(\mathbb{T}^d; \mathbb{R}^d)$, $m \mapsto \mathcal{H}[m, p]$ is strictly convex in $\mathbb{P} \cap \mathcal{C}(\mathbb{T}^d)$, then for all $v \in \mathcal{C}(\mathbb{T}^d; \mathbb{R}^d)$, $m \mapsto \int_{\mathbb{T}^d} m(x) f[m](x, v(x)) dx$ is strictly convex in $\mathbb{P} \cap \mathcal{C}(\mathbb{T}^d)$.*

Proof. Take $\lambda_1 > 0$ and $\lambda_2 > 0$ such that $\lambda_1 + \lambda_2 = 1$ and $m_1 \neq m_2$ in $\mathbb{P} \cap \mathcal{C}(\mathbb{T}^d)$. From (3.14),

$$\begin{aligned} & \int_{\mathbb{T}^d} (\lambda_1 m_1(x) + \lambda_2 m_2(x)) f[\lambda_1 m_1 + \lambda_2 m_2](x, v(x)) dx \\ &= \int_{\mathbb{T}^d} \max_{q \in \mathbb{R}^d} (\lambda_1 m_1(x) + \lambda_2 m_2(x)) (H[\lambda_1 m_1 + \lambda_2 m_2](x, q) - qv(x)) dx. \end{aligned}$$

If for all $x \in \mathbb{T}^d$, the maximum in the latter integrand is achieved by $q^*(x)$, then $x \mapsto q^*(x)$ is a continuous function (from the continuity of v) and we have

$$\begin{aligned} & \int_{\mathbb{T}^d} (\lambda_1 m_1(x) + \lambda_2 m_2(x)) f[\lambda_1 m_1 + \lambda_2 m_2](x, v(x)) dx \\ &= \max_{q \in \mathcal{C}(\mathbb{T}^d; \mathbb{R}^d)} \int_{\mathbb{T}^d} (\lambda_1 m_1(x) + \lambda_2 m_2(x)) (H[\lambda_1 m_1 + \lambda_2 m_2](x, q(x)) - q(x)v(x)) dx. \end{aligned}$$

From this and the convexity of $m \mapsto \mathcal{H}[m, p]$, we deduce that

$$\begin{aligned} & \int_{\mathbb{T}^d} (\lambda_1 m_1(x) + \lambda_2 m_2(x)) f[\lambda_1 m_1 + \lambda_2 m_2](x, v(x)) dx \\ &< \max_{q \in \mathcal{C}(\mathbb{T}^d; \mathbb{R}^d)} \left(\begin{aligned} & \lambda_1 \int_{\mathbb{T}^d} (m_1(x) H[m_1](x, q(x)) - q(x)v(x)) dx + \\ & \lambda_2 \int_{\mathbb{T}^d} (m_2(x) H[m_2](x, q(x)) - q(x)v(x)) dx \end{aligned} \right) \\ &\leq \lambda_1 \max_{q \in \mathcal{C}(\mathbb{T}^d; \mathbb{R}^d)} \int_{\mathbb{T}^d} (m_1(x) H[m_1](x, q(x)) - q(x)v(x)) dx \\ &\quad + \lambda_2 \max_{q \in \mathcal{C}(\mathbb{T}^d; \mathbb{R}^d)} \int_{\mathbb{T}^d} (m_2(x) H[m_2](x, q(x)) - q(x)v(x)) dx \\ &= \lambda_1 \int_{\mathbb{T}^d} m_1(x) f[m_1](x, v(x)) dx + \lambda_2 \int_{\mathbb{T}^d} m_2(x) f[m_2](x, v(x)) dx. \end{aligned}$$

□

3.2. Back to the control of McKean-Vlasov dynamics. As in the end of the previous paragraph, we assume that $n = d$ and $g[m](x, v) = v$. The control of McKean-Vlasov dynamics can be written as a control problem with linear constraints by making the change of variables $z = mv$: it consists of minimizing

$$\tilde{J}(z, m_z) = \int_Q f[m_z(t, \cdot)] \left(x, \frac{z(t, x)}{m_z(t, x)} \right) m_z(t, x) dx dt + \int_{\mathbb{T}^d} u_T(x) m_z(T, x) dx, \quad (3.15)$$

subject to the linear constraints

$$\frac{\partial m_z}{\partial t}(t, x) - \nu \Delta m_z(t, x) + \operatorname{div} z(t, x) = 0 \quad t \in (0, T], x \in \mathbb{T}^d, \quad (3.16)$$

with the initial condition

$$m_z(0, x) = m_0(x), \quad x \in \mathbb{T}^d. \quad (3.17)$$

For simplicity, we assume that f depends locally on m , i.e.

$$f[m](x, v(x)) = \tilde{f}(x, v(x), m(x)), \quad \forall x \in \mathbb{T}^d.$$

We are going to look for sufficient conditions for $(z, m) \mapsto m \tilde{f}(x, \frac{z}{m}, m)$ be a convex function. This condition will thus yield the uniqueness for the above control problem.

Assuming that all the following differentiations are allowed, we see that the Hessian of the latter function is

$$\begin{aligned} \Theta(x, v, m) = & \begin{pmatrix} \frac{1}{m^3} z \cdot D_{vv}^2 \tilde{f} \left(\frac{z}{m}, m \right) z & -\frac{1}{m^2} z \cdot D_{vv}^2 \tilde{f} \left(\frac{z}{m}, m \right) \\ -\frac{1}{m^2} D_{vv}^2 \tilde{f} \left(\frac{z}{m}, m \right) z & \frac{1}{m} D_{vv}^2 \tilde{f} \left(\frac{z}{m}, m \right) \end{pmatrix} \\ & + \begin{pmatrix} 2 \frac{\partial \tilde{f}}{\partial m} \left(\frac{z}{m}, m \right) + m \frac{\partial^2 \tilde{f}}{\partial m^2} \left(\frac{z}{m}, m \right) - 2 \frac{z}{m} \cdot \frac{\partial \nabla_v \tilde{f}}{\partial m} \left(\frac{z}{m}, m \right) & \frac{\partial \nabla_v^T \tilde{f}}{\partial m} \left(\frac{z}{m}, m \right) \\ \frac{\partial \nabla_v \tilde{f}}{\partial m} \left(\frac{z}{m}, m \right) & 0 \end{pmatrix} \end{aligned}$$

where we have omitted the dependency on x for brevity. This is better understood when expressed in terms of (v, m) :

$$\Theta(x, v, m) = \begin{pmatrix} \frac{\partial^2}{\partial m^2} \left(m \tilde{f}(x, v, m) \right) & m \frac{\partial \nabla_v^T \tilde{f}}{\partial m} (x, v, m) \\ m \frac{\partial \nabla_v \tilde{f}}{\partial m} (x, v, m) & m D_{vv}^2 \tilde{f}(x, v, m) \end{pmatrix}. \quad (3.18)$$

We have proved the following

Proposition 2. *We assume that $n = d$ and $g[m](x, v) = v$, and that $f[m](x, v(x)) = \tilde{f}(x, v(x), m(x))$, for all $x \in \mathbb{T}^d$, where \tilde{f} is a smooth function. A sufficient condition for the uniqueness of a minimum (z^*, m^*) such that $m^* > 0$ is that $\Theta(x, v, m)$ be positive definite for all $x \in \mathbb{T}^d$, $m > 0$ and $v \in \mathbb{R}^d$.*

Proposition 3. *We make the same assumptions as in Proposition 2. The positive definiteness of $\Theta(x, v, m)$ for all $x \in \mathbb{T}^d$, $m > 0$ and $v \in \mathbb{R}^d$ implies the sufficient conditions on \tilde{H} in Corollary 2.*

Proof. We observe first that the positive definiteness of Θ implies that $D_{vv}^2 \tilde{f}(x, v, m)$ is positive definite for all $x \in \mathbb{T}^d$, $m > 0$ and $v \in \mathbb{R}^d$.

Let us call $v^* \in \mathbb{R}^d$ the vector achieving $\tilde{H}(x, p, m) = p \cdot v^* + \tilde{f}(x, v^*, m)$. We know that $\nabla_p \tilde{H}(x, p, m) = v^*$. Differentiating the optimality condition for v^* with respect to p , we find that

$$D_{p,p}^2 \tilde{H}(x, p, m) = - \left(D_{v,v}^2 \tilde{f}(x, v^*, m) \right)^{-1}. \quad (3.19)$$

Note that (3.19) implies the strict concavity of $p \mapsto \tilde{H}(x, p, m)$ which is the first desired condition on \tilde{H} . The second condition on \tilde{H} will be a consequence of the implicit function theorem: differentiating \tilde{H} with respect to m , we find that

$$\frac{\partial \tilde{H}}{\partial m}(x, p, m) = \frac{\partial \tilde{f}}{\partial m}(x, v^*, m) + \nabla_v \tilde{f}(x, v^*, m) \cdot \frac{\partial v^*}{\partial m} + p \cdot \frac{\partial v^*}{\partial m} = \frac{\partial \tilde{f}}{\partial m}(x, v^*, m), \quad (3.20)$$

where the last identity comes from the definition of v^* . Differentiating once more with respect to m , we find that

$$\frac{\partial^2 \tilde{H}}{\partial m^2}(x, p, m) = \frac{\partial^2 \tilde{f}}{\partial m^2}(x, v^*, m) + \frac{\partial \nabla_v \tilde{f}}{\partial m}(x, v^*, m) \cdot \frac{\partial v^*}{\partial m}. \quad (3.21)$$

Then the implicit function theorem applied to the optimality condition for v^* yields that

$$\frac{\partial v^*}{\partial m} = - \left(D_{v,v}^2 \tilde{f}(x, v^*, m) \right)^{-1} \frac{\partial \nabla_v \tilde{f}}{\partial m}(x, v^*, m). \quad (3.22)$$

From (3.19)- (3.22), we see that

$$\begin{aligned} \frac{\partial^2}{\partial m^2} \left(m \tilde{H}(x, p, m) \right) &= \frac{\partial^2}{\partial m^2} \left(m \tilde{f}(x, \cdot, m) \right) (v^*) \\ &- \left(m \frac{\partial \nabla_v \tilde{f}}{\partial m}(x, v^*, m) \right) \cdot \left(m D_{v,v}^2 \tilde{f}(x, v^*, m) \right)^{-1} \left(m \frac{\partial \nabla_v \tilde{f}}{\partial m}(x, v^*, m) \right). \end{aligned} \quad (3.23)$$

Hence, $\frac{\partial^2}{\partial m^2} \left(m \tilde{H}(x, p, m) \right)$ is a Schur complement of $\Theta(x, v^*, m)$. Therefore, it is positive definite and we have proved the second condition on \tilde{H} . \square

4. Numerical Simulations. Here we model a situation in which a crowd of pedestrians is driven to leave a given square hall (whose side is 50 meters long) containing rectangular obstacles: one can imagine for example a situation of panic in a closed building, in which the population tries to reach the exit doors. The chosen geometry is represented on Figure 1. The aim is to compare the evolution of the density in




FIGURE 1. Left: the geometry. Right: the density at $t = 0$

two models:

1. Mean field games: we choose $\nu = 0.012$ and the Hamiltonian to be of the form (3.13), i.e. which takes congestion effects into account and depends locally on m ; more precisely:

$$\tilde{H}(x, p, m) = - \frac{8|p|^2}{(1+m)^{\frac{3}{4}}} + \frac{1}{3200}.$$

The system (1.10)- (1.11) becomes

$$\frac{\partial u}{\partial t} + 0.012 \Delta u - \frac{8}{(1+m)^{\frac{3}{4}}} |\nabla u|^2 = -\frac{1}{3200}, \quad (4.1)$$

$$\frac{\partial m}{\partial t} - 0.012 \Delta m - 16 \operatorname{div} \left(\frac{m \nabla u}{(1+m)^{\frac{3}{4}}} \right) = 0. \quad (4.2)$$

The horizon T is $T = 50$ minutes. There is no terminal cost.

There are two exit doors, see Figure 1. The part of the boundary corresponding to the doors is called Γ_D . The boundary conditions at the exit doors are chosen as follows: there is a Dirichlet condition for u on Γ_D , corresponding to an exit cost; in our simulations, we have chosen $u = 0$ on Γ_D . For m , we may assume that $m = 0$ outside the domain, so we also get the Dirichlet condition $m = 0$ on Γ_D .

The boundary Γ_N corresponds to the solid walls of the hall and of the obstacles. A natural boundary condition for u on Γ_N is a homogeneous Neumann boundary condition, i.e. $\frac{\partial u}{\partial n} = 0$ which says that the velocity of the pedestrians is tangential to the walls. The natural condition for the density m is that $\nu \frac{\partial m}{\partial n} + m \frac{\partial \tilde{H}}{\partial p}(\cdot, \nabla u, m) \cdot n = 0$, therefore $\frac{\partial m}{\partial n} = 0$ on Γ_N .

2. Mean field type control: this is the situation where pedestrians or robots use the same feedback law (we may imagine that they follow the strategy decided by a leader); we keep the same Hamiltonian, and the HJB equation becomes

$$\frac{\partial u}{\partial t} + 0.012 \Delta u - \left(\frac{2}{(1+m)^{\frac{3}{4}}} + \frac{6}{(1+m)^{\frac{7}{4}}} \right) |\nabla u|^2 = -\frac{1}{3200}. \quad (4.3)$$

while (4.2) and the boundary condition are unchanged.

The initial density m_0 is piecewise constant and takes two values 0 and 4 people/m², see Figure 1. At $t = 0$, there are 3300 people in the hall.

We use the finite difference method originally proposed in [3], see [1] for some details on the implementation and [2] for convergence results.

On Figure 2, we plot the density m obtained by the simulations for the two models, at $t = 1, 2, 5$ and 15 minutes. With both models, we see that the pedestrians rush towards the narrow corridors leading to the exits, at the left and right sides of the hall, and that the density reaches high values at the intersections of corridors; then congestion effects explain why the velocity is low (the gradient of u) in the regions where the density is high. On the figure, we see that the mean field type control leads to a slower exit of the hall, with lower peaks of density.

Acknowledgements. We warmly thank A. Bensoussan for helpful discussions. The first author was partially funded by the ANR projects ANR-12-MONU-0013 and ANR-12-BS01-0008-01. The second author was partially funded by the Research Grants Council of HKSAR (CityU 500113).

REFERENCES

1. Y. Achdou, *Finite difference methods for mean field games*, Hamilton-Jacobi equations: approximations, numerical analysis and applications (P. Loreti and N. A. Tchou, eds.), Lecture Notes in Math., vol. 2074, Springer, Heidelberg, 2013, pp. 1–47.
2. Y. Achdou, F. Camilli, and I. Capuzzo-Dolcetta, *Mean field games: convergence of a finite difference method*, SIAM J. Numer. Anal. **51** (2013), no. 5, 2585–2612.
3. Y. Achdou and I. Capuzzo-Dolcetta, *Mean field games: numerical methods*, SIAM J. Numer. Anal. **48** (2010), no. 3, 1136–1162.

4. A. Bensoussan and J. Frehse, *Control and Nash games with mean field effect*, Chin. Ann. Math. Ser. B **34** (2013), no. 2, 161–192.
5. A. Bensoussan, J. Frehse, and P. Yam, *Mean field games and mean field type control theory*, Springer Briefs in Mathematics, Springer, New York, 2013.
6. P. Cardaliaguet, J.-M. Lasry, P.-L. Lions, and A. Porretta, *Long time average of mean field games*, Netw. Heterog. Media **7** (2012), no. 2, 279–301.
7. R. Carmona and F. Delarue, *Mean field forward-backward stochastic differential equations*, Electron. Commun. Probab. **18** (2013), no. 68, 15.
8. R. Carmona, F. Delarue, and A. Lachapelle, *Control of McKean-Vlasov dynamics versus mean field games*, Math. Financ. Econ. **7** (2013), no. 2, 131–166.
9. D. A. Gomes and J. Saúde, *Mean field games models—a brief survey*, Dyn. Games Appl. **4** (2014), no. 2, 110–154.
10. O. A. Ladyženskaja, V. A. Solonnikov, and N. N. Ural'ceva, *Linear and quasilinear equations of parabolic type*, Translated from the Russian by S. Smith. Translations of Mathematical Monographs, Vol. 23, American Mathematical Society, Providence, R.I., 1968.
11. J.-M. Lasry and P.-L. Lions, *Jeux à champ moyen. I. Le cas stationnaire*, C. R. Math. Acad. Sci. Paris **343** (2006), no. 9, 619–625.
12. ———, *Jeux à champ moyen. II. Horizon fini et contrôle optimal*, C. R. Math. Acad. Sci. Paris **343** (2006), no. 10, 679–684.
13. ———, *Mean field games*, Jpn. J. Math. **2** (2007), no. 1, 229–260.
14. G.M. Lieberman, *Second order parabolic differential equations*, World Scientific Publishing Co., Inc., River Edge, NJ, 1996.
15. P.-L. Lions, *Cours du Collège de France*, http://www.college-de-france.fr/default/EN/all/equ_der/, 2007–2011.
16. H. P. McKean, Jr., *A class of Markov processes associated with nonlinear parabolic equations*, Proc. Nat. Acad. Sci. U.S.A. **56** (1966), 1907–1911.
17. A. Porretta, *Weak solutions to Fokker-Planck equations and mean field games*, to appear.
18. ———, *On the planning problem for a class of mean field games*, C. R. Math. Acad. Sci. Paris **351** (2013), no. 11–12, 457–462.
19. ———, *On the planning problem for the mean field games system*, Dyn. Games Appl. **4** (2014), no. 2, 231–256.
20. A.-S. Sznitman, *Topics in propagation of chaos*, École d'Été de Probabilités de Saint-Flour XIX—1989, Lecture Notes in Math., vol. 1464, Springer, Berlin, 1991, pp. 165–251.

E-mail address: achdou@ljl1.univ-paris-diderot.fr

E-mail address: mathieu.lauriere@gmail.com




FIGURE 2. The density computed with the two models at different dates. Left: Mean field game. Right: Mean field type control. The scales vary from one date to the other