



# Stabilization for the semilinear wave equation with geometric control condition

Romain Joly, Camille Laurent

## ► To cite this version:

Romain Joly, Camille Laurent. Stabilization for the semilinear wave equation with geometric control condition. IF\_PREPUB. 2012. <hal-00696147v3>

**HAL Id: hal-00696147**

**<https://hal.archives-ouvertes.fr/hal-00696147v3>**

Submitted on 2 Dec 2013

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Stabilization for the semilinear wave equation with geometric control condition

Romain JOLY\* & Camille LAURENT<sup>†‡</sup>

December 2, 2013

## Abstract

In this article, we prove the exponential stabilization of the semilinear wave equation with a damping effective in a zone satisfying the geometric control condition only. The nonlinearity is assumed to be subcritical, defocusing and analytic. The main novelty compared to previous results, is the proof of a unique continuation result in large time for some undamped equation. The idea is to use an asymptotic smoothing effect proved by Hale and Raugel in the context of dynamical systems. Then, once the analyticity in time is proved, we apply a unique continuation result with partial analyticity due to Robbiano, Zuily, Tataru and Hörmander. Some other consequences are also given for the controllability and the existence of a compact attractor.

KEY WORDS: damped wave equation, stabilization, analyticity, unique continuation property, compact attractor.

AMS SUBJECT CLASSIFICATION: 35B40, 35B60, 35B65, 35L71, 93D20, 35B41.

## Résumé

Dans cet article, on prouve la décroissance exponentielle de l'équation des ondes semilinéaires avec un amortissement actif dans une zone satisfaisant seulement la condition de contrôle géométrique. La nonlinéarité est supposée sous-critique, défocalisante et analytique. La principale nouveauté par rapport aux résultats précédents est la preuve d'un résultat de prolongement unique en grand temps pour une solution non amortie. L'idée est d'utiliser un effet régularisant asymptotique

---

\*Institut Fourier - UMR5582 CNRS/Université de Grenoble - 100, rue des Maths - BP 74 - F-38402 St-Martin-d'Hères, France, email: [romain.joly@ujf-grenoble.fr](mailto:romain.joly@ujf-grenoble.fr)

<sup>†</sup>CNRS, UMR 7598, Laboratoire Jacques-Louis Lions, F-75005, Paris, France

<sup>‡</sup>UPMC Univ Paris 06, UMR 7598, Laboratoire Jacques-Louis Lions, F-75005, Paris, France, email: [laurent@ann.jussieu.fr](mailto:laurent@ann.jussieu.fr)

prouvé par Hale et Raugel dans le contexte des systèmes dynamiques. Ensuite, une fois l'analyticité en temps prouvée, on applique un théorème de prolongement unique avec analyticité partielle dû à Robbiano, Zuily, Tataru et Hörmander. Des applications à la contrôlabilité et à l'existence d'attracteur global compact pour l'équation des ondes sont aussi données.

## 1 Introduction

In this article, we consider the semilinear damped wave equation

$$\begin{cases} \square u + \gamma(x)\partial_t u + \beta u + f(u) = 0 & (t, x) \in \mathbb{R}_+ \times \Omega, \\ u(t, x) = 0 & (t, x) \in \mathbb{R}_+ \times \partial\Omega \\ (u, \partial_t u) = (u_0, u_1) \in H_0^1(\Omega) \times L^2(\Omega) \end{cases} \quad (1.1)$$

where  $\square = \partial_{tt}^2 - \Delta$  with  $\Delta$  being the Laplace-Beltrami operator with Dirichlet boundary conditions. The domain  $\Omega$  is a connected  $C^\infty$  three-dimensional Riemannian manifold with boundaries, which is either:

- i) compact.
- ii) a compact perturbation of  $\mathbb{R}^3$ , that is  $\mathbb{R}^3 \setminus D$  where  $D$  is a bounded smooth domain, endowed with a smooth metric equal to the euclidean one outside of a ball.
- iii) or a manifold with periodic geometry (cylinder,  $\mathbb{R}^3$  with periodic metric etc.).

The nonlinearity  $f \in C^1(\mathbb{R}, \mathbb{R})$  is assumed to be defocusing, energy subcritical and such that 0 is an equilibrium point. More precisely, we assume that there exists  $C > 0$  such that

$$f(0) = 0, \quad sf(s) \geq 0, \quad |f(s)| \leq C(1 + |s|)^p \quad \text{and} \quad |f'(s)| \leq C(1 + |s|)^{p-1} \quad (1.2)$$

with  $1 \leq p < 5$ .

We assume  $\beta \geq 0$  to be such that  $\Delta - \beta$  is a negative definite operator, that is that we have a Poincaré inequality  $\int_\Omega |\nabla u|^2 + \beta|u|^2 \geq C \int_\Omega |u|^2$  with  $C > 0$ . In particular, it may require  $\beta > 0$  if  $\partial\Omega = \emptyset$  or if  $\Omega$  is unbounded.

The damping  $\gamma \in L^\infty(\Omega)$  is a non-negative function. We assume that there exist an open set  $\omega \subset \Omega$ ,  $\alpha \in \mathbb{R}$ ,  $x_0 \in \Omega$  and  $R \geq 0$  such that

$$\forall x \in \omega, \quad \gamma(x) \geq \alpha > 0 \quad \text{and} \quad \Omega \setminus B(x_0, R) \subset \omega. \quad (1.3)$$

Moreover, we assume that  $\omega$  satisfies the geometric control condition introduced in [44] and [7]

(GCC) There exists  $L > 0$  such that any generalized geodesic of  $\Omega$  of length  $L$  meets the set  $\omega$  where the damping is effective.

The associated energy  $E \in \mathcal{C}^0(X, \mathbb{R}_+)$  is given by

$$E(u) := E(u, \partial_t u) = \frac{1}{2} \int_{\Omega} (|\partial_t u|^2 + |\nabla u|^2 + \beta |u|^2) + \int_{\Omega} V(u) , \quad (1.4)$$

where  $V(u) = \int_0^u f(s) ds$ . Due to Assumption (1.2) and the Sobolev embedding  $H^1(\Omega) \hookrightarrow L^6(\Omega)$ , this energy is well defined and moreover, if  $u$  solves (1.1), we have, at least formally,

$$\partial_t E(u(t)) = - \int_{\Omega} \gamma(x) |\partial_t u(x, t)|^2 dx \leq 0 . \quad (1.5)$$

The system is therefore dissipative. We are interested in the exponential decay of the energy of the nonlinear damped wave equation (1.1), that is the following property:

(ED) For any  $E_0 \geq 0$ , there exist  $K > 0$  and  $\lambda > 0$  such that, for all solutions  $u$  of (1.1) with  $E(u(0)) \leq E_0$ ,

$$\forall t \geq 0 , \quad E(u(t)) \leq K e^{-\lambda t} E(u(0))$$

Property (ED) means that the damping term  $\gamma \partial_t u$  stabilizes any solution of (1.1) to zero, which is an important property from the dynamical and control points of view.

Our main theorem is as follows.

**Theorem 1.1.** *Assume that the damping  $\gamma$  satisfies (1.3) and the geometric control condition (GCC). If  $f$  is real analytic and satisfies (1.2), then the exponential decay property (ED) holds.*

Theorem 1.1 applies for nonlinearities  $f$  which are globally analytic. Of course, the nonlinearities  $f(u) = |u|^{p-1}u$  are not analytic if  $p \notin \{1, 3\}$ , but we can replace these usual nonlinearities by similar ones as  $f(u) = (u/\text{th}(u))^{p-1}u$ , which are analytic for all  $p \in [1, 5)$ . Note that the estimates (1.2) are only required for  $s \in \mathbb{R}$ , so that it does not imply that  $f$  is polynomial. Moreover, we enhance that (ED) holds in fact for almost all the nonlinearities  $f$  satisfying (1.2), including non-analytic ones.

More precisely, we set

$$\mathfrak{C}^1(\mathbb{R}) = \{ f \in \mathcal{C}^1(\mathbb{R}) \text{ such that there exist } C > 0 \text{ and } p \in [1, 5) \text{ such that (1.2) holds} \} \quad (1.6)$$

endowed with Whitney topology (or any other reasonable topology). We recall that Whitney topology is the topology generated by the neighbourhoods

$$\mathcal{N}_{f, \delta} = \{ g \in \mathfrak{C}^1(\mathbb{R}) \mid \forall u \in \mathbb{R} , \quad \max(|f(u) - g(u)|, |f'(u) - g'(u)|) < \delta(u) \} \quad (1.7)$$

where  $f$  is any function in  $\mathcal{C}^1(\mathbb{R})$  and  $\delta$  is any positive continuous function. The set  $\mathcal{C}^1(\mathbb{R})$  is a Baire space, which means that any generic set, that is any set containing a countable intersection of open and dense sets, is dense in  $\mathcal{C}^1(\mathbb{R})$  (see Proposition 7.1). Baire property ensures that the genericity of a set in  $\mathcal{C}^1(\mathbb{R})$  is a good notion for “the set contains almost all non-linearity  $f$ ”.

**Theorem 1.2.** *Assume that the damping  $\gamma$  satisfies (1.3) and the geometric control condition (GCC). There exists a generic set  $\mathfrak{G} \subset \mathcal{C}^1(\mathbb{R})$  such that the exponential decay property (ED) holds for all  $f \in \mathfrak{G}$ .*

The statements of both theorems lead to some remarks.

- Of course, our results and their proofs should easily extend to any space dimension  $d \geq 3$  if the exponent  $p$  of the nonlinearity satisfies  $p < (d + 2)/(d - 2)$ .
- Actually, it may be possible to get  $\lambda > 0$  in (ED) uniform with respect to the size of the data. We can take for instance  $\lambda = \tilde{\lambda} - \varepsilon$  where  $\tilde{\lambda}$  is the decay rate of the linear equation. The idea is that once we know the existence of a decay rate, we know that the solution is close to zero for a large time. Then, for small solutions, the nonlinear term can be neglected to get almost the same decay rate as the linear equation. We refer for instance to [36] in the context of KdV equation. Notice that the possibility to get the same result with a constant  $K$  independent on  $E_0$  is an open problem.
- The assumption on  $\beta$  is important to ensure some coercivity of the energy and to preclude the spatially constant functions to be undamped solutions for the linear equation. It has been proved in [14] for  $\mathbb{R}^3$  and in [38] for a compact manifold that exponential decay can fail without this term  $\beta$ .
- The geometric control condition is known to be not only sufficient but also necessary for the exponential decay of the linear damped equation. The proof of the optimality uses some sequences of solutions which are asymptotically concentrated outside of the damping region. We can use the same idea in our nonlinear stabilization context. First, the observability for a certain time eventually large is known to be equivalent to the exponential decay of the energy. This was for instance noticed in [14] Proposition 2, in a similar context, see also Proposition 2.5 of this paper. Then, we take as initial data the same sequence that would give a counterexample for the linear observability. The linearizability property (see [19]) allows to obtain that the nonlinear solution is asymptotically close to the linear one. This contradicts the observability property for the nonlinear solution as it does for the linear case. Hence, the geometric control condition is also necessary for the exponential decay of the nonlinear equation.
- Our geometrical hypotheses on  $\Omega$  may look strange, however they are only assumed for sake of simplicity. In fact, our results should apply more generally for any smooth manifold with bounded geometry, that is that  $\Omega$  can be covered by a set of  $\mathcal{C}^\infty$ -charts  $\alpha_i : U_i \mapsto \alpha_i(U_i) \subset \mathbb{R}^3$  such that  $\alpha_i(U_i)$  is equal either to  $B(0, 1)$  or to  $B_+(0, 1) = \{x \in B(0, 1), x_1 > 0\}$  (in the case with boundaries) and such that, for any  $r \geq 0$  and  $s \in [1, \infty]$ , the  $W^{r,s}$ -norm of a function  $u$  in  $W^{r,s}(\Omega, \mathbb{R})$  is equivalent to the norm  $(\sum_{i \in \mathbb{N}} \|u \circ \alpha_i^{-1}\|_{W^{r,s}(\alpha_i(U_i))}^s)^{1/s}$ .

The stabilization property (ED) for Equation (1.1) has been studied in [28], [53], [54] and [13] for  $p < 3$ . For  $p \in [3, 5)$ , our main reference is the work of Dehman, Lebeau and Zuazua [17]. This work is mainly concerned with the stabilization problem previously described, on the Euclidean space  $\mathbb{R}^3$  with flat metric and stabilization active outside of a ball. The main purpose of this paper is to extend their result to a non flat geometry where multiplier methods cannot be used or do not give the optimal result with respect to the geometry. Other stabilization results for the nonlinear wave equation can be found in [2] and the references therein. Some works have been done in the difficult critical case  $p = 5$ , we refer to [14] and [38].

The proofs of these articles use three main ingredients:

- (i) the exponential decay of the linear equation, which is equivalent to the geometric control condition (GCC),
- (ii) a more or less involved compactness argument,
- (iii) a unique continuation result implying that  $u \equiv 0$  is the unique solution of

$$\begin{cases} \square u + \beta u + f(u) &= 0 \\ \partial_t u &= 0 \end{cases} \quad \text{on } [-T, T] \times \omega . \quad (1.8)$$

The results are mainly of the type: “geometric control condition” + “unique continuation” implies “exponential decay”. This type of implication is even stated explicitly in some related works for the nonlinear Schrödinger equation [15] and [37].

In the subcritical case  $p < 5$ , the less understood point is the unique continuation property (iii). In the previous works as [17], the authors use unique continuation results based on Carleman estimates. The resulting geometric assumptions are not very natural and are stronger than (GCC). Indeed, the unique continuation was often proved with some Carleman estimates that required some strong geometric conditions. For instance for a flat metric, the usual geometric assumption that appear are often of “multiplier type” that is  $\omega$  is a neighbourhood of  $\{x \in \partial\Omega \mid (x - x_0) \cdot n(x) > 0\}$  which are known to be stronger than the geometric control condition (see [41] for a discussion about the links between these assumptions). Moreover, on curved spaces, this type of condition often needs to be checked by hand in each situation, which is mostly impossible.

Our main improvement in this paper is the proof of unique continuation in infinite time under the geometric control condition only. We show that, if the nonlinearity  $f$  is analytic (or generic), then one can use the result of Robbiano and Zuily [47] to obtain a unique continuation property (iii) for infinite time  $T = +\infty$  with the geometric control condition (GCC) only.

The central argument of the proof of our main result, Theorem 1.1, is the unique continuation property of [47] (see Section 3). This result applies for solutions  $u$  of (1.8) being smooth in space and analytic in time. If  $f$  is analytic, then the solutions of (1.1) are of course not necessarily analytic in time since the damped wave equations are not smoothing

in finite time. However, the damped wave equations admit an asymptotic smoothing effect, i.e. are smoothing in infinite time. Hale and Raugel have shown in [25] that, for compact trajectories, this asymptotic smoothing effect also concerns the analyticity (see Section 5). In other words, combining [47] and [25] shows that the unique solution of (1.8) is  $u \equiv 0$  if  $f$  is analytic and if  $T = +\infty$ . This combination has already been used by dynamicians in [26] and [32] for  $p < 3$ .

One of the main interests of this paper is the use of arguments coming from both the dynamical study and the control theory of the damped wave equations. The reader familiar with the control theory could find interesting the use of the asymptotic smoothing effect to get unique continuation property with smooth solutions. The one familiar with the dynamical study of PDEs could be interested in the use of Strichartz estimates to deal with the case  $p \in [3, 5)$ . The main part of the proof of Theorem 1.1 is written with arguments coming from the dynamical study of PDEs. They are simpler than the corresponding ones of control theory, but far less accurate since they do not give any estimation for the time of observability. Anyway, such accuracy is not important here since we use the unique continuation property for (1.8) with  $T = +\infty$ . We briefly recall in Section 8 how these propagation of compactness and regularity properties could have been proved with some arguments more usual in the control theory.

Moreover, we give two applications of our results in both contexts of control theory and dynamical systems. First, as it is usual in control theory, some results of stabilization can be coupled with local control theorems to provide global controllability in large time.

**Theorem 1.3.** *Assume that  $f$  satisfies the conditions of Theorem 1.1 or belongs to the generic set  $\mathfrak{G}$  defined by Theorem 1.2. Let  $R_0 > 0$  and  $\omega$  satisfying the geometric control condition. Then, there exists  $T > 0$  such that for any  $(u_0, u_1)$  and  $(\tilde{u}_0, \tilde{u}_1)$  in  $H_0^1(\Omega) \times L^2(\Omega)$ , with*

$$\|(u_0, u_1)\|_{H^1 \times L^2} \leq R_0; \quad \|(\tilde{u}_0, \tilde{u}_1)\|_{H^1 \times L^2} \leq R_0$$

*there exists  $g \in L^\infty([0, T], L^2(\Omega))$  supported in  $[0, T] \times \omega$  such that the unique strong solution of*

$$\begin{cases} \square u + \beta u + f(u) &= g & \text{on } [0, T] \times \Omega \\ (u(0), \partial_t u(0)) &= (u_0, u_1). \end{cases}$$

*satisfies  $(u(T), \partial_t u(T)) = (\tilde{u}_0, \tilde{u}_1)$ .*

The second application of our results concerns the existence of a compact global attractor. A compact global attractor is a compact set, which is invariant by the flow of the PDE and which attracts the bounded sets. The existence of such an attractor is an important dynamical property because it roughly says that the dynamics of the PDE may be reduced

to dynamics on a compact set, which is often finite-dimensional. See for example [24] and [45] for a review on this concept. Theorems 1.1 and 1.2 show that  $\{0\}$  is a global attractor for the damped wave equation (1.1). Of course, it is possible to obtain a more complex attractor by considering an equation of the type

$$\begin{cases} \partial_{tt}^2 u + \gamma(x)\partial_t u = \Delta u - \beta u - f(x, u) & (x, t) \in \Omega \times \mathbb{R}_+, \\ u(x, t) = 0 & (x, t) \in \partial\Omega \times \mathbb{R}_+ \\ (u, \partial_t u) = (u_0, u_1) \in H_0^1 \times L^2 \end{cases} \quad (1.9)$$

where  $f \in C^\infty(\Omega \times \mathbb{R}, \mathbb{R})$  is real analytic with respect to  $u$  and satisfies the following properties. There exist  $C > 0$ ,  $p \in [1, 5)$  and  $R > 0$  such that for all  $(x, u) \in \Omega \times \mathbb{R}$ ,

$$|f(x, u)| \leq C(1 + |u|)^p, \quad |f'_x(x, u)| \leq C(1 + |u|)^p, \quad |f'_u(x, u)| \leq C(1 + |u|)^{p-1} \quad (1.10)$$

$$(x \notin B(x_0, R) \text{ or } |u| \geq R) \implies f(x, u)u \geq 0. \quad (1.11)$$

where  $x_0$  denotes a fixed point of the manifold.

**Theorem 1.4.** *Assume that  $f$  is as above. Then, the dynamical system generated by (1.9) in  $H_0^1(\Omega) \times L^2(\Omega)$  is gradient and admits a compact global attractor  $\mathcal{A}$ .*

Of course, we would get the same result for  $f$  in a generic set similar to the one of Theorem 1.2.

We begin this paper by setting our main notations and recalling the basic properties of Equation (1.1) in Section 2. We recall the unique continuation property of Robbiano and Zuily in Section 3, whereas Sections 4 and 5 are concerned by the asymptotic compactness and the asymptotic smoothing effect of the damped wave equation. The proofs of our main results, Theorem 1.1 and 1.2, are given in Sections 6 and 7 respectively. An alternative proof, using more usual arguments from control theory, is sketched in Section 8. Finally, Theorems 1.3 and 1.4 are discussed in Section 9.

**Acknowledgements:** the second author was financed by the ERC grant *GeCoMethods* during part of the redaction of this article. Moreover, both authors benefited from the fruitful atmosphere of the conference Partial differential equations, optimal design and numerics in Benasque. We also would like to thank Mathieu Léautaud for his remarks about the optimality of Hypothesis (GCC) in the nonlinear context and Geneviève Raugel for her help for removing a non-natural hypothesis of Theorem 1.4.

## 2 Notations and basic properties of the damped wave equation

In this paper, we use the following notations:

$$U = (u, u_t), \quad F = (0, f), \quad A = \begin{pmatrix} 0 & Id \\ \Delta - \beta & -\gamma \end{pmatrix}.$$



In this setting, (1.1) becomes

$$\partial_t U(t) = AU(t) + F(U) .$$

We set  $X = H_0^1(\Omega) \times L^2(\Omega)$  and for  $s \in [0, 1]$ ,  $X^s$  denotes the space  $D((-\Delta + \beta)^{(s+1)/2}) \times D((-\Delta + \beta)^{s/2}) = (H^{1+s}(\Omega) \cap H_0^1(\Omega)) \times H^s(\Omega)$ . Notice that  $X^0 = X$  and  $X^1 = D(A)$  (even if  $\gamma$  is only in  $L^\infty$ ).

We recall that  $E$  denotes the energy defined by (1.4). We also emphasize that (1.2) and the invertibility of  $\Delta - \beta$  implies that a set is bounded in  $X$  if and only if its energy  $E$  is bounded. Moreover, for all  $E_0 \geq 0$ , there exists  $C > 0$  such that

$$\forall (u, v) \in X, E(u, v) \leq E_0 \implies \frac{1}{C} \|(u, v)\|_X^2 \leq E(u, v) \leq C \|(u, v)\|_X^2 \quad (2.1)$$

To simplify some statements in the proofs, we assume without loss of generality that  $3 < p < 5$ . It will avoid some meaningless statements with negative Lebesgue exponents since  $p = 3$  is the exponent where Strichartz estimates are no more necessary and can be replaced by Sobolev embeddings.

We recall that  $\Omega$  is endowed with a metric  $g$ . We denote by  $d$  the distance on  $\Omega$  defined by

$$d(x, y) = \inf \{l(c) \mid c \in C^\infty([0, 1], \Omega) \text{ with } c(0) = x \text{ and } c(1) = y\}$$

where  $l(c)$  is the length of the path  $c$  according to the metric  $g$ . A ball  $B(x, R)$  in  $\Omega$  is naturally defined by

$$B(x, R) = \{y \in \Omega, d(x, y) < R\} .$$

For instance, if  $\Omega = \mathbb{R}^3 \setminus B_{\mathbb{R}^3}(0, 1)$ , the distance between  $(0, 0, 1)$  and  $(0, 0, -1)$  is  $\pi$  (and not 2) and the ball  $B((0, 0, 1), \pi)$  has nothing to do with the classical ball  $B_{\mathbb{R}^3}((0, 0, 1), \pi)$  of  $\mathbb{R}^3$ .

## 2.1 Cauchy problem

The global existence and uniqueness of solutions of the subcritical wave equation (1.1) with  $\gamma \equiv 0$  has been studied by Ginibre and Velo in [21] and [22]. Their method also applies for  $\gamma \neq 0$  since this term is linear and well defined in the energy space  $X$ . Moreover, their argument to prove uniqueness also yields the continuity of the solutions with respect to the initial data.

The central argument is the use of Strichartz estimates.

**Theorem 2.1** (Strichartz estimates).

Let  $T > 0$  and  $(q, r)$  satisfying

$$\frac{1}{q} + \frac{3}{r} = \frac{1}{2}, \quad q \in [7/2, +\infty]. \quad (2.2)$$

There exists  $C = C(T, q) > 0$  such that for every  $G \in L^1([0, T], L^2(\Omega))$  and every  $(u_0, u_1) \in X$ , the solution  $u$  of

$$\begin{cases} \square u + \gamma(x)\partial_t u &= G(t) \\ (u, \partial_t u)(0) &= (u_0, u_1) \end{cases}$$

satisfies the estimate

$$\|u\|_{L^q([0, T], L^r(\Omega))} \leq C \left( \|u_0\|_{H^1(\Omega)} + \|u_1\|_{L^2(\Omega)} + \|G\|_{L^1([0, T], L^2(\Omega))} \right).$$

The result was stated in the Euclidean space  $\mathbb{R}^3$  by Strichartz [48] and Ginibre and Velo with  $q \in (2, +\infty]$ . Kapitanski extended the result to variable coefficients in [33]. On a bounded domain, the first estimates were proved by Burq, Lebeau and Planchon [12] for  $q \in [5, +\infty]$  and extended to a larger range by Blair, Smith and Sogge in [8]. Note that, thanks to the counterexamples of Ivanovici [31], we know that we cannot expect some Strichartz estimates in the full range of exponents in the presence of boundaries.

From these results, we deduce the estimates for the damped wave equation by absorption for  $T$  small enough. We can iterate the operation in a uniform number of steps. Actually, for the purpose of the semilinear wave equation, it is sufficient to consider the Strichartz estimate  $L^{\frac{2p}{p-3}}([0, T], L^{2p}(\Omega))$  which gives  $u^p \in L^{\frac{2}{p-3}}([0, T], L^2(\Omega)) \subset L^1([0, T], L^2(\Omega))$  because  $1 < \frac{2}{p-3} < +\infty$ .

**Theorem 2.2** (Cauchy problem).

Let  $f$  satisfies (1.2). Then, for any  $(u_0, u_1) \in X = H_0^1(\Omega) \times L^2(\Omega)$  there exists a unique solution  $u(t)$  of the subcritical damped wave equation (1.1). Moreover, this solution is defined for all  $t \in \mathbb{R}$  and its energy  $E(u(t))$  is non-increasing in time.

For any  $E_0 \geq 0$ ,  $T \geq 0$  and  $(q, r)$  satisfying (2.2), there exists a constant  $C$  such that, if  $u$  is a solution of (1.1) with  $E(u(0)) \leq E_0$ , then

$$\|u\|_{L^q([0, T], L^r(\Omega))} \leq C \left( \|u_0\|_{H^1(\Omega)} + \|u_1\|_{L^2(\Omega)} \right).$$

In addition, for any  $E_0 \geq 0$  and  $T \geq 0$ , there exists a constant  $C$  such that, if  $u$  and  $\tilde{u}$  are two solutions of (1.1) with  $E(u(0)) \leq E_0$  and  $E(\tilde{u}(0)) \leq E_0$ , then

$$\sup_{t \in [-T, T]} \|(u, \partial_t u)(t) - (\tilde{u}, \partial_t \tilde{u})(t)\|_X \leq C \|(u, \partial_t u)(0) - (\tilde{u}, \partial_t \tilde{u})(0)\|_X.$$

**Proof:** The existence and uniqueness for small times is a consequence of Strichartz estimates and of the subcriticality of the nonlinearity, see [22]. The solution can be globalized backward and forward in time thanks to the energy estimates (1.5) for smooth solutions. Indeed,

$$E(t) \leq E(s) + C \int_t^s E(\tau) d\tau$$

and thus, Gronwall inequality for  $t \leq s$  and the decay of energy for  $t \geq s$  show that the energy does not blow up in finite time. This allows to extend the solution for all times since the energy controls the norm of the space  $X$  by (2.1).

For the uniform continuity estimate, we notice that  $w = u - \tilde{u}$  is solution of

$$\begin{cases} \square w + \beta w + \gamma(x)\partial_t w &= -wg(u, \tilde{u}) \\ (w, \partial_t w)(0) &= (u, \partial_t u)(0) - (\tilde{u}, \partial_t \tilde{u})(0) \end{cases}$$

where  $g(s, \tilde{s}) = \int_0^1 f'(s + \tau(\tilde{s} - s)) d\tau$  fulfills  $|g(s, \tilde{s})| \leq C(1 + |s|^{p-1} + |\tilde{s}|^{p-1})$ . Let  $q = \frac{2p}{p-3}$ , Strichartz and Hölder estimates give

$$\begin{aligned} \|(w, \partial_t w)(t)\|_{L^\infty([0, T], X) \cap L^q([0, T], L^{2p})} &\leq C\|(w, \partial_t w)(0)\|_X + C\|wg(u, \tilde{u})\|_{L^1([0, T], L^2)} \\ &\leq C\|(w, \partial_t w)(0)\|_X + CT\|w\|_{L^\infty([0, T], L^2)} \\ &\quad + T^\theta \|w\|_{L^q([0, T], L^{2p})} \left( \|u\|_{L^q([0, T], L^{2p})}^{p-1} + \|\tilde{u}\|_{L^q([0, T], L^{2p})}^{p-1} \right) \end{aligned}$$

with  $\theta = \frac{5-p}{2} > 0$ . We get the expected result for  $T$  small enough by absorption since we already know a uniform bound (depending on  $E_0$ ) for the Strichartz norms of  $u$  and  $\tilde{u}$ . Then, we iterate the operation to get the result for large  $T$ .  $\square$

## 2.2 Exponential decay of the linear semigroup

In this paper, we will strongly use the exponential decay for the linear semigroup in the case where  $\gamma$  may vanish but satisfies the geometric assumptions of this paper. In this case, (1.3) enables to control the decay of energy outside a large ball and the geometric control condition (GCC) enables to control the energy trapped in this ball.

**Proposition 2.3.** *Assume that  $\gamma \in L^\infty(\Omega)$  satisfies (1.3) and (GCC). There exist two positive constants  $C$  and  $\lambda$  such that*

$$\forall s \in [0, 1], \forall t \geq 0, \quad \|e^{At}\|_{\mathcal{L}(X^s)} \leq Ce^{-\lambda t}.$$

The exponential decay of the damped wave equation under the geometric control condition is well known since the works of Rauch and Taylor on a compact manifold [44] and Bardos, Lebeau and Rauch [7, 6] on a bounded domain. Yet, we did not find any reference for unbounded domain ([3] and [35] concern unbounded domains but local energy only). It is noteworthy that the decay of the linear semigroup in unbounded domains seems not to have been extensively studied for the moment.

We give a proof of Proposition 2.3 using microlocal defect measure as done in Lebeau [39] or Burq [10] (see also [11] for the proof of the necessity). The only difference with

respect to these results is that the manifold that we consider may be unbounded. Since microlocal defect measure only reflects the local propagation, we thus have to use the property of equipartition of the energy to deal with the energy at infinity and to show a propagation of compactness (see [17] for the flat case).

**Lemma 2.4.** *Let  $T > L$  where  $L$  is given by (GCC). Assume that  $(U_{n,0}) \subset X$  is a bounded sequence, which weakly converges to 0 and assume that  $U_n(t) = (u_n(t), \partial_t u_n(t)) = e^{At} U_{n,0}$  satisfies*

$$\int_0^T \int_{\Omega} \gamma(x) |\partial_t u_n|^2 \rightarrow 0. \quad (2.3)$$

Then,  $(U_{n,0})$  converges to 0 strongly in  $X$ .

**Proof:** Let  $\mu$  be a microlocal defect measure associated to  $(u_n)$  (see [18], [50] or [9] for the definition). Note that (2.3) implies that  $\mu$  can also be associated to the solution of the wave equation without damping, so the weak regularity of  $\gamma$  is not problematic for the propagation and we get that  $\mu$  is concentrated on  $\{\tau^2 - |\xi|_x^2 = 0\}$  where  $(\tau, \xi)$  are the dual variables of  $(t, x)$ . Moreover, (2.3) implies that  $\gamma \tau^2 \mu = 0$  and so  $\mu \equiv 0$  on  $S^*(]0, T[ \times \omega)$ . Then, using the propagation of the measure along the generalized bicharacteristic flow of Melrose-Sjöstrand and the geometric control condition satisfied by  $\omega$ , we obtain  $\mu \equiv 0$  everywhere. We do not give more details about propagation of microlocal defect measure and refer to the Appendix [39] or Section 3 of [9] (see also [20] for some close propagation results in a different context). Since  $\mu \equiv 0$ , we know that  $U_n \rightarrow 0$  on  $H^1 \times L^2(]0, T[ \times B(x_0, R))$  for every  $R > 0$ .

To finish the proof, we need the classical equipartition of the energy to get the convergence to 0 in the whole manifold  $\Omega$ . Since  $\gamma$  is uniformly positive outside a ball  $B(x_0, R)$ , (2.3) and the previous arguments imply that  $\partial_t u_n \rightarrow 0$  in  $L^2([0, T] \times \Omega)$ . Let  $\varphi \in C_0^\infty(]0, T[)$  with  $\varphi \geq 0$  and  $\varphi(t) = 1$  for  $t \in [\varepsilon, T - \varepsilon]$ . We multiply the equation by  $\varphi(t)u_n$  and we obtain

$$\begin{aligned} 0 &= - \iint_{[0, T] \times \Omega} \varphi(t) |\partial_t u_n|^2 - \iint_{[0, T] \times \Omega} \varphi'(t) \partial_t u_n u_n + \iint_{[0, T] \times \Omega} \varphi(t) |\nabla u_n|^2 \\ &\quad + \iint_{[0, T] \times \Omega} \varphi(t) \beta |u_n|^2 + \iint_{[0, T] \times \Omega} \varphi(t) \gamma(x) \partial_t u_n u_n. \end{aligned}$$

The  $L^2$ -norm of  $u_n(t)$  is bounded, while  $\partial_t u_n \rightarrow 0$  in  $L^2([0, T] \times \Omega)$ , so the first, second and fifth terms converge to zero. Then, the above equation yields

$$\iint_{[0, T] \times \Omega} \varphi(t) (\beta |u_n|^2 + |\nabla u_n|^2) \rightarrow 0.$$

Finally, notice that the energy identity  $\|U_{n,0}\|_X^2 = \|U_n(t)\|_X^2 + \int_0^T \int_{\Omega} \gamma(x) |\partial_t u_n|^2$  shows that

$$\iint_{[0, T] \times \Omega} \varphi(t) (\beta |u_n|^2 + |\nabla u_n|^2) \sim \|U_{n,0}\|_X^2 \int_0^T \varphi(t)$$

and thus that  $\|U_{n,0}\|_X$  goes to zero.  $\square$

**Proof of Proposition 2.3:** Once Lemma 2.4 is established, the proof follows the arguments of the classical case, where  $\Omega$  is bounded. We briefly recall them.

We first treat the case  $s = 0$ . As in Proposition 2.5, the exponential decay of the energy is equivalent to the observability estimate, that is the existence of  $C > 0$  and  $T > 0$  such that, for any trajectory  $U(t) = e^{At}U_0$  in  $X$ ,

$$\int_0^T \int_{\Omega} \gamma(x) |\partial_t u|^2 \geq C \|U(0)\|_X^2 .$$

We argue by contradiction: assume that (2.4) does not hold for any positive  $T$  and  $C$ . Then, there exists a sequence of initial data  $U_n(0)$  with  $\|U_n(0)\|_X = 1$  and such that

$$\int_0^n \int_{\Omega} \gamma(x) |\partial_t u_n(t, x)|^2 dt dx \xrightarrow{n \rightarrow +\infty} 0 ,$$

where  $(u_n, \partial_t u_n)(t) = U_n(t) = e^{At}U_n(0)$ . Let  $\tilde{U}_n = U_n(n/2 + \cdot)$ . We have

$$\int_{-n/2}^{n/2} \int_{\Omega} \gamma(x) |\partial_t \tilde{u}_n(t, x)|^2 dt dx \xrightarrow{n \rightarrow \infty} 0 ,$$

and, for any  $t \in [-n/2, n/2]$ ,

$$\|\tilde{U}_n(t)\|_X^2 = \|\tilde{U}_n(-n/2)\|_X^2 - \int_{-n/2}^t \int_{\Omega} \gamma(x) |\partial_t \tilde{u}_n(s, x)|^2 ds dx \xrightarrow{n \rightarrow \infty} 1 .$$

Therefore, we can assume that  $U_n(0)$  converges to  $U_{\infty}(0) \in X$ , weakly in  $X$ . Moreover, for any  $T > 0$ ,  $U_n(t)$  and  $\partial_t U_n(t)$  are bounded in  $L^{\infty}([-T, T], X)$  and  $L^{\infty}([-T, T], L^2(\Omega) \times H^{-1}(\Omega))$  respectively. Thus, using Ascoli's Theorem, we may also assume that  $U_n(t)$  strongly converges to  $U_{\infty}(t)$  in  $L^{\infty}([-T, T], L^2(K) \times H^{-1}(K))$  where  $K$  is any compact of  $\Omega$ . Hence,  $(u_{\infty}, \partial_t u_{\infty})(t) = U_{\infty}(t) = e^{At}U_{\infty}(0)$  is a solution of

$$\begin{cases} \square u_{\infty} + \beta u_{\infty} = 0 & \text{on } \mathbb{R} \times \Omega \\ \partial_t u_{\infty} = 0 & \text{on } \mathbb{R} \times \omega. \end{cases} \quad (2.4)$$

in  $L^2 \times H^{-1}$ . Since  $U_{\infty}(0) \in X$  belongs to  $X$ , we deduce that, in fact,  $U_{\infty}(t)$  solves (2.4) in  $X$ .

To finish the proof of Proposition 2.3, we have to show that  $U_{\infty} \equiv 0$ . Indeed, applying Lemma 2.4, we would get that  $U_n$  converges strongly to 0, which contradicts the hypothesis  $\|U_n(0)\|_X = 1$ . Note that  $U_{\infty} \equiv 0$  is a direct consequence of a unique continuation property as Corollary 3.2. However, Corollary 3.2 requires  $\Omega$  to be smooth, whereas Proposition

2.3 could be more general. Therefore, we recall another classical argument to show that  $U_\infty \equiv 0$ .

Denote  $N$  the set of function  $U_\infty(0) \in X$  satisfying (2.4), which is obviously a linear subspace of  $X$ . We will prove that  $N = \{0\}$ . Since  $\gamma(x)|\partial_t u_\infty|^2 \equiv 0$  for functions  $u_\infty$  in  $N$  and since  $N$  is a closed subspace, Lemma 2.4 shows that any weakly convergent subsequence of  $N$  is in fact strongly convergent. By Riesz Theorem,  $N$  is therefore finite dimensional. For any  $t \in \mathbb{R}$ ,  $e^{tA}$  applies  $N$  into itself and thus  $A|_N$  is a bounded linear operator. Assume that  $N \neq \{0\}$ , then  $A|_N$  admits an eigenvalue  $\lambda$  with eigenvector  $Y = (y_0, y_1) \in N$ . This means that  $y_1 = \lambda y_0$  and that  $(\Delta - \beta)y_0 = \lambda^2 y_0$ . Moreover, we know that  $y_1 = 0$  on  $\omega$  and so, if  $\lambda \neq 0$ , that  $y_0 = 0$  on  $\omega$ . This implies  $y_0 \equiv 0$  by the unique continuation property of elliptic operators. Finally, if  $\lambda = 0$ , we have  $(\Delta - \beta)y_0 = 0$  and  $y_0 = 0$ , because, by assumption,  $\Delta - \beta$  is a negative definite operator.

So we have proved  $N = \{0\}$  and therefore  $U_\infty = 0$ , that is  $\tilde{U}_n(0)$  converges to 0 weakly in  $X$ . We can then apply Lemma 2.4 on any interval  $[-n/2, -n/2 + T]$  where  $L$  is the time in the geometric control condition (GCC) and obtain a contradiction to  $\|U_n(0)\|_X = 1$ .

Let us now consider the cases  $s \in (0, 1]$ . The basic semigroup properties (see [43]) shows that, if  $U \in X^1 = D(A)$ , then  $e^{At}U$  belongs to  $D(A)$  and

$$\begin{aligned} \|e^{At}U\|_{X^1} &= \|Ae^{At}U\|_X + \|e^{At}U\|_X = \|e^{At}AU\|_X + \|e^{At}U\|_X \\ &\leq Ce^{-\lambda t} (\|AU\|_X + \|U\|_X) = Ce^{-\lambda t} \|U\|_{D(A)}. \end{aligned}$$

This shows Proposition 2.3 for  $s = 1$ . Notice that we do not have to require any regularity for  $\gamma$  to obtain this result. Then, Proposition 2.3 for  $s \in (0, 1)$  follows by interpolating between the cases  $s = 0$  and  $s = 1$  (see [49]).  $\square$

## 2.3 First nonlinear exponential decay properties

Theorem 2.2 shows that the energy  $E$  is non-increasing along the solutions of (1.1). The purpose of this paper is to obtain the exponential decay of this energy in the sense of property (ED) stated above. We first recall the well-known criterion for exponential decay.

**Proposition 2.5.** *The exponential decay property (ED) holds if and only if, there exist  $T$  and  $C$  such that*

$$E(u(0)) \leq C(E(u(0)) - E(u(T))) = C \int \int_{[0, T] \times \Omega} \gamma(x) |\partial_t u(x, t)|^2 dt dx \quad (2.5)$$

for all solutions  $u$  of (1.1) with  $E(u(0)) \leq E_0$ .

**Proof:** If (ED) holds then obviously (2.5) holds for  $T$  large enough since  $E(u(0)) - E(u(T)) \geq (1 - Ke^{-\lambda T})E(u(0))$ . Conversely, if (2.5) holds, using  $E(u(T)) \leq E(u(0))$ , we

get  $E(u(T)) \leq C/(C+1)E(u(0))$  and thus  $E(u(kT)) \leq (C/(C+1))^k E(u(0))$ . Using again the decay of the energy to fill the gaps  $t \in (kT, (k+1)T)$ , this shows that (ED) holds.  $\square$

First, we prove exponential decay in the case of positive damping, which will be helpful to study what happens outside a large ball since (1.3) is assumed in the whole paper. Note that the fact that  $-\Delta + \beta$  is positive is necessary to avoid for instance the constant undamped solutions.

**Proposition 2.6.** *Assume that  $\omega = \Omega$ , that is that  $\gamma(x) \geq \alpha > 0$  everywhere. Then (ED) holds.*

**Proof:** We recall here the classical proof. We introduce a modified energy

$$\tilde{E}(u) = \int_{\Omega} \frac{1}{2} (|\partial_t u|^2 + |\nabla u|^2 + \beta|u|^2) + V(u) + \varepsilon u \partial_t u$$

with  $\varepsilon > 0$ . Since  $\int_{\Omega} |\nabla u|^2 + \beta|u|^2$  controls  $\|u\|_{L^2}^2$ ,  $\tilde{E}$  is equivalent to  $E$  for  $\varepsilon$  small enough and it is sufficient to obtain the exponential decay of the auxiliary energy  $\tilde{E}$ . Using  $\gamma \geq \alpha > 0$  and  $uf(u) \geq 0$ , a direct computation shows for  $\varepsilon$  small enough that

$$\begin{aligned} \tilde{E}(u(T)) - \tilde{E}(u(0)) &= \int_0^T \int_{\Omega} -\gamma(x)|\partial_t u|^2 + \varepsilon|\partial_t u|^2 + \varepsilon\gamma(x)u\partial_t u - \varepsilon(|\nabla u|^2 + \beta|u|^2) - \varepsilon uf(u) \\ &\leq -C \int_0^T \|(u, \partial_t u)\|_{H^1 \times L^2}^2 \\ &\leq -C \int_0^T \tilde{E}(t) dt \\ &\leq -CT\tilde{E}(T), \end{aligned}$$

where  $C$  is some positive constant, not necessarily the same from line to line. Thus,  $\tilde{E}(u(0)) - \tilde{E}(u(T)) \geq CT\tilde{E}(u(T))$  with  $CT > 0$  and therefore  $\tilde{E}(u(0)) \geq \mu\tilde{E}(u(T))$  with  $\mu > 1$ . As in the proof of Proposition 2.3, this last property implies the exponential decay of  $\tilde{E}$  and thus the one of  $E$ .  $\square$

### 3 A unique continuation result for equations with partially holomorphic coefficients

Comparatively to previous articles on the stabilization of the damped wave equations as [17], one of the main novelties of this paper is the use of a unique continuation theorem

requiring partially analyticity of the coefficients, but very weak geometrical assumptions as shown in Corollary 3.2. We use here the result of Robbiano and Zuily in [47]. This result has also been proved independently by Hörmander in [30] and has been generalised by Tataru in [52]. Note that the idea of using partial analyticity for unique continuation was introduced by Tataru [51] but it requires some global analyticity assumptions that are not fulfilled in our case. All these results use very accurate microlocal analysis and hold in a much more general framework than the one of the wave equation. However, for sake of simplicity, we restrict the statement to this case.

**Theorem 3.1. Robbiano-Zuily, Hörmander (1998)**

Let  $d \geq 1$ ,  $(x_0, t_0) \in \mathbb{R}^d \times \mathbb{R}$  and let  $\mathcal{U}$  be a neighbourhood of  $(x_0, t_0)$ . Let  $(A_{i,j}(x, t))_{i,j=1,\dots,d}$ ,  $b(x, t)$ ,  $(c_i(x, t))_{i=1,\dots,d}$  and  $d(x, t)$  be bounded coefficients in  $\mathcal{C}^\infty(\mathcal{U}, \mathbb{R})$ . Let  $v$  be a strong solution of

$$\partial_{tt}^2 v = \operatorname{div}(A(x, t)\nabla v) + b(x, t)\partial_t v + c(x, t) \cdot \nabla v + d(x, t)v \quad (x, t) \in \mathcal{U} \subset \mathbb{R}^d \times \mathbb{R}. \quad (3.1)$$

Let  $\varphi \in \mathcal{C}^2(\mathcal{U}, \mathbb{R})$  such that  $\varphi(x_0, t_0) = 0$  and  $(\nabla\varphi, \partial_t\varphi)(x, t) \neq 0$  for all  $(x, t) \in \mathcal{U}$ . Assume that:

- (i) the coefficients  $A$ ,  $b$ ,  $c$  and  $d$  are analytic in time,
  - (ii)  $A(x_0, t_0)$  is a symmetric positive definite matrix,
  - (iii) the hypersurface  $\{(x, t) \in \mathcal{U}, \varphi(x, t) = 0\}$  is not characteristic at  $(x_0, t_0)$  that is that we have  $|\partial_t\varphi(x_0, t_0)|^2 \neq \langle \nabla\varphi(x_0, t_0) | A(x_0, t_0)\nabla\varphi(x_0, t_0) \rangle$
  - (iv)  $v \equiv 0$  in  $\{(x, t) \in \mathcal{U}, \varphi(x, t) \leq 0\}$ .
- Then,  $v \equiv 0$  in a neighbourhood of  $(x_0, t_0)$ .

**Proof:** We only have to show that Theorem 3.1 is a direct translation of Theorem A of [47] in the framework of the wave equation. To use the notations of [47], we set  $x_a$  to be the time variable and  $x_b$  the space variable and we set  $(x_0, t_0) = x^0 = (x_b^0, x_a^0)$ . Equation (3.1) corresponds to the differential operator

$$P = \xi_a^2 - {}^t\xi_b A(x_b, x_a)\xi_b - b(x_b, x_a)\xi_a - c(x_b, x_a)\xi_b - d(x_b, x_a)$$

with principal symbol  $p_2 = \xi_a^2 - {}^t\xi_b A(x_b, x_a)\xi_b$ .

All the statement of Theorem 3.1 is an obvious translation of Theorem A of [47], except maybe for the fact that Hypothesis (iii) implies the hypothesis of pseudo-convexity of [47]. We compute  $\{p_2, \varphi\} = 2\xi_a\varphi'_a - 2{}^t\xi_b A(x_a, x_b)\varphi'_b$ . Let us set  $\zeta = (x_a^0, x_b^0, i\varphi'_a(x^0), \xi_b + i\varphi'_b(x^0))$ , then  $\{p_2, \varphi\}(\zeta) = 0$  if and only if

$$i(\varphi'_a(x^0))^2 - i{}^t\varphi'_b(x^0)A(x^0)\varphi'_b(x^0) - {}^t\xi_b A(x^0)\varphi'_b(x^0) = 0.$$

This is possible only if  $(\varphi'_a(x^0))^2 = {}^t\varphi'_b(x^0)A(x^0)\varphi'_b(x^0)$ , that is if the hypersurface  $\varphi = 0$  is characteristic at  $(x_0, t_0)$ . Thus, if this hypersurface is not characteristic, then the pseudo-convexity hypothesis of Theorem A of [47] holds.  $\square$



The previous theorem allows to prove some unique continuation result with some optimal time and geometric assumption. This allows to prove unique continuation where the geometric condition is only, roughly speaking, that we do not contradict the finite speed of propagation.

**Corollary 3.2.** *Let  $T > 0$  (or  $T = +\infty$ ) and let  $b, (c_i)_{i=1,2,3}$  and  $d$  be smooth coefficients in  $\mathcal{C}^\infty(\Omega \times [0, T], \mathbb{R})$ . Assume moreover that  $b, c$  and  $d$  are analytic in time and that  $v$  is a strong solution of*

$$\partial_{tt}^2 v = \Delta v + b(x, t) \partial_t v + c(x, t) \cdot \nabla v + d(x, t) v \quad (x, t) \in \Omega \times (-T, T) . \quad (3.2)$$

Let  $\mathcal{O}$  be a non-empty open subset of  $\Omega$  and assume that  $v(x, t) = 0$  in  $\mathcal{O} \times (-T, T)$ . Then  $v(x, 0) \equiv 0$  in  $\mathcal{O}_T = \{x_0 \in \Omega, d(x_0, \mathcal{O}) < T\}$ .

As consequences:

- a) if  $T = +\infty$ , then  $v \equiv 0$  everywhere,
- b) if  $v \equiv 0$  in  $\mathcal{O} \times (-T, T)$  and  $\overline{\mathcal{O}}_T = \Omega$ , then  $v \equiv 0$  everywhere.

**Proof:** Since  $\Omega$  is assumed to be connected, both consequences are obvious from the first statement.

Let  $x_0$  be given such that  $d(x_0, \mathcal{O}) < T$ . There is a point  $x_* \in \mathcal{O}$  linked to  $x_0$  by a smooth curve of length  $l < T$ , which stays away from the boundary. We introduce a sequence of balls  $B(x_0, r), \dots, B(x_K, r)$  with  $r \in (0, T/K)$ ,  $x_{k-1} \in B(x_k, r)$  and  $x_K = x_*$ , such that  $B(x_k, r)$  stays away from the boundary and is small enough such that it is diffeomorphic to an open set of  $\mathbb{R}^3$  via the exponential map. Note that such a sequence of balls exists because the smooth curve linking  $x_0$  to  $x_K$  is compact and of length smaller than  $T$ . We also notice that it is sufficient to prove Corollary 3.2 in each ball  $B(x_k, r)$ . Indeed, this would enable us to apply Corollary 3.2 in  $B(x_K, r) \times (-T, T)$  to obtain that  $v$  vanishes in a neighbourhood of  $x_{K-1}$  for  $t \in (-T+r, T-r)$  and then to apply it recursively in  $B(x_{K-1}, r) \times (-T+r, T-r), \dots, B(x_1, r) \times (-T+(K-1)r, T-(K-1)r)$  to obtain that  $v(x_0, 0) = 0$ .

From now on, we assume that  $x_0 \in B(x_*, r)$  and that  $v$  vanishes in a neighbourhood  $\mathcal{O}$  of  $x_*$  for  $t \in (-r, r)$ . Since  $d(x_0, x_*) < r$ , we can introduce a non-negative function  $h \in \mathcal{C}^\infty([-r, r], \mathbb{R})$  such that  $h(0) > d(x_0, x_*)$ ,  $h(\pm r) = 0$  and  $|h'(t)| < 1$  for all  $t \in [-r, r]$ . We set  $\mathcal{U} = B(x_*, r) \times (-r, r)$  and for any  $\lambda \in [0, 1]$ , we define

$$\varphi_\lambda(x, t) = d(x, x_*)^2 - \lambda h(t)^2 .$$

Since  $r$  is assumed to be smaller than the radius of injectivity of the exponential map,  $\varphi_\lambda$  is a smooth well-defined function. We prove Corollary 3.2 by contradiction. Assume that  $v(x_0, 0) \neq 0$ . We denote by  $V_\lambda$  the volume  $\{(x, t) \in \mathcal{U}, \varphi_\lambda(x, t) \leq 0\}$ . We notice that  $V_{\lambda_1} \subset V_{\lambda_2}$  if  $\lambda_1 < \lambda_2$ , that for small  $\lambda$ ,  $V_\lambda$  is included in  $\mathcal{O} \times (-r, r)$  where  $v$  vanishes, and that  $V_1$  contains  $(x_0, 0)$  where  $v$  does not vanish. Thus

$$\lambda_0 = \sup\{\lambda \in [0, 1], \forall (x, t) \in V_\lambda, v(x, t) = 0\}$$

is well defined and belongs to  $(0, 1)$ . For  $t$  close to  $-r$  or  $r$ ,  $h(t)$  is small and the section  $\{x, (x, t) \in V_{\lambda_0}\}$  of  $V_{\lambda_0}$  is contained in  $\mathcal{O}$  where  $v$  vanishes. Therefore, by compactness, the hypersurface  $S_{\lambda_0} = \partial V_{\lambda_0}$  must touch the support of  $v$  at some point  $(x_1, t_1) \in \mathcal{U}$  (see Figure 1).




Figure 1: *The proof of Corollary 3.2*

In local coordinates,  $\Delta$  can be written  $\text{div}(A(x)\nabla\cdot) + c(x)\cdot\nabla$ . Moreover,  $\langle \nabla\varphi_\lambda | A\nabla\varphi_\lambda \rangle = |\nabla_g d(\cdot, x_*)|_g^2 = 1$  where the index  $g$  means that the gradient and norm are taken according to the metric. Therefore, the hypersurface  $S_{\lambda_0}$  is non-characteristic at  $(x_1, t_1)$  in the sense of Hypothesis (iii) of Theorem 3.1 since  $|\partial_t\varphi_\lambda(x, t)| = |\lambda h'(t_1)| < 1$ . Thus, we can apply Theorem 3.1 with  $\varphi = \varphi_{\lambda_0}$  at the point  $(x_1, t_1)$ , mapping everything in the 3d-euclidean frame via the exponential chart. We get that  $v$  must vanish in a neighbourhood of  $(x_1, t_1)$ . This is obviously a contradiction since  $(x_1, t_1)$  has been taken in the support of  $v$ .  $\square$

## 4 Asymptotic compactness

As soon as  $t$  is positive, a solution  $u(t)$  of a parabolic PDE becomes smooth and stays in a compact set. The smoothing effect in finite time of course fails for the damped wave equations. However, these PDEs admit in some sense a smoothing effect in infinite time. This effect is called *asymptotic compactness* if one is interested in extracting asymptotic subsequences as in Proposition 4.3, or *asymptotic smoothness* if one uses the regularity of globally bounded solutions as in Proposition 4.4. For the reader interested in these notions,

we refer for example to [24]. The proof of this asymptotic smoothing effect is based on the variation of constant formula  $U(t) = e^{At}U_0 + \int_0^t e^{A(t-s)}F(U(s))ds$  and two properties:

- the exponential decay of the linear group (Proposition 2.3), which implies that the linear part  $e^{At}U_0$  asymptotically disappears,
- the regularity of the nonlinearity  $F$  implying the compactness of the nonlinear term  $\int_0^t e^{A(t-s)}F(U(s))ds$  (Corollary 4.2 below). Note that the subcriticality of  $f$  is the key point of this property and that our arguments cannot be extended as they stand to the critical case  $p = 5$ .

The purpose of this section is to prove some compactness and regularity results about undamped solutions as (1.8). Note that these results could also have been obtained with a more “control theoretic” proof (see section 8 for a sketch of the alternative proof) based on propagation results or observability estimates. Here, we have chosen to give a different one using asymptotic regularization, more usual in dynamical system. The spirit of the proof remains quite similar: prove that the nonlinearity is more regular than it seems *a priori* and use some properties of the damped linear equation.

## 4.1 Regularity of the nonlinearity

Since  $f$  is subcritical, it is shown in [17] that the nonlinear term of 1.1 yields a gain of smoothness.

### Theorem 4.1. Dehman, Lebeau and Zuazua (2003)

Let  $\chi \in C_0^\infty(\mathbb{R}^3, \mathbb{R})$ ,  $R > 0$  and  $T > 0$ . Let  $s \in [0, 1)$  and let  $\varepsilon = \min(1 - s, (5 - p)/2, (17 - 3p)/14) > 0$  with  $p$  and  $f$  as in (1.2). There exist  $(q, r)$  satisfying (2.2) and  $C > 0$  such that the following property holds. If  $v \in L^\infty([0, T], H^{1+s}(\mathbb{R}^3))$  is a function with finite Strichartz norms  $\|v\|_{L^q([0, T], L^r(\mathbb{R}^3))} \leq R$ , then  $\chi(x)f(v) \in L^1([0, T], H^{s+\varepsilon}(\mathbb{R}^3))$  and moreover

$$\|\chi(x)f(v)\|_{L^1([0, T], H^{s+\varepsilon}(\mathbb{R}^3))} \leq C\|v\|_{L^\infty([0, T], H^{1+s}(\mathbb{R}^3))}.$$

The constant  $C$  depends only on  $\chi$ ,  $s$ ,  $T$ ,  $(q, r)$ ,  $R$  and the constant in Estimate (1.2).

Theorem 4.1 is a copy of Theorem 8 of [17], except for two points.

First, we would like to apply the result to a solution  $v$  of the damped wave equation on a manifold possibly with boundaries, where not all Strichartz exponents are available. This leads to the constraint  $q \geq 7/2$  for the Strichartz exponents  $(q, r)$  of (2.2) (see Theorem 2.2). In the proof of Theorem 8 of [17], the useful Strichartz estimate corresponds to  $r = \frac{3(p-1)}{1-\varepsilon}$  and  $q = \frac{2(p-1)}{p-3+2\varepsilon}$  and it is required that  $q \geq p - 1$ , which yields  $\varepsilon \leq (5 - p)/2$ . In this paper, we require also that  $q \geq 7/2$  which yields in addition  $\varepsilon \leq (17 - 3p)/14$ . Notice that  $p < 5$  and thus both bounds are positive.

The second difference is that, in [17], the function  $f$  is assumed to be of class  $C^3$  and to satisfy

$$|f''(u)| \leq C(1 + |u|)^{p-2} \quad \text{and} \quad |f^{(3)}(u)| \leq C(1 + |u|)^{p-3} \quad (4.1)$$

in addition of (1.2). Since Theorem 4.1 concerns the  $L^1(H^{s'})$ -norm of  $\chi(x)f(v)$  for  $s' = s + \varepsilon \leq 1$ , we can omit Assumption (4.1). Actually, we make the assumption  $\varepsilon \leq 1 - s$  which is not present in [17] and a careful study of their proof shows that (1.2) is not necessary under that assumption.

Indeed, let  $\tilde{f}(u) = \text{th}^3(u)|u|^p$ . The function  $\tilde{f}$  is of class  $\mathcal{C}^3$  and satisfies (1.2) and (4.1). Hence, Theorem 8 of [17] can be applied to  $\tilde{f}$  and we can bound the  $L^1(H^{s'})$ -norm of  $\tilde{f}$  as in Theorem 4.1. On the other hand, we notice that  $|\tilde{f}(u)| \underset{\pm\infty}{\sim} |u|^p$ ,  $\tilde{f}'(u) \underset{\pm\infty}{\sim} p|u|^{p-1}$  and  $\tilde{f}'(u) \geq 0$ . Therefore, since  $f$  satisfies (1.2), there exists  $C > 0$  such that  $|f(u)| \leq C(1 + |\tilde{f}(u)|)$  and  $|f'(u)| \leq C(1 + \tilde{f}'(u))$ . Thus, if say  $v > u$  for fixing the notations,

$$\begin{aligned} |f(v) - f(u)| &\leq (v - u) \int_0^1 |f'(u + \tau(v - u))| d\tau \\ &\leq C(v - u) + C(v - u) \int_0^1 \tilde{f}'(u + \tau(v - u)) d\tau = C(v - u) + C(\tilde{f}(v) - \tilde{f}(u)) \\ &\leq C|v - u| + C|\tilde{f}(v) - \tilde{f}(u)| . \end{aligned}$$

For  $0 < s < 1$ , using the above inequalities and the definition of the  $H^{s'}$ -norm as

$$\|\chi f(u)\|_{H^{s'}}^2 = \|\chi f(u)\|_{L^2}^2 + \iint_{\mathbb{R}^6} \frac{|\chi(x)f(u(x)) - \chi(y)f(u(y))|^2}{|x - y|^{2s'}} dx dy ,$$

we obtain

$$\|\chi f(u)\|_{L^1(H^{s'})} \leq C\|u\|_{L^\infty(H^1)} + C\|\tilde{\chi}\tilde{f}(u)\|_{L^1(H^{s'})}$$

where  $\tilde{\chi}$  is another cut-off function with larger support. Hence, for  $0 < s < 1$ , the conclusion of Theorem 4.1 holds not only for  $\tilde{f}$  but also for  $f$ . If  $s' = 1$ , we just apply the chain rule and the proof is easier.

Note that the above arguments show that the constant  $C$  depends on  $f$  through Estimate 1.2 only. Notice in addition that, since  $f$  is only  $\mathcal{C}^1$ , we cannot expect  $\chi f(v)$  to be more regular than  $H^1$  and that is why we also assume  $\varepsilon \leq 1 - s$ .

In this paper, we use a generalisation of Theorem 8 of [17] to non-compact manifolds with boundaries.

**Corollary 4.2.** *Let  $R > 0$  and  $T > 0$ . Let  $s \in [0, 1)$  and let  $\varepsilon = \min(1 - s, (5 - p)/2, (17 - 3p)/14) > 0$  with  $p$  as in (1.2). There exist  $(q, r)$  satisfying (2.2) and  $C > 0$  such that the following property holds. If  $v \in L^\infty([0, T], H^{1+s}(\Omega) \cap H_0^1(\Omega))$  is a function with finite Strichartz norms  $\|v\|_{L^q([0, T], L^r(\Omega))} \leq R$ , then  $f(v) \in L^1([0, T], H_0^{s+\varepsilon}(\Omega))$  and moreover*

$$\|f(v)\|_{L^1([0, T], H_0^{s+\varepsilon}(\Omega))} \leq C\|v\|_{L^\infty([0, T], H^{s+1}(\Omega) \cap H_0^1(\Omega))} .$$

The constant  $C$  depends only on  $\Omega$ ,  $(q, r)$ ,  $R$  and the constant in Estimate (1.2).

**Proof:** Since we assumed that  $\Omega$  has a bounded geometry in the sense that  $\Omega$  is compact or a compact perturbation of a manifold with periodic metric,  $\Omega$  can be covered by a set of  $\mathcal{C}^\infty$ -charts  $\alpha_i : U_i \mapsto \alpha_i(U_i) \subset \mathbb{R}^3$  such that  $\alpha_i(U_i)$  is equal either to  $B(0, 1)$  or to  $B_+(0, 1) = \{x \in B(0, 1), x_1 > 0\}$  and such that, for any  $s \geq 0$  the norm of a function  $u \in H^s(\Omega)$  is equivalent to the norm

$$\left( \sum_{i \in \mathbb{N}} \|u \circ \alpha_i^{-1}\|_{H^s(\alpha_i(U_i))}^2 \right)^{1/2} .$$

Moreover, the Strichartz norm  $L^q([0, T], L^r(\alpha_i(U_i)))$  of  $v \circ \alpha_i^{-1}$  is uniformly controlled from above by the Strichartz norm  $L^q([0, T], L^r(U_i))$  of  $v$  which is bounded by  $R$ .

Therefore, it is sufficient to prove that Corollary 4.2 holds for  $\Omega$  being either  $B(0, 1)$  or  $B_+(0, 1)$ . Say that  $\Omega = B_+(0, 1)$ , the case  $\Omega = B(0, 1)$  being simpler. To apply Theorem 4.1, we extend  $v$  in a neighbourhood of  $B_+(0, 1)$  as follows. For  $x \in B_+(0, 2)$ , we use the radial coordinates  $x = (r, \sigma)$  and we set

$$\tilde{v}(x) = \tilde{v}(r, \sigma) = 5v(1 - r, \sigma) - 20v(1 - r/2, \sigma) + 16v(1 - r/4, \sigma) .$$

Then, for  $x = (x_1, x_2, x_3) \in B_-(0, 2)$ , we set

$$\tilde{v}(x) = 5v(-x_1, x_2, x_3) - 20v(-x_1/2, x_2, x_3) + 16v(-x_1/4, x_2, x_3) .$$

Notice that  $\tilde{v}$  is an extension of  $v$  in  $B(0, 2)$ , which preserves the  $\mathcal{C}^2$ -regularity, and that the  $H^s$ -norm for  $s \leq 2$  as well as the Strichartz norms of  $\tilde{v}$  are controlled by the corresponding norms of  $v$ . Let  $\chi \in \mathcal{C}_0^\infty(\mathbb{R}^3)$  be a cut-off function such that  $\chi \equiv 1$  in  $B_+(0, 1)$  and  $\chi \equiv 0$  outside  $B(0, 2)$ . Applying Theorem 4.1 to  $\chi(x)f(\chi(x)\tilde{v})$  yields a control of  $\|f(v)\|_{L^1([0, T], H^{s+\varepsilon}(B_+(0, 1)))}$  by  $\|v\|_{L^\infty([0, T], H^{s+1}(\Omega))}$ . Finally, notice that  $f(0) = 0$  and thus, the Dirichlet boundary condition on  $v$  naturally implies the one on  $f(v)$ .  $\square$

## 4.2 Asymptotic compactness and regularization effect

As explained in the beginning of this section, using Duhamel formula  $U(t) = e^{At}U_0 + \int_0^t e^{A(t-s)}F(U(s))ds$  and Corollary 4.2, we obtain two propositions related to the asymptotic smoothing effect of the damped wave equations.

**Proposition 4.3.** *Let  $f \in \mathcal{C}^1(\mathbb{R})$  satisfying (1.2), let  $(u_0^n, u_1^n)$  be a sequence of initial data which is bounded in  $X = H_0^1(\Omega) \times L^2(\Omega)$  and let  $(u_n)$  be the corresponding solutions of the damped wave equation (1.1). Let  $(t_n) \in \mathbb{R}$  be a sequence of times such that  $t_n \rightarrow +\infty$  when  $n$  goes to  $+\infty$ .*

*Then, there exist subsequences  $(u_{\phi(n)})$  and  $(t_{\phi(n)})$  and a global solution  $u_\infty$  of (1.1) such that*

$$\forall T > 0, \quad (u_{\phi(n)}, \partial_t u_{\phi(n)})(t_{\phi(n)} + \cdot) \longrightarrow (u_\infty, \partial_t u_\infty)(\cdot) \quad \text{in } \mathcal{C}^0([-T, T], X) .$$

**Proof:** We use the notations of Section 2. Due to the equivalence between the norm of  $X$  and the energy given by (2.1) and the fact that the energy is decreasing in time, we know that  $U_n(t)$  is uniformly bounded in  $X$  with respect to  $n$  and  $t \geq 0$ . So, up to taking a subsequence, it weakly converges to a limit  $U_\infty(0)$  which gives a global solution  $U_\infty$ . We notice that, due to the continuity of the Cauchy problem with respect to the initial data stated in Theorem 2.2, it is sufficient to show that  $U_{\phi(n)}(t_{\phi(n)}) \rightarrow U_\infty(0)$  for some subsequence  $\phi(n)$ . We have

$$\begin{aligned}
U_n(t_n) &= e^{At_n}U_n(0) + \int_0^{t_n} e^{sA}F(U_n(t_n - s)) ds \\
&= e^{At_n}U_n(0) + \sum_{k=0}^{\lfloor t_n \rfloor - 1} e^{kA} \int_0^1 e^{sA}F(U_n(t_n - k - s)) ds + \int_{\lfloor t_n \rfloor}^{t_n} e^{sA}F(U_n(t_n - s)) ds \\
&= e^{At_n}U_n(0) + \sum_{k=0}^{\lfloor t_n \rfloor - 1} e^{kA}I_{k,n} + I_n
\end{aligned} \tag{4.2}$$

Theorem 2.2 shows that the Strichartz norms  $\|u_n(t_n - k - \cdot)\|_{L^q([0,1],L^r(\Omega))}$  are uniformly bounded since the energy of  $U_n$  is uniformly bounded. Therefore, Corollary 4.2 and Proposition 2.3 show that the terms  $I_{n,k} = \int_0^1 e^{sA}F(U_n(t_n - k - s)) ds$ , as well as  $I_n$ , are bounded by some constant  $M$  in  $H^{1+\varepsilon}(\Omega) \times H^\varepsilon(\Omega)$  uniformly in  $n$  and  $k$ . Using Proposition 2.3 again and summing up, we get that the last terms of (4.2) are bounded in  $H^{1+\varepsilon}(\Omega) \times H^\varepsilon(\Omega)$  uniformly in  $n$  by

$$\left\| \sum_{k=0}^{\lfloor t_n \rfloor - 1} e^{kA}I_{k,n} + I_n \right\|_{X^\varepsilon} \leq \sum_{k=0}^{\lfloor t_n \rfloor - 1} C e^{-\lambda k} M + M \leq M \left( 1 + \frac{C}{1 - e^{-\lambda}} \right).$$

Moreover, Proposition 2.3 shows that  $e^{At_n}U_n(0)$  goes to zero in  $X$  when  $n$  goes to  $+\infty$ . Therefore, by a diagonal extraction argument and Rellich Theorem, we can extract a subsequence  $U_{\phi(n)}(t_{\phi(n)})$  that converges to  $U_\infty(0)$  in  $H_0^1(B) \times L^2(B)$  for all bounded set  $B$  of  $\Omega$ .

To finish the proof of Proposition 4.3, we have to show that this convergence holds in fact in  $X$  and not only locally. Let  $\eta > 0$  be given. Let  $T > 0$  and let  $\tilde{U}_n$  be the solution of (1.1) with  $\tilde{U}_n(0) = U_n(t_n - T)$  and with  $\gamma$  being replaced by  $\tilde{\gamma}$ , where  $\tilde{\gamma}(x) \equiv \gamma(x)$  for large  $x$  and  $\tilde{\gamma} \geq \alpha > 0$  everywhere. By Proposition 2.6,  $\|\tilde{U}_n(T)\|_X \leq \eta$  if  $T$  is chosen sufficiently large and if  $n$  is large enough so that  $t_n - T > 0$ . Since the information propagates at finite speed in the wave equation,  $U_n(t_n) \equiv \tilde{U}_n(T)$  outside a large enough bounded set and thus  $U_{\phi(n)}(t_{\phi(n)})$  has a  $X$ -norm smaller than  $\eta$  outside this bounded set. On the other hand, we can assume that the norm of  $U_\infty(0)$  is also smaller than  $\eta$  outside the bounded set. Then, choosing  $n$  large enough,  $\|U_{\phi(n)}(t_{\phi(n)}) - U_\infty(0)\|_X$  becomes smaller than  $3\eta$ .  $\square$

The trajectories  $U^\infty$  appearing in Proposition 4.3 are trajectories which are bounded in  $X$  for all times  $t \in \mathbb{R}$ . The following result shows that these special trajectories are more regular than the usual trajectories of the damped wave equation.

**Proposition 4.4.** *Let  $f \in \mathcal{C}^1(\mathbb{R})$  satisfying (1.2) and let  $E_0 \geq 0$ . There exists a constant  $M$  such that if  $u$  is a solution of (1.1), which exists for all times  $t \in \mathbb{R}$  and satisfies  $\sup_{t \in \mathbb{R}} E(u(t)) \leq E_0$ , then  $t \mapsto U(t) = (u(t), \partial_t u(t))$  is continuous from  $\mathbb{R}$  into  $D(A)$  and*

$$\sup_{t \in \mathbb{R}} \|(u(t), \partial_t u(t))\|_{D(A)} \leq M .$$

In addition,  $M$  depends only on  $E_0$  and the constants in (1.2).

**Proof:** We use a bootstrap argument. For any  $t \in \mathbb{R}$  and  $n \in \mathbb{N}$ ,

$$U(t) = e^{nA}U(t-n) + \sum_{k=0}^{n-1} e^{kA} \int_0^1 e^{sA} F(U(t-k-s)) ds .$$

Using Proposition 2.3, when  $n$  goes to  $+\infty$ , we get

$$U(t) = \sum_{k=0}^{+\infty} e^{kA} \int_0^1 e^{sA} F(U(t-k-s)) ds . \quad (4.3)$$

Moreover, arguing exactly as in the proof of Proposition 4.3, we show that Proposition 2.3 and Corollary 4.2 imply that Equality (4.3) also holds in  $X^\varepsilon$ . Hence,  $U(t)$  is uniformly bounded in  $X^\varepsilon$ . Then, using again Proposition 2.3 and Corollary 4.2, (4.3) also holds in  $X^{2\varepsilon}$  etc. Repeating the arguments and noting that, until the last step,  $\varepsilon$  only depends on  $p$ , we obtain that  $U(t)$  is uniformly bounded in  $X^1 = D(A)$ .

Since the constant  $C$  of Corollary 4.2 only depends on  $f$  through Estimate (1.2), the same holds for the bound  $M$  here.  $\square$

**Proposition 4.5.** *The Sobolev embedding  $H^2(\Omega) \hookrightarrow \mathcal{C}^0(\Omega)$  holds and there exists a constant  $\mathcal{K}$  such that*

$$\forall u \in H^2(\Omega) , \quad \sup_{x \in \Omega} |u(x)| \leq \mathcal{K} \|u\|_{H^2} .$$

In particular, the solution  $u$  in the statement of Proposition 4.4 belongs to  $\mathcal{C}^0(\overline{\Omega} \times \mathbb{R}, \mathbb{R})$  and  $\sup_{(x,t) \in \overline{\Omega} \times \mathbb{R}} |u(x,t)| \leq \mathcal{K}M$ .

**Proof:** Proposition 4.5 follows directly from the fact that  $\Omega$  has a bounded geometry and from the classical Sobolev embedding  $H^2 \hookrightarrow \mathcal{C}^0$  in the ball  $B(0,1)$  of  $\mathbb{R}^3$ .  $\square$



## 5 Smoothness and uniqueness of non-dissipative complete solutions

In this section, we consider only a non-dissipative complete solution, that is a solution  $u^*$  existing for all times  $t \in \mathbb{R}$  for which the energy  $E$  is constant. In other words,  $u^*(t)$  solves

$$\begin{cases} \partial_{tt}^2 u^* = \Delta u^* - \beta u^* - f(u^*) & (x, t) \in \Omega \times \mathbb{R} , \\ u^*(x, t) = 0 & (x, t) \in \partial\Omega \times \mathbb{R} \\ \partial_t u^*(x, t) = 0 & (x, t) \in \text{supp}(\gamma) \times \mathbb{R} \end{cases} \quad (5.1)$$

Since the energy  $E$  is not dissipated by  $u^*(t)$ , we can write  $E(u^*)$  instead of  $E(u^*(t))$ . Yet, an interesting fact that will be used several times in the sequel is that such  $u^*$  is, at the same time, solution of both damped and undamped equations.

The purpose of this section is:

- First show that  $u^*$  is analytic in time and smooth in space. The central argument is to use a theorem of J.K. Hale and G. Raugel in [25].
- Then use the unique continuation result of L. Robbiano and C. Zuily stated in Corollary 3.2 to show that  $u^*$  is necessarily an equilibrium point of (1.1).
- Finally show that the assumption  $sf(s) \geq 0$  imply that  $u^* \equiv 0$ .

We enhance that the first two steps are valid and very helpful in a more general framework than the one of our paper.

### 5.1 Smoothness and partial analyticity of $u^*$

First, we recall here the result of Section 2.2 of [25], adapting the statement to suit our notations.

#### Theorem 5.1. Hale and Raugel (2003)

Let  $Y$  be a Banach space. Let  $P_n \in \mathcal{L}(Y)$  be a sequence of continuous linear maps and let  $Q_n = Id - P_n$ . Let  $A : D(A) \rightarrow Y$  be the generator of a continuous semigroup  $e^{tA}$  and let  $G \in \mathcal{C}^1(Y)$ . We assume that  $V$  is a complete mild solution in  $Y$  of

$$\partial_t V(t) = AV(t) + G(V(t)) \quad \forall t \in \mathbb{R} .$$

We further assume that

- (i)  $\{V(t), t \in \mathbb{R}\}$  is contained in a compact set  $K$  of  $Y$ .
- (ii) for any  $y \in Y$ ,  $P_n y$  converges to  $y$  when  $n$  goes  $+\infty$  and  $(P_n)$  and  $(Q_n)$  are sequences of  $\mathcal{L}(Y)$  bounded by  $K_0$ .
- (iii) the operator  $A$  splits in  $A = A_1 + B_1$  where  $B_1$  is bounded and  $A_1$  commutes with  $P_n$ .
- (iv) there exist  $M$  and  $\lambda > 0$  such that  $\|e^{At}\|_{\mathcal{L}(Y)} \leq Me^{-\lambda t}$  and  $\|e^{(A_1 + Q_n B_1)t}\|_{\mathcal{L}(Q_n Y, Y)} \leq Me^{-\lambda t}$  for all  $t \geq 0$ .



(v)  $G$  is analytic in the ball  $B_Y(0, r)$ , where  $r$  is such that  $r \geq 4K_0 \sup_{t \in \mathbb{R}} \|V(t)\|_Y$ . More precisely, there exists  $\rho > 0$  such that  $G$  can be extended to an holomorphic function of  $B_Y(0, r) + iB_Y(0, \rho)$ .

(vi)  $\{DG(V(t))V_2 \mid t \in \mathbb{R}, \|V_2\|_Y \leq 1\}$  is a relatively compact set of  $Y$ .

Then, the solution  $V(t)$  is analytic from  $t \in \mathbb{R}$  into  $Y$ .

More precisely, Theorem 5.1 is Theorem 2.20 (which relates to Theorem 2.12) of [25] applied with Hypothesis (H3mod) and (H5).

Proposition 4.4 shows that  $u^*$  is continuous in both space and time variables. We apply Theorem 5.1 to show that, because  $f$  is analytic,  $u^*$  is also analytic with respect to the time.

**Proposition 5.2.** *Let  $f \in \mathcal{C}^1(\mathbb{R})$  satisfying (1.2) and let  $E_0 \geq 0$ . Let  $\mathcal{K}$  and  $M$  be the constants given by Propositions 4.4 and 4.5. Assume that  $f$  is analytic in  $[-4\mathcal{K}M, 4\mathcal{K}M]$ . Then for any non-dissipative complete solution  $u^*(t)$  solving (5.1) and satisfying  $E(u^*) \leq E_0$ ,  $t \mapsto u^*(\cdot, t)$  is analytic from  $\mathbb{R}$  into  $X^\alpha$  with  $\alpha \in (1/2, 1)$ . In particular, for all  $x \in \Omega$ ,  $u^*(x, t)$  is analytic with respect to the time.*

**Proof:** Theorem 5.1 uses strongly some compactness properties. Therefore, we need to truncate our solution to apply the theorem on a bounded domain (of course, this is not necessary and easier if  $\Omega$  is already bounded).

Let  $\chi \in \mathcal{C}_0^\infty(\bar{\Omega})$  be such that  $\frac{\partial \chi}{\partial \nu} = 0$  on  $\partial\Omega$ ,  $\chi \equiv 1$  in  $\{x \in \Omega, \gamma(x) = 0\}$  and  $\text{supp}(\chi)$  is included in a smooth bounded subdomain  $\mathcal{O}$  of  $\Omega$ . Since Proposition 4.4 shows that  $u^* \in \mathcal{C}^0(\mathbb{R}, D(A))$  and since  $u^*$  is constant with respect to the time in  $\text{supp}(\gamma)$ ,  $(1 - \chi)u^*$  is obviously analytic from  $\mathbb{R}$  into  $D(A)$ . It remains to obtain the analyticity of  $\chi u^*$ .

In this proof, the damping  $\gamma$  needs to be more regular than just  $L^\infty(\Omega)$ . We replace  $\gamma$  by a damping  $\tilde{\gamma} \in C^\infty(\Omega)$ , which has the same geometrical properties (GCC) and (1.3) and which vanishes where  $\gamma$  does. Notice that  $\gamma \partial_t u^* \equiv 0 \equiv \tilde{\gamma} \partial_t u^*$ , therefore replacing  $\gamma$  by  $\tilde{\gamma}$  has no consequences here.

Let  $v = \chi u^*$ , we have

$$\begin{cases} \partial_{tt}^2 v + \tilde{\gamma}(x) \partial_t v = \Delta v - \beta v + g(x, v) & (x, t) \in \mathcal{O} \times \mathbb{R}_+, \\ v(x, t) = 0 & (x, t) \in \partial\mathcal{O} \times \mathbb{R}_+ \end{cases} \quad (5.2)$$

with  $g(x, v) = -\chi(x)f(v + (1 - \chi)u^*(x)) - 2(\nabla\chi \cdot \nabla u^*)(x) - (u^* \Delta\chi)(x)$ . We apply Theorem 5.1 with the following setting. Let  $Y = X^\alpha = H^{1+\alpha}(\mathcal{O}) \cap H_0^1(\mathcal{O}) \times H_0^\alpha(\mathcal{O})$  with  $\alpha \in (1/2, 1)$ . Let  $V = (v, \partial_t v)$  and let  $G(v) = (0, g(\cdot, v))$ . We set

$$A = A_1 + B_1 = \begin{pmatrix} 0 & Id \\ \Delta - \beta & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 0 & -\tilde{\gamma} \end{pmatrix}.$$

Let  $(\lambda_k)_{k \geq 1}$  be the negative eigenvalues of the Laplacian operator on  $\mathcal{O}$  with Dirichlet boundary conditions and let  $(\varphi_k)$  be corresponding eigenfunctions. We set  $P_n$  to be the canonical projections of  $X$  on the subspace generated by  $((\varphi_k, 0))_{k=1 \dots n}$  and  $((0, \varphi_k))_{k=1 \dots n}$ .

To finish the proof of Proposition 5.2, we only have to check that the hypotheses of Theorem 5.1 hold.

The trajectory  $V$  is compact since we know by Proposition 4.4 that it is bounded in  $X^1$ , which gives (i).

Hypothesis (ii) and (iii) hold with  $K_0 = 1$  by construction of  $P_n$  and because  $B_1$  is bounded in  $Y$  since  $\tilde{\gamma}$  belongs to  $C^\infty(\Omega)$ .

The first part of Hypothesis (iv) follows from Proposition 2.3. The second estimate  $\|e^{(A_1+Q_n B_1)t}\|_{\mathcal{L}(Q_n Y, Y)} \leq M e^{-\lambda t}$  means that the restriction  $Q_n A Q_n$  of  $A$  to the high frequencies of the wave operator also generates a semigroup satisfying an exponential decay. By a result of Haraux in [29], we know that this also holds, see Section 2.3.2 of [25] for the detailed arguments.

We recall that  $u^*(x, \cdot)$  is constant outside  $\chi^{-1}(1)$  and belongs locally to  $H^{1+\alpha}$  since  $u^* \in D(A)$ . Therefore, the terms  $(1 - \chi)u^*(x)$ ,  $\nabla\chi \cdot \nabla u^*$  and  $u^* \Delta\chi$  appearing in the definition of  $g$  are in  $H^1$ . Moreover, they satisfy Dirichlet boundary condition on  $\partial\Omega$  since  $u^* \equiv 0$  and  $\partial_\nu \chi \equiv 0$  there. Of course, they also satisfy Dirichlet boundary condition on the other parts of  $\partial\mathcal{O}$  since  $\chi \equiv 0$  outside  $\mathcal{O}$ . Notice that  $\alpha > 1/2$  and thus  $H^{1+\alpha}(\mathcal{O}) \cap H_0^1(\mathcal{O})$  is an algebra included in  $\mathcal{C}^0$ . Therefore, (1.2) shows that  $G$  is of class  $\mathcal{C}^1$  in the bounded sets of  $Y$ . Since  $u \in [-4\mathcal{K}M, 4\mathcal{K}M] \mapsto f(u) \in \mathbb{R}$  is analytic, it can be extended to a holomorphic function in  $[-4\mathcal{K}M, 4\mathcal{K}M] + i[-\rho, \rho]$  for small  $\rho > 0$ . Using again the embedding  $H^{1+\alpha}(\mathcal{O}) \hookrightarrow \mathcal{C}^0(\mathcal{O})$  and the definitions of  $\mathcal{K}$  and  $M$ , we deduce that (v) holds.

Finally, for  $V_2 = (v_2, \partial_t v_2)$  with  $\|V_2\|_Y \leq 1$   $DG(V(t))V_2 = (0, -\chi(x)f'(v(t)) + (1 - \chi)u^*(x))v_2)$  is relatively compact in  $Y$  since  $v(t)$  is bounded in  $H^2 \cap H_0^1$  due to Proposition 4.4 and therefore  $v_2 \in H^{1+\alpha} \mapsto \chi(x)f'(v(t)) + (1 - \chi)u^*(x)v_2 \in H^\alpha$  is a compact map. This yields (vi).  $\square$

Once the time-regularity of  $u^*$  is proved, the space-regularity follows directly.

**Proposition 5.3.** *Let  $f$  and  $u^*$  be as in Proposition 5.2, then  $u^* \in \mathcal{C}^\infty(\Omega \times \mathbb{R})$ .*

**Proof:** Proposition 5.2 shows that  $u^*$  and all its time-derivatives belongs to  $X^\alpha$  with  $\alpha \in (1/2, 1)$ . Due to the Sobolev embeddings, this implies that any time-derivative of  $u^*$  is Hölder continuous. Writing

$$\Delta u^* = \partial_{tt}^2 u^* + \beta u^* + f(u^*) \tag{5.3}$$

and using the local elliptic regularity properties (see [42] and the references therein for example), we get that  $u^*$  is locally of class  $\mathcal{C}^{2,\lambda}$  in space for some  $\lambda \in (0, 1)$ . Thus,  $u^*$  is of class  $\mathcal{C}^{2,\lambda}$  in both time and space. Then, we can use a bootstrap argument in (5.3) to show that  $u^*$  is of class  $\mathcal{C}^{2k,\lambda}$  for all  $k \in \mathbb{N}$ .  $\square$

## 5.2 Identification of $u^*$

The smoothness and the partial analyticity of  $u^*$  shown in Propositions 5.2 and 5.3 enable us to use the unique continuation result of [47].

**Proposition 5.4.** *Let  $f$  and  $u^*$  be as in Proposition 5.2, then  $u^*$  is constant in time, i.e.  $u^*$  is an equilibrium point of the damped wave equation (1.1).*

**Proof:** Setting  $v = \partial_t u^*$ , we get  $\partial_{tt}^2 v = \Delta v - \beta v - f'(u^*)v$ . Propositions 5.2 and 5.3 show that  $u^*$  is smooth and analytic with respect to the time and moreover  $v \equiv 0$  in  $\text{supp}(\gamma)$ . Thus, the unique continuation result stated in Corollary 3.2 yields  $v \equiv 0$  everywhere.  $\square$

The sign assumption on  $f$  directly implies that 0 is the only possible equilibrium point of (1.1).

**Corollary 5.5.** *Let  $f \in \mathcal{C}^1(\mathbb{R})$  satisfying (1.2) and let  $E_0 \geq 0$ . Let  $\mathcal{K}$  and  $M$  be the constants given by Propositions 4.4 and 4.5 and assume that  $f$  is analytic in  $[-4\mathcal{K}M, 4\mathcal{K}M]$ . Then, the unique solution  $u^*$  of (5.1) with  $E(u^*) \leq E_0$  is  $u^* \equiv 0$ .*

**Proof:** Due to Proposition 5.4,  $u^*$  is solution of  $\Delta u^* - \beta u^* = f(u^*)$ . By multiplying by  $u^*$  and integrating by part, we obtain  $\int_{\Omega} |\nabla u^*|^2 + \beta |u^*|^2 dx = - \int_{\Omega} u^* f(u^*) dx$ , which is non-positive due to Assumption (1.2). Since  $\beta \geq 0$  is such that  $\Delta - \beta$  is negative definite, this shows that  $u^* \equiv 0$ .  $\square$

## 6 Proof of Theorem 1.1

Due to Proposition 2.5, Theorem 1.1 directly follows from the following result.

**Proposition 6.1.** *Let  $f \in \mathcal{C}^1(\mathbb{R})$  satisfying (1.2) and let  $E_0 \geq 0$ . Let  $\mathcal{K}$  and  $M$  be the constants given by Propositions 4.4 and 4.5. Assume that  $f$  is analytic in  $[-4\mathcal{K}M, 4\mathcal{K}M]$  and that  $\gamma$  is as in Theorem 1.1. Then, there exist  $T > 0$  and  $C > 0$  such that for any  $u$  solution of (1.1) with  $E(u)(0) \leq E_0$  satisfies*

$$E(u)(0) \leq C \iint_{[0, T] \times \Omega} \gamma(x) |\partial_t u|^2 dt dx.$$

**Proof:** We argue by contradiction: we assume that there exists a sequence  $(u_n)$  of solutions of (1.1) and a sequence of times  $(T_n)$  converging to  $+\infty$  such that

$$\iint_{[0, T_n] \times \Omega} \gamma(x) |\partial_t u_n|^2 dt dx \leq \frac{1}{n} E(u_n)(0) \leq \frac{1}{n} E_0. \quad (6.1)$$

Denote  $\alpha_n = (E(u_n(0)))^{1/2}$ . Since  $\alpha \in [0, \sqrt{E_0}]$ , we can assume that  $\alpha_n$  converges to a limit  $\alpha$  when  $n$  goes to  $+\infty$ . We distinguish two cases:  $\alpha > 0$  and  $\alpha = 0$ .

- First case:  $\alpha_n \rightarrow \alpha > 0$

Notice that, due to (2.1),  $\|(u_n, \partial_t u_n)(0)\|_X$  is uniformly bounded from above and from below by positive numbers. We set  $u_n^* = u_n(T_n/2 + \cdot)$ . Due to the asymptotic compactness property stated in Proposition 4.3, we can assume that  $u_n^*$  converges to a solution  $u^*$  of (1.1) in  $\mathcal{C}^0([-T, T], X)$  for all time  $T > 0$ . We notice that

$$E(u_n(0)) \geq E(u_n^*(0)) = E(u_n(0)) - \iint_{[0, T_n/2] \times \Omega} \gamma(x) |\partial_t u_n|^2 \geq (1 - 1/n)E(u_n(0))$$

and thus  $E(u^*(0)) = \alpha^2 > 0$ . Moreover, (6.1) shows that  $\gamma \partial_t u_n^*$  converges to zero in  $L^2([-T, T], L^2(\Omega))$  for any  $T > 0$  and thus  $\partial_t u^* \equiv 0$  in  $\text{supp}(\gamma)$ . In other words,  $u^*$  is a non-dissipative solution of (1.1), i.e. a solution of (5.1) with  $E(u^*) = \alpha^2 \leq E_0$ . Corollary 5.5 shows that  $u^* \equiv 0$ , which contradicts the positivity of  $E(u^*(0))$ .

- Second case:  $\alpha_n \rightarrow 0$

The assumptions on  $f$  allow to write  $f(s) = f'(0)s + R(s)$  with

$$|R(s)| \leq C(|s|^2 + |s|^p) \quad \text{and} \quad |R'(s)| \leq C(|s| + |s|^{p-1}). \quad (6.2)$$

Let us make the change of unknown  $w_n = u_n/\alpha_n$ . Then,  $w_n$  solves

$$\square w_n + \gamma(x) \partial_t w_n + (\beta + f'(0))w_n + \frac{1}{\alpha_n} R(\alpha_n w_n) = 0 \quad (6.3)$$

and

$$\iint_{[0, T_n] \times \Omega} \gamma(x) |\partial_t w_n|^2 \, dt dx \leq \frac{1}{n}. \quad (6.4)$$

Denote  $W_n = (w_n, \partial_t w_n)$ . Due to the equivalence between norm and energy given by (2.1), the scaling  $w_n = u_n/\alpha_n$  implies that  $\|(w_n(0), \partial_t w_n(0))\|_X$  is uniformly bounded from above and from below by positive numbers. Moreover, (6.1) implies

$$\|W_n(t)\|_X = \frac{\|(U_n(t))\|_X}{\alpha_n} \geq C \frac{E(u_n(t))^{1/2}}{\alpha_n} \geq C \frac{(E(u_n(0)) - \alpha_n^2/n)^{1/2}}{\alpha_n} \geq \frac{C}{2} > 0 \quad (6.5)$$

for any  $t \in [0, T_n]$  and  $n$  large enough.

We set  $f_n = 1/\alpha_n R(u_n)$  and  $F_n = (0, f_n)$ . The stability estimate of Proposition 2.2 implies that  $\|u_n\|_{L^q([k, k+1], L^r)} \leq C\alpha_n$  uniformly for  $n, k \in \mathbb{N}$ . In particular, combined with (6.2), this gives

$$\|f_n\|_{L^1([k, k+1], L^2)} = \left\| \frac{1}{\alpha_n} R(\alpha_n w_n) \right\|_{L^1([k, k+1], L^2)} \leq C(\alpha_n + \alpha_n^{p-1})$$

We can argue as in Proposition 4.3 and write

$$\begin{aligned}
W_n(T_n) &= e^{\tilde{A}T_n}W_n(0) + \sum_{k=0}^{\lfloor T_n \rfloor - 1} e^{\tilde{A}(T_n - k)} \int_0^1 e^{-\tilde{A}s} F_n(k + s) ds \\
&\quad + e^{\tilde{A}(T_n - \lfloor T_n \rfloor)} \int_0^{T_n - \lfloor T_n \rfloor} e^{-\tilde{A}s} F_n(\lfloor T_n \rfloor + s) ds .
\end{aligned} \tag{6.6}$$

where  $\tilde{A}$  is the modified damped wave operator

$$\tilde{A} = \begin{pmatrix} 0 & Id \\ \Delta - \beta - f'(0) & -\gamma \end{pmatrix} .$$

Notice that  $\tilde{A}$  satisfies an exponential decay as  $A$  in Proposition 2.3 since (1.2) implies  $f'(0) \geq 0$ . By summing up as in Proposition 4.3, we get

$$\|W_n(T_n)\|_X \leq C e^{-\lambda T_n} + C(\alpha_n + \alpha_n^{p-1})$$

which goes to zero, in a contradiction with (6.5).  $\square$

As a direct consequence of Proposition 6.1, we obtain a unique continuation property for nonlinear wave equations. Notice that the time of observation  $T$  required for the unique continuation is not explicit. Thus, this result is not so convenient as a unique continuation property. But it may be useful for other nonlinear stabilization problems as  $\square u + \gamma(x)g(\partial_t u) + f(u) = 0$ .

**Corollary 6.2.** *Let  $f \in C^1(\mathbb{R})$  satisfying (1.2) and let  $E_0 \geq 0$ . Assume that  $f$  is analytic in  $\mathbb{R}$  and that  $\omega$  is an open subset of  $\Omega$  satisfying (GCC). Then, there exist  $T > 0$  such that the only solution  $u$  of*

$$\begin{cases} \square u + \beta u + f(u) = 0 & \text{on } [-T, T] \times \Omega \\ \partial_t u \equiv 0 & \text{on } [-T, T] \times \omega . \end{cases} \tag{6.7}$$

with  $E(u)(0) \leq E_0$  is  $u \equiv 0$ .

**Proof:** Corollary 6.2 is a straightforward consequence of Proposition 6.1 since we can easily construct a smooth damping  $\gamma$  supported in  $\omega$  and such that  $Supp(\gamma)$  satisfies (GCC). We only have to remark that  $u$  solution of (6.7) is also solution of (1.1).  $\square$

## 7 Proof of Theorem 1.2

Before starting the proof of Theorem 1.2 itself, we prove that  $\mathfrak{C}^1(\mathbb{R})$  is a Baire space, that is that any countable intersection of open dense sets is dense. This legitimizes the genericity in  $\mathfrak{C}^1(\mathbb{R})$  as a good notion of large subsets of  $\mathfrak{C}^1(\mathbb{R})$ . We recall that  $\mathfrak{C}^1(\mathbb{R})$  is defined by (1.6) and endowed by Whitney topology, the open sets of which are generated by the neighbourhoods  $\mathcal{N}_{f,\delta}$  defined by (1.7).

**Proposition 7.1.** *The space  $\mathfrak{C}^1(\mathbb{R})$  endowed with Whitney topology is a Baire space.*

**Proof:** The set  $\mathfrak{C}^1(\mathbb{R})$  is not an open set of  $\mathcal{C}^1(\mathbb{R})$ , and neither a submanifold. It is a closed subset of  $\mathcal{C}^1(\mathbb{R})$ , but  $\mathcal{C}^1(\mathbb{R})$  endowed with Whitney topology is not a completely metrizable space, since it is not even metrizable (the neighbourhoods of a function  $f$  are not generated by a countable subset of them). Therefore, we have to come back to the basic proof of Baire property as in [23].

Let  $\mathcal{U}$  be an open set of  $\mathfrak{C}^1(\mathbb{R})$  and let  $(\mathcal{O}_n)_{n \in \mathbb{N}}$  be a sequence of open dense sets of  $\mathfrak{C}^1(\mathbb{R})$ . By density, there exists a function  $f_0 \in \mathfrak{C}^1(\mathbb{R})$  in  $\mathcal{U} \cap \mathcal{O}_0$  and by openness, there exists a positive continuous function  $\delta_0$  such that the neighbourhood  $\mathcal{N}_{f_0, \delta_0}$  is contained in  $\mathcal{U} \cap \mathcal{O}_0$ . By choosing  $\delta_0$  small enough, one can also assume that  $\mathcal{N}_{f_0, 2\delta_0} \subset \mathcal{U} \cap \mathcal{O}_0$  and that  $\sup_{u \in \mathbb{R}} |\delta_0(u)| \leq 1/2^0$ . By recursion, one constructs similar balls  $\mathcal{N}_{f_n, \delta_n} \subset \mathcal{N}_{f_{n-1}, \delta_{n-1}} \subset \mathcal{N}_{f_0, \delta_0}$  such that  $\mathcal{N}_{f_n, 2\delta_n} \subset \mathcal{U} \cap \mathcal{O}_n$  and that  $\sup_{u \in \mathbb{R}} |\delta_n(u)| \leq 1/2^n$ . Since  $\mathcal{C}^1([-m, m], \mathbb{R})$  endowed with the uniform convergence topology is a complete metric space, the sequence  $(f_i)$  converges to a function  $f \in \mathcal{C}^1(\mathbb{R}, \mathbb{R})$  uniformly in any compact set of  $\mathbb{R}$ . By construction, the limit  $f$  satisfies

$$\forall n \in \mathbb{N}, \forall u \in \mathbb{R}, \quad \max(|f(u) - f_n(u)|, |f'(u) - f'_n(u)|) \leq \delta_n(u) < 2\delta_n(u), \quad (7.1)$$

as well as  $f(0) = 0$  and  $uf(u) \geq 0$  since any  $f_n$  satisfies (1.2). Moreover, there exist  $C > 0$  and  $p \in [1, 5)$  such that  $f_0$  satisfies

$$|f_0(u)| \leq C(1 + |u|)^p \quad \text{and} \quad |f'_0(u)| \leq C(1 + |u|)^{p-1}. \quad (7.2)$$

Since  $\max(|f(u) - f_0(u)|, |f'(u) - f'_0(u)|) \leq \delta_0(u) \leq 1$ ,  $f$  also satisfies (7.2) with a constant  $C' = C + 1$ . Therefore,  $f$  satisfies (1.2) and thus belongs to  $\mathfrak{C}^1(\mathbb{R})$ . In addition,  $f$  satisfying (7.1) and  $\mathcal{N}_{f_n, 2\delta_n}$  being contained in  $\mathcal{U} \cap \mathcal{O}_n$ , we get  $f \in \mathcal{U} \cap \mathcal{O}_n$  for all  $n$ . This shows that  $\bigcap_{n \in \mathbb{N}} \mathcal{O}_n$  intersects any open set  $\mathcal{U}$  and therefore is dense in  $\mathfrak{C}^1(\mathbb{R})$ .  $\square$

**Proof of Theorem 1.2:** We denote by  $\mathfrak{G}_n$  the set of functions  $f \in \mathfrak{C}^1(\mathbb{R})$  such that the exponential decay property (ED) holds for  $E_0 = n$ . Obviously,  $\mathfrak{G} = \bigcap_{n \in \mathbb{N}} \mathfrak{G}_n$  and hence it is sufficient to prove that  $\mathfrak{G}_n$  is an open dense subset of  $\mathfrak{C}^1(\mathbb{R})$ . We sketch here the main arguments to prove this last property.

**$\mathfrak{G}_n$  is a dense subset:** let  $\mathcal{N}$  be a neighbourhood of  $f_0 \in \mathfrak{C}^1(\mathbb{R})$ . Up to choosing  $\mathcal{N}$  smaller, we can assume that the constant in (1.2) is independent of  $f \in \mathcal{N}$ . Due to Propositions 4.4 and 4.5, there exist constants  $\mathcal{K}$  and  $M$  such that, for all  $f \in \mathcal{N}$ , all the global non-dissipative trajectories  $u$  of (1.1) with  $E(u) \leq n$  are such that  $\|u\|_{L^\infty(\Omega \times \mathbb{R})} \leq \mathcal{K}M$ . We claim that we can choose  $f \in \mathcal{N}$  as close to  $f_0$  as wanted such that  $f$  is analytic on  $[-4\mathcal{K}M, 4\mathcal{K}M]$  and still satisfies (1.2). Then, Proposition 6.1 shows that  $f$  satisfies (ED) with  $E_0 = n$  i.e. that  $f \in \mathfrak{G}_n$ .

To obtain this suitable function  $f$ , we proceed as follows. First, we set  $a = 4\mathcal{K}M$  and notice that it is sufficient to explain how we construct  $f$  in  $[-a, a]$ . Indeed, one can easily extend a perturbation  $f$  of  $f_0$  in  $[-a, a]$  satisfying  $f(s)s \geq 0$  to a perturbation  $\tilde{f}$  of  $f_0$  in  $\mathbb{R}$ , equal to  $f_0$  outside of  $[-a - 1, a + 1]$  and such that  $f(s)s \geq 0$  in  $[-a - 1, a + 1]$ . We construct  $f$  in  $[-a, a]$  as follows. Since  $f_0(s)s \geq 0$ , we have that  $f'_0(0) \geq 0$ . We perturb  $f_0$  to  $f_1$  such that  $f_1(0) = 0$ ,  $f'_1(s) \geq \varepsilon > 0$  in a small interval  $[-\eta, \eta]$  and  $sf_1(s) \geq 2\varepsilon$  in  $[-a, -\eta] \cup [\eta, a]$ , where  $\varepsilon$  could be chosen as small as needed. Then we perturb  $f_1$  to obtain a function  $f_2$  which is analytic in  $[-a, a]$  and satisfies  $f'_2(s) > 0$  in  $[-\eta, \eta]$ ,  $sf_2(s) \geq \varepsilon$  in  $[-a, -\eta] \cup [\eta, a]$  and  $|f_2(0)| < \varepsilon/a$ . Finally, we set  $f(s) = f_2(s) - f_2(0)$  and check that  $f$  is analytic and satisfies  $sf(s) \geq 0$  in  $[-a, a]$ . Moreover, up to choosing  $\varepsilon$  very small,  $f$  is as close to  $f_0$  as wanted.

**$\mathfrak{G}_n$  is an open subset:** let  $f_0 \in \mathfrak{G}_n$ . Proposition 2.3 shows the existence of a constant  $C$  and a time  $T$  such that, for all solution  $u$  of (1.1),

$$E(u(0)) \leq E_0 \implies E(u(0)) \leq C \int_0^T \int_{\Omega} \gamma(x) |\partial_t u(x, t)|^2 dx dt . \quad (7.3)$$

The continuity of the trajectories in  $X$  with respect to  $f \in \mathfrak{C}^1(\mathbb{R})$  is not difficult to obtain: using the strong control of  $f$  given by Whitney topology, the arguments are the same as the ones of the proof of the continuity with respect to the initial data, stated in Theorem 2.2. Thus, (7.3) holds also for any  $f$  in a neighbourhood  $\mathcal{N}$  of  $f_0$ , replacing the constant  $C$  by a larger one. Therefore, Proposition 2.3 shows that  $\mathcal{N} \subset \mathfrak{G}_n$  and hence that  $\mathfrak{G}_n$  is open.  $\square$

## 8 A proof of compactness and regularity with the usual arguments of control theory

In this section, we give an alternative proof of the compactness and regularity properties of Propositions 4.3 and 4.4. We only give its outline since it is redundant with previous results of the article. Moreover, it is quite similar to the arguments of [17]. Yet, the arguments



of this section are interesting because they do not require any asymptotic arguments and they show a regularization effect through an observability estimate with a finite time  $T$ , which can be explicit. However, for the moment, it seems impossible to obtain an analytic regularity similar to Proposition 5.2 with this kind of arguments.

Instead of using a Duhamel formula with an infinite interval of time  $(-\infty, t)$  as in (4.3), the main idea is to use as a black-box an observability estimate for  $T$  large enough,  $T$  being the time of geometric control condition,

$$\|U_0\|_{X^s}^2 \leq C \|Be^{tA}U_0\|_{L^2([0,T],X^s)}^2 \quad (8.1)$$

where

$$A = \begin{pmatrix} 0 & Id \\ \Delta - \beta & -\gamma \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 0 & 0 \\ 0 & -\gamma \end{pmatrix} .$$

The first aim is to prove that a solution of (5.1), globally bounded in energy, is also globally bounded in  $X^s$  for  $s \in [0, 1]$ . We proceed step by step. First, let us show that it is bounded in  $X^\varepsilon$ .

- we fix  $T >$  large enough to get the observability estimate (8.1). By the existence theory on each  $[t_0, t_0 + T]$ ,  $u|_{[t_0, t_0 + T]}$  is bounded in Strichartz norms, uniformly for  $t_0 \in \mathbb{R}$ . Since the nonlinearity is subcritical, Corollary 4.2 gives that  $f(u)$  is globally bounded in  $L^1([t_0, t_0 + T], H^{1+\varepsilon})$ .
- we decompose the solution into its linear and nonlinear part by the Duhamel formula

$$U(t) = e^{A(t-t_0)}U(t_0) + \int_{t_0}^t e^{A(t_0-\tau)}f(U(\tau))d\tau = U_{lin} + U_{Nlin}.$$

Since  $f(u)$  is bounded in  $L^1([t_0, t_0 + T], H^{1+\varepsilon})$ ,  $U_{Nlin}$  is uniformly bounded in  $C([t_0, t_0 + T], X^\varepsilon)$ .

- we will now use the linear observability estimate (8.1) with  $s = \varepsilon$ , applying it to  $U_{lin}$ :

$$\|U(t_0)\|_{X^\varepsilon}^2 = \|U_{lin}(t_0)\|_{X^\varepsilon}^2 \leq C \int_{t_0}^{t_0+T} \|\gamma(x)\partial_t u_{lin}\|_{H^\varepsilon}^2. \quad (8.2)$$

Then, using triangular inequality, we get

$$\begin{aligned} \int_{t_0}^{t_0+T} \|\gamma(x)\partial_t u_{lin}\|_{H^\varepsilon}^2 &\leq 2 \int_{t_0}^{t_0+T} \|\gamma(x)\partial_t u\|_{H^\varepsilon}^2 + 2 \int_{t_0}^{t_0+T} \|\gamma(x)\partial_t u_{Nlin}\|_{H^\varepsilon}^2 \\ &\leq 2 \int_{t_0}^{t_0+T} \|\gamma(x)\partial_t u_{Nlin}\|_{H^\varepsilon}^2 \leq C \end{aligned}$$

where we have used  $\partial_t u \equiv 0$  on  $\omega$  and  $U_{Nlin}$  is bounded in  $C([t_0, t_0 + T], X^\varepsilon)$ . Combining this with (8.2) for any  $t_0 \in \mathbb{R}$ , we obtain that  $U$  is uniformly bounded in  $X^\varepsilon$  on  $\mathbb{R}$ .



Repeating the arguments, we show that  $u$  is bounded in  $X^{2\varepsilon}$ ,  $X^{3\varepsilon}$  and so on... until  $X^1$ . Similar ideas allows to prove some theorem of propagation of compactness in finite time, replacing the asymptotic compactness property of Proposition 4.3.

As said above, an advantage of this method, compared to the one used in Propositions 4.3 and 4.4, is that it allows to propagate the regularity or the compactness on some finite interval of fixed length. Yet, it seems that such propagation results are not available in the analytic setting. Indeed, it seems that, for nonlinear equations, the propagation of analytic regularity or of nullity in finite time is much harder to prove. We can for instance refer to the weaker (with respect to the geometry) result of Alinhac-Métivier [1] or the negative result of Métivier [40].

## 9 Applications

### 9.1 Control of the nonlinear wave equation

In this subsection, we give a short proof of Theorem 1.3, which states the global controllability of the nonlinear wave equation. The first step consists in a local control theorem.

**Theorem 9.1** (Local control). *Let  $\omega$  satisfying the geometric control condition for a time  $T$ . Then, there exists  $\delta$  such that for any  $(u_0, u_1)$  in  $H_0^1(\Omega) \times L^2(\Omega)$ , with*

$$\|(u_0, u_1)\|_{H_0^1 \times L^2} \leq \delta$$

*there exists  $g \in L^\infty([0, T], L^2)$  supported in  $[0, T] \times \omega$  such that the unique strong solution of*

$$\begin{cases} \square u + \beta u + f(u) &= g & \text{on } [0, T] \times \Omega \\ (u(0), \partial_t u(0)) &= (u_0, u_1). \end{cases}$$

*satisfies  $(u(T), \partial_t u(T)) = (0, 0)$ .*

**Proof:** The proof is exactly the same as Theorem 3 of [17] or Theorem 3.2 of [38]. The main argument consists in seeing the problem as a perturbation of the linear controllability which is known to be true in our setting.  $\square$

Now, as it is very classical, we can combine the local controllability with our stabilization theorem to get global controllability.

**Sketch of the proof of Theorem 1.3:** In a first step, we choose as a control  $g = -\gamma(x)\partial_t \tilde{u}$  where  $\tilde{u}$  is solution of (1.1) with initial data  $(u_0, u_1)$ . By uniqueness of solutions, we have  $u = \tilde{u}$ . Therefore, thanks to Theorem 1.1, for a large time  $T_1$ , only depending on

$R_0$ , we have  $\|(u(T_1), \partial_t u(T_1))\|_{H^1 \times L^2} \leq \delta$ . Then, Theorem 9.1 allows to find a control that brings  $(u(T_1), \partial_t u(T_1))$  to 0. In other words, we have found a control  $g$  supported in  $\omega$  that brings  $(u_0, u_1)$  to 0. We obtain the same result for  $(\tilde{u}_0, \tilde{u}_1)$  and conclude, by reversibility of the equation, that we can also bring 0 to  $(\tilde{u}_0, \tilde{u}_1)$ .  $\square$

## 9.2 Existence of a compact global attractor

In this subsection, we give the modification of the proofs of this paper necessary to get Theorem 1.4 about the existence of a global attractor.

The energy associated to (1.9) in  $X = H_0^1(\Omega) \times L^2(\Omega)$  is given by

$$E(u, v) = \int_{\Omega} \frac{1}{2} (|\nabla u|^2 + |v|^2) + V(x, u) dx ,$$

where  $V(x, u) = \int_0^u f(x, \xi) d\xi$ .

The existence of a compact global attractor for (1.9) is well known for the Sobolev subcritical case  $p < 3$ . The first proofs in this case go back to 1985 ([24] and [27]), see [45] for other references. The case  $p = 3$  as been studied in [5] and [4]. For  $p \in (3, 5)$ , Kapitanski proved in [34] the existence of a compact global attractor for (1.9) if  $\Omega$  is a compact manifold without boundary and if  $\gamma(x) = \gamma$  is a constant damping. Using the same arguments as in the proof of our main result, we can partially deal with the case  $p \in (3, 5)$  with a localized damping  $\gamma(x)$  and with unbounded manifold with boundaries.

Assume that  $f$  satisfies the assumptions of Theorem 1.4. We first claim that we can assume in addition that  $f(x, 0) = 0$  on  $\partial\Omega$  in order to guarantee the Dirichlet boundary condition for  $f(x, u)$  if  $u \in H_0^1(\Omega)$ . Indeed, let  $\varphi$  be the solution of  $\Delta\varphi - \beta\varphi = f(x, 0)$  with Dirichlet boundary conditions, which is well defined and smooth, since by assumptions  $f(x, 0)$  is smooth and compactly supported. Let  $\chi$  be a smooth compactly supported cut-off function such that  $\frac{\partial\chi}{\partial\nu} = 0$  on  $\partial\Omega$  and  $\chi \equiv 1$  in the ball  $B(x_0, R)$  of Hypothesis (1.11). If  $f(x, 0) \neq 0$  on  $\partial\Omega$ , we consider the new variable  $\tilde{u} = u - \chi\varphi$  and the new equation

$$\partial_{tt}^2 \tilde{u} + \gamma(x) \partial_t \tilde{u} = \Delta \tilde{u} - \beta \tilde{u} - \tilde{f}(x, \tilde{u}) \tag{9.1}$$

where  $\tilde{f}(x, \tilde{u}) = f(x, \tilde{u} + \chi\varphi(x)) - \chi f(x, 0) + 2\nabla\varphi \cdot \nabla\chi + \varphi\Delta\chi$ . One directly checks that  $u$  solves (1.9) if and only if  $\tilde{u}$  solves (9.1). Moreover  $\tilde{f}$  is smooth, analytic in  $\tilde{u}$  and satisfies Hypotheses (1.10) and (1.11) and in addition  $\tilde{f}(x, 0) = 0$  on the boundary. Clearly, since our change of variables is a simple translation, obtaining a compact global attractor for (9.1) is equivalent to obtaining a compact global attractor for (1.9). In what follows, we may thus assume that  $f(x, 0) = 0$  in addition to the hypotheses of Theorem 1.4.

The arguments of this paper show the following properties.

- i) **The positive trajectories of bounded sets are bounded.** Indeed, (1.11) implies that, for  $x \notin B(x_0, R)$ , we have  $V(x, u) = \int_0^u f(x, \xi) d\xi \geq 0$ . Moreover, for  $x \in B(x_0, R)$ ,  $V(x, \cdot)$  is non-increasing on  $(-\infty, -R)$  and non-decreasing on  $(R, \infty)$ . Thus,  $V(x, u)$  is bounded from below for  $x \in B(x_0, R)$  and

$$\forall (u, v) \in X, \quad E(u, v) \geq \frac{1}{2} \|(u, v)\|_X^2 + \text{vol}(B(x_0, R)) \inf V.$$

The Sobolev embeddings  $H^1(\Omega) \hookrightarrow L^{p+1}(\Omega)$  shows that the bounded sets of  $X$  have a bounded energy. Since the energy  $E$  is non increasing along the trajectories of (1.9), we get that the trajectory of a bounded set is bounded.

- ii) **The dynamical system is asymptotically smooth.** The asymptotic compactness exactly corresponds to the statement of Proposition 4.3. Let us briefly explain why it can be extended to the case where  $f$  depends on  $x$ . The key point is the extension of Corollary 4.2. First notice that, since we have assumed that  $f(x, 0) = 0$  on  $\partial\Omega$ ,  $f(x, u)$  satisfies Dirichlet boundary condition if  $u$  does. Then, it is not difficult to see that the discussion following Theorem 4.1 can be extended to the case  $f$  depending on  $x$  by using estimates (1.10). Corollary 4.2 follows then, except for a small change: since it is possible that  $f(x, 0) \neq 0$  for some  $x \in \Omega$ , the conclusion of Corollary 4.2 should be replaced by

$$\|f(x, v)\|_{L^1([0, T], H_0^{s+\varepsilon}(\Omega))} \leq C(1 + \|v\|_{L^\infty([0, T], H^{s+1}(\Omega) \cap H_0^1(\Omega))}).$$

Then the proof of Proposition 4.3 is based on Corollary 4.2, the boundedness of the positive trajectories of bounded sets (both could be extended to the case where  $f$  depends on  $x$  as noticed above) and an application of Proposition 2.6 outside of a large ball. We conclude by noticing that, for  $x$  large,  $f(x, u)u \geq 0$  and  $\gamma(x) \geq \alpha > 0$  and thus Proposition 2.6 can still be applied exactly as in the proof of Proposition 4.3.

- iii) **The dynamical system generated by (1.9) is gradient**, that is that the energy  $E$  is non-increasing in time and is constant on a trajectory  $u$  if and only if  $u$  is an equilibrium point of (1.9). This last property is shown in Proposition 5.4 for  $f$  independent of  $x$  but can be easily generalised for  $f = f(x, u)$ . Notice that the proof of this property is the one, where the analyticity of  $f$  is required since the unique continuation property of Section 3 is used. Finally, we enhance that the gradient structure of (1.9) is interesting from the dynamical point of view since it implies that any trajectory  $u(t)$  converges when  $t$  goes to  $+\infty$  to the set of equilibrium points.
- iv) **The set of equilibrium points is bounded.** The argument is similar to the one of Corollary 5.5: if  $e$  is an equilibrium point of (1.9) then (1.11) implies that

$$\int \frac{1}{2} |\nabla e|^2 + \beta |e|^2 = - \int_{\Omega} f(x, e) e dx \leq -\text{vol}(B(x_0, R)) \inf_{(x, u) \in \Omega \times \mathbb{R}} f(x, u)u,$$

where we have bounded  $f(x, u)u$  from below exactly as we have done for  $V(x, u)$  in i).

It is well known (see [24] or Theorem 4.6 of [45] for examples) that Properties i)-iv) yield the existence of a compact global attractor. Hence, we obtain the conclusion of Theorem 1.4.

## References

- [1] S. Alinhac and G. Métivier, *Propagation de l'analyticité des solutions de systèmes hyperboliques non-linéaires*, Inventiones Mathematicae n°75(2) (1984), pp. 189-204.
- [2] L. Aloui, S. Ibrahim and K. Nakanishi, *Exponential energy decay for damped Klein-Gordon equation with nonlinearities of arbitrary growth*, Communication in Partial Differential Equations n°36 (2011), pp. 797-818.
- [3] L. Aloui, M. Khenissi, *Stabilisation pour l'équation des ondes dans un domaine extérieur*, Rev. Mat. Iberoamericana 18 n°1 (2002), pp. 1-16.
- [4] J. Arrieta, A. Carvalho and J.K. Hale, *A damped hyperbolic equation with critical exponent*, Communication in PDE n°17 (1992), pp. 841-866.
- [5] A.V. Babin and M.I. Vishik, *Attractors of Evolutionary Equations*, North-Holland, 1989.
- [6] C. Bardos, G. Lebeau and J. Rauch, *Un exemple d'utilisation des notions de propagation pour le contrôle et la stabilisation de problèmes hyperboliques*, Rend. Sem. Mat. Univ. Politec. Torino, Rendiconti del Seminario Matematico (già "Conferenze di Fisica e di Matematica"). Università e Politecnico di Torino (1988), pp. 11-31.
- [7] C. Bardos, G. Lebeau and J. Rauch, *Sharp sufficient conditions for the observation, control and stabilization of waves from the boundary*, SIAM Journal on Control and Optimization n°30 (1992), pp. 1024-1065.
- [8] M.D. Blair, H.F. Smith and C.D. Sogge, *Strichartz estimates for the wave equation on manifolds with boundary*, Annales de l'Institut Henri Poincaré (C) Non Linear Analysis n°26 (2009), pp. 1817-1829.
- [9] N. Burq, *Mesures semi-classiques et mesures de défaut. (French) [Semiclassical measures and defect measures]* Séminaire Bourbaki, Vol. 1996/97. Astérisque n°245 (1997), Exp. n°826, 4, pp. 167-195.
- [10] N. Burq, *Contrôlabilité exacte des ondes dans des ouverts peu réguliers. (French) [Exact controllability of waves in nonsmooth domains]* Asymptotic Analysis n°14 (1997), pp. 157-191.

- [11] N. Burq and P. Gérard. Condition nécessaire et suffisante pour la contrôlabilité exacte des ondes. *Comptes rendus de l'Académie des sciences. Série 1, Mathématique*, 325(7):749–752, 1997.
- [12] N. Burq, G. Lebeau and F. Planchon, *Global existence for energy critical waves in 3-D domains*, Journal of the American Mathematical Society n°21 (2008), pp. 831-845.
- [13] B. Dehman, *Stabilisation pour l'équation des ondes semi-linéaire*, Asymptotic Analysis n°27 (2001), pp. 171-181.
- [14] B. Dehman and P. Gérard, *Stabilization for the Nonlinear Klein Gordon Equation with critical Exponent*, preprint.
- [15] B. Dehman, P. Gérard and G. Lebeau *Stabilization and control for the nonlinear Schrödinger equation on a compact surface*, Mathematische Zeitschrift n°254 (2006), pp. 729-749.
- [16] B. Dehman and G. Lebeau, *Analysis of the HUM Control Operator and Exact Controllability for Semilinear Waves in Uniform Time*, SIAM Journal on Control and Optimization n°48 (2009), pp. 521-550.
- [17] B. Dehman, G. Lebeau and E. Zuazua, *Stabilization and control for the subcritical semilinear wave equation*, Annales Scientifiques de l'École Normale Supérieure n°36 (2003), pp. 525-551.
- [18] P. Gérard. Microlocal Defect Measures. *Comm. Partial Diff. eq.*, 16:1762–1794, 1991.
- [19] P. Gérard, *Oscillations and Concentration Effects in Semilinear Dispersive Wave Equations*, Journal of Functional Analysis n°141 (1996), pp. 60-98.
- [20] P. Gérard and E. Leichtnam *Ergodic properties of eigenfunctions for the Dirichlet problem*, Duke Mathematical Journal n°71 (1993), pp. 559-607.
- [21] J. Ginibre et G. Velo, *The global Cauchy problem for the nonlinear Klein-Gordon equation*, Mathematische Zeitschrift n°189 (1985), pp. 487-505.
- [22] J. Ginibre et G. Velo, *The global Cauchy problem for the nonlinear Klein-Gordon equation. II*. Annales de l'Institut Henri Poincaré, Analyse Non Linéaire n°6 (1989), pp. 15-35.
- [23] M. Golubitsky and V. Guillemin, *Stable mapping and their singularities*, Graduate Texts in Mathematics n°14, Springer-Verlag, New York-Heidelberg, 1973.
- [24] J.K. Hale, *Asymptotic behavior of dissipative systems*, Mathematical Survey n°25, American Mathematical Society, 1988.

- [25] J.K. Hale and G. Raugel, *Regularity, determining modes and Galerkin methods*, Journal de Mathématiques Pures et Appliquées n°82 (2003), pp. 1075-1136.
- [26] J.K. Hale and G. Raugel, manuscript.
- [27] A. Haraux, *Two remarks on hyperbolic dissipative problems*. Nonlinear partial differential equations and their applications. Collège de France seminar, Vol. VII (Paris, 1983-1984), pp. 161-179. Res. Notes in Math. 122. Pitman, Boston, MA, 1985.
- [28] A. Haraux, *Stabilization of trajectories for some weakly damped hyperbolic equations*, Journal of Differential Equations n°59 (1985), pp. 145-154.
- [29] A. Haraux, *Une remarque sur la stabilisation de certains systèmes du deuxième ordre en temps*, Portugaliae Mathematica n°46 (1989), pp. 246-257.
- [30] L. Hörmander, *On the uniqueness of the Cauchy problem under partial analyticity assumptions*, Geometrical optics and related topics (Cortona, 1996), Progr. Nonlinear Differential Equations Appl., Birkhäuser Boston n°32 (1997), pp. 179-219.
- [31] O. Ivanovici, *Counter-examples to the Strichartz estimates for the wave equation in general domains with boundary*, Journal of the European Mathematical Society, n°14(5) (2012).
- [32] R. Joly, *New examples of damped wave equations with gradient-like structure*, Asymptotic Analysis n°53 (2007), pp. 237-253.
- [33] L.V. Kapitanski, *Some generalizations of the Strichartz-Brenner inequality*, Leningrad Mathematical Journal n°1 (1990), pp. 693-726.
- [34] L.V. Kapitanski, *Minimal compact global attractor for a damped semilinear wave equation*, Communications in Partial Differential Equations n°20 (1995), pp. 1303-1323.
- [35] M. Khenissi, *Equation des ondes amorties dans un domaine extérieur. (French. English and French summary) [Dissipative wave equation in an exterior domain]*, Bulletin de la Société Mathématique de France n°131 (2003), pp. 211-228.
- [36] C. Laurent, L. Rosier and B. Y. Zhang, *Control and stabilization of the Korteweg-de Vries equation on a periodic domain*, Communications in Partial Differential Equations n°35(4) (2010), pp. 707-744.
- [37] C. Laurent, *Global controllability and stabilization for the nonlinear Schrödinger equation on some compact manifolds of dimension 3*, SIAM Journal on Mathematical Analysis n°42 (2010), pp. 785-832.

- [38] C. Laurent, *On stabilization and control for the critical Klein Gordon equation on 3-D compact manifolds*, Journal of Functional Analysis n°260 (2011), pp. 1304-1368.
- [39] G. Lebeau. *Equation des ondes amorties*. In *Algebraic and geometric methods in mathematical physics: proceedings of the Kaciveli Summer School, Crimea, Ukraine, 1993*, page 73. Springer, 1996.
- [40] G. Métivier, *Counterexamples to Hölmgren's uniqueness for analytic non linear Cauchy problems*, Inventiones Mathematicae n°112(1) (1993), pp. 217-222.
- [41] L. Miller, *Escape function conditions for the observation, control, and stabilization of the wave equation*, SIAM Journal on Control and Optimization n°41 (2002), pp. 1554-1566.
- [42] C. Miranda, *Partial differential equations of elliptic type*. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 2. Springer-Verlag, New York-Berlin, 1970.
- [43] A. Pazy, *Semigroups of linear operators and applications to partial differential equations*, Applied Mathematical Sciences n°44 (1983), Springer-Verlag.
- [44] J. Rauch and M. Taylor, *Exponential decay of solutions to hyperbolic equations in bounded domains*, Indiana University Mathematical Journal n°24 (1974), pp. 79-86.
- [45] G. Raugel, *Handbook of dynamical systems*, chapter 17 of volume 2. Edited by B.Fiedler, Elsevier Science, 2002.
- [46] L. Robbiano, *Théorème d'unicité adapté au contrôle des solutions des problèmes hyperboliques*, Communications in Partial Differential Equations n°16 (1991), pp. 789-800.
- [47] L. Robbiano and C. Zuily, *Uniqueness in the Cauchy problem for operators with partially holomorphic coefficients*, Inventiones Mathematicae n°131 (1998), pp. 493-539.
- [48] R. Strichartz, *Restriction of Fourier transform to quadratic surfaces and decay of solutions to the wave equation*, Duke Mathematical Journal n°44 (1977), pp. 705-714.
- [49] L. Tartar, *An Introduction to Sobolev Spaces and Interpolation*. Lecture Notes of the Unione Matematica Italiana n°3. Springer, Berlin, 2007.
- [50] L. Tartar. H-measures, a new approach for studying homogenisation, oscillations and concentration effects in partial differential equations. *Proceedings of the Royal Society of Edinburgh. Section A. Mathematics*, 115(3-4):193–230, 1990.
- [51] D. Tataru, *Unique continuation for solutions to PDE's; between Hörmander's theorem and Holmgren's theorem*, Communication in Partial Differential Equations. n°20(5-6) (1995), pp. 855-884.



- [52] D. Tataru, *Unique continuation for operators with partially analytic coefficients*, Journal de Mathématiques Pures et Appliquées. n°78(5) (1999), pp. 505-521.
- [53] E. Zuazua, *Exponential decay for semilinear wave equations with localized damping*, Communications in Partial Differential Equations n°15 (1990), pp. 205-235.
- [54] E. Zuazua, *Exponential decay for the semilinear wave equation with localized damping in unbounded domains*, Journal de Mathématiques Pures et Appliquées n°70 (1992), pp. 513-529.