



Trapping colloids near chemical stripes via critical Casimir forces

Matthias Troendle, Olga Zvyagolskaya, Andrea Gambassi, Dominik Vogt, Ludger Harnau, Clemens Bechinger, Siegfried Dietrich

► To cite this version:

Matthias Troendle, Olga Zvyagolskaya, Andrea Gambassi, Dominik Vogt, Ludger Harnau, et al.. Trapping colloids near chemical stripes via critical Casimir forces. *Molecular Physics*, Taylor & Francis, 2011, pp.1. <10.1080/00268976.2011.553639>. <hal-00686243>


HAL Id: hal-00686243

<https://hal.archives-ouvertes.fr/hal-00686243>

Submitted on 9 Apr 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Trapping colloids near chemical stripes via critical Casimir forces

Journal:	<i>Molecular Physics</i>
Manuscript ID:	TMPH-2010-0470
Manuscript Type:	Special Issue paper - In honour of Bob Evans
Date Submitted by the Author:	30-Nov-2010
Complete List of Authors:	Troendle, Matthias; Max Planck Institute for Metals Research, Dietrich; Universität Stuttgart, Institut für Theoretische und Angewandte Physik Zvyagolskaya, Olga; Universität Stuttgart, 2. Physikalisches Institut Gambassi, Andrea; SISSA — International School for Advanced Studies and INFN; Laboratoire de Physique Théorique et Hautes Energies, UMR 7589, Université Pierre et Marie Curie Vogt, Dominik; Universität Stuttgart, 2. Physikalisches Institut Harnau, Ludger; Max Planck Institute for Metals Research, Dietrich; Universität Stuttgart, Institut für Theoretische und Angewandte Physik Bechinger, Clemens; Universität Stuttgart, 2. Physikalisches Institut; Max Planck Institute for Metals Research, Dietrich Dietrich, Siegfried; Max Planck Institute for Metals Research; Universität Stuttgart, Institut für Theoretische und Angewandte Physik
Keywords:	Solvation forces, critical Casimir effect, colloidal suspension, chemically patterned substrates
<p>Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.</p>	

troendlesub.zip

SCHOLARONE™
Manuscripts

For Peer Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INVITED ARTICLE

Trapping colloids near chemical stripes via critical Casimir forces

Matthias Tröndle^{a,b*}, Olga Zvyagolskaya^c, Andrea Gambassi^{d,e}, Dominik Vogt^c, Ludger Harnau^{a,b}, Clemens Bechinger^{a,c} and Siegfried Dietrich^{a,b}

^aMax-Planck-Institut für Metallforschung, Heisenbergstr.3, 70569 Stuttgart, Germany;

^bInstitut für Theoretische und Angewandte Physik, Universität Stuttgart, Pfaffenwaldring 57, 70569 Stuttgart, Germany;

^c2. Physikalisches Institut, Universität Stuttgart, Pfaffenwaldring 57, 70569 Stuttgart, Germany;

^dSISSA — International School for Advanced Studies and INFN, via Bonomea 265, 34136 Trieste, Italy;

^eLaboratoire de Physique Théorique et Hautes Energies, UMR 7589, Université Pierre et Marie Curie - Paris VI, 4 Place Jussieu, 75252 Paris Cedex 05, France

(Received 00 Month 200x; final version received 00 Month 200x)

We study theoretically and experimentally the solvent-mediated critical Casimir force acting on colloidal particles immersed in a binary liquid mixture of water and 2,6-lutidine and close to substrates which are chemically patterned with periodically alternating stripes of antagonistic adsorption preferences. These patterns are experimentally realized via microcontact printing. Upon approaching the critical demixing point of the solvent, normal and lateral critical Casimir forces generate laterally confining effective potentials for the colloids. We analyze in detail the rich behavior of the spherical colloids close to such substrates. For all patterned substrates we investigated, our measurements of these effective potentials agree with the corresponding theoretical predictions. Since both the directions and the strengths of the critical Casimir forces can be tuned by minute temperature changes, this provides a new mechanism for controlling colloids as model systems, opening encouraging perspectives for applications.

Keywords: Solvation forces, critical Casimir effect, colloidal suspension, chemically patterned substrates

1. Introduction

The structures and the dynamics of colloidal suspensions are to a large extent determined by the effective interactions among the mesoscopic colloids which are mediated by the solvent surrounding them. A rather intriguing example of such solvation forces is the critical Casimir effect [1] which occurs close to a critical point of the solvent. As a soft-matter analogue of the quantum-electrodynamic Casimir effect [2, 3], the confinement of the concentration fluctuations within a near-critical binary liquid mixture by the surfaces of the colloidal particles results in an attractive or repulsive effective interaction [4–6]. Upon approaching the bulk critical demixing point of the mixture at temperature $T = T_c$ the critical fluctuations become long-ranged so that the resulting effective interaction among colloids can extend well beyond the molecular scale of the solvent; it may act even at distances comparable to the size of the colloids. The strength and the spatial range of the critical Casimir force depend sensitively on temperature because the correlation length ξ , which characterizes the extension of the spatial correlations of the critical fluctuations and

*Corresponding author. Email: troendle@mf.mpg.de

1 therefore sets the interaction range, diverges as $\xi \simeq \xi_0 |(T - T_c)/T_c|^{-\nu}$ upon approaching
2 the critical point, $T \rightarrow T_c$, where $\nu \simeq 0.63$ is a standard bulk critical exponent¹ and ξ_0
3 is a solvent-specific microscopic length. The critical Casimir effect is *universal* in the
4 sense that all quantities related to it, such as the various components of the force and the
5 associated potentials, can be suitably expressed in terms of scaling functions which are
6 identical for all those fluids undergoing the same kind of transition, irrespective of their
7 molecular details and material properties, which actually enter only via two independent
8 non-universal amplitudes, such as ξ_0 and the value of T_c .

9 Besides some indirect quantitative experimental determinations of the critical Casimir
10 force acting within wetting films [7–9], its *direct* effect on a spherical colloid close to a
11 substrate and immersed in a critical water-lutidine mixture was measured only recently
12 [10]. These experiments clearly demonstrated that, upon approaching the critical point,
13 the critical Casimir force can modify the effective interaction between the colloid and
14 the substrate by several multiples of $k_B T$. Moreover, depending on the preference of the
15 confining surfaces for lutidine or water — denoted as $(+)$ and $(-)$ effective bound-
16 ary conditions (BC) for the order parameter, respectively — the critical Casimir force
17 between the colloid and the substrate is attractive [for (\pm, \pm) boundary conditions] or re-
18 pulsive [for $(+, -)$ boundary conditions] [10]. In addition, these critical Casimir forces
19 depend strongly on the geometrical shape of the confining surfaces so that topographically
20 structured surfaces experience also a component of the force which is parallel to the walls
21 [11], whereas nonspherical particles are subject to a torque [12].

22 One can combine these features of the critical Casimir effect by using surfaces which
23 are *chemically patterned* on the micron-scale in order to obtain also *lateral* critical
24 Casimir forces [13]. The critical Casimir interaction between a colloid and a substrate
25 which is chemically patterned with stripes of alternating adsorption preferences results
26 in a laterally confining potential for the colloid, which can be controlled and reversibly
27 switched on and off by varying the temperature of the near-critical solvent [14–16]. Re-
28 cently, such a substrate has been realized by removing via a focused ion beam a regular
29 pattern of equally spaced stripes from the hydrophobic HMDS-coating of a glass sur-
30 face [14]. The effective interaction between this chemically patterned substrate and a col-
31 loidal particle has then been measured by monitoring via digital video microscopy the
32 behaviour of a dilute solution of colloidal particles as the demixing critical point of the
33 binary mixture of water and lutidine is approached [14]. In comparing these data with
34 the corresponding theoretical predictions [15] it turned out that the critical Casimir poten-
35 tial resulting from a chemical pattern depends rather strongly on its geometrical details
36 and it actually provides a sensitive tool to probe these microscopic features, which might
37 not be easily accessible otherwise [15]. Indeed agreement between the theoretical predic-
38 tions of Ref. [15] and the experimental data of Ref. [14] for the spatially varying effective
39 potential was found only by assuming that the chemical steps between two subsequent
40 stripes within the pattern were not microscopically sharp due to fluctuations inherent to
41 the preparation process of the structures [15]. However, in spite of this improvement in
42 the comparison between the *shapes* of the theoretical and experimental potentials, the
43 analysis of the temperature dependence of the corresponding potential *depths* renders an
44 amplitude ξ_0 of the correlation length ξ (treated as a fitting parameter) which is about
45 twice as large [15] as the one known from independent previous measurements. This per-
46 sisting discrepancy, together with the necessity to test independently the aforementioned
47 assumption used in Ref. [15], calls for a more detailed experimental study of the critical
48 Casimir force acting in the presence of patterned substrates.

49 Colloidal suspensions are not only relevant as model systems in soft-matter physics

50
51
52
53
54
55
56
57
58 ¹Here we quote the value of ν belonging to the Ising universality class, which is relevant for the interpretation of experi-
59 mental data concerning classical binary liquid mixtures.

1 but they may find also applications in integrated micro- and nano-devices. Since critical
2 Casimir forces provide a tool to exert active control over the strength and the direction
3 of interactions between colloids in such devices, a thorough quantitative understanding of
4 the underlying physics is essential. Recently, it was theoretically predicted that for suitable
5 geometrical parameters colloids can even levitate in a stable position above chemically
6 patterned substrates [16]. Such a “critical Casimir levitation” may even help in overcoming
7 the problem of stiction which hampers the functioning of currently available micro-
8 and nano-electromechanical devices. Therefore, in order to benefit from this wide range of
9 possibilities, a theoretical understanding of the phenomenon and the experimental demon-
10 stration of its applicability are essential.

11 Here, we report on experimental studies of the critical Casimir interaction between a
12 colloid and substrates which are chemically structured by microcontact printing of alkane-
13 thiols. We shall see below that, in the present context the substrates prepared by this tech-
14 nique turn out to exhibit higher chemical and geometrical resolution than those prepared
15 via focused ion beam, which was used in previous experiments [14]. A dilute suspension
16 of hydrophilic spherical colloids of radius R which impose $(-)$ boundary conditions to
17 a near-critical water-lutidine mixture is exposed to patterned substrates consisting of al-
18 ternating and periodically repeating stripes of width L_{\pm} with boundary conditions (\pm) so
19 that the periodicity is $P = L_{+} + L_{-}$ along the x -direction, whereas the pattern is trans-
20 lationally invariant along the orthogonal y -direction. In order to extend previous inves-
21 tigations [14, 15] to a wider range of chemical patterns, we consider here sequences of
22 stripes with various periodicities P and stripe widths L_{-} . In particular this allows us to
23 investigate the case in which the effects of neighboring chemical steps interfere, which
24 strongly affects the resulting forces acting on the colloid. As in Ref. [14] we measure
25 the laterally varying equilibrium spatial distribution of the colloids by digital video mi-
26 croscopy and define an effective potential on the basis of the equilibrium number density
27 of the colloids projected along the vertical z direction (orthogonal to x and y). We analyze
28 theoretically this potential and the ensuing surface-to-surface distance z of the colloid
29 from the substrate and compare our findings with the experimental data. Our theoretical
30 analysis provides accurate information also about the vertical probability distribution of
31 the positions of the colloidal particles which, however, cannot be resolved by the kind of
32 video microscopy used in the current experiment. For all stripe widths and temperatures
33 studied we find quantitative agreement between the measurements and the theoretical pre-
34 dictions. Both the shapes and the depths of the measured potentials agree very well with
35 the theoretical analysis. Moreover, the uncertainty of the actual position of the subsequent
36 chemical steps is significantly reduced by using the present microcontact printing in order
37 to structure the substrate. The amplitude ξ_0 of the bulk correlation length, which we esti-
38 mate by comparing the depth of the measured potentials with the theoretical predictions,
39 turns out to be in quantitative agreement with previous, independent estimates. Thus, our
40 detailed comparison of experimental data and theory demonstrates that indeed the full
41 benefits of the critical Casimir force can be reaped reliably and may be utilized in poten-
42 tial applications, opening a new route for using colloids as model systems or in micro-
43 and nano-devices.

51 2. Theory

52 2.1. Critical Casimir potential

53 According to renormalization group theory, in the vicinity of the critical point at $T = T_c$
54 the normal and lateral critical Casimir forces as well as the corresponding potential can
55 be described by *universal scaling functions* (see, e.g., Refs. [4, 5, 17]). These scaling
56 functions are universal in the sense that they do not depend on molecular details but only
57
58
59
60

1 on the gross features of the system and of the confining surfaces. These gross features,
 2 among which there are the spatial dimension d , the symmetries of the system, and the
 3 geometry of the boundaries, determine the so-called bulk [4, 5] and surface [4, 18–20]
 4 universality classes of the associated critical point. The relevant thermodynamic proper-
 5 ties which emerge upon approaching the point of the continuous phase transition can be
 6 understood and analyzed in terms of the fluctuations of the so-called order parameter ϕ
 7 of the phase transition. For the consolute point of phase segregation of a binary liquid
 8 mixture ϕ is given by the difference between the local and the mean concentration of one
 9 of the two components of the mixture; thus, it is a scalar quantity and the bulk univer-
 10 sality class is of the so-called Ising type (see, e.g., Refs. [10, 21]). For classical binary
 11 liquid mixtures, the surfaces generically exhibit preferential adsorption for one of the two
 12 species of the mixture, which results in a local enhancement of the order parameter ϕ
 13 close to the surface. This enhancement is effectively described by symmetry-breaking
 14 surface fields [4, 18–20] and it is usually denoted by (+) and (−) boundary conditions
 15 (BC) corresponding to having a preference for $\phi > 0$ and $\phi < 0$, respectively, at the sur-
 16 face. For the water-lutidine mixture we are interested in, one conventionally indicates the
 17 preferential adsorption for lutidine and water as (+) and (−) BC, respectively (see also
 18 Refs. [10, 21]). In the experimental setup described below, the lower critical demixing
 19 point of the water-lutidine mixture is always approached from the homogeneous (mixed)
 20 phase at temperatures $T < T_c$ upon increasing T at *fixed critical composition* of the mix-
 21 ture. From the experimental point of view it is rather difficult to quantify the strength of
 22 the adsorption preference exhibited by the different portions of the surfaces. Therefore
 23 the comparison with theoretical predictions requires assumptions, which can be verified a
 24 posteriori. In the present case we have qualitative experimental evidence that the chemical
 25 treatment of the surfaces results in rather pronounced adsorption preferences and therefore
 26 we shall assume that the surfaces are characterized by the so-called *strong critical adsorp-*
 27 *tion* fixed point [18, 19]. We consider here neither the opposite case of weak adsorption
 28 at the surfaces (see, e.g., Refs. [22, 23]) nor the effects due to off-critical compositions of
 29 the mixture, which might lead to a bridging transition analogous to capillary condensation
 30 (see, e.g., Refs. [21, 24–29]).

31 For the film geometry — in which the binary liquid mixture is confined between two
 32 parallel, planar, and macroscopically extended walls at a distance L — the normal critical
 33 Casimir force per unit area is given by [17]

$$34 \quad f_{(+,\pm)}(L, T) = k_B T \frac{1}{L^d} k_{(+,\pm)}(L/\xi), \quad (1)$$

35 where the index $(+, \pm)$ denotes the combination of BC (+) and (−) at the two walls. (In
 36 the absence of a symmetry-breaking bulk field within the film, which would correspond to
 37 an off-critical concentration of the mixture, one has $f_{(-,-)} = f_{(+,+)$.) The universal scaling
 38 functions $k_{(+,\pm)}$ depend only on the scaled variable L/ξ , i.e., on the film thickness L
 39 in units of the bulk correlation length ξ . The functions $k_{(+,\pm)}$ have been calculated exactly
 40 in $d = 2$ [30], for $d < 4$ using perturbative field-theoretical methods [31] or effective
 41 theories [32], and in $d = 3$ numerically via Monte Carlo simulations [33–35]. At present,
 42 a quantitatively reliable theoretical determination of both $k_{(+,+)}$ and $k_{(+,-)}$ in $d = 3$ is
 43 provided only by Monte Carlo simulations. For a chemically structured wall opposing
 44 another planar wall forming a film geometry, the critical Casimir force has been obtained
 45 within mean-field theory (MFT) [13] and from Monte Carlo simulations [36, 37].

46 For a spherical colloid with (−) BC opposite to a chemically patterned substrate with
 47 alternating (+) and (−) BC the critical Casimir potential has been calculated both nu-
 48 merically within full mean-field theory (corresponding to $d = 4$) and by resorting to the
 49 so-called Derjaguin approximation exploiting the full knowledge of $k_{(+,\pm)}$ in the film ge-
 50 ometry for $d = 3$ [16]. Accordingly, the critical Casimir force and the critical Casimir
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60

potential for a colloid close to a chemically patterned wall can be expressed in terms of universal scaling functions which take the geometry into account and which depend on the following scaled quantities:

- $\Theta = z/\xi$, where z is the surface-to-surface distance between the colloid and the substrate,
- $\Delta = z/R$, where R is the radius of the spherical colloid,
- $\Xi = x/\sqrt{Rz}$, where x is the lateral coordinate of the center of the colloid such that $x = 0$ corresponds to the colloid being located opposite to the center of a stripe with $(-)$ BC,
- $\Pi = P/\sqrt{Rz}$, where P is the periodicity of the stripe pattern along the x -direction,
- $\lambda = L_-/P$, where L_- is the width of the stripes with $(-)$ BC.

In particular in $d = 3$ the critical Casimir potential Φ_C can be written as [16]

$$\Phi_C(L_-, P, x, z, R, T) = k_B T \frac{R}{z} \vartheta(\lambda, \Pi, \Xi, \Theta, \Delta), \quad (2)$$

where ϑ is a universal scaling function. The realization of the present geometrical setup of a sphere facing a plane still represents a challenge for lattice based Monte Carlo simulations. Accordingly, up to now it is not possible to obtain accurate numerical data for this three-dimensional geometry and therefore one has to rely on the Derjaguin approximation (DA), based on the assumption of additivity, in order to calculate approximately the critical Casimir potential (see Ref. [16]). Within the Derjaguin approximation, the scaling function ϑ [Eq. (2)] can be expressed in terms of the known scaling functions $k_{(\pm,-)}$ for the film geometry [16]:

$$\vartheta(\lambda, \Pi, \Xi, \Theta, \Delta \rightarrow 0) = \frac{\vartheta_{(+,-)}(\Theta) + \vartheta_{(-,-)}(\Theta)}{2} + \frac{\vartheta_{(+,-)}(\Theta) - \vartheta_{(-,-)}(\Theta)}{2} \omega(\lambda, \Pi, \Xi, \Theta), \quad (3)$$

where

$$\vartheta_{(\pm,-)}(\Theta) = 2\pi \int_1^\infty d\beta (\beta - 1) \beta^{-3} k_{(\pm,-)}(\beta\Theta) \quad (4)$$

are the scaling functions of the critical Casimir potential of a colloid in front of a homogeneous wall as calculated within the Derjaguin approximation [10], and

$$\omega(\lambda, \Pi, \Xi, \Theta) = 1 + \sum_{n=-\infty}^{\infty} \left\{ \omega_s(\Xi + \Pi(n + \frac{\lambda}{2}), \Theta) - \omega_s(\Xi + \Pi(n - \frac{\lambda}{2}), \Theta) \right\} \quad (5)$$

with ω_s given by

$$\omega_s(\Xi \geq 0, \Theta) = \mp 1 \pm \frac{\Xi^4 \int_1^\infty ds \frac{s^{\text{sarccos}(s^{-1/2}) - \sqrt{s-1}}}{(1 + \Xi^2 s/2)^d} \Delta k(\Theta[1 + \Xi^2 s/2])}{\vartheta_{(+,-)}(\Theta) - \vartheta_{(-,-)}(\Theta)}, \quad (6)$$

where $\Delta k(\Theta) = k_{(+,-)}(\Theta) - k_{(-,-)}(\Theta)$. For $\Theta = 0$ and $\Theta \gg 1$ there are analytic expressions for ω_s (see Eqs. (6) and (7) in Ref. [15], respectively). The accuracy of the Derjaguin approximation has been checked numerically within mean-field theory and it has turned out that this approximation describes quantitatively the actual behavior of those numerical data which correspond to $\Theta \gtrsim 4$ for $\Delta \lesssim 1$ and $\Pi \gtrsim 0.5$, as well as to $0 \leq \Theta \lesssim 4$ with $\Delta \lesssim 0.3$ and $\Pi \gtrsim 2$ [16]. In the experiments discussed further below the corresponding values of Θ , Δ , and Π vary within these ranges so that, assuming that the previous quantitative conclusions extend to $d = 3$, we expect the Derjaguin approximation to be quantitatively reliable.

2.2. Background forces

In addition to the critical Casimir force due to the critical fluctuations of the solvent, the colloidal particles of the suspension are subjected to additional effective forces which are characterized by a smooth and rather mild dependence on temperature. Typical background forces acting within the colloidal suspensions of present interest are due to (screened) electrostatic and dispersion interactions and to the gravitational field. In a first approximation, which neglects possible mutual influences of these forces, their total potential is given by the sum of the corresponding contributions: (i) the electrostatic potential Φ_{el} , (ii) the gravitational potential Φ_g , (iii) the van der Waals interaction Φ_{vdW} , and (iv) the effective critical Casimir potential Φ_C .

Electrostatics — This interaction originates from the fact that, due to the formation of charge double-layers, the surface of the colloids and of the substrate acquire a surface charge once immersed in the liquid solvent. As a result, the polystyrene colloids of radius $R = 1.2\mu\text{m}$ and (–) BC immersed in water-lutidine mixtures experience an electrostatic repulsion from the substrate. The screened electrostatic potential of the colloid at a surface-to-surface distance z from a *homogeneous* substrate with (\pm) BC is well approximated by

$$\Phi_{el,\pm}(z)/(k_B T) = \exp\{-\kappa(z - z_0^\pm)\}, \tag{7}$$

where κ^{-1} is the screening length and z_0^\pm describes the strength of the electrostatic repulsion from the substrate with (\pm) BC. Although the values of these parameters are determined by the surface charge of the colloid, the dielectric constant of the mixture etc. [38], here they will be treated as fitting parameters of the actual experimental data for $\Phi_{el,\pm}$. In experimental conditions similar to the present ones as far as the mixture and the colloids are concerned, one finds $\kappa^{-1} \simeq 12\text{nm}$ [10, 21, 23] and $z_0^\pm \simeq 0.1 - 0.2\mu\text{m}$ [10, 15, 21, 23] as typical values at $T \simeq T_c \simeq 307\text{K}$. In view of a possible difference between z_0^+ and z_0^- , the resulting electrostatic potential of a colloid close to a *patterned* substrate such as the one described above depends on the lateral position x of the colloid. Such a dependence can be accounted for theoretically within the Derjaguin approximation (by repeating the calculations described in Appendices A.2, B, and C in Ref. [16] for Φ_C), which is expected to be particularly accurate due to the exponential decay of $\Phi_{el,\pm}$ in Eq. (7) as a function of the distance z from the substrate. For this electrostatic potential $\Phi_{el}(x, z)$ one therefore finds (for a colloid facing the center of a (–) stripe at $x = 0$)

$$\Phi_{el}(x, z) = \frac{\Phi_{el,+}(z) + \Phi_{el,-}(z)}{2} + \frac{\Phi_{el,+}(z) - \Phi_{el,-}(z)}{2} \Omega(x), \tag{8}$$

where

$$\Omega(x) = 1 + \sum_{n=-\infty}^{\infty} \{\theta([x + nP + L_-/2]/\Lambda) - \theta([x + nP - L_-/2]/\Lambda)\}, \tag{9}$$

with $\theta(u) = -\text{erf}(u)$ and $\Lambda = \sqrt{2R\kappa^{-1}} \simeq 0.17\mu\text{m}$. Ω and Φ_{el} depend on the geometric parameters L_- and P describing the pattern.

Dispersion forces — For the particular choice of materials and conditions used in the present experiments, van der Waals forces turn out to be negligible compared with the other contributions [10, 15, 21, 39]. In addition, the temperature dependence of the dielectric permittivity of the water-lutidine mixture is negligible (see footnote 4 in Ref. [15]) and therefore no significant changes are expected to occur in Φ_{vdW} and Φ_{el} within the range of temperatures explored experimentally.

Gravity — Due to buoyancy the colloid immersed in the solvent above the patterned

substrate experiences a gravitational potential given by

$$\Phi_g(z) = (\rho_{\text{PS}} - \rho_{\text{WL}})g \frac{4\pi}{3}R^3 z \equiv Gz, \quad (10)$$

where $\rho_{\text{PS}} \simeq 1.055\text{g/cm}^3$ and $\rho_{\text{WL}} \simeq 0.988\text{g/cm}^3$ [40] are the mass densities of the polystyrene colloid and of the water-lutidine mixtures at the critical composition and near T_c , respectively, and $g \simeq 9.81\text{m/s}^2$ is the gravitational acceleration. Accordingly, at $T \simeq 307\text{K}$ one has $G \simeq 1.12k_B T/\mu\text{m}$ and therefore, compared to the other contributions, it turns out that the gravitational potential depends rather mildly on the distance z because it varies only over a few $k_B T$ on the relevant length scale of a few microns. The expression in Eq. (10) assumes that the colloidal particle of mass density ρ_{PS} is floating in a homogeneous medium of mass density ρ_{WL} . However, the laterally varying adsorption preferences of the substrate induce the formation of alternating water-rich or lutidine-rich regions close to the surface of the patterned substrate, which laterally alter the resulting mass density of the solvent as a consequence of water and lutidine having different mass densities. This implies that the effective gravitational constant G acquires a dependence on x . In addition, the preferential adsorption of the colloid, with the ensuing formation of an adsorption profile around it, can lead to a modification of the effective density ρ_{PS} of the colloid itself. However, on the basis of our estimates, we expect all these effects to be negligible for the present experimental conditions [15].

2.3. Total potential

The total potential Φ of the sum of the forces acting on the colloid is given by

$$\Phi(x, z, T) = \Phi_C(L_-, P, x, z, R, T) + \Phi_{\text{el}}(x, z) + \Phi_g(z), \quad (11)$$

where the theoretical expressions for the individual contributions are given by Eqs. (2), (8), and (10). Here and in the following we do not indicate the dependence of Φ on L_- , P , and R , because the values of these parameters are fixed for each individual experiment. Figure 1 shows the total potential $\Phi(x, z, T)$ of a single colloid with $(-)$ BC opposite to a chemically patterned substrate, as a function of both x and z and for three values of the temperature T close to the critical value T_c . These three values correspond to different correlation lengths ξ , as indicated in the figure. The gray area in the x - z plane indicates the vertical projection of the stripe with $(-)$ BC, the center of which corresponds to $x = 0$. In Fig. 1 the white part of the x - z plane corresponds to the projection of the stripe with $(+)$ BC, the center of which is located at $x = P/2 = 0.9\mu\text{m}$. The potential of the forces acting on the colloid is translationally invariant along the y -direction which is not shown in Fig. 1. $\Phi(x, z, T)$ in Fig. 1 has been calculated by using geometrical and interaction parameters which mimic the actual experimental conditions and by using values of the correlation length ξ which are experimentally available. As anticipated above, Fig. 1 clearly shows that the gravitational tail of the potential, which characterizes $\Phi(x, z, T)$ at large values of z , is indeed rather flat on the scales of $k_B T$ and of hundreds of nm. As a consequence of thermal fluctuations — which cause the colloid to explore a region of space within which the total potential Φ differs from its minimum by a few $k_B T$ — the particle is expected to display large fluctuations $\Delta z \simeq k_B T/G$ along the vertical direction. At small particle-substrate distances electrostatic forces are responsible for the strong repulsion of the colloid from the substrate. In Fig. 1 we allowed for a lateral inhomogeneity of the electrostatic potential (i.e., $z_0^+ \neq z_0^-$), which might occur due to different surface charges on the different stripes. This is clearly visible in panel (a) of Fig. 1, which corresponds to a rather small value of the correlation length ξ so that, within the range $z \simeq 0.1\mu\text{m}$




Figure 1. Total potential $\Phi(x, z, T)$ of a colloid with radius $R = 1.2\mu\text{m}$ opposite to a chemically patterned substrate with $L_- = 0.9\mu\text{m}$ and $P = 1.8\mu\text{m}$ for three temperatures corresponding to (a) $\xi = 5\text{nm}$, (b) $\xi = 22\text{nm}$, and (c) $\xi = 26\text{nm}$, respectively. The electrostatic potential corresponds to $z_0^- = 0.12\mu\text{m}$, $z_0^+ = 0.08\mu\text{m}$, and $\kappa^{-1} = 12\text{nm}$ (see the main text for details). At small separations $z \lesssim z_0^\pm$, the colloid is strongly repelled from the substrate due to electrostatics. The effective gravitational potential associated with buoyancy is rather flat, with a spatial slope $G \simeq 1.12k_B T/\mu\text{m}$. Accordingly, the surface-to-surface particle-substrate distance z exhibits large thermal fluctuations of the order of $k_B T/G \simeq 1\mu\text{m}$ as long as ξ is small. Upon approaching the critical point [from (a) to (c)], ξ increases and a deep, local potential minimum arises rapidly as a function of ξ above that part of the substrate (indicated by the shaded area) with the same preferential adsorption as the colloid. The colloid is eventually confined in this deep potential well at a distance $z \simeq z_0^-$ with almost no vertical fluctuations. The black lines correspond to cuts through the potential surface at constant values of z and x , respectively.

Fig. 1(a) refers to, the contributions of the critical Casimir force are negligible. In this case, the x -dependent electrostatic contribution dominates at small values of z , whereas the laterally homogeneous gravitational potential dominates at larger distances. However, upon approaching the critical point [Figs. 1(b) and 1(c)], the correlation length ξ increases and the critical Casimir force acting on the colloid builds up; it is repulsive within the region with (+) BC whereas it is attractive within the region with (-) BC. In the latter case the behavior of the colloid is eventually determined by the electrostatic repulsion and the attractive critical Casimir force, whereas in the former case it is determined by gravitation and the repulsive Casimir effect. Above a certain threshold value of ξ , which depends on the specific choice of the various geometrical and physical parameters, a very deep and steep potential well develops rapidly close to the stripe with (-) BC, which therefore confines the vertical motion of the colloid at much smaller values of z than before with very limited thermal fluctuations of the particle-wall distance. In contrast, the vertical repulsive critical Casimir force, which the colloid experiences above the stripe with (+) BC, pushes it further away from the surface, but the corresponding fluctuations of the vertical position z (still limited only by the gravitational tail) are not significantly affected. Thus, the full theoretical analysis of the various forces at play reveals a rather interesting energy landscape which is strongly temperature dependent.

2.4. Measured potential

In order to measure experimentally the total potential Φ of the forces acting on a colloid a very effective approach consists in monitoring the equilibrium Brownian motion of a *single* particle and inferring from the sampled probability distribution function $\bar{P}(x, y, z) \propto \exp(-\Phi(x, z, T)/(k_B T))$ the potential as $\Phi/(k_B T) = -\ln \bar{P} + \text{const}$, where x and y are the lateral coordinates of the projection of the colloid center onto the substrate surface. This approach forms the basis of total internal reflection microscopy, which has been used in Refs. [10, 21, 23] to study critical Casimir forces. Alternatively, one can study a colloidal *suspension* which is sufficiently dilute so that the inter-particle interaction is negligible. In this case, the mean equilibrium number density $\rho(x, y, z)$ of the colloids at position (x, y, z) is proportional to the single-colloid probability distribution function \bar{P} and therefore it is again given by $\rho(x, y, z) \propto \exp(-\Phi(x, z, T)/k_B T)$. In the experimental setup described below, the positions of the centers of the colloids are monitored via a digital video camera positioned below the substrate. Accordingly, the surface-to-surface

distances z of the colloids from the substrate are not resolved and the camera records only the projected number density $\rho_P(x, y) \equiv \int_0^\infty dz \rho(x, y, z)$. Due to the translational invariance of the chemical pattern along the y direction of length $l \gg P, R$, the density $\rho_P(x, y)$ can be conveniently projected further onto the x -axis, resulting in an effective number density $\hat{\rho}(x) = l^{-1} \int_0^l dy \rho_P(x, y) = l^{-1} \int_0^l dy \int_0^\infty dz \rho(x, y, z)$, which depends only on x . This projection increases the statistics and therefore the accuracy with which this projected density can be determined experimentally. Subsequently, an effective potential $\hat{V}(x)$ (up to an irrelevant additive constant) can be associated with $\hat{\rho}(x)$ such that $\hat{\rho}(x) \propto \exp(-\hat{V}(x)/(k_B T))$. (Note that due to the thermal fluctuations of the colloids along the vertical direction, even if one knows the average distance $z_{\text{avg}}(x)$ of the colloid from the substrate at a certain lateral position x , the effective potential $\hat{V}(x)$ is not simply given by $\Phi(x, z_{\text{avg}}(x), T) + \text{const}$, as it was implicitly assumed in Ref. [14].) The measured potential

$$\delta\hat{V}(x) = \hat{V}(x) - \hat{V}(P/2) = -k_B T \ln(\hat{\rho}(x)/\hat{\rho}(P/2)) \quad (12)$$

is eventually defined such that it vanishes for a colloid opposite to the center of a repulsive (+) stripe at $x = P/2$. We emphasize that $\delta\hat{V}(x)$ contains universal ingredients stemming from the scaling function associated with Φ_C (see Eq. (11)) as well as nonuniversal contributions due to Φ_{el} and Φ_{g} .

2.5. Non-ideal stripe patterns

Due to the preparation process (see below) the actual position $x = x_s(y)$ of each chemical step separating two adjacent stripes might vary smoothly along the y -axis. This variation affects the measured effective potential \hat{V} of the colloids as long as it occurs on a length scale which is comparable or smaller than the typical distance ℓ_{msd} along the y -axis which each particle explores during the acquisition of the images by the camera. The images acquired during the experiments allow one to estimate such a mean-square displacement ℓ_{msd} to be of the order of tens of μm [14, 15, 41]. The projection along the y -axis, which yields the density $\hat{\rho}(x)$, effectively causes a broadening of $\hat{\rho}(x)$ compared to the case of straight (ideal) chemical steps with $x_s(y) = \text{const}$. In addition, locally a smooth intrinsic chemical gradient of the step leads to such an effect, too. For illustration purposes, we first consider a single chemical step, which is ideally located at $x = 0$ and which generates a potential $\Phi(x, z, T)$. (This reasoning can be extended to the periodic chemical pattern we are presently interested in by assuming additivity of the forces, see Ref. [16] for details.) In order to estimate the effect of these variations of the actual position of the step along the y -axis, we assume that the local position $x_s(y)$ of the step does not change significantly along the y -axis on the scale of the radius R of the colloid, so that on this scale it can be considered as ideal and therefore generates a potential $\Phi(x - x_s(y), z, T)$. For $x_s(y)$ we assume an effective Gaussian distribution $p(x_s)$ along the y -axis, with zero average and standard deviation Δx . Accordingly, the projection $l^{-1} \int_0^l dy$ along the y -axis turns into $\int_{-\infty}^{\infty} dx_s p(x_s)$ and affects the resulting projected density $\hat{\rho}(x)$ and the resulting potential $\hat{V}(x)$ [see Eq. (12)].

2.6. Particle-substrate distance

The theoretical knowledge of the total potential $\Phi(x, z, T)$ [see Eq. (11)] allows one to predict the particle-substrate distance z as a function of the lateral variable x and temperature, a quantity which is not accessible to the experiments discussed below. As anticipated above, the rather small value of G in Eq. (10) is responsible for rather large fluctuations of the particle-substrate distance z around the position $z = z_{\text{min}}(x)$ at which the potential $\Phi(x, z, T)$ has a minimum as a function of z for a fixed lateral position x of the particle

and which corresponds to the position of mechanical equilibrium. In the presence of these large fluctuations it is convenient to consider the x -dependent mean particle-substrate distance $z_{\text{avg}}(x)$, which is determined by the probability distribution function \bar{P} of the colloid, i.e., by the total potential as²

$$z_{\text{avg}}(x) = \frac{1}{N(x)} \int_0^\infty dz z \exp\{-\Phi(x, z, T)/(k_B T)\}, \quad (13)$$

where $N(x) = \int_0^\infty dz \exp\{-\Phi(x, z, T)/(k_B T)\}$ is the normalization. In order to describe the thermal fluctuations of the vertical position of the colloid it is convenient to consider the probability $\bar{P}_<(z; x)$ that for a fixed lateral position x the colloid has a surface-to-surface distance from the substrate smaller than a given z :

$$\bar{P}_<(z; x) = \frac{1}{N(x)} \int_0^z dz' \exp\{-\Phi(x, z', T)/(k_B T)\}. \quad (14)$$

In order to generalize the notion of “standard deviation” to the present case of an asymmetric distribution of the particle-substrate distances at fixed lateral position, we define a lower value $z_{\text{low}}(x)$ and an upper value $z_{\text{upp}}(x)$ of the particle-substrate distances such that

$$\bar{P}_<(z_{\text{low}}(x); x) = 0.159 \quad \text{and} \quad \bar{P}_<(z_{\text{upp}}(x); x) = 1 - 0.159, \quad (15)$$

so that the probability of the colloid to be at a distance z with $z_{\text{low}}(x) < z < z_{\text{upp}}(x)$ is $\simeq 68\%$, whereas the probability to find it at distances smaller (larger) than $z_{\text{low}}(x)$ ($z_{\text{upp}}(x)$) is $\simeq 16\%$; these two properties define the standard deviation for a Gaussian distribution.

Figure 2 shows the behavior of the colloid with (–) BC above a chemically patterned substrate for the two lateral positions (a) $x = 0$ and (b) $x = P/2$ at which the colloid is floating above the center of a (–) and of a (+) stripe, respectively, as a function of the correlation length ξ . In Fig. 2 the values of the geometrical parameters (P, R, L_-) are chosen to correspond to the actual experimental conditions, whereas the parameters z_0^\pm governing the electrostatic repulsion from the substrate are varied within a range which was determined by previous independent experiments. Panel (a) clearly demonstrates that the fluctuations of z for a colloid opposite to an attractive stripe decrease significantly upon increasing the correlation length ξ above a certain threshold value ξ^* which depends on z_0^- . This reflects the emergence of the deep potential well shown in Fig. 1 which results from the competition between an increasingly attractive critical Casimir force and a repulsive electrostatic repulsion, the former being always overwhelmed by the latter around $z = z_0^-$. Indeed, for small values of ξ the average position $z_{\text{avg}}(x = 0)$ is typically determined by the competition between the electrostatic repulsion and the gravitational part, such that $z_{\text{avg}}(x = 0) \simeq z_0^- + k_B T/G \simeq 1 \mu\text{m}$ with fluctuations of the order of $k_B T/G \simeq 1 \mu\text{m}$. On the other hand, for larger values of ξ , one has $z_{\text{upp,avg,low,min}}(x = 0) \simeq z_0^-$ and there are only very small thermal fluctuations of the particle-substrate distance, at most a few tens of nm. Depending on the relative strength of the electrostatic repulsion and the critical Casimir attraction, one can have $z_{\text{min}} < z_{\text{low}}$ if the former dominates the latter, i.e., at small values of ξ or the opposite at large values of ξ . From Fig. 2(a) one concludes that the choice of z_0^- strongly affects the behavior of the particle, both in determining the values of

²Note that the Derjaguin approximation holds only for distances which are small on the scale of the particle size (a detailed analysis of its applicability for the system under consideration is given in Ref. [16]). However, in Eq. (13) also large values of z occur. But at these large particle-wall distances the critical Casimir force as well as the electrostatic force are negligibly small compared to the gravitational force, so that using this approximation is nonetheless not detrimental.




Figure 2. Distances of the colloid from the substrate, for a fixed lateral position (a) $x = 0$ and (b) $x = P/2$, as a function of the bulk correlation length ξ . Here, $P = 1.8 \mu\text{m}$, $R = 1.2 \mu\text{m}$, $L_- = 0.9 \mu\text{m}$, whereas z_{upp} , z_{avg} , z_{low} , and z_{min} indicate the upper, average, lower, and potential-minimum distances of the colloid, respectively (see the main text). The solid lines correspond to the choice $z_0^- = z_0^+ = 0.09 \mu\text{m}$, the dashed lines to $z_0^- = 0.12 \mu\text{m}$ and $z_0^+ = 0.09 \mu\text{m}$, and the dotted lines to $z_0^- = 0.09 \mu\text{m}$ and $z_0^+ = 0.15 \mu\text{m}$. As expected, the fluctuations of the particle-substrate distance z for a colloid opposite to an attractive stripe [$x = 0$, panel (a)] decrease significantly upon increasing the correlation length ξ , due to the fact that the particle is mainly localized around the deep potential well which forms as a consequence of the action of the critical Casimir force. In fact, $z_{\text{upp,avg,low,min}} \simeq z_0^-$ for large ξ . The actual value of z_0^- strongly affects the behavior of the particle, both in determining the values of $z_{\text{upp,avg,low,min}}$ close to the critical point and in setting the threshold value ξ^* of ξ at which one observes such a sharp transition towards strong spatial confinement for $\xi > \xi^*$. From panel (a) one infers, e.g., that $\xi^* \simeq 17 \text{nm}$ and $\xi^* \simeq 27 \text{nm}$ for the solid and dashed curves, respectively. On the other hand, the dependence on z_0^+ is not pronounced and indeed in panel (a) the dotted lines practically coincide with the solid ones. For a colloid opposite to a repulsive stripe [$x = P/2$, panel (b)], due to the repulsive nature of both the electrostatic and the critical Casimir force and the very weak gravitational attraction one has $z_{\text{min}} < z_{\text{low}}$ and the average position is typically of the order of $1 \mu\text{m}$ with fluctuations of the same order. As anticipated, also in this case the actual value of z_0^+ is not very relevant, in particular for large values of ξ . Analogously, the behavior of the colloid at $x = P/2$ is not affected by the choice of z_0^- and in panel (b) the dashed lines practically coincide with the solid ones.

$z_{\text{upp,avg,low,min}}$ and in setting the threshold value ξ^* of ξ above which the particle becomes strongly confined, whereas the dependence on z_0^+ is negligible for the behavior at $x = 0$, because z_0^+ controls the electrostatic interaction with the adjacent stripe. Analogously, the behavior of a colloid at $x = P/2$ as shown in Fig. 2(b), i.e., opposite to a repulsive stripe is not affected by the choice of z_0^- . However, for this configuration, also the actual value of z_0^+ does not affect significantly the resulting behavior of the particle-substrate distance at $x = P/2$, in particular for large values of ξ . Indeed, due to the repulsive nature of both the electrostatic and the critical Casimir force and the weak gravitational attraction, the average position is typically of the order of $z_{\text{avg}}(x = P/2) \simeq z_0^+ + k_B T/G \simeq k_B T/G$ (with an additional linear contribution $\propto \xi$ for large values of ξ) allowing fluctuations of the order of $k_B T/G \simeq 1 \mu\text{m}$.

In Fig. 3 the particle-substrate distance (characterized via $z_{\text{avg,upp,low}}$) and the position of mechanical equilibrium z_{min} are reported as functions of the lateral coordinate of the colloid within the period P and for two different temperatures, i.e., two different values of ξ . In Fig. 3, the choice of the parameters corresponds to actually experimentally accessible values. Figure 3 clearly shows that for $\xi = 10 \text{nm}$ (dashed lines) the particle-substrate distance is laterally constant and the colloid does not react to the presence of the chemical pattern on the substrate, apart for a possible effect due to a change in the electrostatic interaction (which here is taken to be the same on the different stripes). On the other hand, for $\xi = 20 \text{nm}$ (solid line) the colloid is strongly attracted to that part of the substrate with the same preferential adsorption (in Fig. 3 indicated by a shaded background), as a consequence of the emerging critical Casimir forces. Indeed for $|x| \lesssim L_-/2$ the particle is abruptly localized at a distance $z \simeq z_0^-$, which is primarily set by the electrostatic repul-




Figure 3. Particle-substrate distance z (characterized via $z_{\text{upp,avg,low}}$, see the main text) and position of mechanical equilibrium z_{min} as functions of the lateral coordinate of the particle for $\xi = 10\text{nm}$ (dashed curves) and $\xi = 20\text{nm}$ (solid curves). Here, $z_0^- = z_0^+ = 0.09\mu\text{m}$, $P = 1.8\mu\text{m}$, $R = 1.2\mu\text{m}$, and $L_- = 0.9\mu\text{m}$. For small values of the correlation length ξ the various characteristic distances are almost independent of the lateral coordinate, whereas above a certain threshold value ξ^* of ξ (see also Fig. 2) the colloid opposite to the attractive stripe is strongly confined close to the wall at a distance $z \simeq z_0^-$ with almost no fluctuations.

sion and which corresponds to the position $z_{\text{min}}(0)$ of the minimum of the potential, with almost no thermal fluctuations. In the region above the repulsive stripes (in Fig. 3 indicated by a white background) and within the temperature range considered here, instead, the repulsive critical Casimir force pushes the colloid only slightly further away from the substrate, by a distance of the order of the increasing correlation length ξ . But the amplitude of the thermal fluctuations of the particle-substrate distance are barely affected by the onset of the repulsive critical Casimir force.

The analysis of the particle-substrate distance shows that the behavior of the colloidal particle and the resulting potential are drastically influenced by the strong attraction of the colloid close to an attractive stripe. Accordingly, the electrostatic repulsion from this stripe, i.e., the value of z_0^- , affects significantly the total potential and has therefore to be considered carefully in the comparison between theoretical predictions with the experimental data. On the contrary, the actual value of z_0^+ as well as the actual value of κ do not significantly affect the resulting behavior of the potentials as long as they are within the range appropriate for the present experiment which can be inferred from previous, independent measurements [21].

3. Experimental setup

In the present context of critical Casimir forces, surfaces with a periodic pattern of alternating stripes with opposite adsorption preferences for the two components of the binary mixture of water and lutidine were obtained by micro-contact printing (μCP) of alkane-thiols. Since there is a vast literature covering this technique [42–44] here we shall describe it only briefly. Stamps were prepared by casting poly(dimethylsiloxane) (PMDS) onto a master which was topographically structured by a lithographic process. After the PMDS is cured one obtains an elastic stamp which exhibits the negative structure of the master. In the next step the stamp is wetted with a 1mM ethanolic solution of nonpolar $\text{HS}(\text{CH}_2)_{17} - \text{CH}_3$ thiol. When the excess liquid is removed, the stamp is brought into mechanical contact for several seconds with a glass substrate coated by a 30nm layer of gold. This results in hydrophobic regions the geometry of which corresponds to the protruding parts of the stamp. Finally, the substrate is dipped into a 1mM ethanolic solution of polar $\text{HS}(\text{CH}_2)_{11} - \text{OH}$ thiol for several minutes in order to render the remaining bare

1 gold surface hydrophilic. With this technique we were able to produce periodic arrays of
2 hydrophilic and hydrophobic stripes of widths between $0.9\mu\text{m}$ and $3\mu\text{m}$ over a typical
3 total extension of 0.5cm^2 . The typical lateral edge resolution of the chemical structures
4 is of the order of $50 - 100\text{nm}$. In previous studies [14] the chemical patterns were cre-
5 ated by removing locally a hydrophobic monolayer from a hydrophilic surfaces with a
6 focused ion beam (FIB). Although under ideal conditions the resolution of FIB is within
7 the range of several nanometers, in the case of non-conducting surfaces we observed dis-
8 tortions/deflections of the ion beam due to an electrostatic charging of the glass surface,
9 leading to deviations of the chemical steps from straight lines. This causes considerable
10 discrepancies between the measured critical Casimir forces acting on particles near such
11 surfaces and the corresponding theoretical predictions, which assume straight and sharp
12 boundaries for the stripes [15]. Since such charging effects are basically absent in μCP ,
13 here we have an improved control of the geometrical structure of the imprinted chemical
14 pattern and therefore much better agreement with theory (see below).
15

16
17 In order to probe the critical Casimir forces which act on a colloidal particle exposed
18 to the patterned substrates prepared as described above, we used polystyrene (PS) probe
19 particles with radius $R = 1.2\mu\text{m}$. Their surface charge, as provided from the manufacturer,
20 is $10\mu\text{C}/\text{cm}^2$ which renders them hydrophilic, realizing $(-)$ boundary conditions. The
21 particles were dissolved in a critical water-2,6-lutidine mixture which has a lower critical
22 demixing point at a lutidine mass fraction of $c_L^c \cong 0.286$ and a critical temperature of $T_c =$
23 307K . Particle positions were monitored by digital video microscopy which allows one
24 to track the projection of their centers onto the substrate plane with a spatial resolution of
25 about 50nm . For each measurement the particle trajectories were recorded for 40 minutes
26 with a frequency of 4 frames per second. Within the field of view of $290\mu\text{m} \times 225\mu\text{m}$
27 around 20 to 40 particles have been monitored, corresponding to a projected density of one
28 particle per square of the size of 25×25 to 17×17 particle diameters. Thus, the particle
29 density was sufficiently small to exclude the presence of critical Casimir forces among
30 neighboring colloids, such that only their interaction with the patterned substrate is probed
31 during the measurements.
32

33 The temperature of the sample was controlled by the procedure described in detail in
34 Ref. [14]. With this setup, temperatures close to T_c could be stabilized within 10mK
35 over several hours. In contrast to temperature *changes*, which could be resolved within
36 mK accuracy, larger errors in the determination of *absolute* temperatures (in particular the
37 measured critical temperature) occur. This is due to the fact, that the metal-resistance ther-
38 mometer (Pt100) could not be placed within the sample cell but it was instead attached
39 outside but close to the field of view. Indeed, the thermometer measures a temperature
40 T^{out} lower than the actual temperature T inside the sample cell. We associate the temper-
41 ature $T^{\text{out}} = T_c^{\text{out}}$ with the critical temperature $T = T_c^{\text{exp}}$ of the solvent in the sample at
42 that particular temperature for which critical opalescence is observed when shining a laser
43 beam into the sample cell. This leads to systematic errors on the absolute temperature of
44 T_c^{out} and T_c^{exp} of the order of 50mK . Using the assumption that $T - T^{\text{out}} = \text{const}$ (this
45 constant could not be determined and is inter alia dependent on the ambient temperature),
46 we have measured the temperature difference $\Delta T = T_c^{\text{out}} - T^{\text{out}}$ and identified it with
47 $T_c^{\text{exp}} - T \equiv \Delta T$. Since the comparison with the theoretical predictions depends crucially
48 on the actual value $T_c \simeq 307\text{K}$ of the critical temperature, in the analysis below we account
49 for such a possible systematic error by considering $\Delta T_c^* = T_c - T_c^{\text{exp}}$ as an additional fit-
50 ting parameter, where $T_c \simeq 307\text{K}$ is the actual critical temperature of the water-lutidine
51 mixture.
52
53
54
55
56
57
58
59
60


4. Comparison of theory with experiments

The total potential Φ of the forces [Eq. (11)] acting on the colloidal particle has been calculated on the basis of the Derjaguin approximation with $k_{(\pm,-)}$ [see Eqs. (1) and (6)] obtained from Monte Carlo simulations. In the following we shall use for the scaling functions $k_{(\pm,-)}$ of the critical Casimir force between two planar walls with $(\pm,-)$ BC the numerical estimate referred to as “approximation (i)” in Figs. 9 and 10 of Ref. [34]. We have checked that different [34] or more recent and accurate [35] estimates for $k_{(\pm,-)}$ actually lead to essentially the same effective potentials, the only difference being a small additional overall shift of the resulting fitted value of the critical temperature (see further below). A detailed analysis in spatial dimension $d = 4$ suggests that the Derjaguin approximation we have used in our theoretical predictions for $d = 3$ should be rather accurate in describing the actual potential of the colloid within the range of parameters experimentally studied here [16]. As anticipated above, in order to predict the effective potential \hat{V} we need to fix also the value of the parameters which determine the electrostatic interaction. For the comparison between theory and experiment we fix the screening length κ^{-1} to the value $\kappa^{-1} = 12\text{nm}$ which has been reported from independent measurements on the same system (see, e.g., Ref. [21]). In order to fit our theoretical predictions to the experimental data, we vary instead the unknown values of the parameters z_0^\pm [Eqs. (7) and (8)] within the range $0.08 - 0.15\mu\text{m}$, which can be reasonably expected on the basis of previous measurements on homogeneous substrates [21]. However, the results for $\delta\hat{V}(0)$ are hardly affected by the particular choice of z_0^+ so that in the following we keep it fixed at $z_0^+ = 0.09\mu\text{m}$. The amplitude ξ_0^+ of the correlation length has been determined by independent experiments as $\xi_0^+ \simeq 0.20 \pm 0.02\text{nm}$ (see, e.g., Tab. III in Ref. [21]) so that we vary ξ_0^+ within the range $0.18 - 0.22\text{nm}$ in order to obtain the best fit to the experimental data by the theoretical scaling functions. Moreover, as mentioned above, the experimental uncertainty in the determination of the absolute value of the critical temperature T_c^{exp} is taken into account by considering as an additional fitting parameter the shift $\Delta T_c^* = T_c - T_c^{\text{exp}}$ (see the end of Sec. 3) of up to $|\Delta T_c^*| \simeq 100\text{mK}$. (The values of ΔT_c^* may be different for the individual stripe widths L_- which characterize the substrates investigated in independent experimental runs.) However, the *relative* uncertainty in the determination of the temperature within a single experimental run for a given stripe width L_- is smaller than $\pm 10\text{mK}$ (see above). Thus, for the comparison carried out below, we are left with ΔT_c^* , z_0^- , ξ_0^+ , and Δx (see Subsec. 2.5) as fitting parameters which, however, are all limited to rather small ranges of values in order to be in accordance with independent and previous experimental results.

4.1. Depth of the potential

The depth of the measured laterally confining potential is given by $\Delta\hat{V} = -\delta\hat{V}(0) > 0$ [see Eq. (12)]. Figure 4(a) shows $\Delta\hat{V}$ as a function of the deviation $\Delta T = T_c^{\text{exp}} - T$ of the temperature from the experimentally determined critical demixing temperature T_c^{exp} , for various stripe widths L_- and measured in independent experimental runs as described above. The periodicities P of the various patterns the stripes belong to are determined from the photolithography mask and are given in Tab. 1. The potential depth $\Delta\hat{V}$ is determined as the difference of the potential between its value at the center of a repulsive stripe and at the center of an attractive stripe, carrying an uncertainty of $\pm 0.15k_B T$ near the critical temperature $T_c \simeq 307\text{K}$. Upon approaching T_c , for a stripe width $L_- \lesssim 1.5\mu\text{m}$ the effects of two adjacent chemical steps interfere, which results in an effectively reduced potential depth compared to the case of a very wide stripe (see the dashed curves in Fig. 4(b); in Fig. 4(a) the experimental uncertainty in determining the critical temperature T_c^{exp} is responsible for the relative displacements among the various curves).

Figure 4. Depth $\Delta\hat{V}$ of the effective potentials of a colloid close to patterned substrates, with periods $P = L_- + L_+$ and $(-)$ stripe widths L_- given in Tab. 1, as a function of the temperature deviation from the critical temperature. The symbols represent the experimental data which are affected by an experimental uncertainty of $\pm 0.15k_B T$ for the potential depth and of $\pm 0.01\text{K}$ for the temperature differences between the various data points belonging to one and the same value for L_- .



(a) $\Delta\hat{V}$ as a function of $\Delta T = T_c^{\text{exp}} - T$. (For better visibility, the data for $L_- = 3.25\mu\text{m}$ are shifted by 0.1K along the ΔT axis, as indicated by the red arrow.) The solid lines represent the best fit of the theory to the experimental data for each *individual* value of L_- with fixed parameters κ and z_0^+ which do not affect the resulting behavior significantly as discussed in the main text. ξ_0^+ has been varied within the reasonable range $0.18\text{nm} \lesssim \xi_0^+ \lesssim 0.22\text{nm}$ known from the literature. The values of all parameters corresponding to the solid lines are given in Tab. 1. Clearly, for each individual L_- the actual critical temperature T_c is shifted by $\Delta T_c^* = T_c - T_c^{\text{exp}}$ from the value T_c^{exp} determined experimentally, with ΔT_c^* being within the experimental accuracy $|\Delta T_c^*| \lesssim 0.1\text{K}$. The dashed curves represent a *common* fit to all data. For this latter fit we have assumed ξ_0^+ and z_0^- to be the same for the various independent experimental runs, independently of the corresponding value of L_- , whereas $\Delta T_c^* = T_c - T_c^{\text{exp}}$ (besides Δx as determined below) is used for adjustment to the data for each individual L_- . The best fit for all these parameters renders $\xi_0^+ = 0.22\text{nm}$, $z_0^- = 0.11\mu\text{m}$, and the values reported in Tab. 2.

(b) $\Delta\hat{V}$ as a function of $\Delta T + \Delta T_c^* = T_c - T$ using the values of ΔT_c^* for the best common fit (see dashed lines in (a)) given in Tab. 2. This plot takes into account the experimental uncertainty of measuring the absolute value of T_c^{exp} . The data points, which have been shifted accordingly, basically fall on top of each other within their error bars. The dashed lines correspond to the dashed ones shown in (a) and represent the theoretical predictions. For $L_- \geq 1.60\mu\text{m}$ the various curves are almost indistinguishable from each other, whereas for $L_- \leq 1.30\mu\text{m}$ the potential depth is effectively reduced due to the interference of the effects of two neighboring steps and due to the fact that the steps are non-ideal.

The effect of non-ideal chemical steps may result in an effectively reduced potential depth as well (see, for example, Fig. 4(b) and, c.f., Fig. 6(b)). Accordingly, in comparing the experimental data for $\Delta\hat{V}$ with the theoretical predictions we allow for an uncertainty in the local position of the boundary between adjacent stripes, as described in Subsec. 2.5. However, as will be shown below, we are able to determine Δx rather precisely from the lateral variation of the potential. The values of Δx , which yield the best agreement and which are used for the comparison shown in Fig. 4, are given in Tab. 1. On the basis of our theoretical analysis, it turns out that the temperature dependence of the potential depth becomes independent of the stripe width when the latter is sufficiently large. Adopting for the geometrical and physical parameters the values corresponding to the present experimental conditions, this is expected to occur for $L_- \gtrsim 2\mu\text{m}$. From Fig. 4 one can see, however, that the experimentally determined data for individual experimental runs are shifted along the temperature axis with respect to each other, which reflects the uncertainty of up to 100mK in measuring the absolute value of the critical temperature. On the other hand, within a single individual experimental run corresponding to a certain stripe width, the temperature difference between the various data points can be measured with the high accuracy of less than 10mK. Upon comparing the experimental data shown as symbols in Fig. 4 with our theoretical predictions we take this into account by introducing, for each value of L_- , a possible shift $\Delta T_c^*(L_-) = T_c - T_c^{\text{exp}}(L_-)$ between the actual critical temperature T_c and the value T_c^{exp} determined in that particular experiment. Accordingly, for the data corresponding to a certain L_- , the actual distance from the critical

Table 1. Values of the parameters for which best agreement is obtained between theory and the data for $\delta\hat{V}(x)$ for each individual experimental run, corresponding to a single value of L_- (see Fig. 4(a)). The values $\kappa^{-1} = 12\text{nm}$ and $z_0^+ = 0.09\mu\text{m}$ are fixed because their choice does not affect significantly the resulting theoretical curves in Fig. 4.

$P[\mu\text{m}]^a$	$L_-^{\text{exp}}[\mu\text{m}]^b$	$L_-[\mu\text{m}]^c$	$\xi_0^+[\text{nm}]$	$z_0^-[\mu\text{m}]$	$\Delta T_c^*[\text{mK}]$	$\Delta x[\mu\text{m}]^d$
6.0	3.0	3.25	0.22	0.103	86	0.15
5.4	2.7	2.25	0.21	0.128	-5	0.10
4.2	2.1	1.60	0.22	0.095	88	0.22
3.6	1.8	1.30	0.19	0.140	-18	0.19
1.8	0.9	0.90	0.20	0.121	-28	0.09

^aMost of the experimental data for the potential $\delta\hat{V}(x)$ discussed here are not influenced by the actual value of P as long as $P \gtrsim 2L_-$. Accordingly, we assume the periodicity to be the one determined by the inscribed photolithographic mask pattern reported here.

^bWidth of the stripes of the photolithographic mask pattern. Due to the μCP stamping process we estimate the uncertainty of the actual width of the thiol stripes to be up to $\pm 0.5\mu\text{m}$.

^cValue of the stripe width for which best agreement between theory and experiment is obtained.

^dValue of Δx for which best agreement between theory and experiment is obtained (see Figs. 5 and 6).

Table 2. Values of the parameters for which best agreement is obtained between theory and all experimental data together (see Fig. 4(b)), so that the values of ξ_0^+ and z_0^- are the same, and ΔT_c^* (in addition to Δx determined from the shape of the potentials; see below) is the only parameter allowed to vary for the various stripe widths L_- . As in Tab. 1, the values of $\kappa^{-1} = 12\text{nm}$ and $z_0^+ = 0.09\mu\text{m}$ are fixed. (For a description of the parameter values P , L_-^{exp} , and L_- see the footnotes in Tab. 1.)

$P[\mu\text{m}]^a$	$L_-^{\text{exp}}[\mu\text{m}]$	$L_-[\mu\text{m}]$	$\xi_0^+[\text{nm}]$	$z_0^-[\mu\text{m}]$	$\Delta T_c^*[\text{mK}]$	$\Delta x[\mu\text{m}]$
6.0	3.0	3.25	0.22	0.110	49	0.15
5.4	2.7	2.25	0.22	0.110	54	0.10
4.2	2.1	1.60	0.22	0.110	0	0.22
3.6	1.8	1.30	0.22	0.110	72	0.19
1.8	0.9	0.90	0.22	0.110	29	0.09

point is given by $T_c - T = \Delta T + \Delta T_c^*(L_-)$.


The dashed curves in Fig. 4 correspond to the *common* fit to all experimental data which leads to the fitting parameters ξ_0^+ and z_0^- taking the same values $\xi_0^+ = 0.22\text{nm}$ and $z_0^- = 0.11\mu\text{m}$ for all L_- as obtained from the least-square method. On the other hand, the individual temperature shifts $\Delta T_c^*(L_-)$ are given in Tab. 2 for each data set corresponding to a single value of L_- . The values of ξ_0^+ and z_0^- are both in agreement with independent previous findings [21]. Figure 4(b) shows the depth of the potentials as a function of $\Delta T + \Delta T_c^*(L_-) = T_c - T$, i.e., shifted along the temperature axis by $\Delta T_c^*(L_-)$ as given in Tab. 2. This accounts for the uncertainty in determining experimentally the absolute value of the temperature (see the discussion at the end of Sec. 3), and indeed, as expected for significantly wide stripes $L_- \gg \xi$, the data overlap with each other within the error bars, reflecting that in this limit $\Delta\hat{V}$ is independent of L_- .

The solid lines in Fig. 4(a) correspond to fits of the theoretical predictions to each *individual* experiment dealing with a specific stripe width. Distinct from the previous common fit, here for each stripe width L_- we allow for a variation of the values of ξ_0^+ and z_0^- in addition to ΔT_c and Δx . On the basis of a least-square fit, best agreement is obtained for the values given in Tab. 1, which agree with those obtained in independent previous studies [21] within the corresponding experimental accuracy.

4.2. Shape of the potentials

Figures 5 and 6 show the total potential of the forces acting on the colloid as a function of its lateral position for various temperatures and for two stripe widths. Symbols represent the experimental data, whereas the solid and dashed lines are the corresponding theoretical

Figure 5. Lateral variation of the effective potential $\delta\hat{V}(x)$ (see Eq. (12)) for a colloid opposite to a chemically patterned substrate and immersed in the water-lutidine binary liquid mixture at its critical concentration for various temperatures $T_c^{\text{exp}} - \Delta T$. Symbols indicate experimental data, whereas the lines are the corresponding theoretical predictions for ideal (dashed lines) and non-ideal (solid lines) stripe patterns. The parameters used to calculate the theoretical curves are given in Tab. 1.




(a) $L_- = 3.25\mu\text{m}$. From top to bottom the critical point is approached and the measured temperature deviations ΔT are 0.3, 0.18, 0.15 (0.153), 0.145 (0.14), 0.13 (0.128), 0.12, 0.11 (0.113), and 0.10 K with an uncertainty of up to $\pm 0.01\text{K}$ with respect to each other. If indicated, the values in brackets are corrected values (but compatible within the experimental inaccuracy) which have been used for evaluating the theoretical predictions.

(b) Same as (a) for $L_- = 2.25\mu\text{m}$ and $\Delta T = 0.175$ (0.165), 0.16 (0.152), 0.145 (0.143), 0.13, 0.115, 0.10 K.

predictions with ξ_0^+ , z_0^- , and ΔT_c^* fixed to the values reported in Tab. 1, which have been determined from the fit of the depth $\Delta\hat{V}$ of the potential. In addition, for a few cases and as indicated in the figure captions, we use the leeway provided by the experimental uncertainty of $\pm 10\text{mK}$ for the temperature value. The dashed lines in Figs. 5 and 6 are based on the assumption that the stripe patterns are *ideal* whereas the solid ones refer to *non-ideal* patterns for which we fitted the parameters Δx (see Subsec. 2.5) in order to obtain the best agreement between theoretical predictions and the set of experimental data at distinct temperatures. The resulting values of Δx for the various widths of the pattern are reported in Tab. 1. In fact, the broadness of the transition regions across the chemical steps of $\delta\hat{V}(x)$ between its extremal values is affected practically exclusively by Δx . We have checked that the other parameters and potential additional effects such as polydispersity and weak adsorption cannot account for the discrepancies of the slopes of the dashed curves and the experimentally determined ones (see also Ref. [15]). Thus, Δx and L_- are determined de facto independently of the choice of ξ_0^+ , z_0^- , and ΔT_c^* and in the following they can be regarded fixed upon variation of the values of these latter parameters. In particular, this is important for narrow stripes as shown in Fig. 6, where even the depth of the potential depends on L_- and Δx , in contrast to broader stripes.

As anticipated, in the case of rather large stripe widths $L_- = 3.25\mu\text{m}$ and $L_- = 2.25\mu\text{m}$ shown in Fig. 5 and for the temperatures considered here the effects of two neighboring chemical steps do not interfere and the resulting potential across the center of the attractive chemical stripe is flat. Accordingly, from the comparison between the experimental data for the shapes of the potentials and the corresponding theoretical prediction we can infer the width L_- of the stripes — now treated as a fitting parameter — independently of its initially assumed value L_-^{exp} determined by the width of the photolithography mask. Indeed, the actual width of the stripe imprinted by the mask may differ by up to $\pm 0.5\mu\text{m}$ from L_-^{exp} characterizing the mask itself. It is only L_-^{exp} which is controlled during the preparation process [41]. On the other hand, the actual value of P is the same as the one of the photolithography mask because the periodicity is not affected by the stamping

Figure 6. Effective potential $\delta\hat{V}(x)$ (see Eq. (12)) for various temperatures $T_c^{\text{exp}} - \Delta T$ and relatively small widths of the attractive stripes. Symbols indicate experimental data, whereas the lines are the corresponding theoretical predictions for ideal (dashed lines) and non-ideal (solid lines) stripe patterns. The parameters used to calculate the theoretical curves are given in Tab. 1.



(a) Same as Fig. 5(a) for $L_- = 1.60\mu\text{m}$ and $\Delta T = 0.3, 0.23$ (0.218), 0.21 (0.198), 0.19 (0.182), 0.17, 0.15 (0.153) K.

(b) Same as (a) for $L_- = 1.30\mu\text{m}$ (left panel) and $L_- = 0.9\mu\text{m}$ (right panel) with $\Delta T = 0.3, 0.12, 0.1, 0.09$ (0.083), 0.08 K for $L_- = 1.30\mu\text{m}$ and $\Delta T = 0.3, 0.18, 0.16, 0.145, 0.13$ K for $L_- = 0.9\mu\text{m}$.

process. (Note that, within the parameter range investigated in Fig. 5, the effects of neighboring chemical steps do not influence each other and therefore the actual value of the periodicity P of the pattern does not affect the determination of L_- .) In agreement with this observation, the comparison with the theoretical predictions leads to fitted values L_- which are within the aforementioned range indicated in Tab. 1. From Fig. 5 we find that the experimental techniques described above lead to rather sharp chemical steps between two stripes with an uncertainty of only $\Delta x \leq 0.15\mu\text{m}$.

The comparison for narrower stripe widths $L_- \lesssim 2\mu\text{m}$ is shown in Fig. 6. Even for these cases, in which the effects of two neighboring chemical steps do interfere, the experimental data agree very well with the theoretical predictions obtained from the Derjaguin approximation. Whereas for $L_- = 1.6\mu\text{m}$ and $L_- = 1.3\mu\text{m}$ we have to take into account an uncertainty Δx of the position and associated with the shape of the chemical steps of about $0.2\mu\text{m}$, for the smallest stripe width $L_- = 0.9\mu\text{m}$ the chemical pattern turns out to be almost ideal with $\Delta x < 0.1\mu\text{m}$ (see Tab. 1).

5. Conclusions and Outlook

5.1. Summary

We have presented a detailed theoretical and experimental study of the effective solvent-mediated forces acting on colloids suspended in a near-critical binary liquid mixture of water and 2.6-lutidine and close to chemically patterned substrates. In contrast to earlier investigations [14] we have obtained a chemical pattern of stripes of different widths and with laterally alternating adsorption preferences for either lutidine [(+) boundary condition] or water [(-) boundary condition] by monolayers of thiols deposited on gold-coated substrates via microcontact printing. The solute colloidal particles — $2.4\mu\text{m}$ diameter polystyrene spheres — were rendered hydrophilic, corresponding to (-) boundary conditions, due to their surface charge. Based on digital video microscopy our main experimental findings are the following.

1. At the fixed critical composition of the water-lutidine mixture, upon raising the temperature T towards its lower critical demixing point at $T = T_c$, lateral forces acting on

1 the colloidal particles arise gradually. These critical Casimir forces cause the colloidal
2 particles to be attracted to those chemical stripes which have the same adsorption pref-
3 erence as the particles and to be repelled from those stripes with the opposite adsorption
4 preference.

5
6 2. These lateral and normal contributions to the total effective forces (compare Sub-
7 sec. 2.2) are negligible at temperatures T more than a few hundred mK away from the
8 critical value T_c but they increase significantly upon approaching T_c . Eventually, they
9 lead to laterally confining potentials for the colloidal particles with potential depths of
10 several $k_B T$ (see Fig. 4). These potentials can be reversibly tuned by minute temperature
11 changes.

12 3. Across a chemical step the critical Casimir potentials vary rather abruptly between
13 two plateau values on a length scale of $\approx 0.8\mu\text{m}$ (see Fig. 5). This indicates that the
14 microscopic structures of the chemical steps formed by imprinted layers of alkanethiols
15 with different endgroups are much sharper than those created previously by a focused
16 ion beam [14].

17
18 4. For rather narrow stripes of widths $\lesssim 2\mu\text{m}$ the effects of two neighboring chemical
19 steps interfere and consequently reduce the potential depth and lead to rounded shapes
20 of the effective potentials (see Fig. 6).

21
22 These experimental observations can be consistently interpreted in terms of the occurrence
23 of the critical Casimir effect, and it is possible to quantitatively compare the measured
24 potentials with the corresponding theoretical predictions. We have derived the effective
25 potentials within the Derjaguin approximation (see Subsec. 2.1 and Ref. [16]) based on
26 universal scaling functions for the critical Casimir force between two plates with (\pm, \pm) or
27 $(+, -)$ boundary conditions as obtained from Monte Carlo simulation data. (Differences
28 in estimates for these scaling functions as obtained from various Monte Carlo simulations
29 do not affect our results significantly.) In contrast to the present experimental measure-
30 ments we are able to theoretically analyze the spatially fully resolved critical Casimir
31 potential acting on a colloid. The resulting energy landscape for a colloidal particle is
32 rather rich due to the interplay of several relevant forces, which we have taken into ac-
33 count (see Fig. 1). Typically, thermal fluctuations lead to large fluctuations of the lateral
34 position and the distance between the colloidal particles and the substrate (see Fig. 2).
35 Upon approaching T_c , however, strong normal and lateral critical Casimir forces abruptly
36 localize the colloids very close to and above stripes exhibiting the same adsorption pref-
37 erence as the particles (see Fig. 3). Calculating the effectively one-dimensional, projected
38 potentials for a single colloid, as they are obtained experimentally, we also take into ac-
39 count the possibility that the chemical pattern is not ideal, in addition to other experimen-
40 tal uncertainties. Not only the theoretical predictions for the potential depths agree with
41 the experimental data (see Fig. 4) but also the correlation length ξ , as determined from
42 our comparison, follows rather well the expected universal power law behavior and the
43 associated non-universal amplitude ξ_0 is in agreement with previous independent experi-
44 mental determinations (see Tab. 1). Moreover, the shapes of the potentials as a function of
45 the lateral position of the colloid show good agreement between the theory and the exper-
46 iments (see Figs. 5 and 6). Since critical Casimir forces probe rather sensitively the details
47 of the geometry of the patterns [15], we find that the chemical steps obtained by the ex-
48 perimental method used here are sharp, with deviations of only a few percent in terms of
49 the stripe width. From the detailed comparison with rather narrow chemical stripes (see
50 Fig. 6(b)) we infer that even for such cases the Derjaguin approximation describes the
51 actual behavior quite well, as expected from a corresponding theoretical analysis within
52 mean-field theory [16]. We conclude that the quantitatively successful comparison be-
53 tween the experimental data and the theoretical predictions demonstrates the significance
54 and reliability of the critical Casimir effect for colloidal suspensions and reveals a new
55 means of using them as model systems in soft-matter physics or in applications in inte-
56
57
58
59
60

1 grated micro- or nano-devices.

2 3 4 5 **5.2. Perspectives**

6 *“After many failures, involving many fluid samples, I did succeed in levitating a hard*
7 *platelet.”*

8 — Robert Evans.

9
10 Since quantum-electrodynamic Casimir forces acting on the scale of nm and μm are typ-
11 ically always attractive, they are responsible for stiction occurring in micro- and nano-
12 mechanical devices. According to a generalization of the Earnshaw theorem, for quan-
13 tum Casimir forces acting on conventional materials in vacuum, stable levitation is not
14 possible [45]. Therefore, in order to overcome the problem of stiction using the quantum-
15 electrodynamic Casimir effect the system must be immersed in a fluid and the bodies have
16 to be designed very specifically [46, 47]. Obviously however, for a system immersed in
17 a fluid it is natural to use solvent mediated forces such as the critical Casimir forces to
18 control its behavior. Since one can obtain attractive and repulsive *critical* Casimir forces
19 by suitable surface treatments as discussed here, a properly chosen combination of chem-
20 ical substrate patterns may easily lead to levitation of a colloidal particle at a stable height
21 above the substrate [16]. One finds that this critical Casimir levitation not only can be
22 achieved without involving any other additional force but that the stable equilibrium dis-
23 tance between the wall and the colloid can be tuned steeply by varying the temperature
24 [16]. Recently [16] it was predicted that the ranges of geometrical parameters for a pattern
25 of laterally alternating stripes (as discussed here), which allow for critical Casimir levi-
26 tation of colloidal particles, are rather large. Moreover, critical Casimir levitation is robust
27 even in the presence of other forces and of thermal fluctuations [16] as they occur typ-
28 ically in colloidal suspensions, so that an experimental realization of this effect appears
29 to be possible. These results show, that chemical patterning of substrates allows one to
30 design the critical Casimir effect over a wide range of properties.

31
32
33 *“Ions are the invention of the devil.”*

34 — Robert Evans.

35
36 By adding salt to the solvent of the colloidal suspension the electrostatic repulsion be-
37 tween the substrate and a colloid is strongly screened; this provides the possibility to ex-
38 plore the spatial variation of critical Casimir forces over a much wider range of distances
39 and correlation lengths of the fluid. Moreover, a rather complex behavior of the resulting
40 forces acting on the colloid can be expected because electrostatic forces induced by ions
41 lead to rather delicate physical features. Recently, preliminary measurements have shown
42 a rich and novel behavior of a colloid immersed in a salt-rich solvent upon approaching
43 the critical point [48]. The onset of strong attractive forces even several K below the crit-
44 ical temperature is observed and, in addition, these forces are found to remain attractive,
45 independent of the boundary conditions, (\pm, \pm) or $(+, -)$, throughout a large tempera-
46 ture range. Currently it is not clear whether these observations can be entirely attributed
47 to critical Casimir forces or whether, e.g., the coupling between electrostatic interactions
48 and critical fluctuations has to be considered. Thus, using these intriguing effects together
49 with a chemical patterning of substrates may open a new field of phenomena with poten-
50 tial applications in colloid science.

51 52 53 54 55 **Acknowledgement**

56
57 It is a great pleasure for the authors to dedicate this paper to Bob Evans on the occasion of
58 his 65th birthday in view of his long lasting inspiration and encouragement across many
59
60

1 fields of statistical physics.
2
3
4

5 References

- 6 [1] M.E. Fisher and P.G. de Gennes, C. R. Acad. Sci., Paris, Ser. B **287**, 207 (1978).
7 [2] H.G.B. Casimir, Proc. K. Ned. Akad. Wet. **51**, 793 (1948).
8 [3] M. Kardar and R. Golestanian, Rev. Mod. Phys. **71**, 1233 (1999).
9 [4] M. Krech, *The Casimir Effect in Critical Systems* (World Scientific, Singapore, 1994).
10 [5] J.G. Brankov, D.M. Danchev and N.S. Tonchev, *Theory of critical phenomena in finite size systems* (World Scientific,
11 Singapore, 2000).
12 [6] A. Gambassi, J. Phys.: Conf. Ser. **161**, 012037 (2009).
13 [7] R. Garcia and M.H.W. Chan, Phys. Rev. Lett. **83**, 1187 (1999); *ibid* **88**, 086101 (2002).
14 [8] A. Ganshin, S. Scheidemantel, R. Garcia and M.H.W. Chan, Phys. Rev. Lett. **97**, 075301 (2006).
15 [9] M. Fukuto, Y.F. Yano and P.S. Pershan, Phys. Rev. Lett. **94**, 135702 (2005).
16 [10] C. Hertlein, L. Helden, A. Gambassi, S. Dietrich and C. Bechinger, Nature **451**, 172 (2008).
17 [11] M. Tröndle, L. Harnau and S. Dietrich, J. Chem. Phys. **129**, 124716 (2008).
18 [12] S. Kondrat, L. Harnau and S. Dietrich, J. Chem. Phys. **131**, 204902 (2009).
19 [13] M. Sprenger, F. Schlesener and S. Dietrich, J. Chem. Phys. **124**, 134703 (2006).
20 [14] F. Soyka, O. Zvyagol'skaya, C. Hertlein, L. Helden and C. Bechinger, Phys. Rev. Lett. **101**, 208301 (2008).
21 [15] M. Tröndle, S. Kondrat, A. Gambassi, L. Harnau and S. Dietrich, EPL **88**, 40004 (2009).
22 [16] M. Tröndle, S. Kondrat, A. Gambassi, L. Harnau and S. Dietrich, J. Chem. Phys. **133**, 074702 (2010).
23 [17] M. Krech and S. Dietrich, Phys. Rev. Lett. **66**, 345 (1991); Phys. Rev. A **46**, 1886 (1992); *ibid* 1922 (1992).
24 [18] K. Binder, Critical Behavior at Surfaces, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and J. L.
25 Lebowitz, Vol. 8 (Academic, London, 1983), p. 1.
26 [19] H.W. Diehl, Field-theoretical Approach to Critical Behaviour at Surfaces, in *Phase Transitions and Critical Phenom-*
27 *ena*, edited by C. Domb and J. L. Lebowitz, Vol. 10 (Academic, London, 1986), p. 75.
28 [20] H.W. Diehl, Int. J. Mod. Phys. B **11**, 3503 (1997).
29 [21] A. Gambassi, A. Maciolek, C. Hertlein, U. Nellen, L. Helden, C. Bechinger and S. Dietrich, Phys. Rev. E **80**, 061143
30 (2009).
31 [22] T.F. Mohry, A. Maciolek and S. Dietrich, Phys. Rev. E **81**, 061117 (2010).
32 [23] U. Nellen, L. Helden and C. Bechinger, EPL **88**, 26001 (2009).
33 [24] F. Schlesener, A. Hanke and S. Dietrich, J. Stat. Phys. **110**, 981 (2003).
34 [25] A. Drzewiński, A. Maciolek and R. Evans, Phys. Rev. Lett. **85**, 3079 (2000).
35 [26] A. Maciolek, A. Drzewiński and R. Evans, Phys. Rev. E **64**, 056137 (2001).
36 [27] R. Evans, J. Phys. Cond. Mat. **2**, 8989 (1990).
37 [28] T. Bieker and S. Dietrich, Physica A **252**, 85 (1998).
38 [29] C. Bauer, T. Bieker and S. Dietrich, Phys. Rev. E **62**, 5324 (2000).
39 [30] R. Evans and J. Stecki, Phys. Rev. B **49**, 8842 (1994).
40 [31] M. Krech, Phys. Rev. E **56**, 1642 (1997).
41 [32] Z. Borjan and P.J. Upton, Phys. Rev. Lett. **101**, 125702 (2008).
42 [33] O. Vasilyev, A. Gambassi, A. Maciolek and S. Dietrich, EPL **80**, 60009 (2007).
43 [34] O. Vasilyev, A. Gambassi, A. Maciolek and S. Dietrich, Phys. Rev. E **79**, 041142 (2009); *ibid* **80**, 039902(E) (2009).
44 [35] M. Hasenbusch, Phys. Rev. B **82**, 104425 (2010).
45 [36] F. Parisen Toldin and S. Dietrich, J. Stat. Mech. P 11003 (2010).
46 [37] F. Parisen Toldin and S. Dietrich, in *Proceedings of the Ninth Conference on Quantum Field Theory Under the Influence*
47 *of External Conditions (QFEXT09)*, edited by K. A. Milton Milton and M. Bordag (World Scientific, Singapore, 2010),
48 p. 355.
49 [38] J. Israelachvili, *Intermolecular and Surface Forces, 2nd ed.* (Academic, New York, 1992).
50 [39] D. Dantchev, F. Schlesener and S. Dietrich, Phys. Rev. E **76**, 011121 (2007).
51 [40] Y. Jayalakshmi, J.S. Van Duijneveldt and D. Beysens, J. Chem. Phys. **100**, 604 (1994).
52 [41] D. Vogt, diploma thesis, University of Stuttgart (2009).
53 [42] D. Qin, Y.N. Xia and G.M. Whitesides, Nature Protocols **5**, 491 (2010).
54 [43] R. Menard, *Springer Handbook of Nanotechnology, Chapter 9: Stamping Techniques for Micro- and Nanofabrication*
55 (Springer, 2004).
56 [44] A. Kumar, H.A. Biebuyck and G.M. Whitesides, Langmuir **10**, 1498 (1994).
57 [45] S.J. Rahi, M. Kardar and T. Emig, Phys. Rev. Lett. **105**, 070404 (2010).
58 [46] J.N. Munday, F. Capasso and V.A. Parsegian, Nature **457**, 170 (2009).
59 [47] A.W. Rodriguez, D. Woolf, A.P. McCauley, F. Capasso, J.D. Joannopoulos and S.G. Johnson, Phys. Rev. Lett. **105**,
60 060401 (2010).
[48] U. Nellen and C. Bechinger (unpublished).