



Energy estimates and cavity interaction for a critical-exponent cavitation model

Sylvia Serfaty, Duvan Henao

► **To cite this version:**

Sylvia Serfaty, Duvan Henao. Energy estimates and cavity interaction for a critical-exponent cavitation model. 2012. <hal-00665991>

HAL Id: hal-00665991

<https://hal.archives-ouvertes.fr/hal-00665991>

Submitted on 3 Feb 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Energy estimates and cavity interaction for a critical-exponent cavitation model

Duvan Henao

Sylvia Serfaty

July 11, 2011

Abstract

We consider the minimization of $\int_{\Omega_\varepsilon} |D\mathbf{u}|^p \, d\mathbf{x}$ in a perforated domain $\Omega_\varepsilon := \Omega \setminus \bigcup_{i=1}^M B_\varepsilon(\mathbf{a}_i)$ of \mathbb{R}^n , among maps $\mathbf{u} \in W^{1,p}(\Omega_\varepsilon, \mathbb{R}^n)$ that are incompressible ($\det D\mathbf{u} \equiv 1$), invertible, and satisfy a Dirichlet boundary condition $\mathbf{u} = \mathbf{g}$ on $\partial\Omega$. If the volume enclosed by $\mathbf{g}(\partial\Omega)$ is greater than $|\Omega|$, any such deformation \mathbf{u} is forced to map the small holes $B_\varepsilon(\mathbf{a}_i)$ onto macroscopically visible cavities (which do not disappear as $\varepsilon \rightarrow 0$). We restrict our attention to the critical exponent $p = n$, where the energy required for cavitation is of the order of $\sum_{i=1}^M v_i |\log \varepsilon|$ and the model is suited, therefore, for an asymptotic analysis (v_1, \dots, v_M denote the volumes of the cavities). We obtain estimates for the “renormalized” energy $\frac{1}{n} \int_{\Omega_\varepsilon} \left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|^p \, d\mathbf{x} - \sum_i v_i |\log \varepsilon|$, showing its dependence on the size and the shape of the cavities, on the initial distance between the cavitation points $\mathbf{a}_1, \dots, \mathbf{a}_M$, and on the distance from these points to the outer boundary $\partial\Omega$. Based on those estimates we conclude, for the case of two cavities, that either the cavities prefer to be spherical in shape and well separated, or to be very close to each other and appear as a single equivalent round cavity. This is in agreement with existing numerical simulations, and is reminiscent of the interaction between cavities in the mechanism of ductile fracture by void growth and coalescence.

1 Introduction

1.1 Motivation

In nonlinear elasticity, cavitation is the name given to the sudden formation of cavities in an initially perfect material, due to its incompressibility (or near-incompressibility), in response to a sufficiently large and triaxial tension. It plays a central role in the initiation of fracture in metals [35, 62, 34, 78, 58] and in elastomers [29, 80, 32, 22, 18] (especially in reinforced elastomers [57, 31, 15, 9, 52]), via the mechanism of void growth and coalescence. It has important applications such as the indirect measurement of mechanical properties [45] or the rubber-toughening of brittle polymers [46, 14, 76, 48]. Mathematically, it constitutes a realistic example of a regular variational problem with singular minimizers, and corresponds to the case when the stored-energy function of the material is not $W^{1,p}$ -quasiconvex [2, 5, 7], the Jacobian determinant is not weakly continuous [7], and important properties such as the invertibility of the deformation may not pass to the weak limit [55, Sect. 11]. The problem has been studied by many authors, beginning with Gent-Lindley

[30] and Ball [4]; see the review papers [29, 41, 25], the variational models of Müller-Spector [55] and Sivaloganathan-Spector [70], and the recent works [38, 49] for further motivation and references.

The standard model in the variational approach to cavitation considers functionals of the form

$$\int_{\Omega} |D\mathbf{u}|^p \, d\mathbf{x}, \quad (1.1)$$

where the deformation $\mathbf{u} : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}^n$ is constrained to be incompressible (i.e. $\det D\mathbf{u} = 1$) and globally invertible, and either a Dirichlet condition $\mathbf{u} = \mathbf{g}$ or a force boundary condition is applied. Unless the boundary condition is exactly compatible with the volume, cavities have to be formed. If $p < n$ this can happen while still keeping a finite energy. A typical deformation creating a cavity of volume $\omega_n A^n$ at the origin (ω_n being the volume of the unit ball in \mathbb{R}^n) is given by

$$\mathbf{u}(\mathbf{x}) = \sqrt[n]{A^n + |\mathbf{x}|^n} \frac{\mathbf{x}}{|\mathbf{x}|}. \quad (1.2)$$

We can easily compute that

$$|D\mathbf{u}|^2 \underset{\mathbf{x}=\mathbf{0}}{\sim} \frac{(n-1)A^2}{|\mathbf{x}|^2}. \quad (1.3)$$

In that situation the origin is called a *cavitation point*, which belongs to the domain space, and its image by \mathbf{u} is the *cavity*, belonging to the target space. Contrarily to the usual, we study the critical case $p = n$ where the cavity behaviour (1.2) just fails to be of finite energy.

This fact is analogous to what happens for \mathbb{S}^1 -valued harmonic maps in dimension 2, which were particularly studied in the context of the Ginzburg-Landau model, see Bethuel-Brezis-Hélein [10]. For \mathbb{S}^1 -valued maps \mathbf{u} from $\Omega \subset \mathbb{R}^2$, the topological degree of \mathbf{u} around a point \mathbf{a} is defined by the following integer

$$d = \frac{1}{2\pi} \int_{\partial B(\mathbf{a}, r)} \frac{\partial \mathbf{u}}{\partial \tau} \times \mathbf{u}.$$

Points around which this is not zero are called vortices. Typical vortices of degree d look like $\mathbf{u} = e^{id\theta}$ (in polar coordinates). If $d \neq 0$ again $|D\mathbf{u}|^2$ just fails to be integrable since for the typical vortex $|D\mathbf{u}|^2 \underset{\mathbf{x}=\mathbf{0}}{\sim} \frac{|d|^2}{|\mathbf{x}|^2}$, just as above (1.3), up to a constant factor. So there is an analogy in that sense between maps from Ω to \mathbb{C} which are constrained to satisfy $|\mathbf{u}| = 1$, and maps from Ω to \mathbb{R}^2 which satisfy the incompressibility constraint $\det D\mathbf{u} = 1$. We see that in this analogy (in dimension 2) the volume of the cavity divided by π plays the role of the absolute value of the degree for \mathbb{S}^1 -valued maps. In this correspondence two important differences appear: the degree is quantized while the cavity volume is not; on the other hand the degree has a sign, which can lead to “cancellations” between vortices, while the cavity volume is always positive.

In the context of \mathbb{S}^1 -valued maps, two possible ways of giving a meaning to $\int_{\Omega} |D\mathbf{u}|^2$ are the following. The first is to relax the constraint $|\mathbf{u}| = 1$ and replace it by a penalization, and study instead

$$\int_{\Omega} |D\mathbf{u}|^2 + \frac{1}{\varepsilon^2} (1 - |\mathbf{u}|^2)^2 \quad (1.4)$$

in the limit $\varepsilon \rightarrow 0$; this is the Ginzburg-Landau approximation. The second is to study the energy with the constraint $|\mathbf{u}| = 1$ but in a punctured domain $\Omega_{\varepsilon} := \Omega \setminus \cup_i B(\mathbf{a}_i, \varepsilon)$ where \mathbf{a}_i 's stand for

the vortex locations:

$$\min_{|\mathbf{u}|=1} \int_{\Omega_\varepsilon} |D\mathbf{u}|^2 \quad (1.5)$$

again in the limit $\varepsilon \rightarrow 0$; this can be called the “renormalized energy approach”. Both of these approaches were followed in [10], where it is proven that the Ginzburg-Landau approach essentially reduces to the renormalized energy approach. More specifically, when there are vortices at \mathbf{a}_i , $|D\mathbf{u}|$ will behave like $|d_i|/|\mathbf{x} - \mathbf{a}_i|$ near each vortex (where d_i is the degree of the vortex) and both energies (1.4) and (1.5) will blow up like $\pi \sum_i d_i^2 \log \frac{1}{\varepsilon}$ as $\varepsilon \rightarrow 0$. It is shown in [10] that when this divergent term is subtracted off (this is the “renormalization” procedure), what remains is a nondivergent term depending on the positions of the vortices \mathbf{a}_i and their degrees d_i (and the domain), called precisely the renormalized energy. That energy is essentially a Coulombian interaction between the points \mathbf{a}_i behaving like charged particles (vortices of same degree repel, those of opposite degrees attract) and it can be written down quite explicitly.

Our goal here is to study cavitation in the same spirit. A first attempt, which would be the analogue of (1.4), would be to relax the incompressibility constraint and study for example

$$\int_{\Omega} |D\mathbf{u}|^2 + \frac{(1 - \det D\mathbf{u})^2}{\varepsilon}. \quad (1.6)$$

We do not however follow this route which seems to present many difficulties (one of them is that this energy in two dimensions is scale invariant, and that contrarily to (1.4) the nonlinearity contains as many order of derivatives as the other term), but it remains a seemingly interesting open problem, which would have good physical sense. Rather we follow the second approach, i.e. that of working in punctured domains while keeping the incompressibility constraint.

For the sake of generality we consider holes which can be of different radii $\varepsilon_1, \dots, \varepsilon_m$, define $\Omega_\varepsilon := \Omega \setminus \cup_{i=1}^m \overline{B}(\mathbf{a}_i, \varepsilon_i)$ and look at

$$\min_{\det D\mathbf{u}=1} \int_{\Omega_\varepsilon} |D\mathbf{u}|^2 \quad (1.7)$$

(or $\min_{\det D\mathbf{u}=1} \int_{\Omega_\varepsilon} |D\mathbf{u}|^n$ in dimension n), in the limit $\varepsilon \rightarrow 0$. This also has a reasonable physical interpretation: it corresponds to studying the incompressible deformation of a body that contains micro-voids which expand under the applied boundary deformation. One may think of the points \mathbf{a}_i as fixed, then they correspond to defects that pre-exist, just as above. Or the model can be seen as a fracture model where we postulate that the body will first break around the most energetically favorable points \mathbf{a}_i (see, e.g., the discussion in [4, 43, 69, 29, 42, 55, 6, 71, 74, 49, 50]). It can also be compared to the core-radius approach in dislocation models [13, 59, 28].

Following the analogy above, we would like to be able to subtract from (1.7) a leading order term proportional to $\log \frac{1}{\varepsilon}$, in order to extract at the next order a “renormalized” term which will tell us how cavities “interact” (attract or repel each other), according to their positions and shapes. This is more difficult than the problem (1.5) because the condition $\det D\mathbf{u} = 1$ is much less constraining than $|\mathbf{u}| = 1$. While the maps with $|\mathbf{u}| = 1$ can be parametrized by lifting in the form $\mathbf{u} = e^{i\varphi}$, to our knowledge no parametrization of that sort exists for incompressible maps. In addition while the only characteristic of a vortex is an integer –its degree–, for incompressible maps, the characteristics of a cavity are more complex –they comprise the volume of the cavity and its

shape-, and there is no quantization. For these reasons we cannot really hope for something as nice and explicit as a complete “renormalized energy” for this toy cavitation model. However we will show that we can obtain, in particular in the case of two cavities, some quantitative information about the cavities interaction that is reminiscent of the renormalized energy.

1.2 Method and main results : energy lower bounds

Our method relies on obtaining general and ansatz-free lower bounds for the energy on the one hand, and on the other hand upper bounds via explicit constructions, which match as much as possible the lower bounds. This is in the spirit of Γ -convergence (however we will not prove a complete Γ -convergence result). For simplicity in this section we present the results in dimension 2, but they carry over in higher dimension.

To obtain lower bounds we use the “ball construction method”, which was introduced in the context of Ginzburg-Landau by Jerrard [44] and Sandier [65, 66]. The crucial estimate for Ginzburg-Landau, or more simply S^1 -valued harmonic maps, is the following simple relation, corollary of Cauchy-Schwarz:

$$\int_{\partial B(\mathbf{a}, r)} |D\mathbf{u}|^2 \geq \frac{1}{2\pi r} \left(\int_{\partial B(\mathbf{a}, r)} \frac{\partial \mathbf{u}}{\partial \tau} \times \mathbf{u} \right)^2 = 2\pi \frac{d^2}{r} \quad (1.8)$$

if d is the degree of the map on $\partial B(\mathbf{a}, r)$. Integrating this relation over r ranging from ε to 1 yields a lower bound for the energy on annuli, with the logarithmic behavior stated above. One sees that the equality case in (1.8) is achieved when \mathbf{u} is exactly radial (which corresponds to $\mathbf{u} = e^{id\theta}$ in polar coordinates), so the least energetically costly vortices are the radial ones. For an arbitrary number of vortices the “ball construction” à la Jerrard and Sandier allows to paste together the lower bounds obtained on disjoint annuli. Previous constructions for bounded numbers of vortices include those of Bethuel-Brezis-Hélein [10] and Han-Shafrir [36]. The ball construction method will be further described in Section 3.1.

For the cavitation model, there is an analogue to the above calculation, which is also our starting point. Assume that \mathbf{u} develops a cavity of volume v around a cavitation point \mathbf{a} in the domain space. By v we really denote the excess of volume created by the cavity (we still refer to it as cavity volume), this way the image of the ball $B(\mathbf{a}, \varepsilon)$ contains a volume $\pi\varepsilon^2 + v$. Using the Cauchy-Schwarz inequality, we may write

$$\int_{\partial B(\mathbf{a}, r)} |D\mathbf{u}|^2 \geq \frac{1}{2\pi r} \left(\int_{\partial B(\mathbf{a}, r)} |D\mathbf{u} \cdot \tau| \right)^2. \quad (1.9)$$

But then one can observe that $\int_{\partial B(\mathbf{a}, r)} |D\mathbf{u} \cdot \tau|$ is exactly the length of the image curve of the circle $\partial B(\mathbf{a}, r)$. We may then use the classical isoperimetric inequality

$$(\text{Per } E(\mathbf{a}, r))^2 \geq 4\pi |E(\mathbf{a}, r)| \quad (1.10)$$

where $|\cdot|$ denotes the volume, and $E(\mathbf{a}, r)$ is the region enclosed by this image curve, which contains the cavity, and has volume $\pi r^2 + v$ by incompressibility. Inserting this into (1.9), we are led to

$$\int_{\partial B(\mathbf{a}, r)} \frac{|D\mathbf{u}|^2}{2} \geq \frac{\text{Per}^2(E(\mathbf{a}, r))}{4\pi r} \geq \frac{|E(\mathbf{a}, r)|}{r} \geq \frac{v}{r} + \pi r. \quad (1.11)$$

This is the building block that we will integrate over r and insert into the ball construction, to obtain our first lower bound, which is proved in Section 3.1. To state it, we will use the notion of weak determinant:

$$\langle \text{Det } D\mathbf{u}, \phi \rangle := -\frac{1}{n} \int_{\Omega} \mathbf{u}(\mathbf{x}) \cdot (\text{cof } D\mathbf{u}(\mathbf{x})) D\phi(\mathbf{x}) \, d\mathbf{x}, \quad \forall \phi \in C_c^\infty(\Omega)$$

whose essential features we recall in Section 2.4; as well as Müller and Spector’s invertibility “condition INV” [55] which is defined in Section 2.3 (Definition 5) and which essentially means that the deformations of the material, in addition to being one-to-one, cannot create cavities which would at the same time be filled by material coming from elsewhere. Even though we have discussed dimension 2, we directly state the result in dimension n .

Proposition 1.1. *Let Ω be an open and bounded set in \mathbb{R}^n , and $\Omega_\varepsilon = \Omega \setminus \cup_{i=1}^m \overline{B}(\mathbf{a}_i, \varepsilon_i)$ where $\mathbf{a}_1, \dots, \mathbf{a}_m \in \Omega$ and the $\overline{B}(\mathbf{a}_i, \varepsilon_i)$ are disjoint. Suppose that $\mathbf{u} \in W^{1,n}(\Omega_\varepsilon, \mathbb{R}^n)$ and that condition INV is satisfied. Suppose, further, that $\text{Det } D\mathbf{u} = \mathcal{L}^n$ in Ω_ε (where \mathcal{L}^n is the Lebesgue measure), and let $v_i := |E(\mathbf{a}_i, \varepsilon_i)| - \omega_n \varepsilon_i^n$ (with $E(\mathbf{a}_i, \varepsilon_i)$ as in (1.10)). Then for any $R > 0$*

$$\frac{1}{n} \int_{\Omega_\varepsilon} \left(\left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|^n - 1 \right) d\mathbf{x} \geq \left(\sum_{i, B(\mathbf{a}_i, R) \subset \Omega} v_i \right) \log \frac{R}{2 \sum_{i=1}^m \varepsilon_i}.$$

Note that $\sum_i v_i = V$ is the total cavity volume, which due to incompressibility is completely determined by the Dirichlet data, in the case of a displacement boundary value problem.

Examining the equality cases in the chain of inequalities (1.9)–(1.11) already tells us that the minimal energy is obtained when “during the ball construction” all circles (at least for r small) are mapped into circles and the cavities are spherical. A more careful examination of (1.9) indicates that the map should at least locally follow the model cavity map (1.2). It is the same argument that has been used by Sigalovanathan and Spector [72, 73] to prove the radial symmetry of minimizers for the model with power $p < n$.

When there is more than one cavity, and two cavities are close together, we can observe that there is a geometric obstruction to all circles “of the ball construction” being mapped into circles. This is true for any number of cavities larger than 1; to quantify it is in principle possible but a bit inextricable for more than 2. For that reason and for simplicity, we restrict to the case of two cavities, and now explain the quantitative point.

Let \mathbf{a}_1 and \mathbf{a}_2 be the two cavitation points with $|\mathbf{a}_1 - \mathbf{a}_2| = d$, small compared to 1. For simplicity of the presentation let us also assume that $\varepsilon_1 = \varepsilon_2 = \varepsilon$. The ball construction is very simple in such a situation: three disjoint annuli are constructed, $B(\mathbf{a}_1, d/2) \setminus B(\mathbf{a}_1, \varepsilon)$, $B(\mathbf{a}_2, d/2) \setminus B(\mathbf{a}_2, \varepsilon)$ and $B(\mathbf{a}, R) \setminus B(\mathbf{a}, d)$, where \mathbf{a} is the midpoint of \mathbf{a}_1 and \mathbf{a}_2 (see Figure 1). These annuli can be seen as a union of concentric circles centred at \mathbf{a}_1 , \mathbf{a}_2 , \mathbf{a} respectively. To achieve the optimality condition above, each of these circles would have to be mapped by \mathbf{u} into a circle. If this were true, the images of $B(\mathbf{a}_1, d/2)$ and $B(\mathbf{a}_2, d/2)$ would be two disjoint balls containing the two cavities, call them E_1 and E_2 . By incompressibility, $|E_1| = v_1 + \pi(d/2)^2$ and $|E_2| = v_2 + \pi(d/2)^2$. Then the image of $B(\mathbf{a}, d)$ would also have to be a ball, call it E , which contains the disjoint union $E_1 \cup E_2$, and by incompressibility

$$|E| = v_1 + v_2 + \pi d^2. \tag{1.12}$$




Figure 1: Ball construction in the reference configuration

If d is small compared to v_1 and v_2 it is easy to check this is geometrically impossible: the radius of the ball E_1 is certainly bigger than $\sqrt{v_1/\pi}$, that of E_2 than $\sqrt{v_2/\pi}$ and since E is a ball that contains their disjoint union, its radius is at least the sum of the two, hence $|E| \geq (\sqrt{v_1} + \sqrt{v_2})^2$. This is incompatible with (1.12) unless $\pi d^2 \geq 2\sqrt{v_1 v_2}$.




Figure 2: Incompressible deformation $\mathbf{u} : B(\mathbf{0}, d) \setminus \{\mathbf{a}_1, \mathbf{a}_2\} \rightarrow \mathbb{R}^2$, $d = |\mathbf{a}_2 - \mathbf{a}_1|$, opening distorted cavities of volumes $v_1 + \pi\varepsilon_1^2$, $v_2 + \pi\varepsilon_2^2$; deformed configuration for increasing values of the displacement load ($\mu := \sqrt{\frac{v_1 + v_2}{\pi d^2}}$). Choice of parameters: $d = 1$, $\frac{v_2}{v_1} = 0.3$.

So in practice, if d is small compared to the volumes, the circles are not all mapped to exact circles, the inclusion and disjointness are preserved, but some distortion in the shape of the images has to be created *either* for the “balls before merging” i.e. E_1 and E_2 – this corresponds to what is sketched on Figure 2 – *or* for the “balls after merging” i.e. E – this corresponds to what is sketched in Figure 3 (the situations of Figures 2 and 3 correspond to the test-maps we will use to get energy upper bounds, see below Section 1.3).

A convenient tool to quantify how much these sets differ from balls, which is what we exactly mean by “distortion”, is the following




Figure 3: Incompressible deformation of $B(\mathbf{0}, d)$, $d := |\mathbf{a}_2 - \mathbf{a}_1|$, for increasing values of $\mu := \sqrt{\frac{v_1 + v_2}{\pi d^2}}$. Final cavity volumes v_1 and v_2 given by $d = 1$, $\frac{v_2}{v_1} = 0.3$.

Definition 1. *The Fraenkel asymmetry of a measurable set $E \subset \mathbb{R}^n$ is defined as*

$$D(E) := \min_{\mathbf{x} \in \mathbb{R}^n} \frac{|E \Delta B(\mathbf{x}, r_E)|}{|E|}, \quad \text{with } r_E \text{ such that } |B(\mathbf{x}, r_E)| = |E|$$

where Δ denotes the symmetric difference between sets.

Note that $D(E)$ is a scale-free quantity which depends not on the size of E , but on its shape.

The following proposition, which we shall prove in Section 3.3, allows to make the observations above quantitative in terms of the distortions.

Proposition 1.2. *Let E , E_1 , and E_2 be sets of positive measure in \mathbb{R}^n , $n \geq 2$ such that $E \supset E_1 \cup E_2$ and $E_1 \cap E_2 = \emptyset$, and assume without loss of generality that $|E_1| \geq |E_2|$. Then*

$$\frac{|E|D(E)^{\frac{n}{n-1}} + |E_1|D(E_1)^{\frac{n}{n-1}} + |E_2|D(E_2)^{\frac{n}{n-1}}}{|E| + |E_1 \cup E_2|} \geq C_n \left(\frac{|E_2|}{|E_1| + |E_2|} \right)^{\frac{n}{n-1}} \left(\frac{(|E_1|^{\frac{1}{n}} + |E_2|^{\frac{1}{n}})^n - |E|}{(|E_1|^{\frac{1}{n}} + |E_2|^{\frac{1}{n}})^n - |E_1 \cup E_2|} \right)^{\frac{n(n+1)}{2(n-1)}}$$

for some constant $C_n > 0$ depending only on n .

The fact that E_1, E_2, E cannot simultaneously be balls is made explicit by the fact that $D(E_1), D(E_2), D(E)$ cannot all vanish unless the right-hand side is negative, which can happen only if $|E|$ is large relative to $|E_1|$ and $|E_2|$. The first factor in the estimate degenerates only when one of the sets is very small compared to the other.

Note that such a geometric constraint is also true for more than two merging balls, so in principle we could treat (with more effort) the case of more than two cavities, however the estimates would degenerate as the number of cavities gets large.

These estimates on the distortions are useful for us thanks to the following improved isoperimetric inequality, precisely expressed in terms of the Fraenkel asymmetry:

Proposition 1.3 (Fusco-Maggi-Pratelli [27]). *For every Borel set $E \subset \mathbb{R}^n$*

$$\text{Per } E \geq n\omega_n^{\frac{1}{n}} |E|^{\frac{n-1}{n}} (1 + CD(E)),$$

where C is a universal constant.

In dimension 2, we thus have the improved isoperimetric inequality

$$(\text{Per } E)^2 \geq 4\pi|E| + C|E|D^2(E), \quad (1.13)$$

for some universal $C > 0$. Inserting (1.13) instead of (1.10) into the basic estimate (1.11) gives us

$$\int_{\partial B(\mathbf{a}, r)} \frac{|D\mathbf{u}|^2}{2} \geq \frac{|E(\mathbf{a}, r)|}{r} + \frac{C}{r} |E(\mathbf{a}, r)| D^2(E(\mathbf{a}, r)) \geq \frac{v}{r} + \pi r + \frac{C}{r} |E(\mathbf{a}, r)| D^2(E(\mathbf{a}, r)). \quad (1.14)$$

This then allows us to get improved estimates when integrating over r (in a ball construction procedure), keeping track of the fact that to achieve equality, all level curves $E(\mathbf{a}, r)$ which are images of circles during the ball construction would have to be circles. This way, after subtracting off the leading order term $\sum_i v_i \log \frac{1}{\sum_i \varepsilon_i}$ we can retrieve a next order ‘‘renormalized’’ term that will account for the cavity interaction. This is expressed in the following main result.

Theorem 1 (Lower bound). *Given $\Omega \subset \mathbb{R}^n$ a bounded open set, let $\Omega_\varepsilon := \Omega \setminus (\overline{B}_{\varepsilon_1}(\mathbf{a}_1) \cup \overline{B}_{\varepsilon_2}(\mathbf{a}_2))$, where $\mathbf{a}_1, \mathbf{a}_2 \in \Omega$, $\varepsilon_1, \varepsilon_2 > 0$, and assume that $B_{\varepsilon_1}(\mathbf{a}_1)$ and $B_{\varepsilon_2}(\mathbf{a}_2)$ are disjoint and contained in Ω . Suppose that $\mathbf{u} \in W^{1,n}(\Omega_\varepsilon, \mathbb{R}^n)$ satisfies condition INV and $\text{Det } D\mathbf{u} = \mathcal{L}^n$ in Ω_ε . Set*

$$\mathbf{a} := \frac{\mathbf{a}_1 + \mathbf{a}_2}{2}, \quad d := |\mathbf{a}_1 - \mathbf{a}_2|, \quad v_1 := |E(\mathbf{a}_1, \varepsilon_1)| - \omega_n \varepsilon_1^n, \quad v_2 := |E(\mathbf{a}_2, \varepsilon_2)| - \omega_n \varepsilon_2^n.$$

Then, for all R such that $B(\mathbf{a}, R) \subset \Omega$,

$$\begin{aligned} \frac{1}{n} \int_{\Omega_\varepsilon \cap B(\mathbf{a}, R)} \left(\left| \frac{D\mathbf{u}(\mathbf{x})}{\sqrt{n-1}} \right|^n - 1 \right) d\mathbf{x} &\geq v_1 \log \frac{R}{2\varepsilon_1} + v_2 \log \frac{R}{2\varepsilon_2} \\ &+ C(v_1 + v_2) \left(\left(\frac{\min\{v_1, v_2\}}{v_1 + v_2} \right)^{\frac{n}{n-1}} - \frac{\omega_n d^n}{v_1 + v_2} \right)_+ \log \min \left\{ \left(\frac{v_1 + v_2}{2^n \omega_n d^n} \right)^{\frac{1}{n^2}}, \frac{R}{d}, \frac{d}{\max\{\varepsilon_1, \varepsilon_2\}} \right\} \end{aligned}$$

for some constant C independent of Ω , \mathbf{a}_1 , \mathbf{a}_2 , d , v_1 , v_2 , ε_1 , and ε_2 (t_+ stands for $\max\{0, t\}$).

Two main differences appear in this lower bound compared to Proposition 1.1. First, the leading order term $(v_1 + v_2) \log \frac{1}{\varepsilon_1 + \varepsilon_2}$ has been improved to $v_1 \log \frac{1}{\varepsilon_1} + v_2 \log \frac{1}{\varepsilon_2}$, which shows that the energy goes to infinity as $\varepsilon_1 \rightarrow 0$ or $\varepsilon_2 \rightarrow 0$, even if $\varepsilon_1 + \varepsilon_2 \not\rightarrow 0$. This term is optimal since it coincides with the leading order term in the upper bound of Theorem 2 below, and in fact it should be possible to replace $\sum_i v_i \log \frac{1}{\sum_i \varepsilon_i}$ with $\sum_i (v_i \log \frac{1}{\varepsilon_i})$ in Proposition 1.1 (however, this would require a more sophisticated ball construction, and it is not immediately clear how to obtain a general result for the case of more than two cavities). Second, and returning to the discussion in dimension two and choosing $\varepsilon_1 = \varepsilon_2 = \varepsilon$, compared to Proposition 1.1 we have gained the new term

$$C(v_1 + v_2) \left(\left(\frac{\min\{v_1, v_2\}}{v_1 + v_2} \right)^2 - \frac{\pi d^2}{v_1 + v_2} \right) \min \left\{ \log \sqrt[4]{\frac{v_1 + v_2}{4\pi d^2}}, \log \frac{R}{d}, \log \frac{d}{\varepsilon} \right\},$$

This term is of course worthless unless $\frac{\pi d^2}{v_1+v_2} < \left(\frac{\min\{v_1, v_2\}}{v_1+v_2}\right)^2$ i.e. $\pi d^2 \leq \frac{\min\{v_1^2, v_2^2\}}{v_1+v_2}$. Under that condition, it expresses an interaction between the two cavities in terms of the distance of the cavitation points relative to the data of v_1, v_2 and ε . As $\frac{\pi d^2}{v_1+v_2} \rightarrow 0$ the interaction tends logarithmically to $+\infty$; this expresses a *logarithmic repulsion* between the cavities, unless the term $\log \frac{d}{\varepsilon}$ is the one that achieves the min above, which can only happen if $\log d$ is comparable to $\log \varepsilon$. This expresses an *attraction of the cavities when they are close compared to the puncture scale*, which we believe means that two cavities thus close would energetically prefer to be merged into one. This suggests that three scenarii are energetically possible:

Scenario (i) the cavities are spherical and the cavitation points are well separated (but not necessarily the cavities themselves), this is the situation of Figure 3

Scenario (ii) the cavitation points are at distance $\ll 1$ but all but one cavity are of very small volume and hence “close up” in the limit $\varepsilon \rightarrow 0$

Scenario (iii) “outer circles” (in the ball construction) are mapped into circles and cavities (as well as cavitation points) are pushed together to form one equivalent round cavity, this is the situation of Figure 2. This seems to correspond to void coalescence (c.f. [81, 47]).

1.3 Method and main results: upper bound

After obtaining this lower bound, we show that it is close to being optimal (at least in scale). To do so we need to construct explicit test maps and evaluate their energy (in terms of the parameters of the problem). The main difficulty is that these test maps have to satisfy the incompressibility condition outside of the cavitation points, and as we mentioned previously, there is no simple parametrization of such incompressible maps. The main known result in that area is the celebrated result of Dacorogna and Moser [20] which provides an existence result for incompressible maps with compatible boundary conditions. Two methods are proposed in their work, one of them constructive, however they are not explicit enough to evaluate the Dirichlet energy of the map.

The question we address can be phrased in the following way: given a domain with a certain number of “round holes” at certain distances from each other, and another domain of same volume, with the same number of holes whose volumes are prescribed but whose positions and shapes are free; can we find an incompressible map that maps one to the other, and can we estimate its energy $\int |D\mathbf{u}|^n$ in terms of the distance of the holes and the cavity volumes?

We answer positively this question, still in the case of two holes, by using two tools:

- (a) a family of explicitly defined incompressible deformations preserving angles, that we introduce
- (b) the construction of incompressible maps of Rivière and Ye [63, 64], which is more tractable than Dacorogna and Moser to obtain energy estimates.

We believe it would be of interest to tackle that question in a more general setting: compute the minimal Dirichlet energy of an incompressible map between two domains with same volume, and the same number of holes, the holes having arbitrary shapes and sizes; and find appropriate

geometric parameters to evaluate it as a function of the domains. This question is beyond the scope of our paper however and we do not attempt to treat it in that much generality.




Figure 4: Transition from round to distorted cavities: $d = 1$, $\sqrt{\frac{v_1+v_2}{\pi d^2}} = 1.5$, $\frac{v_2}{v_1} = 0.3$.

Our main result (proved in Section 4.1) is the following.

Theorem 2. *Let $\mathbf{a}_1, \mathbf{a}_2 \in \mathbb{R}^n$, $v_1 \geq v_2 \geq 0$, and suppose that $d := |\mathbf{a}_1 - \mathbf{a}_2| > \varepsilon_1 + \varepsilon_2$. Then, for every $\delta \in [0, 1]$ there exists \mathbf{a}^* in the line segment joining \mathbf{a}_1 and \mathbf{a}_2 , and a piecewise smooth map $\mathbf{u} \in C(\mathbb{R}^n \setminus \{\mathbf{a}_1, \mathbf{a}_2\}, \mathbb{R}^n)$ satisfying condition INV, such that $\text{Det } D\mathbf{u} = \mathcal{L}^n + v_1\delta_{\mathbf{a}_1} + v_2\delta_{\mathbf{a}_2}$ in \mathbb{R}^n and for all $R > 0$*

$$\int_{B(\mathbf{a}^*, R) \setminus (B_{\varepsilon_1}(\mathbf{a}_1) \cup B_{\varepsilon_2}(\mathbf{a}_2))} \frac{1}{n} \left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|^n dx \leq C_1(v_1 + v_2 + \omega_n R^n) + v_1 \left(\log \frac{R}{\varepsilon_1} \right)_+ + v_2 \left(\log \frac{R}{\varepsilon_2} \right)_+ \\ + C_2(v_1 + v_2) \left((1 - \delta) \left(\log \frac{R}{d} \right)_+ + \delta \left(\sqrt[n]{\frac{v_2}{v_1}} \log \frac{d}{\varepsilon_1} + \sqrt[2n]{\frac{v_2}{v_1}} \log \frac{d}{\varepsilon_2} \right) \right)$$

(C_1 and C_2 are universal constants depending only on n).

If we are not preoccupied with boundary conditions but just wish to build a test configuration with cavities of prescribed volumes and cavitation points at distance d , then the above result suffices. This is obtained by our construction of an explicit family of incompressible maps, which contains parameters allowing for all possible cavitation points distances d and cavity volumes v_1, v_2 . The feature of this construction is that it allows for our almost optimal estimates, as the shapes of the cavities are automatically adjusted to the optimal scenario according to the ratio between $d, \varepsilon, \sqrt{v_1}, \sqrt{v_2}$, their logs, etc, as in the three scenarii of the end of the previous subsection. In other words, the construction builds cavities which, when d is comparable to ε , are distorted and form one equivalent round cavity while the deformation rapidly becomes radially symmetric (as in Scenario (iii)); and cavities which are more and more round as d gets large compared to ε (as in Scenario (i)). For the extreme cases $\delta = 1$ and $\delta = 0$, the maps are those that were presented in Figures 2 and 3 respectively. The result for intermediate values of δ is shown in Figure 4.




Figure 5: Geometric construction of domains Ω_1, Ω_2 satisfying $\frac{|\Omega_1|}{|\Omega_2|} = \frac{v_1}{v_2}$.

The idea of the construction is the following. Take two intersecting balls $B(\tilde{\mathbf{a}}_1, \rho_1)$ and $B(\tilde{\mathbf{a}}_2, \rho_2)$ such that the width of their union is exactly $2d$ and the width of their intersection is $2d\delta$, and let Ω_1 and Ω_2 be as in Figure 5 (the precise definition is given in (4.4)). As will be proved in Section 4.1, for every $\delta \in [0, 1]$ there are unique ρ_1 and ρ_2 such that $\frac{|\Omega_1|}{|\Omega_2|} = \frac{v_1}{v_2}$. The cavitation points \mathbf{a}_1 and \mathbf{a}_2 are suitably placed in Ω_1 and Ω_2 , respectively, in such a way that $|\mathbf{a}_1 - \mathbf{a}_2| = d$. It is always possible to choose \mathbf{a}^* between \mathbf{a}_1 and \mathbf{a}_2 such that $\overline{\Omega_1 \cup \Omega_2}$ is *star-shaped* with respect to \mathbf{a}^* . In order to define \mathbf{u} in $\mathbb{R}^n \setminus \overline{\Omega_1 \cup \Omega_2}$ we choose \mathbf{a}^* as the origin and look for an *angle-preserving* map

$$\mathbf{u}(\mathbf{x}) = \lambda \mathbf{a}^* + f(\mathbf{x}) \frac{\mathbf{x} - \mathbf{a}^*}{|\mathbf{x} - \mathbf{a}^*|}, \quad \lambda^n - 1 := \frac{v_1 + v_2}{|\Omega_1 \cup \Omega_2|} = \frac{v_1}{|\Omega_1|} = \frac{v_2}{|\Omega_2|}.$$

By so doing, we can solve the incompressibility equation $\det D\mathbf{u} = 1$ explicitly, since for angle-preserving maps the equation has the same form as in the radial case,

$$\det D\mathbf{u}(\mathbf{x}) = \frac{f^{n-1}(\mathbf{x}) \frac{\partial f}{\partial r}(\mathbf{x})}{r^{n-1}} \equiv 1, \quad r = |\mathbf{x} - \mathbf{a}^*|,$$

which we will see can be solved as

$$f^n(\mathbf{x}) = |\mathbf{x} - \mathbf{a}^*|^n + A \left(\frac{\mathbf{x} - \mathbf{a}^*}{|\mathbf{x} - \mathbf{a}^*|} \right)^n,$$

where the function $A : \mathbb{S}^{n-1} \rightarrow \mathbb{R}$ is completely determined if we prescribe \mathbf{u} on $\partial\Omega_1 \cup \partial\Omega_2$. Inside Ω_1 and Ω_2 the deformation \mathbf{u} is defined analogously, taking \mathbf{a}_1 and \mathbf{a}_2 as the corresponding origins. The resulting map creates cavities at \mathbf{a}_1 and \mathbf{a}_2 with the desired volumes, and with exactly the same shape as $\partial\Omega_1$ and $\partial\Omega_2$. For compatibility we impose $\mathbf{u}(\mathbf{x}) = \lambda \mathbf{x}$ on $\partial\Omega_1 \cup \partial\Omega_2$.

In the energy estimate, $(1 - \delta) \log \frac{R}{d}$ is the excess energy due to the distortion of the ‘outer’ curves $\mathbf{u}(\partial B(\mathbf{a}^*, r))$, $r \in (d, R)$, and $\delta \left(n \sqrt{\frac{v_2}{v_1}} \log \frac{d}{\varepsilon_1} + 2n \sqrt{\frac{v_2}{v_1}} \log \frac{d}{\varepsilon_2} \right)$ is that due to the distortion of the curves $\mathbf{u}(\partial B(\mathbf{a}_i, r))$, $r \in (\varepsilon_i, d)$, $i = 1, 2$ near the cavities. When $\delta = 0$, $\overline{\Omega_1}$ and $\overline{\Omega_2}$ are tangent balls, the cavities are spherical, and the second term in the estimate vanishes. The outer curves are distorted because their shape depends on that of $\partial(\Omega_1 \cup \Omega_2)$, hence a price of the order of

$(v_1 + v_2) \log \frac{R}{d}$ is felt in the energy. When $\delta = 1$, at the opposite end, $\Omega_1 \cup \Omega_2$ is a ball of radius d , the deformation is radially symmetric outside $\Omega_1 \cup \Omega_2$, and no extra price for the outer curves is paid. In contrast, the cavities are “D-shaped” (they are copies of $\partial\Omega_1$ and $\partial\Omega_2$), and a price of order $(v_1 + v_2) \sqrt[2n]{\frac{v_2}{v_1}} \log \frac{d}{\varepsilon}$ is obtained as a consequence (in this case the excess energy vanishes as $\frac{v_2}{v_1} \rightarrow 0$, in agreement with the prediction of Theorem 1).

Since the last term of the energy estimate is linear in δ , by taking¹ either $\delta = 0$ or $\delta = 1$ (and assuming $R > d$) the estimate becomes

$$C(v_1 + v_2) \min \left\{ \log \frac{R}{d}, \sqrt[n]{\frac{v_2}{v_1}} \log \frac{d}{\varepsilon_1} + \sqrt[2n]{\frac{v_2}{v_1}} \log \frac{d}{\varepsilon_2} \right\}.$$

Comparing it against the corresponding term for the lower bound, namely²,

$$C(v_1 + v_2) \min \left\{ \left(\frac{v_2}{v_1} \right)^{\frac{n}{n-1}} \log \frac{R}{d}, \left(\frac{v_2}{v_1} \right)^{\frac{n}{n-1}} \log \frac{d}{\max\{\varepsilon_1, \varepsilon_2\}} \right\},$$

we observe that there are still some qualitative differences. First of all, in the case when $\varepsilon_1 \ll \varepsilon_2$, a term of the form $\log \frac{d}{\varepsilon_1} + \log \frac{d}{\varepsilon_2}$ is much larger than $\log \frac{d}{\max\{\varepsilon_1, \varepsilon_2\}}$. We believe that the expression in the lower bound quantifies more accurately the effect of the distortion of the cavities, and that the obstacle for obtaining a comparable expression in the upper bound is that the domains Ω_1 and Ω_2 in our explicit constructions are required to be star-shaped. For example, in the case $d \sim \varepsilon_2$, an energy minimizing deformation \mathbf{u} would try to create a spherical cavity at \mathbf{a}_1 (so as to prevent a term of order $\log \frac{d}{\varepsilon_1}$ from appearing in the energy due to the distortion of the first cavity), and, at the same time, to rapidly become radially symmetric (because of the price of order $\log \frac{R}{d}$ due to the distortion of the ‘outer’ circles). Therefore, for values of $\pi\varepsilon_2^2 \ll v_1 + v_2$, the second cavity would be of the form $B \setminus B_1$ for some balls B_1 and B such that $B_1 \subset B$, $|B_1| = v_1$, and $|B| = v_1 + v_2$. In other words, \mathbf{u} must create “moon-shaped” cavities, which cannot be obtained if \mathbf{u} is angle-preserving.

In the second place, the interaction term in the lower bound vanishes as $\frac{v_2}{v_1} \rightarrow 0$ regardless of whether the minimum is achieved at $\log \frac{R}{d}$ or at $\log \frac{d}{\varepsilon}$, whereas in the upper bound this vanishing effect is obtained only for the case of distorted cavities (when $\log \frac{d}{\varepsilon}$ is the smallest). This is because when $\delta = 0$ and $v_1 \gg v_2$, the circular sector³ $\{\mathbf{a}^* + de^{i\theta}, \theta \in (\frac{\pi}{2}, \frac{3\pi}{2})\}$ is mapped to a curve $\lambda\mathbf{a}^* + f(\varphi)e^{i\varphi}$ with polar angles φ ranging almost from 0 to 2π . This “angular distortion” necessarily produces a strict inequality in (1.9), so in principle it could be possible to quantify its effect in the lower bound. It is not clear, however, whether for a minimizer an interaction term of the form $(v_1 + v_2) \log \frac{R}{d}$ will always be present (in the case when $\frac{v_2}{v_1} \rightarrow 0$), or if the fact that such a term appears in the upper bound is a limitation of the method used for the explicit constructions.

Finally, the factor $\frac{v_2}{v_1}$ in front of $\log \frac{d}{\varepsilon_1}$ and $\log \frac{d}{\varepsilon_2}$ is raised to a different exponent in each term, the reason being that Ω_1 and Ω_2 play different roles in the upper bound construction. Provided $\delta > 0$, when $\frac{v_2}{v_1} \rightarrow 0$ the first subdomain is becoming more and more like a circle (its height and its width tend to be equal, and the distortion of the first cavity tends to vanish) whereas Ω_2 becomes

¹When considering boundary conditions, not all values of δ can be chosen, see the discussion below.

²we assume, e.g., that $v_1 + v_2 < 4\pi R^2$, in order to illustrate the main point

³we state this in two dimensions for simplicity

increasingly distorted (the ratio between its height and its width tends to infinity). The factor $2^n \sqrt{\frac{v_2}{v_1}}$ in front of $\log \frac{d}{\varepsilon_2}$ is only due to the fact that the effect in the energy of the distortion of the cavities also depends on the size of the cavity.

Dirichlet boundary conditions

If we want our maps to satisfy specific Dirichlet boundary conditions, then they need to be “completed” outside of the ball $B(\mathbf{a}^*, R)$ of the previous theorem. For that we use the method of Rivière and Ye, and show how to obtain explicit Dirichlet energy estimates from it. We consider the radially symmetric loading of a ball, but other boundary conditions could also be handled. Let \mathbf{a}^* , δ , ρ_1 , ρ_2 , Ω_1 , Ω_2 be as before. We are to find R_1 , R_2 , and an incompressible diffeomorphism $\mathbf{u} : \{R_1 < |\mathbf{x} - \mathbf{a}^*| < R_2\} \rightarrow \mathbb{R}^n$ such that

- i) $\Omega_1 \cup \Omega_2 \subset B(\mathbf{a}^*, R_1)$ and $\mathbf{u}|_{\partial B(\mathbf{a}^*, R_1)}$ coincides with the map of Theorem 2
- ii) $\mathbf{u}|_{\partial B(\mathbf{a}^*, R_2)}$ is radially symmetric.




Figure 6: Transition to a radially symmetric map. A larger initial domain is necessary in order to create spherical cavities. Parameters: $\Omega = B(\mathbf{0}, R_2)$, $\sqrt{\frac{v_1 + v_2}{\pi d^2}} = 1.5$, $\frac{v_2}{v_1} = 0.3$, $d = 1$, $R_1 \approx d$.

Not all values of R_1 and R_2 are suitable for the existence of a solution, since the reference configuration $\{R_1 \leq |\mathbf{x} - \mathbf{a}^*| \leq R_2\}$ must contain enough material to fill the space between $\mathbf{u}(\partial B(\mathbf{a}^*, R_2))$ (with shape prescribed by the Dirichlet data) and $\mathbf{u}(\partial B(\mathbf{a}^*, R_1))$ (whose shape is determined by Theorem 2, see Figure 6). In the case of a radially symmetric loading, the farther $\Omega_1 \cup \Omega_2$ is from being a ball, the larger the reference configuration has to be. If $\delta = 1$ nothing has to be imposed; if $\delta < 1$, we must have that

$$\omega_n(R_2^n - R_1^n) \geq C(v_1 + v_2)(1 - \delta)$$

for some constant C (see Lemma 4.5). It turns out that the above necessary condition is also sufficient, as we show in the following theorem:

Theorem 3. *Suppose that $\mathbf{a}_1, \mathbf{a}_2 \in \mathbb{R}^n$ and $d := |\mathbf{a}_1 - \mathbf{a}_2| > \varepsilon_1 + \varepsilon_2$. Let $\delta \in [0, 1]$, $v_1 \geq v_2 \geq 0$,*

$$V_\delta := 2^{2n+1}n(v_1 + v_2)(1 - \delta), \quad R_1 \geq \max \left\{ \sqrt[n]{\frac{V_\delta}{\omega_n}}, 2d \right\}, \quad R_2 := \sqrt[n]{R_1^n + \frac{V_\delta}{\omega_n}}. \quad (1.15)$$

Then there exists \mathbf{a}^ in the segment joining \mathbf{a}_1 and \mathbf{a}_2 and a piecewise smooth homeomorphism $\mathbf{u} \in W^{1,\infty}(\mathbb{R}^n \setminus \{\mathbf{a}_1, \mathbf{a}_2\}, \mathbb{R}^n)$ such that $\text{Det } D\mathbf{u} = \mathcal{L}^n + v_1\delta_{\mathbf{a}_1} + v_2\delta_{\mathbf{a}_2}$ in \mathbb{R}^n , $\mathbf{u}|_{\mathbb{R}^n \setminus B(\mathbf{a}^*, R_2)}$ is radially symmetric, and for all $R \geq R_1$*

$$\begin{aligned} \frac{1}{n} \int_{B(\mathbf{a}^*, R) \setminus (B_{\varepsilon_1}(\mathbf{a}_1) \cup B_{\varepsilon_2}(\mathbf{a}_2))} \left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|^n \mathrm{d}\mathbf{x} &\leq C_1(v_1 + v_2 + \omega_n R^n) + v_1 \log \frac{R}{\varepsilon_1} + v_2 \log \frac{R}{\varepsilon_2} \\ &+ C_2(v_1 + v_2) \left((1 - \delta) \left(\log \sqrt[n]{\frac{V_\delta}{\omega_n d^n}} \right)_+ + \delta \left(\sqrt[n]{\frac{v_2}{v_1}} \log \frac{d}{\varepsilon_1} + \sqrt[2n]{\frac{v_2}{v_1}} \log \frac{d}{\varepsilon_2} \right) \right). \end{aligned}$$

The main differences with respect to Theorem 2 are that \mathbf{u} is now radially symmetric in $\mathbb{R}^n \setminus B(\mathbf{a}^*, R_2)$ and that $\log \frac{R}{d}$ has been replaced with $\log \sqrt[n]{\frac{V_\delta}{\omega_n d^n}} = C + \log \sqrt[n]{\frac{(v_1 + v_2)(1 - \delta)}{\omega_n d^n}}$ in the interaction term. The proof is presented in Section 4.2. As a consequence we finally obtain

Corollary 1. *Let Ω be a ball of radius $R \geq 2d$, with $d > \varepsilon_1 + \varepsilon_2 > 0$. Then, for every $v_1 \geq v_2 \geq 0$ there exist $\mathbf{a}_1, \mathbf{a}_2 \in \Omega$ with $|\mathbf{a}_1 - \mathbf{a}_2| = d$, and a Lipschitz homeomorphism $\mathbf{u} : \Omega \setminus \{\mathbf{a}_1, \mathbf{a}_2\} \rightarrow \mathbb{R}^n$, such that $\text{Det } D\mathbf{u} = \mathcal{L}^n + v_1\delta_{\mathbf{a}_1} + v_2\delta_{\mathbf{a}_2}$ in Ω , $\mathbf{u}|_{\partial\Omega} \equiv \lambda \mathbf{id}$ (with $\lambda^n - 1 := \frac{v_1 + v_2}{|\Omega|}$), and*

$$\begin{aligned} \frac{1}{n} \int_{\Omega \setminus (B_{\varepsilon_1}(\mathbf{a}_1) \cup B_{\varepsilon_2}(\mathbf{a}_2))} \left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|^n \mathrm{d}\mathbf{x} &\leq C_1(v_1 + v_2 + \omega_n R^n) + v_1 \log \frac{R}{\varepsilon_1} + v_2 \log \frac{R}{\varepsilon_2} \\ &+ C_2(v_1 + v_2) \min_{\delta \in [\delta_0, 1]} \left((1 - \delta) \left(\log \frac{(v_1 + v_2)(1 - \delta)}{\omega_n d^n} \right)_+ + \delta \left(\sqrt[n]{\frac{v_2}{v_1}} \log \frac{d}{\varepsilon_1} + \sqrt[2n]{\frac{v_2}{v_1}} \log \frac{d}{\varepsilon_2} \right) \right) \end{aligned}$$

with $\delta_0 := \max \left\{ 0, 1 - \frac{|\Omega| - 2^n \omega_n d^n}{4^{n+1} n \omega_n d^n} \right\}$.

The value of δ_0 is such that $\delta \geq \delta_0$ if and only if $\omega_n R^n \geq \omega_n R_1^n + V_\delta$, with $\omega_n R_1^n := V_\delta + \omega_n (2d)^n$; the idea is to be able to use Theorem 3 and obtain a final energy estimate depending only on $v_1, v_2, d, \varepsilon_1, \varepsilon_2$ and the size $|\Omega|$ of the domain.

1.4 Convergence results

Once we have upper and lower bounds, we are able to show that for “almost-minimizers” one of the three scenarii described after Theorem 1 holds in the limit $\varepsilon \rightarrow 0$.

Theorem 4. *Let Ω be an open and bounded set in \mathbb{R}^n , $n \geq 2$. Let $\varepsilon_j \rightarrow 0$ be a sequence, that we will denote in the sequel simply by ε . Let $\{\Omega_\varepsilon\}_\varepsilon$ be a corresponding sequence of domains of the form $\Omega_\varepsilon = \Omega \setminus \bigcup_{i=1}^m \overline{B}_\varepsilon(\mathbf{a}_{i,\varepsilon})$, with $m \in \mathbb{N}$, $\mathbf{a}_{1,\varepsilon}, \dots, \mathbf{a}_{m,\varepsilon} \in \Omega$ and ε such that the balls $B_\varepsilon(\mathbf{a}_{1,\varepsilon}), \dots, B_\varepsilon(\mathbf{a}_{m,\varepsilon})$ are disjoint. Assume that for each $i = 1, \dots, m$ the sequence $\{\mathbf{a}_{i,\varepsilon}\}_\varepsilon$ is compactly contained in Ω . Suppose, further, that there exists $\mathbf{u}_\varepsilon \in W^{1,n}(\Omega_\varepsilon, \mathbb{R}^n)$ satisfying condition INV, $\text{Det } D\mathbf{u}_\varepsilon = \mathcal{L}^n$ in Ω_ε , $\sup_\varepsilon \|\mathbf{u}_\varepsilon\|_{L^\infty(\Omega_\varepsilon)} < \infty$ and*

$$\frac{1}{n} \int_{\Omega_\varepsilon} \left| \frac{D\mathbf{u}_\varepsilon(\mathbf{x})}{\sqrt{n-1}} \right|^n d\mathbf{x} \leq \left(\sum_{i=1}^m v_{i,\varepsilon} \right) \log \frac{\text{diam } \Omega}{\varepsilon} + C \left(|\Omega| + \sum_{i=1}^m v_{i,\varepsilon} \right), \quad (1.16)$$

where⁴ $v_{i,\varepsilon} := |E(\mathbf{a}_{i,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon)| - \omega_n \varepsilon^n$ and C is a universal constant.

Then (extracting a subsequence) the limits $\mathbf{a}_i = \lim_{\varepsilon \rightarrow 0} \mathbf{a}_{i,\varepsilon}$ and $v_i = \lim_{\varepsilon \rightarrow 0} v_{i,\varepsilon}$, $i = 1, \dots, m$ are well defined, and there exists $\mathbf{u} \in \bigcap_{1 \leq p < n} W^{1,p}(\Omega, \mathbb{R}^n) \cap W_{\text{loc}}^{1,n}(\Omega \setminus \{\mathbf{a}_1, \dots, \mathbf{a}_m\}, \mathbb{R}^n)$ such that

- $\mathbf{u}_\varepsilon \rightharpoonup \mathbf{u}$ in $W_{\text{loc}}^{1,n}(\Omega \setminus \{\mathbf{a}_1, \dots, \mathbf{a}_m\}, \mathbb{R}^n)$
- $\text{Det } D\mathbf{u}_\varepsilon \xrightarrow{*} \text{Det } D\mathbf{u}$ in $\Omega \setminus \{\mathbf{a}_1, \dots, \mathbf{a}_m\}$ locally in the sense of measures
- $\text{Det } D\mathbf{u} = \sum_{i=1}^m v_i \delta_{\mathbf{a}_i} + \mathcal{L}^n$ in Ω .

When $m = 2$, one of the following holds:

i) if $\mathbf{a}_1 \neq \mathbf{a}_2$ and $v_1, v_2 > 0$ (assume without l.o.g. $v_1 \geq v_2$), then

- the cavities $\text{im}_T(\mathbf{u}, \mathbf{a}_1)$ and $\text{im}_T(\mathbf{u}, \mathbf{a}_2)$ (as defined in (2.3)) are balls of volume v_1, v_2
- $|E(\mathbf{a}_{i,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon) \Delta \text{im}_T(\mathbf{u}, \mathbf{a}_i)| \rightarrow 0$ as $\varepsilon \rightarrow 0$ for $i = 1, 2$
- under the additional assumption that $v_1 + v_2 < 2^n \omega_n (\text{dist}(\frac{\mathbf{a}_1 + \mathbf{a}_2}{2}, \partial\Omega))^n$,

$$\frac{\omega_n |\mathbf{a}_2 - \mathbf{a}_1|^n}{v_1 + v_2} \geq C_1 \exp \left(-C_2 \left(1 + \frac{|\Omega|}{v_1 + v_2} + \log \frac{\omega_n (\text{diam } \Omega)^n}{v_1 + v_2} \right) \right) / \left(\frac{v_2}{v_1 + v_2} \right)^{\frac{n}{n-1}}$$

for some universal constants C_1 and C_2 depending only on n ;

ii) if $\min\{v_1, v_2\} = 0$ (say $v_2 = 0$), then $\text{im}_T(\mathbf{u}, \mathbf{a}_1)$ (the only cavity opened by \mathbf{u}) is spherical;

iii) if $\mathbf{a}_1 = \mathbf{a}_2$ and $v_1, v_2 > 0$ (assume $v_1 \geq v_2$), then

- $\text{im}_T(\mathbf{u}, \mathbf{a}_1)$ is a ball of volume $v_1 + v_2$
- $|\mathbf{a}_{2,\varepsilon} - \mathbf{a}_{1,\varepsilon}| = O(\varepsilon)$ as $\varepsilon \rightarrow 0$

⁴Now we write $E(\mathbf{a}_{i,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon)$, and not just $E(\mathbf{a}_{i,\varepsilon}, \varepsilon)$, to highlight the dependence on \mathbf{u}_ε . It corresponds to the cavity opened by \mathbf{u}_ε at $\mathbf{a}_{i,\varepsilon}$ (compare with (1.10) and (2.3)).

- *the cavities must be distorted in the following sense (C_n being as in Proposition 1.2):*

$$\liminf_{\varepsilon \rightarrow 0} \frac{v_1 D(E(\mathbf{a}_{1,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon))^{\frac{n}{n-1}} + v_2 D(E(\mathbf{a}_{2,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon))^{\frac{n}{n-1}}}{v_1 + v_2} > C_n \left(\frac{v_2}{v_1 + v_2} \right)^{\frac{n}{n-1}}. \quad (1.17)$$

In the situation of two cavities, the three cases above correspond to the three scenarii of the end of Section 1.2 in the same order.

The main ingredients for the proof are the comparison of the upper bound (1.16) with the lower bounds Proposition 1.1 and Theorem 1, standard compactness arguments, and an argument introduced by Struwe [77] in the context of Ginzburg-Landau which allows to deduce from the energy bounds sufficient compactness of \mathbf{u}_ε .

1.5 Additional comments and remarks

We note first that our analysis works provided that the distance of the cavitation points to the boundary does not get small (thus the domain cannot be too thin either). It is an interesting question to better understand what happens when they do get close to the boundary, as well as the effect of the boundary conditions.

Second, it follows from our work that it is always necessary to compare quantities in the reference configuration with quantities in the deformed configuration, due to the scale-invariance in elasticity. For example, we have shown that a large price needs to be paid (in terms of elastic energy) in order to open spherical cavities whenever the distance between the cavitation points is small *compared to the final size of the cavities* ($\omega_n d^n \ll v_1 + v_2$). If we only know that the cavitation points are becoming closer and closer to each other, from this alone we cannot conclude that the cavities will interact and that the total elastic energy will go to infinity, as the following argument shows. Suppose that \mathbf{u} is an incompressible map defined on the unit cube $Q \subset \mathbb{R}^n$, opening a cavity, and satisfying affine boundary conditions of the form $\mathbf{u}(\mathbf{x}) \equiv \mathbf{A}\mathbf{x}$ on ∂Q , $\mathbf{A} \in \mathbb{R}^{n \times n}$. Then, by rescaling \mathbf{u} and reproducing it periodically, it is possible to construct a sequence of incompressible maps creating an increasingly large number of cavities, at cavitation points that are closer and closer to each other, in such a way that all the deformations in the sequence have exactly the same elastic energy (cf. Ball & Murat [7]; see also [60, 49, 50]). This is possible because the cavities themselves are also becoming increasingly smaller, with radii decaying at the same rate as the distance between neighbouring cavitation points. This example also shows that the strategy of filling the material with an arbitrarily large number of small cavities is, in a sense, equivalent to forming a single big cavity (there is no interaction between the singularities). Here we complement that result by showing that if it is not possible to create an infinite number of cavities, then the interaction effects in the energy do become noticeable, and under some circumstances can even be quantified.

Third, we mention that the idea of partitioning the domain and using angle-preserving maps inside the resulting subdomains (as described in Section 1.3) can be used to produce test maps that are incompressible and open any prescribed number of cavities (for example by dividing the initial domain in angular sectors, as in Figure 7). The relative size of the cavities can be controlled by specifying the volume ratios of the subdomains in the partition; the cavity shapes will also be




Figure 7: Incompressible maps creating multiple cavities of arbitrary sizes.

determined by the shape of those subdomains. The deformations thus constructed may be relevant for future work on the subject, for instance when obtaining energy estimates.

Finally, we discuss the case $p \neq n$. It is not clear how to extend the analysis to this case, the main reason being that the energy is no longer conformally invariant while the “ball-construction method” is only suited for such cases. To see this in a simple way, let us consider the case of two cavities, assuming incompressibility, letting $\varepsilon_1 = \varepsilon_2 \rightarrow 0$, and let us try to reproduce the steps (1.8) and (1.11) with (1.14). The p -equivalent of (1.14) obtained by Hölder’s inequality (and by relating $|D\mathbf{u}|^{n-1}$ to the area element $|(\text{cof } D\mathbf{u})\boldsymbol{\nu}|$, see Lemma 3.1) is

$$\int_{\partial B(\mathbf{a}, r)} \left| \frac{D\mathbf{u}(\mathbf{x})}{\sqrt{n-1}} \right|^p d\mathcal{H}^{n-1}(\mathbf{x}) \geq \frac{\text{Per}(E(\mathbf{a}, r))^{\frac{p}{n-1}}}{(n\omega_n r^{n-1})^{\frac{p}{n-1}-1}} \geq n\omega_n^{\frac{n-p}{n}} \frac{|E(\mathbf{a}, r)|^{\frac{p}{n}}}{r^{1-(n-p)}} \left(1 + CD(E(\mathbf{a}, r))^{\frac{p}{n-1}} \right).$$

According to this, when $p \neq n$ we may bound from below the energy in $B(\mathbf{a}_1, \frac{d}{2}) \cup B(\mathbf{a}_2, \frac{d}{2})$ (with $d = |\mathbf{a}_2 - \mathbf{a}_1|$) by

$$\int_{B(\mathbf{a}_1, \frac{d}{2}) \cup B(\mathbf{a}_2, \frac{d}{2})} \frac{\omega_n^{\frac{p-n}{n}}}{n} \left| \frac{D\mathbf{u}(\mathbf{x})}{\sqrt{n-1}} \right|^p \geq \left(v_1^{\frac{p}{n}} + v_2^{\frac{p}{n}} \right) \left(\frac{d}{2} \right)^{n-p} + C(v_1 + v_2)^{\frac{p}{n}} \int_0^{\frac{d}{2}} \langle D(E(\mathbf{a}_i, r))^{\frac{p}{n-1}} \rangle r^{n-p-1},$$

where $\langle D(E(\mathbf{a}_i, r))^{\frac{p}{n-1}} \rangle$ stands for the average distortion

$$\langle D(E(\mathbf{a}_i, r))^{\frac{p}{n-1}} \rangle := \left(v_1^{\frac{p}{n}} D(E(\mathbf{a}_1, r))^{\frac{p}{n-1}} + v_2^{\frac{p}{n}} D(E(\mathbf{a}_2, r))^{\frac{p}{n-1}} \right) (v_1 + v_2)^{-\frac{p}{n}}.$$

Analogously, we can bound the energy in $B(\mathbf{a}, R) \setminus \bar{B}(\mathbf{a}, d)$ (with $\mathbf{a} = \frac{\mathbf{a}_1 + \mathbf{a}_2}{2}$) by

$$\int_{B(\mathbf{a}, R) \setminus \bar{B}(\mathbf{a}, d)} \frac{\omega_n^{\frac{p-n}{n}}}{n} \left| \frac{D\mathbf{u}(\mathbf{x})}{\sqrt{n-1}} \right|^p \geq (v_1 + v_2)^{\frac{p}{n}} \int_d^R r^{n-p-1} + C(v_1 + v_2)^{\frac{p}{n}} \int_d^R D(E(\mathbf{a}, r))^{\frac{p}{n-1}} r^{n-p-1}$$

and obtain:

$$\begin{aligned} \int_{B(\mathbf{a}, R)} \frac{\omega_n^{\frac{p-n}{n}}}{n} \left| \frac{D\mathbf{u}(\mathbf{x})}{\sqrt{n-1}} \right|^p &\geq (v_1 + v_2)^{\frac{p}{n}} \left(\int_0^{\frac{d}{2}} + \int_d^R \right) r^{n-p-1} + \underbrace{\left(v_1^{\frac{p}{n}} + v_2^{\frac{p}{n}} - (v_1 + v_2)^{\frac{p}{n}} \right) \left(\frac{d}{2} \right)^{n-p}}_{II} \\ &+ \underbrace{C(v_1 + v_2)^{\frac{p}{n}} \left[\int_0^{\frac{d}{2}} \langle D(E(\mathbf{a}_i, r))^{\frac{p}{n-1}} \rangle r^{n-p-1} + \int_d^R D(E(\mathbf{a}, r))^{\frac{p}{n-1}} r^{n-p-1} \right]}_{III}. \end{aligned}$$

Assume that $v_1 + v_2$ is fixed (as is the case in the Dirichlet problem). Let us first consider the case $p < n$. Since the limit $\varepsilon \rightarrow 0$ is not singular in this case (contrarily to $p = n$), the problem cannot be analyzed by asymptotic analysis. If we guide ourselves only by the second and third terms (II and III), when $p < n$ we can say the following. The factor $v_1^{\frac{p}{n}} + v_2^{\frac{p}{n}} - (v_1 + v_2)^{\frac{p}{n}}$ in II is minimized when $\min\{v_1, v_2\} = 0$, hence it motivates the creation of just one cavity (the same can be said for the problem with M cavities, because $v_1^{\frac{p}{n}} + \dots + v_M^{\frac{p}{n}}$ is concave and the restriction $v_1 + \dots + v_M = \text{const.}$ is linear). If the above difference has to be positive, the factor $\left(\frac{d}{2}\right)^n$ suggests that the two cavitation points would want to be arbitrarily close, and that the cavities will tend to act as a single cavity. This is consistent with the prediction for III; indeed, consider the corresponding estimate for $p = n$:

$$\begin{aligned} \frac{1}{n} \int_{\Omega_\varepsilon \cap B(\mathbf{a}, R)} \left| \frac{D\mathbf{u}(\mathbf{x})}{\sqrt{n-1}} \right|^n d\mathbf{x} &\geq (v_1 + v_2) \left(\int_\varepsilon^{\frac{d}{2}} + \int_d^R \right) \frac{dr}{r} \\ &+ C(v_1 + v_2) \left[\int_\varepsilon^{\frac{d}{2}} \langle D(E(\mathbf{a}_i, r))^{\frac{n}{n-1}} \rangle \frac{dr}{r} + \int_d^R D(E(\mathbf{a}, r))^{\frac{n}{n-1}} \frac{dr}{r} \right]. \end{aligned}$$

Under a logarithmic cost, it is much more important to minimize the distortions $D(E(\mathbf{a}_i, r))$ of the circles $\mathbf{u}(\partial B(\mathbf{a}_i, r))$, $i = 1, 2$, $\varepsilon < r < \frac{d}{2}$ near the cavities, rather than the distortion of the outer circles $D(E(\mathbf{a}, r))$, $r > d$. As was discussed before, this leads either to the case of well-separated and spherical cavities (scenario (i) in p. 9), or to the conclusion that if outer circles are mapped to circles (scenario (iii)) then the distance between cavitation points must be of order ε (Theorem 4iii). In contrast, When $p < n$, in the presence of the weight r^{n-p-1} , minimizing the distortions $D(E(\mathbf{a}, r))$, $r > d$ gains more relevance compared to the distortion near the cavities.

For the previous reasons, we believe that the deformations of scenario (i) will not be global minimizers, instead the body will prefer to open a single cavity. If multiple cavities have to be created, then the cavitation points will try to be close to each other, and the deformation will try to rapidly become radially symmetric. The cavities will be distorted and try to act as a single cavity (as in scenario (iii), which creates a state of strain potentially leading to fracture by coalescence), at distances between the cavitation points that are of order 1 (not of order ε). This, in fact, is what has been observed numerically [81, 47].

Let us now turn to $p > n$. The lower bound reads

$$\int_{\Omega_\varepsilon \cap B(\mathbf{a}, R)} \frac{\omega_n^{\frac{p-n}{n}}}{n} \left| \frac{D\mathbf{u}(\mathbf{x})}{\sqrt{n-1}} \right|^p d\mathbf{x} + \frac{(v_1 + v_2)^{\frac{p}{n}} R^{n-p}}{p-n} \geq \underbrace{(v_1^{\frac{p}{n}} + v_2^{\frac{p}{n}}) \int_\varepsilon^{\frac{d}{2}} r^{n-p-1}}_I + \underbrace{(v_1 + v_2)^{\frac{p}{n}} d^{n-p}}_{II} \\ + C(v_1 + v_2)^{\frac{p}{n}} \left[\int_\varepsilon^{\frac{d}{2}} \langle D(E(\mathbf{a}_i, r))^{\frac{p}{n-1}} \rangle r^{n-p-1} + \int_d^R D(E(\mathbf{a}, r))^{\frac{p}{n-1}} r^{n-p-1} \right].$$

This time the limit $\varepsilon \rightarrow 0$ is singular, even more so than for $p = n$. The factor $v_1^{\frac{p}{n}} + v_2^{\frac{p}{n}}$ is now minimized when the cavities have equal volumes. Regarding d , the first term prefers small distances ($d = 2\varepsilon$) while the second prefers $d \rightarrow \infty$; since $(v_1 + v_2)^{\frac{p}{n}} > v_1^{\frac{p}{n}} + v_2^{\frac{p}{n}}$, it can be said that II has a stronger influence, hence d large should be preferred⁵. With respect to the third term, it is now much more vital to create spherical cavities (so as to minimize the first of the two integrals) than when $p = n$. This implies that it is scenario (i), rather than (ii) or (iii), which should be observed.

The case $p < n$, therefore, should favour a single cavity and coalescence, $p > n$ should favour many cavities and splitting, and both situations are possible in the borderline case that we have studied: $p = n$.

1.6 Plan of the paper

In Section 2 we describe our notation and recall the notions of perimeter, reduced boundary, topological image, distributional determinant, and the invertibility condition INV. In Section 3 we begin by extending (1.14) to the case of an arbitrary power p and space dimension n (Lemma 3.1). In Section 3.1 we prove the lower bound for an arbitrary number of cavities using the ball construction method (Proposition 1.1). In Section 3.2, we prove the main lower bound (Theorem 1) and postpone the proof of our estimate on the distortions (Proposition 1.2) to Section 3.3. The energy estimates for the angle-preserving ansatz are presented in Section 4.1 and proved in Section 4.3. In Section 4.2 we show how to complete the maps away from the cavitation points so as to fulfil the boundary conditions, and in Section 4.4 we comment briefly on the numerical computations presented in this paper based on the constructive method of Dacorogna & Moser [20]. Finally, the proof of the main compactness result and of the fact that in the limit only one of the three scenarii holds (Theorem 4) is given in Section 5.

1.7 Acknowledgements

We give special thanks to G. Francfort for his interest and his involvement in this project. We also thank J.-F. Babadjian, J. Ball, Y. Brenier, A. Contreras, R. Kohn, R. Lecaros, G. Mingione, C. Mora-Corral, S. Müller, T. Rivière, and N. Rougerie for useful discussions.

⁵although in order to be sure it would be necessary to compute the energy in the transition region $B(\mathbf{a}, d) \setminus (B(\mathbf{a}_1, \frac{d}{2}) \cup B(\mathbf{a}_2, \frac{d}{2}))$

2 Notation and preliminaries

2.1 General notation

Let n denote the space dimension. Vector-valued and matrix-valued quantities will be written in bold face. The set of unit vectors in \mathbb{R}^n is denoted by \mathbb{S}^{n-1} . Given a set $E \subset \mathbb{R}^n$, $\lambda \geq 0$ and $\mathbf{h} \in \mathbb{R}^n$, we define $\lambda E := \{\lambda \mathbf{x} : \mathbf{x} \in E\}$ and $E + \mathbf{h} := \{\mathbf{x} + \mathbf{h} : \mathbf{x} \in E\}$. The interior and the closure of E are denoted by $\text{Int } E$ and \overline{E} , and the symmetric difference of two sets E_1 and E_2 by $E_1 \triangle E_2$. If E_1 is compactly contained in E_2 , we write $E_1 \subset\subset E_2$. The notations $B(\mathbf{x}, R)$, $B_R(\mathbf{x})$ are used for the open ball of radius R centred at \mathbf{x} , and $\overline{B}(\mathbf{a}, R)$, $\overline{B}_R(\mathbf{a})$ for the corresponding closed ball. The distance from a point \mathbf{x} to a set E is denoted by $\text{dist}(\mathbf{x}, E)$, the distance between sets by $\text{dist}(E_1, E_2)$, and the diameter of a set by $\text{diam } E$.

Given \mathbf{A} an $n \times n$ matrix, \mathbf{A}^T will be its transpose, $\det \mathbf{A}$ its determinant, and $\text{cof } \mathbf{A}$ its cofactor matrix (defined by $\mathbf{A}^T \text{cof } \mathbf{A} = (\det \mathbf{A})\mathbf{1}$, where $\mathbf{1}$ stands for the $n \times n$ identity matrix). The adjugate matrix of \mathbf{A} is $\text{adj } \mathbf{A} = (\text{cof } \mathbf{A})^T$.

The Lebesgue and the k -dimensional Hausdorff measure are denoted by \mathcal{L}^n and \mathcal{H}^k , respectively. If E is a measurable set, $\mathcal{L}^n(E)$ is also written $|E|$ (as well as $|I|$ for the length of an interval I). The measure of the k -dimensional unit ball is ω_k (accordingly, $\mathcal{H}^{n-1}(\partial B(\mathbf{x}, r)) = n\omega_n r^{n-1}$). The exterior product of $1 \leq k \leq n$ vectors $\mathbf{a}_1, \dots, \mathbf{a}_k \in \mathbb{R}^n$ is denoted by $\mathbf{a}_1 \wedge \dots \wedge \mathbf{a}_k$ or $\bigwedge_{i=1}^k \mathbf{a}_i$. It is k -linear, antisymmetric, and such that $|\mathbf{a}_1 \wedge \dots \wedge \mathbf{a}_k|$ is the k -dimensional measure of the k -prism formed by $\mathbf{a}_1, \dots, \mathbf{a}_k$ (see, e.g., [24, 75, 33, 1]). In particular, $|\mathbf{x}|^2 = |\mathbf{x} \cdot \mathbf{e}|^2 + |\mathbf{x} \wedge \mathbf{e}|^2$ for all $\mathbf{x} \in \mathbb{R}^n$ and $\mathbf{e} \in \mathbb{S}^{n-1}$. With a slight abuse of notation, when $k = n$ the expression $\mathbf{a}_1 \wedge \dots \wedge \mathbf{a}_n$ is used to denote the determinant (in the standard basis) of the matrix with column vectors $\mathbf{a}_1, \dots, \mathbf{a}_n \in \mathbb{R}^n$.

The characteristic function of a set E is referred to as χ_E , and the restriction of \mathbf{u} to E as $\mathbf{u}|_E$. The sign function $\text{sgn} : \mathbb{R} \rightarrow \{-1, 0, 1\}$ is given by $\text{sgn } x = x/|x|$ if $x \neq 0$, $\text{sgn } 0 = 0$. The notation \mathbf{id} is used for the identity function $\mathbf{id}(\mathbf{x}) \equiv \mathbf{x}$. The symbol f_E stands for the integral average $\frac{1}{|E|} \int_E f$. The support of a function f is represented by $\text{spt } f$.

The space of infinitely differentiable functions with compact support is denoted by $C_c^\infty(\Omega)$, and the L^p norm of a function f by $\|f\|_{L^p}$. Sobolev spaces are denoted by $W^{1,p}(\Omega, \mathbb{R}^n)$, as usual. The Hilbert space $W^{1,2}(\Omega, \mathbb{R}^n)$ is denoted by $H^1(\Omega, \mathbb{R}^n)$. The weak derivative (the linear transformation) of a map $\mathbf{u} \in W^{1,p}(\Omega, \mathbb{R}^n)$ at a point $\mathbf{x} \in \mathbb{R}^n$ is identified with the gradient $D\mathbf{u}(\mathbf{x})$ (the matrix of weak partial derivatives).

Use will be made of the coarea formula (see, e.g., [24, 23, 1]): if $E \subset \mathbb{R}^n$ is measurable and $\phi : E \rightarrow \mathbb{R}$ is Lipschitz, then for all $f \in L^1(E)$

$$\int_E f(\mathbf{x}) |D\phi(\mathbf{x})| \, d\mathbf{x} = \int_{-\infty}^{\infty} \left(\int_{\{\mathbf{x} \in E : \phi(\mathbf{x}) = t\}} f(\mathbf{x}) \, d\mathcal{H}^{n-1}(\mathbf{x}) \right) dt.$$

2.2 Perimeter and reduced boundary

Definition 2. *The perimeter of a measurable set $E \subset \mathbb{R}^n$ is defined as*

$$\text{Per } E := \sup \left\{ \int_E \text{div } \mathbf{g}(\mathbf{y}) \, d\mathbf{y} : \mathbf{g} \in C_c^1(\mathbb{R}^n, \mathbb{R}^n), \|\mathbf{g}\|_\infty \leq 1 \right\}.$$

Definition 3. Given $\mathbf{y}_0 \in \mathbb{R}^n$ and a non-zero vector $\boldsymbol{\nu} \in \mathbb{R}^n$, we define

$$H^+(\mathbf{y}_0, \boldsymbol{\nu}) := \{\mathbf{y} \in \mathbb{R}^n : (\mathbf{y} - \mathbf{y}_0) \cdot \boldsymbol{\nu} \geq 0\}, \quad H^-(\mathbf{y}_0, \boldsymbol{\nu}) := \{\mathbf{y} \in \mathbb{R}^n : (\mathbf{y} - \mathbf{y}_0) \cdot \boldsymbol{\nu} \leq 0\}.$$

The reduced boundary of a measurable set $E \subset \mathbb{R}^n$, denoted by $\partial^* E$, is defined as the set of points $\mathbf{y} \in \mathbb{R}^n$ for which there exists a unit vector $\boldsymbol{\nu} \in \mathbb{R}^n$ such that

$$\lim_{r \rightarrow 0^+} \frac{|E \cap H^-(\mathbf{y}, \boldsymbol{\nu}) \cap B(\mathbf{y}, r)|}{|B(\mathbf{y}, r)|} = \frac{1}{2} \quad \text{and} \quad \lim_{r \rightarrow 0^+} \frac{|E \cap H^+(\mathbf{y}, \boldsymbol{\nu}) \cap B(\mathbf{y}, r)|}{|B(\mathbf{y}, r)|} = 0.$$

If $\mathbf{y} \in \partial^* E$ then $\boldsymbol{\nu}$ is uniquely determined and is called the unit outward normal to E .

The definition of perimeter coincides precisely with the \mathcal{H}^{n-1} -measure of the reduced boundary, as follows from the well-known results of Federer, Fleming and De Giorgi (see, e.g., [24, 83, 23, 1])⁶.

2.3 Degree and topological image

We begin by recalling the notion of topological degree for maps \mathbf{u} that are only weakly differentiable [56, 26, 12, 17].

If $\mathbf{u} \in W^{1,p}(\Omega, \mathbb{R}^n)$ and $\mathbf{x} \in \mathbb{R}^n$, then, for a.e. $r \in (0, \infty)$ with $\partial B(\mathbf{x}, r) \subset \Omega$,

(R1) $\mathbf{u}(\mathbf{z})$ and $D\mathbf{u}(\mathbf{z})$ are defined at \mathcal{H}^{n-1} -a.e. $\mathbf{z} \in \partial B(\mathbf{x}, r)$

(R2) $\mathbf{u}|_{\partial B(\mathbf{x}, r)} \in W^{1,p}(\partial B(\mathbf{x}, r), \mathbb{R}^n)$

(R3) $D(\mathbf{u}|_{\partial B(\mathbf{x}, r)})(\mathbf{z}) = (D\mathbf{u}(\mathbf{z}))|_{T_{\mathbf{z}}(\partial B(\mathbf{x}, r))}$ (the n -dimensional and the tangential weak derivatives coincide; $T_{\mathbf{z}}(\partial B(\mathbf{x}, r))$ denotes the tangent plane) for \mathcal{H}^{n-1} -a.e. $\mathbf{z} \in \partial B(\mathbf{x}, r)$

(this follows by approximating by C^∞ maps and using the coarea formula). If, moreover, $p > n - 1$, then, by Morrey's inequality, there exists a unique map $\bar{\mathbf{u}} \in C^0(\partial B(\mathbf{x}, r))$ that coincides with $\mathbf{u}|_{\partial B(\mathbf{x}, r)}$ \mathcal{H}^{n-1} -a.e. With an abuse of notation we write $\mathbf{u}(\partial B(\mathbf{x}, r))$ to denote $\bar{\mathbf{u}}(\partial B(\mathbf{x}, r))$.

If $p > n - 1$ and (R2) is satisfied, for every $\mathbf{y} \in \mathbb{R}^n \setminus \mathbf{u}(\partial B(\mathbf{x}, r))$ we define $\deg(\mathbf{u}, \partial B(\mathbf{x}, r), \mathbf{y})$ as the classical Brouwer degree [68, 26] of $\mathbf{u}|_{\partial B(\mathbf{x}, r)}$ with respect to \mathbf{y} . The degree $\deg(\mathbf{u}, \partial B(\mathbf{x}, r), \cdot)$ is the only $L^1(\mathbb{R}^n)$ map [56, 12] such that

$$\int_{\mathbb{R}^n} \deg(\mathbf{u}, \partial B(\mathbf{x}, r), \mathbf{y}) \operatorname{div} \mathbf{g}(\mathbf{y}) \, d\mathbf{y} = \int_{\partial B(\mathbf{x}, r)} \mathbf{g}(\mathbf{u}(\mathbf{z})) \cdot (\operatorname{cof} D\mathbf{u}(\mathbf{z})) \boldsymbol{\nu}(\mathbf{z}) \, d\mathcal{H}^{n-1}(\mathbf{z}) \quad (2.1)$$

for every $\mathbf{g} \in C^1(\mathbb{R}^n, \mathbb{R}^n)$, $\boldsymbol{\nu}(\mathbf{z})$ being the outward unit normal to $\partial B(\mathbf{x}, r)$.

For a map $\mathbf{u} \in W^{1,p}(\Omega, \mathbb{R}^n)$ that is invertible, orientation-preserving, and regular except for the creation of a finite number of cavities, $\deg(\mathbf{u}, \partial B(\mathbf{x}, r), \mathbf{y})$ is equal to 1, roughly speaking, only at those points \mathbf{y} enclosed by $\mathbf{u}(\partial B(\mathbf{x}, r))$. Because of this, the degree is useful for the study of cavitation, since we can detect a cavity by looking at the set of points where the degree is 1, but which do not belong to the image of \mathbf{u} (they are not part of the deformed body). This gave rise to Šverák's notion of topological image [79].

⁶When $\operatorname{Per} E = \infty$, the result is true if we consider the *measure-theoretic* boundary, as defined in [23, Th. 5.11.1]. For sets of finite perimeter the two notions of boundary coincide \mathcal{H}^{n-1} -a.e., thanks to a result of Federer [24] (also available in [1, Th. 3.61], [23, Lemma 5.8.1], or [83, Sect. 5.6]).

Definition 4. Let $\mathbf{u} \in W^{1,p}(\partial B(\mathbf{x}, r), \mathbb{R}^n)$ for some $\mathbf{x} \in \mathbb{R}^n$, $r > 0$, and $p > n - 1$. Then

$$\text{im}_{\mathbb{T}}(\mathbf{u}, B(\mathbf{x}, r)) := \{\mathbf{y} \in \mathbb{R}^n : \deg(\mathbf{u}, \partial B(\mathbf{x}, r), \mathbf{y}) \neq 0\}.$$

It was pointed out by Müller-Spector [55, Sect. 11] that Sobolev maps may create cavities in some part of the body, and subsequently fill them with material from somewhere else (even if they are one-to-one a.e. [3]). In order to avoid this pathological behaviour, they defined a stronger invertibility condition, based on the topological image⁷.

Definition 5. Let $\mathbf{u} \in W^{1,p}(\Omega, \mathbb{R}^n)$ with $p > n - 1$. We say that \mathbf{u} satisfies condition INV if

i) $\mathbf{u}(\mathbf{z}) \in \text{im}_{\mathbb{T}}(\mathbf{u}, B(\mathbf{x}, r))$ for a.e. $\mathbf{z} \in B(\mathbf{x}, r) \cap \Omega$

ii) $\mathbf{u}(\mathbf{z}) \in \mathbb{R}^n \setminus \text{im}_{\mathbb{T}}(\mathbf{u}, B(\mathbf{x}, r))$ for a.e. $\mathbf{z} \in \Omega \setminus B(\mathbf{x}, r)$

for every $\mathbf{x} \in \mathbb{R}^n$ and a.e. $r \in (0, \infty)$ such that $\mathbf{u}|_{\partial B(\mathbf{x}, r)} \in W^{1,p}(\partial B(\mathbf{x}, r), \mathbb{R}^n)$.

In the following proposition we summarize some of the main virtues of condition INV. We add a sketch of the proof to make it easier for the interested reader to compile the different ideas and conciliate the different notation in [79], [55, Lemmas 2.5, 3.5 and 7.3], [17, Lemmas 3.8 and 3.10], [39, Lemma 2], and [40, Prop. 6 and Lemma 15].

Proposition 2.1. Let $\mathbf{u} \in W^{1,p}(\Omega, \mathbb{R}^n)$ with $p > n - 1$ satisfy $\det D\mathbf{u} > 0$ a.e. and condition INV. Then, for every $\mathbf{x} \in \mathbb{R}^n$ there exists a full- \mathcal{L}^1 -measure subset $R_{\mathbf{x}}$ of $\{r \in (0, \infty) : \partial B(\mathbf{x}, r) \subset \Omega\}$ for which (R1)–(R3), conditions i)–ii) of Definition 5, and the following properties are satisfied:

i) $\deg(\mathbf{u}, \partial B(\mathbf{x}, r), \mathbf{y}) \in \{0, 1\}$ for every $\mathbf{y} \in \mathbb{R}^n \setminus \mathbf{u}(\partial B(\mathbf{x}, r))$

ii) $\partial^* \text{im}_{\mathbb{T}}(\mathbf{u}, B(\mathbf{x}, r)) = \mathbf{u}(\partial B(\mathbf{x}, r))$ up to \mathcal{H}^{n-1} -null sets

iii) $\text{Per}(\text{im}_{\mathbb{T}}(\mathbf{u}, B(\mathbf{x}, r))) = \int_{\partial B(\mathbf{x}, r)} |(\text{cof } D\mathbf{u}(\mathbf{z}))\boldsymbol{\nu}(\mathbf{z})| d\mathcal{H}^{n-1}(\mathbf{z})$

iv) $|\text{im}_{\mathbb{T}}(\mathbf{u}, B(\mathbf{x}, r))| = \frac{1}{n} \int_{\partial B(\mathbf{x}, r)} \mathbf{u}(\mathbf{z}) \cdot (\text{cof } D\mathbf{u}(\mathbf{z}))\boldsymbol{\nu}(\mathbf{z}) d\mathcal{H}^{n-1}(\mathbf{z})$.

Moreover, for every $\mathbf{x}, \mathbf{x}' \in \mathbb{R}^n$ and every $r \in R_{\mathbf{x}}$, $r' \in R_{\mathbf{x}'}$

v) $\text{im}_{\mathbb{T}}(\mathbf{u}, B(\mathbf{x}, r)) \subset \text{im}_{\mathbb{T}}(\mathbf{u}, B(\mathbf{x}', r'))$ if $B(\mathbf{x}, r) \subset B(\mathbf{x}', r')$

vi) $\text{im}_{\mathbb{T}}(\mathbf{u}, B(\mathbf{x}, r)) \cap \text{im}_{\mathbb{T}}(\mathbf{u}, B(\mathbf{x}', r')) = \emptyset$ if $B(\mathbf{x}, r) \cap B(\mathbf{x}', r') = \emptyset$.

⁷The original definition of condition INV in [55, Sect. 3] required that i) and ii) were satisfied only for a.e. $r \in (0, \infty)$ such that $B(\mathbf{x}, r) \subset \Omega$. Here we impose i) and ii) for a.e. $r \in (0, \infty)$ such that $\partial B(\mathbf{x}, r) \subset \Omega$. As explained in [37], this modification is necessary when considering perforated domains, due to Sivaloganathan & Spector's example of leakage between cavities [74, Sect. 6].

Proof. Call Ω_0 the set of $\mathbf{x} \in \Omega$ for which there exist $\mathbf{w} \in C^1(\mathbb{R}^n, \mathbb{R}^n)$ and a compact set $K \subset \Omega$ such that

$$\lim_{r \rightarrow 0^+} \frac{|K \cap B(\mathbf{x}, r)|}{|B(\mathbf{x}, r)|} = 1, \quad \mathbf{u}|_K = \mathbf{w}|_K, \quad \text{and} \quad D\mathbf{u}|_K = D\mathbf{w}|_K. \quad (2.2)$$

Since $\mathbf{u} \in W^{1,p}(\Omega, \mathbb{R}^n)$, it is possible to find (combining Federer's approximation of approximately differentiable maps by Lipschitz functions, Rademacher's theorem, and Whitney's extension theorem, see, e.g., [23, Cor. 6.6.3.2], [24, Thms. 3.1.8 and 3.1.16], [55, Prop. 2.4], [39, Lemma 1]) an increasing sequence of compact sets $\{K_j\}_{j \in \mathbb{N}}$ contained in Ω , and a sequence $\{\mathbf{w}_j\}_{j \in \mathbb{N}}$ of maps in $C^1(\mathbb{R}^n, \mathbb{R}^n)$, such that $\mathbf{u}|_{K_j} = \mathbf{w}_j|_{K_j}$, $\nabla \mathbf{u}|_{K_j} = D\mathbf{w}_j|_{K_j}$, and $|\Omega \setminus K_j| < \frac{1}{j}$ for each $j \in \mathbb{N}$. By Lebesgue's differentiation theorem, $|K_j \setminus K'_j| = 0$ where $K'_j := \{\mathbf{x} \in K_j : \lim_{r \rightarrow 0^+} (r^{-n} |B(\mathbf{x}, r) \setminus K|) = 0\}$. Since $\Omega_0 \supset \bigcup_{j \in \mathbb{N}} K'_j$, it follows that $|\Omega \setminus \Omega_0| = 0$.

Define $R_{\mathbf{x}}$ as the subset of $\{r \in (0, \infty) : \partial B(\mathbf{x}, r) \subset \Omega\}$ for which (R1)–(R3), conditions i)-ii) of Definition 5, and the following properties are satisfied:

$$(R4) \quad \mathcal{H}^{n-1}(\partial B(\mathbf{x}, r) \setminus \Omega_0) = 0$$

$$(R5) \quad \det D\mathbf{u}(\mathbf{z}) > 0 \text{ for } \mathcal{H}^{n-1}\text{-a.e. } \mathbf{z} \in \partial B(\mathbf{x}, r).$$

The fact that $|\{r \in (0, \infty) : \partial B(\mathbf{x}, r) \subset \Omega\} \setminus R_{\mathbf{x}}| = 0$ is a consequence of the coarea formula and of the discussion before Definition 4. For this choice of $R_{\mathbf{x}}$ we have that the properties listed in the proposition are satisfied for all (not only for a.e.) $r \in R_{\mathbf{x}}$. This follows from (2.1), the fact that $\mathbf{u}|_{\Omega_0}$ is one to one (by [55, Lemmas 3.4 and 2.5]; only minor modifications are required, see [39, Lemma 2] if necessary), and a careful inspection of the proofs of [55, Lemmas 2.5, 3.5 and 7.3]. \square

By Proposition 2.1v) the topological image of $B(\mathbf{x}, r)$ can be defined for all $\mathbf{x} \in \mathbb{R}^n$ and all $r \geq 0$ such that $\{\mathbf{z} : r < |\mathbf{z}| < r + \delta\} \subset \Omega$ for some $\delta > 0$ (not only for radii $r \in R_{\mathbf{x}}$). Indeed, since the sequence $\{\text{im}_{\mathbb{T}}(\mathbf{u}, B(\mathbf{x}, r)) : r \in R_{\mathbf{x}}\}$ is increasing for every $\mathbf{x} \in \mathbb{R}^n$, we may define

$$E(\mathbf{x}, r) := \bigcap_{\substack{r' > r \\ r' \in R_{\mathbf{x}}}} \text{im}_{\mathbb{T}}(\mathbf{u}, B(\mathbf{x}, r')). \quad (2.3)$$

Whenever explicit mention of \mathbf{u} is necessary (such as in Theorem 4 where sequences of deformations are considered), we write $E(\mathbf{a}, r; \mathbf{u})$. Finally, if a point $\mathbf{a} \in \mathbb{R}^n$ is such that $B(\mathbf{a}, \delta) \setminus \{\mathbf{a}\} \subset \Omega$ for some $\delta > 0$, we define its topological image as $E(\mathbf{a}, 0)$, and denote it by $\text{im}_{\mathbb{T}}(\mathbf{u}, \mathbf{a})$.

2.4 The distributional determinant

It is well known that the Jacobian determinant of a C^2 vector-valued map $\mathbf{u} : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}^n$ has a divergence structure. When $n = 2$ or $n = 3$, this is

$$\det D\mathbf{u} = u_{1,1}u_{2,2} - u_{2,1}u_{1,2} = (u_1u_{2,2})_{,1} - (u_1u_{2,1})_{,2}$$

$$\begin{aligned} \det D\mathbf{u} &= u_{1,1} \begin{vmatrix} u_{2,2} & u_{2,3} \\ u_{3,2} & u_{3,3} \end{vmatrix} + u_{1,2} \begin{vmatrix} u_{2,3} & u_{2,1} \\ u_{3,3} & u_{3,1} \end{vmatrix} + u_{1,3} \begin{vmatrix} u_{2,1} & u_{2,2} \\ u_{3,1} & u_{3,2} \end{vmatrix} \\ &= \left(u_1 \begin{vmatrix} u_{2,2} & u_{2,3} \\ u_{3,2} & u_{3,3} \end{vmatrix} \right)_{,1} + \left(u_1 \begin{vmatrix} u_{2,3} & u_{2,1} \\ u_{3,3} & u_{3,1} \end{vmatrix} \right)_{,2} + \left(u_1 \begin{vmatrix} u_{2,1} & u_{2,2} \\ u_{3,1} & u_{3,2} \end{vmatrix} \right)_{,3}, \end{aligned}$$

where $u_{i,j}$ denotes the j -th partial derivative of the i -th component of \mathbf{u} . In higher dimensions, we may write $\det D\mathbf{u} = \operatorname{Div}((\operatorname{adj} D\mathbf{u}) \frac{\mathbf{u}}{n})$.

One of the main ideas in Ball's theory for nonlinear elasticity [2] is that if the divergence is taken in the sense of distributions, the right-hand side of the above expressions is well defined for maps that are only weakly differentiable. This motivated his definition of the *distributional determinant* of a map $\mathbf{u} \in W^{1,n-1}(\Omega, \mathbb{R}^n) \cap L_{\operatorname{loc}}^\infty(\Omega, \mathbb{R}^n)$ as the distribution $\operatorname{Det} D\mathbf{u} \in \mathcal{D}'(\Omega)$ given by

$$\langle \operatorname{Det} D\mathbf{u}, \phi \rangle := -\frac{1}{n} \int_{\Omega} \mathbf{u}(\mathbf{x}) \cdot (\operatorname{cof} D\mathbf{u}(\mathbf{x})) D\phi(\mathbf{x}) \, d\mathbf{x}, \quad \phi \in C_c^\infty(\Omega) \quad (2.4)$$

(see also [54, 16, 10, 67, 21, 11] and references therein for subsequent developments and for the role of $\operatorname{Det} D\mathbf{u}$ in compensated compactness, homogenization, liquid crystals, and superconductivity). If a map $\mathbf{u} \in W^{1,p}(\Omega, \mathbb{R}^n)$, $p > n - 1$, satisfies condition INV, then $\mathbf{u}(\mathbf{z})$ is contained in the region enclosed by $\mathbf{u}(\partial B(\mathbf{x}, r))$ for every $\mathbf{x} \in \mathbb{R}^n$, a.e. $\mathbf{z} \in \Omega \cap B(\mathbf{x}, r)$, and a.e. $r > 0$ such that $\partial B(\mathbf{x}, r) \subset \Omega$. Consequently, $\mathbf{u} \in L_{\operatorname{loc}}^\infty(\Omega, \mathbb{R}^n)$, and the distributional determinant is well defined.

Proposition 2.2 (cf. [55], Lemma 8.1). *Let $\mathbf{u} \in W^{1,p}(\Omega, \mathbb{R}^n)$, $p > n - 1$, satisfy $\det D\mathbf{u} > 0$ a.e. and condition INV. Then*

- i) $\operatorname{Det} D\mathbf{u} = (\det D\mathbf{u})\mathcal{L}^n + \mu^s$, where μ^s is singular with respect to \mathcal{L}^n
- ii) $|E(\mathbf{x}, r) \setminus \operatorname{im}_T(\mathbf{u}, B(\mathbf{x}, r))| = 0$ for every $\mathbf{x} \in \mathbb{R}^n$ and $r \in R_{\mathbf{x}}$
- iii) $|E(\mathbf{x}, r_2) \setminus E(\mathbf{x}, r_1)| = \operatorname{Det} D\mathbf{u}(A_{r_1, r_2})$ for all $r_1 \geq 0$ and $r_2 > 0$ such that the annulus $A_{r_1, r_2} := \{\mathbf{x} \in \mathbb{R}^n : r_1 < |\mathbf{x}| < r_2\}$ is contained in Ω .

Proof. Let $\mathbf{x} \in \mathbb{R}^n$ and set $S := \{r \in (0, \infty) : \partial B(\mathbf{x}, r) \subset \Omega\}$. The map

$$\omega(r) := \frac{1}{n} \int_{\partial B(\mathbf{x}, r)} \mathbf{u}(\mathbf{z}) \cdot (\operatorname{cof} D\mathbf{u}(\mathbf{z})) \boldsymbol{\nu}(\mathbf{z}) \, d\mathcal{H}^{n-1}(\mathbf{z}), \quad r \in R_{\mathbf{x}}$$

belongs to $L^1(S)$. Suppose $[r_1, r_2] \subset S$ for some $r_1, r_2 \in R_{\mathbf{x}}$. For $\delta > 0$ let $\phi_\delta(\mathbf{z}) := \psi_\delta(|\mathbf{z} - \mathbf{x}|)$, where $\psi_\delta \in C_c^\infty([0, \infty))$ is such that $\psi_\delta = 1$ in $(r_1 + \delta, r_2 - \delta)$, $\psi_\delta = 0$ in $[0, r_1] \cup [r_2, \infty)$, and $\delta \|\psi'_\delta\|_\infty \leq 2$. It is clear that $\phi_\delta \rightarrow \chi_{A_{r_1, r_2}}$ pointwise as $\delta \rightarrow 0^+$, and that

$$\langle \operatorname{Det} D\mathbf{u}, \phi_\delta \rangle = \omega(r_2) - \omega(r_1) + \int_{r_1}^{r_1 + \delta} \delta \psi'_\delta(r) (\omega(r_1) - \omega(r)) + \int_{r_2 - \delta}^{r_2} \delta \psi'_\delta(r) (\omega(r_2) - \omega(r)).$$

The proof follows from [55, Lemma 8.1], Proposition 2.1iv)–v), and Lebesgue's differentiation theorem applied to ω . \square

3 Lower bounds

The following is the basic estimate that allows us to relate the elastic energy to the volume and distortion of the cavities. It extends (1.14) to an arbitrary exponent p and dimension n .

Lemma 3.1. *Suppose that $\mathbf{u} \in W^{1,p}(\Omega, \mathbb{R}^n)$, $p > n - 1$, satisfies $\det D\mathbf{u} > 0$ a.e. and condition INV. Then, for every $\mathbf{x} \in \Omega$ and $r \in R_{\mathbf{x}}$ (as defined in Proposition 2.1),*

$$\int_{\partial B(\mathbf{x}, r)} \left| \frac{D(\mathbf{u}|_{\partial B(\mathbf{x}, r)})(\mathbf{x})}{\sqrt{n-1}} \right|^p d\mathcal{H}^{n-1}(\mathbf{x}) \geq \left(\frac{|E(\mathbf{x}, r)|}{|B(\mathbf{x}, r)|} \right)^{\frac{p}{n}} (1 + CD(E(B(\mathbf{x}, r))))^{\frac{p}{n-1}}.$$

Equality is attained only if $\mathbf{u}|_{\partial B(\mathbf{x}, r)}$ is radially symmetric.

Proof. Given $\mathbf{x} \in \mathbb{R}^n$, $r > 0$ and $\mathbf{z} \in \partial B(\mathbf{x}, r)$ such that $D\mathbf{u}(\mathbf{z})$ is well defined, we have that

$$\begin{aligned} |(\operatorname{cof} D\mathbf{u}(\mathbf{z}))\boldsymbol{\nu}(\mathbf{z})| &= |(D\mathbf{u}(\mathbf{z}))\mathbf{e}_1 \wedge \cdots \wedge (D\mathbf{u}(\mathbf{z}))\mathbf{e}_{n-1}| \leq |(D\mathbf{u})\mathbf{e}_1| \cdots |(D\mathbf{u})\mathbf{e}_{n-1}| \\ &\leq (n-1)^{\frac{1-n}{2}} (|(D\mathbf{u})\mathbf{e}_1|^2 + \cdots + |(D\mathbf{u})\mathbf{e}_{n-1}|^2)^{\frac{n-1}{2}}, \end{aligned}$$

$\{\mathbf{e}_1, \dots, \mathbf{e}_{n-1}, \boldsymbol{\nu}(\mathbf{z})\}$ being an orthonormal basis of \mathbb{R}^n with $\boldsymbol{\nu}(\mathbf{z}) := (\mathbf{z} - \mathbf{x})/r$. Equality holds only if $|(D\mathbf{u})\mathbf{e}_i| = |(D\mathbf{u})\mathbf{e}_j|$ and $(D\mathbf{u})\mathbf{e}_i \perp (D\mathbf{u})\mathbf{e}_j$ for $i \neq j$, as in Sivaloganathan-Spector [72, 73]. If $r \in R_{\mathbf{x}}$, by Propositions 2.1iii), 2.2ii), and 1.3, we obtain

$$\int_{\partial B(\mathbf{x}, r)} \left| \frac{D(\mathbf{u}|_{\partial B(\mathbf{x}, r)})(\mathbf{x})}{\sqrt{n-1}} \right|^{n-1} d\mathcal{H}^{n-1} \geq \left(\frac{|E(\mathbf{x}, r)|}{\omega_n r^n} \right)^{\frac{n-1}{n}} (1 + CD(E(\mathbf{x}, r))).$$

The conclusion follows by Jensen's inequality. \square

3.1 Ball constructions, the case of multiple cavities

In this Section we prove Proposition 1.1 (our first lower bound, valid for an arbitrary number of cavities). We start by introducing the necessary notation, and by recalling the ball construction method in Ginzburg-Landau theory, following the presentation in [67].

Collections of balls will be denoted by expressions with \mathcal{B} . If B is a ball, $r(B)$ denotes its radius. If \mathcal{B} is a collection of balls, then $r(\mathcal{B}) = \sum_{B \in \mathcal{B}} r(B)$. If $\lambda \geq 0$, $\lambda\mathcal{B} := \{\lambda B : B \in \mathcal{B}\}$. We use $\bigcup \mathcal{B}$ to denote the union $\bigcup_{B \in \mathcal{B}} B$ of a collection of balls. Given a measurable set A and a collection of balls \mathcal{B} , we denote $\{B \cap A : B \in \mathcal{B}\}$ by $A \cap \mathcal{B}$. Given $\mathcal{F} : \mathbb{R}^n \times (0, \infty) \rightarrow \mathbb{R}$, we regard \mathcal{F} as a function defined on the set of all balls (cf. [67, Def. 4.1]), and write $\mathcal{F}(B)$ for $\mathcal{F}(\mathbf{x}, r)$ if $B = B(\mathbf{x}, r)$ (or $\bar{B}(\mathbf{x}, r)$). Also, we write $\mathcal{F}(\mathcal{B})$ for $\sum_{B \in \mathcal{B}} \mathcal{F}(B)$ if \mathcal{B} is a collection of balls.

Proposition 3.2 (cf. [67], Th. 4.2). *Let \mathcal{B}_0 be a finite collection of disjoint closed balls and let $t_0 := r(\mathcal{B}_0)$. There exists a family $\{\mathcal{B}(t) : t \geq t_0\}$ of collections of disjoint closed balls such that $\mathcal{B}(t_0) = \mathcal{B}_0$ and*

- i) *For every $s \geq t \geq t_0$, $\bigcup \mathcal{B}(t) \subset \bigcup \mathcal{B}(s)$.*
- ii) *There exists a finite set T such that if $[t_1, t_2] \subset [t_0, \infty) \setminus T$, then $\mathcal{B}(t_2) = \frac{t_2}{t_1} \mathcal{B}(t_1)$.*

iii) $r(\mathcal{B}(t)) = t$ for every $t \geq t_0$.

We point out that we chose a different parametrization from the one in [67, Th. 4.2]. Here t corresponds to e^t there.

Definition 6 ([67], Def. 4.1). *We say that a function $\mathcal{F} : \mathbb{R}^n \times (0, \infty) \rightarrow \mathbb{R}$ is monotonic (when regarded as a function defined in the set of balls) if $\mathcal{F}(\mathbf{x}, r)$ is continuous with respect to r and $\mathcal{F}(\mathcal{B}) \leq \mathcal{F}(\mathcal{B}')$ for any families of disjoint closed balls $\mathcal{B}, \mathcal{B}'$ such that $\bigcup \mathcal{B} \subset \bigcup \mathcal{B}'$.*

Proposition 3.3 (cf. [67], Prop. 4.1). *Let $\mathcal{F} : \mathbb{R}^n \times (0, \infty) \rightarrow \mathbb{R}$ be monotonic in the sense of Definition 6. Let \mathcal{B}_0 and $\{\mathcal{B}(t) : t \geq t_0\}$ satisfy the conditions of Proposition 3.2. Then,*

$$\mathcal{F}(\mathcal{B}(s)) - \mathcal{F}(\mathcal{B}_0) \geq \int_{t_0}^s \sum_{B(\mathbf{x}, r) \in \mathcal{B}(t)} r \frac{\partial \mathcal{F}}{\partial r}(\mathbf{x}, r) \frac{dt}{t} \quad (3.1)$$

for every $s \geq t_0$, and for every $B \in \mathcal{B}(s)$

$$\mathcal{F}(B) - \mathcal{F}(\mathcal{B}_0 \cap B) \geq \int_{t_0}^s \sum_{B(\mathbf{x}, r) \in \mathcal{B}(t) \cap B} r \frac{\partial \mathcal{F}}{\partial r}(\mathbf{x}, r) \frac{dt}{t}. \quad (3.2)$$

Lemma 3.1 applied to $\mathcal{F}(\mathbf{x}, r) = \int_{B(\mathbf{x}, r)} \left(\left| \frac{D\mathbf{u}(\mathbf{x})}{\sqrt{n-1}} \right|^p - 1 \right) d\mathbf{x}$ and Proposition 3.3 immediately imply the following result (stated without proof).

Proposition 3.4. *Suppose that $\mathbf{u} \in W^{1,p}(\Omega, \mathbb{R}^n)$ with $p > n - 1$ satisfies $\det D\mathbf{u} > 0$ a.e. and condition INV. Suppose, further, that \mathcal{B}_0 and $\{\mathcal{B}(t) : t \geq t_0\}$ satisfy the conditions of Proposition 3.2. Then, for every $s > t_0$ such that $\Omega_s := \bigcup \mathcal{B}(s) \setminus \bigcup \mathcal{B}_0 \subset \Omega$,*

$$\frac{1}{n} \int_{\Omega_s} \left(\left| \frac{D\mathbf{u}(\mathbf{x})}{\sqrt{n-1}} \right|^p - 1 \right) d\mathbf{x} \geq \int_{t_0}^s \sum_{B \in \mathcal{B}(t)} |B| \left(\frac{|E_B|_n^{\frac{p}{n}}}{|B|_n^{\frac{p}{n}}} (1 + CD(E_B))^{\frac{p}{n-1}} - 1 \right) \frac{dt}{t},$$

where E_B denotes $E(\mathbf{x}, r)$ for $B = \overline{B}(\mathbf{x}, r)$. Analogously, for every $B \in \mathcal{B}(s)$

$$\frac{1}{n} \int_{B \setminus \bigcup \mathcal{B}_1} \left(\left| \frac{D\mathbf{u}(\mathbf{x})}{\sqrt{n-1}} \right|^p - 1 \right) d\mathbf{x} \geq \int_{t_0}^s \sum_{B' \in \mathcal{B}(t) \cap B} |B'| \left(\frac{|E_{B'}|_n^{\frac{p}{n}}}{|B'|_n^{\frac{p}{n}}} (1 + CD(E_{B'}))^{\frac{p}{n-1}} - 1 \right) \frac{dt}{t}.$$

Proposition 1.1 finally follows from Proposition 3.4 and the incompressibility constraint:

Proof of Proposition 1.1. Let $A := \{i : B(\mathbf{a}_i, R) \subset \Omega\}$, $t_0 := r(\mathcal{B}_0) = \sum_{i \in A} \varepsilon_i$, and $\mathcal{B}_0 := \bigcup_{i \in A} \overline{B}_{\varepsilon_i}(\mathbf{a}_i)$. Let $\{\mathcal{B}(t) : t \geq t_0\}$ be the family obtained by applying Proposition 3.2 to \mathcal{B}_0 . Then, applying Proposition 3.4, if $\bigcup \mathcal{B}(s) \subset \Omega$,

$$\frac{1}{n} \int_{\Omega_\varepsilon \cap \bigcup \mathcal{B}(s)} \left(\left| \frac{D\mathbf{u}(\mathbf{x})}{\sqrt{n-1}} \right|^n - 1 \right) d\mathbf{x} \geq \int_{t_0}^s \sum_{B \in \mathcal{B}(t)} \left((|E_B| - |B|) + C|E_B|D(E_B)^{\frac{n}{n-1}} \right) \frac{dt}{t}. \quad (3.3)$$

Proceeding as in the proof of Proposition 2.2 and using incompressibility we obtain

$$\left| E_B \setminus \bigcup_{\mathbf{a}_i \in B} E(\mathbf{a}_i, \varepsilon_i) \right| = \text{Det } D\mathbf{u} \left(B \setminus \bigcup_{\mathbf{a}_i \in B} \overline{B}_{\varepsilon_i}(\mathbf{a}_i) \right) = |B| - \sum_{\mathbf{a}_i \in B} \omega_n \varepsilon_i^n,$$

hence, by the definition of v_i in the statement of the proposition,

$$|E_B| - |B| = \left| \bigcup_{\mathbf{a}_i \in B} E(\mathbf{a}_i, \varepsilon_i) \right| - \sum_{\mathbf{a}_i \in B} \omega_n \varepsilon_i^n = \sum_{\mathbf{a}_i \in B} v_i. \quad (3.4)$$

Combining (3.3) and (3.4) we obtain

$$\frac{1}{n} \int_{\Omega_\varepsilon \cap \bigcup \mathcal{B}(s)} \left(\left| \frac{D\mathbf{u}(\mathbf{x})}{\sqrt{n-1}} \right|^n - 1 \right) d\mathbf{x} \geq \left(\sum_{i, B(\mathbf{a}_i, R) \subset \Omega_\varepsilon} v_i \right) \log \frac{s}{t_0} + C \int_{t_0}^s \left(\sum_{B \in \mathcal{B}(t)} |E_B| D(E_B)^{\frac{n}{n-1}} \right) \frac{dt}{t}.$$

Let $s_0 := \sup\{s \in [t_0, R) : \bigcup \mathcal{B}(s) \subset \Omega\}$. If $s_0 = R$, the claim is proved. Otherwise, from Proposition 3.2 we deduce that there exists a ball $B(\mathbf{a}, r) \in \mathcal{B}(s_0)$, of radius $r \leq s_0$, containing at least one \mathbf{a}_i , $i \in A$, such that $\overline{B}(\mathbf{a}, r) \cap \partial\Omega \neq \emptyset$. The proof is completed by observing that

$$R < \text{dist}(\mathbf{a}_i, \partial\Omega) \leq |\mathbf{a}_i - \mathbf{a}| + \text{dist}(\mathbf{a}, \partial\Omega) < 2s_0.$$

□

3.2 The case of two cavities: proof of Theorem 1

In this section, we prove Theorem 1 assuming Proposition 1.2, whose proof is postponed to Section 3.3.

We will need the following lemma.

Lemma 3.5 (Modulus of continuity of the distortion). *Let $E, E' \subset \mathbb{R}^n$ be measurable. Then*

- i) $||E|D(E) - |E'|D(E')| \leq 2|E \Delta E'|$
- ii) $\left| |E|D(E)^{\frac{n}{n-1}} - |E'|D(E')^{\frac{n}{n-1}} \right| \leq 2^{\frac{n}{n-1}} \frac{n+1}{n-1} |E \Delta E'|.$

Proof. Let B' be a ball such that $|B'| = |E'|$ and $|E'|D(E') = |E' \Delta B'|$. For all measurable sets B

$$|E \Delta B| - |E'|D(E') = \|\chi_E - \chi_B\|_{L^1} - \|\chi_{E'} - \chi_{B'}\|_{L^1} \leq \|\chi_E - \chi_{E'}\|_{L^1} + \|\chi_B - \chi_{B'}\|_{L^1}.$$

Testing with concentric balls, and taking the minimum over all balls B with $|B| = |E|$, yields

$$|E|D(E) - |E'|D(E') \leq \|\chi_E - \chi_{E'}\|_{L^1} + ||E| - |E'||$$

($\|\chi_B - \chi_{B'}\|_{L^1} = ||E| - |E'|$ since B and B' are concentric). Combining this with the fact that $||E| - |E'|| = ||\chi_E\|_{L^1} - \|\chi_{E'}\|_{L^1}| \leq \|\chi_E - \chi_{E'}\|_{L^1}$, we obtain i).

Property ii) follows from i), the mean value theorem, and the fact that $D(E) \leq 2$ for all E (a direct consequence of its definition). To be more precise, suppose that $|E| > |E'|$, then

$$\begin{aligned}
& \left| |E|D(E)^{\frac{n}{n-1}} - |E'|D(E')^{\frac{n}{n-1}} \right| \\
&= \left| |E|^{-\frac{1}{n-1}} (|E|D(E))^{\frac{n}{n-1}} - |E'|^{-\frac{1}{n-1}} (|E'|D(E'))^{\frac{n}{n-1}} \right| \\
&\leq |E|^{-\frac{1}{n-1}} \left| (|E|D(E))^{\frac{n}{n-1}} - (|E'|D(E'))^{\frac{n}{n-1}} \right| + (|E'|D(E'))^{\frac{n}{n-1}} \left| |E|^{-\frac{1}{n-1}} - |E'|^{-\frac{1}{n-1}} \right| \\
&\leq \frac{2n}{n-1} |E|^{-\frac{1}{n-1}} (\max\{|E|D(E), |E'|D(E')\})^{\frac{1}{n-1}} |E \Delta E'| + \frac{2^{\frac{n}{n-1}}}{n-1} ||E| - |E'||,
\end{aligned}$$

completing the proof. \square

We now proceed to the proof of Theorem 1. As in (3.4), by Proposition 2.2 we have that $|E(B)| = |B| + \sum_{i: \mathbf{a}_i \in B} v_i$ for all balls B with $\partial B \subset \Omega_\varepsilon$. Hence, Lemma 3.1 implies that

$$\frac{1}{n} \int_{\partial B(\mathbf{x}, r)} \left(\left| \frac{D\mathbf{u}(\mathbf{x})}{\sqrt{n-1}} \right|^n - 1 \right) d\mathcal{H}^{n-1}(\mathbf{x}) \geq \left(\sum_{i: \mathbf{a}_i \in B(\mathbf{x}, r)} v_i + C|E(\mathbf{x}, r)|D(E(\mathbf{x}, r))^{\frac{n}{n-1}} \right) \frac{1}{r} \quad (3.5)$$

for all $\mathbf{x} \in \mathbb{R}^n$ and all $r \in R_{\mathbf{x}}$. Given $R > d$ such that $B(\mathbf{a}, R) \subset \Omega$, let

$$A_1 := B_{d/2}(\mathbf{a}_1) \setminus \overline{B_{\varepsilon_1}(\mathbf{a}_1)}, \quad A_2 := B_{d/2}(\mathbf{a}_2) \setminus \overline{B_{\varepsilon_2}(\mathbf{a}_2)}, \quad A_3 := B_R(\mathbf{a}) \setminus \overline{B_d(\mathbf{a})}.$$

By considering that $\Omega_\varepsilon \cap B(\mathbf{a}, R) \supset A_1 \cup A_2 \cup A_3$ and integrating successively in each annulus, we obtain

$$\begin{aligned}
\frac{1}{n} \int_{\Omega_\varepsilon \cap B(\mathbf{a}, R)} \left(\left| \frac{D\mathbf{u}(\mathbf{x})}{\sqrt{n-1}} \right|^n - 1 \right) d\mathbf{x} &\geq v_1 \log \frac{d}{2\varepsilon_1} + v_2 \log \frac{d}{2\varepsilon_2} + (v_1 + v_2) \log \frac{R}{d} \\
&+ C \int_{\varepsilon_1}^{d/2} |E(\mathbf{a}_1, r)|D(E(\mathbf{a}_1, r))^{\frac{n}{n-1}} \frac{dr}{r} + C \int_{\varepsilon_2}^{d/2} |E(\mathbf{a}_2, r)|D(E(\mathbf{a}_2, r))^{\frac{n}{n-1}} \frac{dr}{r} \\
&+ C \int_d^R |E(\mathbf{a}, r)|D(E(\mathbf{a}, r))^{\frac{n}{n-1}} \frac{dr}{r}.
\end{aligned} \quad (3.6)$$

Proposition 1.2 applied to $E_1 = E(\mathbf{a}_1, \frac{d}{2})$, $E_2 = E(\mathbf{a}_2, \frac{d}{2})$, and $E = E(\mathbf{a}, r)$, $r \in (d, R)$ gives

$$\begin{aligned}
& |E(\mathbf{a}, r)|D(E(\mathbf{a}, r))^{\frac{n}{n-1}} \\
&\geq C(v_1 + v_2) \left(\frac{(|E_1|^{\frac{1}{n}} + |E_2|^{\frac{1}{n}})^n - |E(\mathbf{a}, r)|}{(|E_1|^{\frac{1}{n}} + |E_2|^{\frac{1}{n}})^n - |E_1 \cup E_2|} \right)^{\frac{n(n+1)}{2(n-1)}} \left(\frac{\min\{|E_1|, |E_2|\}}{|E_1| + |E_2|} \right)^{\frac{n}{n-1}} \\
&\quad - |E(\mathbf{a}_1, d/2)|D(E(\mathbf{a}_1, d/2))^{\frac{n}{n-1}} - |E(\mathbf{a}_2, d/2)|D(E(\mathbf{a}_2, d/2))^{\frac{n}{n-1}}.
\end{aligned} \quad (3.7)$$

Define $g(\beta_1, \beta_2) := (\beta_1^{\frac{1}{n}} + \beta_2^{\frac{1}{n}})^n - (\beta_1 + \beta_2)$ (when $n = 2$, $g(\beta_1, \beta_2) = 2\sqrt{\beta_1\beta_2}$). Using that $|E_i| = v_i + \frac{\omega_n d^n}{2^n}$, $i = 1, 2$ we may write

$$(|E_1|^{\frac{1}{n}} + |E_2|^{\frac{1}{n}})^n = g(|E_1|, |E_2|) + (|E_1| + |E_2|) = g(|E_1|, |E_2|) + 2 \cdot \frac{\omega_n d^n}{2^n} + v_1 + v_2. \quad (3.8)$$

Estimate (3.7) is meaningful if $|E(\mathbf{a}, r)| \leq (|E_1|^{\frac{1}{n}} + |E_2|^{\frac{1}{n}})^n$, i.e. if

$$\omega_n d^n \leq \omega_n r^n \leq g\left(v_1 + \frac{\omega_n d^n}{2^n}, v_2 + \frac{\omega_n d^n}{2^n}\right) + \frac{\omega_n d^n}{2^{n-1}} \quad (3.9)$$

(since g is increasing in β_1 and β_2 and $g(\beta, \beta) = (2^n - 2)\beta$, the inequality holds at least for $r = d$). Define ρ as the radius for which $\omega_n r^n$ is in the middle of the two extremes in (3.9),

$$\omega_n \rho^n := (2^{n-1} + 1) \frac{\omega_n d^n}{2^n} + \frac{1}{2} g\left(v_1 + \frac{\omega_n d^n}{2^n}, v_2 + \frac{\omega_n d^n}{2^n}\right). \quad (3.10)$$

For all $r \in (d, \min\{\rho, R\})$ we have that $E(\mathbf{a}, r) \subset E(\mathbf{a}, \rho)$, hence

$$|E(\mathbf{a}, r)| < \omega_n \rho^n + v_1 + v_2 = \frac{1}{2} g(|E_1|, |E_2|) + (2^{n-1} + 1) \frac{\omega_n d^n}{2^n} + v_1 + v_2. \quad (3.11)$$

Noticing that g is 1-homogeneous, combining (3.8) and (3.11) we obtain

$$\frac{(|E_1|^{\frac{1}{n}} + |E_2|^{\frac{1}{n}})^n - |E(\mathbf{a}, r)|}{(|E_1|^{\frac{1}{n}} + |E_2|^{\frac{1}{n}})^n - |E_1 \cup E_2|} \geq \frac{\frac{1}{2} g(|E_1|, |E_2|) - (2^{n-1} + 1 - 2) \frac{\omega_n d^n}{2^n}}{g(|E_1|, |E_2|)} = \frac{1}{2} - \frac{2^{n-1} - 1}{g\left(\frac{2^n |E_1|}{\omega_n d^n}, \frac{2^n |E_2|}{\omega_n d^n}\right)}.$$

Without loss of generality, assume that $\omega_n d^n < v_1 + v_2$. Estimate $g\left(\frac{2^n |E_1|}{\omega_n d^n}, \frac{2^n |E_2|}{\omega_n d^n}\right)$ by

$$\begin{aligned} g(1+x, 1+y) &= \sum_{k=1}^{n-1} \binom{n}{k} \left((1+x)^k (1+y)^{n-k} \right)^{\frac{1}{n}} \geq \sum_{k=1}^{n-1} \binom{n}{k} (1+kx)^{\frac{1}{n}} (1+(n-k)y)^{\frac{1}{n}} \\ &\geq \sum_{k=1}^{n-1} \binom{n}{k} (1+x)^{\frac{1}{n}} (1+y)^{\frac{1}{n}} \geq (2^n - 2) (1+x+y)^{\frac{1}{n}} \end{aligned} \quad (3.12)$$

(with $x = \frac{2^n |E_1|}{\omega_n d^n} - 1 = \frac{2^n v_1}{\omega_n d^n}$ and $y = \frac{2^n v_2}{\omega_n d^n}$) to obtain

$$\left(\frac{(|E_1|^{\frac{1}{n}} + |E_2|^{\frac{1}{n}})^n - |E(\mathbf{a}, r)|}{(|E_1|^{\frac{1}{n}} + |E_2|^{\frac{1}{n}})^n - |E_1 \cup E_2|} \right)^{\frac{n(n+1)}{2(n-1)}} \geq \left(\frac{1}{2} - \frac{2^{n-1} - 1}{(2^n - 2) \left(1 + 2^n \frac{v_1 + v_2}{\omega_n d^n}\right)^{\frac{1}{n}}} \right)^{\frac{n(n+1)}{2(n-1)}} \geq 4^{-\frac{n(n+1)}{2(n-1)}}.$$

On the other hand, $|E_1 \cup E_2| < 2(v_1 + v_2)$ (because $\omega_n d^n < v_1 + v_2$), and since $|E_1| \geq v_1$ and $|E_2| \geq v_2$, we can substitute $\frac{\min\{|E_1|, |E_2|\}}{|E_1| + |E_2|}$ with $\frac{\min\{v_1, v_2\}}{v_1 + v_2}$ in (3.7). Hence, for all $r \in (d, \min\{\rho, R\})$, all $s_1 \in (\varepsilon_1, d/2)$ and all $s_2 \in (\varepsilon_2, d/2)$,

$$\begin{aligned} &|E(\mathbf{a}, r)| D(E(\mathbf{a}, r))^{\frac{n}{n-1}} + |E(\mathbf{a}_1, s_1)| D(E(\mathbf{a}_1, s_1))^{\frac{n}{n-1}} + |E(\mathbf{a}_2, s_2)| D(E(\mathbf{a}_1, s_1))^{\frac{n}{n-1}} \\ &\geq C(n) (v_1 + v_2) \left(\frac{\min\{v_1, v_2\}}{v_1 + v_2} \right)^{\frac{n}{n-1}} - \sum_{i=1}^2 \left| |E(\mathbf{a}_i, s_i)| D(E(\mathbf{a}_i, s_i))^{\frac{n}{n-1}} - |E(\mathbf{a}_i, \frac{d}{2})| D(E(\mathbf{a}_i, \frac{d}{2}))^{\frac{n}{n-1}} \right|. \end{aligned} \quad (3.13)$$

Denoting $E(\mathbf{a}_1, s_1)$, $E(\mathbf{a}_2, s_2)$, and $E(\mathbf{a}, r)$ by E_{s_1} , E_{s_2} , and E_r , from (3.6) we obtain

$$\begin{aligned} &\frac{1}{n} \int_{\Omega_\varepsilon \cap B(\mathbf{a}, R)} \left(\left| \frac{D\mathbf{u}(\mathbf{x})}{\sqrt{n-1}} \right|^n - 1 \right) d\mathbf{x} \geq v_1 \log \frac{R}{2\varepsilon_1} + v_2 \log \frac{R}{2\varepsilon_2} \\ &+ C \inf_{\substack{r \in (d, \min\{\rho, R\}) \\ s_i \in (\varepsilon_i, d/2)}} \left(|E_r| D(E_r)^{\frac{n}{n-1}} + |E_{s_1}| D(E_{s_1})^{\frac{n}{n-1}} + |E_{s_2}| D(E_{s_2})^{\frac{n}{n-1}} \right) \log \min \left\{ \frac{\rho}{d}, \frac{R}{d}, \frac{d}{\varepsilon} \right\}, \end{aligned} \quad (3.14)$$

with $\varepsilon = \max\{\varepsilon_1, \varepsilon_2\}$. In order to estimate $\log \frac{\rho}{d}$, from (3.10) and (3.12) we find that

$$\frac{\rho^n}{d^n} \geq 2^{-(n+1)} g \left(1 + \frac{2^n v_1}{\omega_n d^n}, 1 + \frac{2^n v_2}{\omega_n d^n} \right) \geq (2^{-1} - 2^{-n}) \left(1 + 2^n \frac{v_1 + v_2}{\omega_n d^n} \right)^{\frac{1}{n}} \geq (1 - 2^{1-n}) \left(\frac{v_1 + v_2}{\omega_n d^n} \right)^{\frac{1}{n}}.$$

The proof is completed by combining (3.13) and (3.14) with Lemma 3.5.

3.3 Estimate on the distortions

This section is devoted to the proof of Proposition 1.2.

Lemma 3.6. *Let $q > 1$ and suppose that E , E_1 , and E_2 are sets of positive measure such that $E \supset E_1 \cup E_2$ and $E_1 \cap E_2 = \emptyset$. Then*

$$\frac{|E|D(E)^q + |E_1|D(E_1)^q + |E_2|D(E_2)^q}{|E| + |E_1 \cup E_2|} \geq \min_{B, B_1, B_2} \left(\frac{\|\chi_B - \chi_{B_1} - \chi_{B_2}\|_{L^1} - (|B| - |B_1| - |B_2|)}{|E| + |E_1 \cup E_2|} \right)^q,$$

where the minimum is taken over all balls B , B_1 , B_2 with $|B| = |E|$, $|B_1| = |E_1|$, $|B_2| = |E_2|$.

Proof. Let B , B_1 , B_2 attain the minimum in the definition of $D(E)$, $D(E_1)$, $D(E_2)$, that is, suppose that $|B| = |E|$, $|B_1| = |E_1|$, $|B_2| = |E_2|$ and

$$|E|D(E) = |E \triangle B|, \quad |E_1|D(E_1) = |E_1 \triangle B_1|, \quad |E_2|D(E_2) = |E_2 \triangle B_2|.$$

Since $\chi_B - \chi_{B_1} - \chi_{B_2} = (\chi_B - \chi_E) + (\chi_E - \chi_{E_1} - \chi_{E_2}) + (\chi_{E_1} - \chi_{B_1}) + (\chi_{E_2} - \chi_{B_2})$, then

$$\|\chi_B - \chi_{B_1} - \chi_{B_2}\|_{L^1} - \|\chi_E - \chi_{E_1} - \chi_{E_2}\|_{L^1} \leq |E|D(E) + |E_1|D(E_1) + |E_2|D(E_2).$$

Also, note that $\|\chi_E - \chi_{E_1} - \chi_{E_2}\|_{L^1} = |E| - |E_1| - |E_2| = |B| - |B_1| - |B_2|$ because $E_1 \cap E_2 = \emptyset$ and $E_1 \cup E_2 \subset E$. The result follows by Jensen's inequality applied to the map $t \mapsto t^q$. \square

Lemma 3.7. *Let B, B_1, B_2 be measurable subsets of \mathbb{R}^n . Then*

$$\|\chi_B - \chi_{B_1} - \chi_{B_2}\|_{L^1} - (|B| - |B_1| - |B_2|) = 2(|B_1| + |B_2| - |B \cap (B_1 \cup B_2)|) \quad (3.15)$$

$$= 2(|B_1 \setminus B| + |B_2 \setminus B| + |B \cap B_1 \cap B_2|). \quad (3.16)$$

Proof. Consider, first, the elementary relations

$$|B_i \setminus B| = |B_i| - |B \cap B_i|, \quad i = 1, 2. \quad (3.17)$$

$$|B \cap (B_1 \cup B_2)| = |B \cap B_1| + |B \cap B_2| - |B \cap B_1 \cap B_2|. \quad (3.18)$$

$$|B \setminus (B_1 \cup B_2)| = |B| - |B \cap (B_1 \cup B_2)|. \quad (3.19)$$

From (3.17) and (3.18) we obtain

$$|B_1 \setminus B| + |B_2 \setminus B| + |B \cap B_1 \cap B_2| = |B_1| + |B_2| - |B \cap (B_1 \cup B_2)|. \quad (3.20)$$

From (3.19) and (3.20) we obtain

$$|B \setminus (B_1 \cup B_2)| = |B| - (|B_1| + |B_2|) + (|B_1 \setminus B| + |B_2 \setminus B| + |B \cap B_1 \cap B_2|). \quad (3.21)$$

Decomposing \mathbb{R}^n as $\bigcup_{\alpha, \alpha_1, \alpha_2 \in \{0,1\}} \{\mathbf{y} : (\chi_B, \chi_{B_1}, \chi_{B_2}) = (\alpha, \alpha_1, \alpha_2)\}$ we find that

$$\begin{aligned} \|\chi_B - \chi_{B_1} - \chi_{B_2}\|_{L_1} = & |B \cap B_1 \cap B_2| + |B \setminus (B_1 \cup B_2)| \\ & + 2|(B_1 \cap B_2) \setminus B| + |(B_1 \setminus B) \setminus B_2| + |(B_2 \setminus B) \setminus B_1|. \end{aligned}$$

Since $|(B_1 \cap B_2) \setminus B|$ can be seen either as $|(B_1 \setminus B) \cap B_2|$ or as $|(B_2 \setminus B) \cap B_1|$,

$$\|\chi_B - \chi_{B_1} - \chi_{B_2}\|_{L_1} = |B \cap B_1 \cap B_2| + |B \setminus (B_1 \cup B_2)| + |B_1 \setminus B| + |B_2 \setminus B|.$$

Using (3.20) and (3.19) we obtain (3.15); from (3.21) we obtain (3.16). \square

From (3.15) we see that the minimization problem in the conclusion of Lemma 3.6 is equivalent to

$$\max\{|B \cap (B_1 \cup B_2)| : B, B_1, B_2 \text{ balls of radii } R, R_1, R_2\}, \quad (3.22)$$

where R, R_1, R_2 are such that $|E| = \omega_n R^n$, $|E_1| = \omega_n R_1^n$, $|E_2| = \omega_n R_2^n$.

Lemma 3.8. *Suppose $0 < R_1, R_2 < R < R_1 + R_2$. Then (3.22) admits a solution, unique up to isometries of the plane, characterized by the facts that:*

- i) *the centres of B, B_1, B_2 are aligned*
- ii) *$\emptyset \neq B_1 \cap B_2 \subset B$, $B_1 \not\subset B$, and $B_2 \not\subset B$*
- iii) *$\partial B \cap \partial B_1$, $\partial B_1 \cap \partial B_2$, and $\partial B_2 \cap \partial B$ are $((n-2)$ -dimensional) circles having the same radius (or, if $n = 2$, the common chords between B and B_1 , B_1 and B_2 , and B_2 and B all three have the same length, see Figure 8a).*

In addition, the solution to (3.22) is such that

$$|B \cap B_1 \cap B_2| \geq \frac{2^{n-1}}{n!} (R_1 + R_2 - R)^{\frac{n+1}{2}} \left(\frac{R_1 R_2}{R_1 + R_2} \right)^{\frac{n-1}{2}}. \quad (3.23)$$

The proof of Lemma 3.8 uses the auxiliary Lemmas 3.9 and 3.10. As mentioned in Section 2, we write $\mathbf{a} \wedge \mathbf{b}$ to denote the exterior product of $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$. In particular, we use that $|\mathbf{a} \wedge \mathbf{b}| = |\mathbf{b}| \operatorname{dist}(\mathbf{a}, \langle \mathbf{b} \rangle)$. The purpose of Lemma 3.9 is to show that $B(\mathbf{p} + h\mathbf{e}, R)$ can be written as the intersection of the two sets in Figure 8b), for all $h \in \mathbb{R}$. We then write the derivative of the area of the sublevel sets with respect to h as a surface integral on $\partial B(\mathbf{p} + h\mathbf{e}, R)$, using the coarea formula (Lemma 3.10).

Lemma 3.9. *Let $R > 0$, $\mathbf{p} \in \mathbb{R}^n$, $\mathbf{e} \in \mathbb{S}^{n-1}$. Define*

$$\begin{aligned} \phi(\mathbf{y}) &:= (\mathbf{y} - \mathbf{p}) \cdot \mathbf{e} - \sqrt{R^2 - |(\mathbf{y} - \mathbf{p}) \wedge \mathbf{e}|^2} \\ \psi(\mathbf{y}) &:= (\mathbf{y} - \mathbf{p}) \cdot \mathbf{e} + \sqrt{R^2 - |(\mathbf{y} - \mathbf{p}) \wedge \mathbf{e}|^2} \end{aligned}$$

in the infinite slab $S := \{\mathbf{y} \in \mathbb{R}^n : |(\mathbf{y} - \mathbf{p}) \wedge \mathbf{e}| < R\}$. Then, for all $h \in \mathbb{R}$,

$$B(\mathbf{p} + h\mathbf{e}, R) = \{\mathbf{y} \in S : \phi(\mathbf{y}) < h\} \cap \{\mathbf{y} \in S : \psi(\mathbf{y}) > h\}.$$




Figure 8: On the left: optimal choice of B , B_1 and B_2 in (3.23), with $h = h_1 = h_2$. On the right: sublevel sets $\{\phi < h\}$ and $\{\psi > h\}$ in the proof of Lemma 3.10 (as h increases the level sets move along the slab S , in the direction of \mathbf{e}).

Proof. By Pithagoras's theorem $|\mathbf{y} - (\mathbf{p} + h\mathbf{e})|^2 = |(\mathbf{y} - \mathbf{p}) \cdot \mathbf{e} - h|^2 + |(\mathbf{y} - \mathbf{p}) \wedge \mathbf{e}|^2$. Then $\mathbf{y} \in B(\mathbf{p} + h\mathbf{e}, R)$ if and only if $\mathbf{y} \in S$ and $|(\mathbf{y} - \mathbf{p}) \cdot \mathbf{e} - h| < \sqrt{R^2 - |(\mathbf{y} - \mathbf{p}) \wedge \mathbf{e}|^2}$, that is, if and only if

$$\begin{aligned} \mathbf{y} \in S, \quad (\mathbf{y} - \mathbf{p}) \cdot \mathbf{e} \geq h \quad \text{and} \quad \phi(\mathbf{y}) < h, \\ \text{or} \quad \mathbf{y} \in S, \quad (\mathbf{y} - \mathbf{p}) \cdot \mathbf{e} \leq h \quad \text{and} \quad \psi(\mathbf{y}) > h. \end{aligned}$$

This proves that $B(\mathbf{p} + h\mathbf{e}, R) \subset \{\phi < h\} \cap \{\psi > h\}$,

$$\begin{aligned} \{\phi < h\} \setminus B(\mathbf{p} + h\mathbf{e}, R) &\subset \{\mathbf{y} \in \mathbb{R}^n : (\mathbf{y} - \mathbf{p}) \cdot \mathbf{e} < h\} \\ \text{and} \quad \{\psi > h\} \setminus B(\mathbf{p} + h\mathbf{e}, R) &\subset \{\mathbf{y} \in \mathbb{R}^n : (\mathbf{y} - \mathbf{p}) \cdot \mathbf{e} > h\}. \end{aligned}$$

From this we see that $\{\phi < h\} \cap \{\psi > h\} \subset B(\mathbf{p} + h\mathbf{e}, R)$, so the conclusion follows. \square

Lemma 3.10. *Let $\mathbf{p} \in \mathbb{R}^n$, $R > 0$, $E \subset \mathbb{R}^n$ measurable, and suppose that*

$$\mathcal{H}^{n-1}(\partial B(\mathbf{p}, R) \cap \partial E) = 0. \quad (3.24)$$

Then the map $\mathbf{y} \mapsto |B(\mathbf{y}, R) \cap E|$ is differentiable at $\mathbf{y} = \mathbf{p}$ with gradient

$$D_{\mathbf{y}}(|B(\mathbf{y}, R) \cap E|)|_{\mathbf{y}=\mathbf{p}} = \int_{\partial B(\mathbf{p}, R) \cap E} \frac{\mathbf{z} - \mathbf{p}}{R} d\mathcal{H}^{n-1}(\mathbf{z}).$$

Proof. Given $\mathbf{e} \in \mathbb{S}^{n-1}$ arbitrary, let ϕ , ψ , and S be as in Lemma 3.9. By definition of ϕ and ψ , we have that $\phi(\mathbf{y}) < (\mathbf{y} - \mathbf{p}) \cdot \mathbf{e} < \psi(\mathbf{y})$ for all $\mathbf{y} \in S$, hence

$$(\mathbf{y} - \mathbf{p}) \cdot \mathbf{e} \leq h \Rightarrow \phi(\mathbf{y}) < h \quad \text{and} \quad (\mathbf{y} - \mathbf{p}) \cdot \mathbf{e} \geq h \Rightarrow \psi(\mathbf{y}) > h$$

for all $h \in \mathbb{R}$. Thus, $\{\phi < h\} \cup \{\psi > h\} = S$ and is independent of h . From the elementary relation $|E \cap S_1 \cap S_2| + |E \cap (S_1 \cup S_2)| = |E \cap S_1| + |E \cap S_2|$ we obtain (first for the case $|E \cap S| < \infty$, then

for all measurable sets)

$$\begin{aligned}
& |E \cap B(\mathbf{p} + h\mathbf{e}, R)| - |E \cap B(\mathbf{p}, R)| \\
&= (|E \cap \{\phi < h\}| + |E \cap \{\psi > h\}| - |E \cap S|) - (|E \cap \{\phi < 0\}| + |E \cap \{\psi > 0\}| - |E \cap S|) \\
&= |E \cap \{0 \leq \phi < h\}| - |E \cap \{0 < \psi \leq h\}|.
\end{aligned}$$

Writing $\mathbf{y} \in S$ as $\mathbf{p} + \lambda\mathbf{e} + \mu\mathbf{e}'$, with $|\mathbf{e}'| = 1$ and $\mathbf{e} \perp \mathbf{e}'$, a direct computation shows that

$$D\phi(\mathbf{y}) = \mathbf{e} - \frac{\mu\mathbf{e}'}{\sqrt{R^2 - \mu^2}} \quad \text{and} \quad D\psi(\mathbf{y}) = \mathbf{e} + \frac{\mu\mathbf{e}'}{\sqrt{R^2 - \mu^2}}.$$

Hence, by the coarea formula and Pithagoras's theorem,

$$\begin{aligned}
|E \cap B(\mathbf{p} + h\mathbf{e}, R)| - |E \cap B(\mathbf{p}, R)| &= \int_0^h \left(\int_{\{\phi=\tau\} \cap E} \frac{d\mathcal{H}^{n-1}(\mathbf{y})}{|D\phi(\mathbf{y})|} - \int_{\{\psi=\tau\} \cap E} \frac{d\mathcal{H}^{n-1}(\mathbf{y})}{|D\psi(\mathbf{y})|} \right) d\tau \\
&= \int_0^h \int_{\partial B(\mathbf{p} + \tau\mathbf{e}, R) \cap E} \operatorname{sgn}(\lambda - \tau) \frac{\sqrt{R^2 - \mu^2}}{R} d\mathcal{H}^{n-1}(\mathbf{y}) d\tau \\
&= \mathbf{e} \cdot \int_0^h \int_{\partial B(\mathbf{p} + \tau\mathbf{e}, R) \cap E} \frac{\mathbf{y} - \mathbf{p} - \tau\mathbf{e}}{R} d\mathcal{H}^{n-1}(\mathbf{y}) d\tau.
\end{aligned}$$

Since h and \mathbf{e} are arbitrary, the above equation expresses that for all $\mathbf{h} \in \mathbb{R}^n$

$$|E \cap B(\mathbf{p} + \mathbf{h}, R)| - |E \cap B(\mathbf{p}, R)| = \mathbf{h} \cdot \int_0^1 \int_{\partial B(\mathbf{p}, R)} \frac{\mathbf{z} - \mathbf{p}}{R} \chi_{E - \tau\mathbf{h}}(\mathbf{z}) d\mathcal{H}^{n-1}(\mathbf{z}) d\tau.$$

Denoting $|\{\tau \in (0, 1) : \mathbf{z} + \tau\mathbf{h} \in E\}|$ by $\alpha(\mathbf{z}, \mathbf{h}, E)$, Fubini's theorem gives

$$\begin{aligned}
& \left| |E \cap B(\mathbf{p} + \mathbf{h}, R)| - |E \cap B(\mathbf{p}, R)| - \mathbf{h} \cdot \int_{\partial B(\mathbf{p}, R) \cap E} \frac{\mathbf{z} - \mathbf{p}}{R} d\mathcal{H}^{n-1}(\mathbf{z}) \right| \\
& \leq |\mathbf{h}| \int_{\partial B(\mathbf{p}, R)} (\chi_E(\mathbf{z}) - \alpha(\mathbf{z}, \mathbf{h}, E)) d\mathcal{H}^{n-1}(\mathbf{z}).
\end{aligned}$$

Due to the connexity of the line segment joining \mathbf{z} and $\mathbf{z} + \mathbf{h}$, if $\operatorname{dist}(\mathbf{z}, \partial E) \geq |\mathbf{h}|$ then either $\mathbf{z} \in \operatorname{Int} E$ and $\alpha(\mathbf{z}, \mathbf{h}, E) = \chi_E(\mathbf{z}) = 1$, or $\mathbf{z} \in \mathbb{R}^n \setminus \bar{E}$ and $\alpha(\mathbf{z}, \mathbf{h}, E) = \chi_E(\mathbf{z}) = 0$. Therefore,

$$\begin{aligned}
& \limsup_{\mathbf{h} \rightarrow 0} |\mathbf{h}|^{-1} \left| |E \cap B(\mathbf{p} + \mathbf{h}, R)| - |E \cap B(\mathbf{p}, R)| - \mathbf{h} \cdot \int_{\partial B(\mathbf{p}, R) \cap E} \frac{\mathbf{z} - \mathbf{p}}{R} d\mathcal{H}^{n-1}(\mathbf{z}) \right| \\
& \leq \lim_{\mathbf{h} \rightarrow 0} \mathcal{H}^{n-1}(\{\mathbf{z} \in \partial B(\mathbf{p}, R) : \operatorname{dist}(\mathbf{z}, \partial E) < |\mathbf{h}|\}) = \mathcal{H}^{n-1}(\partial B(\mathbf{p}, R) \cap \partial E),
\end{aligned}$$

completing the proof. \square

Remark 1. The example $\mathbf{p} = \mathbf{0}$, $R = 1$, $E = (-1, 1)^n \setminus B(\mathbf{0}, 1)$ shows that $|B(\mathbf{y}, R) \cap E|$ is not always differentiable with respect to \mathbf{y} if (3.24) is not satisfied. However, this condition holds in the situations to be considered in the sequel, namely, when E is a ball, the union of balls, or the intersection of balls of radii different from R .

Proof of Lemma 3.8. The existence of solutions to (3.22) can be easily deduced from the continuity of $|B \cap (B_1 \cup B_2)|$ with respect to the centres of B , B_1 , and B_2 . Let (B, B_1, B_2) be one such solution. We divide the proof of i)-iii) in the following steps.

Step 1: one of the following possibilities occur

$$\text{dist}(B_1 \cap B_2, B) > 0, \quad \text{dist}(B_1 \cap B_2, \mathbb{R}^n \setminus B) > 0, \quad \text{or} \quad B_1 \cap B_2 = \emptyset. \quad (3.25)$$

Suppose, looking for a contradiction, that neither $\overline{B_1 \cap B_2} \cap \overline{B} = \emptyset$ nor $\overline{B_1 \cap B_2} \subset B$. Then, by the connexity of $\overline{B_1 \cap B_2}$, there exists $\mathbf{x}_0 \in \overline{B_1 \cap B_2} \cap \partial B$. Let $B = B(\mathbf{p}, R)$, $\mathbf{e} := \frac{\mathbf{x}_0 - \mathbf{p}}{|\mathbf{x}_0 - \mathbf{p}|}$, and consider the following parametrization of $\partial B(\mathbf{p}, R)$ using spherical coordinates

$$\mathbf{f}(\theta, \boldsymbol{\xi}) := \mathbf{p} + (R \cos \theta) \mathbf{e} + (R \sin \theta) \boldsymbol{\xi}, \quad \theta \in [0, \pi], \quad \boldsymbol{\xi} \in \mathbb{S}_{\mathbf{e}}^{n-2} := \mathbb{S}^{n-1} \cap \langle \mathbf{e} \rangle^\perp.$$

Applying Lemma 3.10 to $E = \overline{B_1} \cup \overline{B_2}$ (see Remark 1)

$$\begin{aligned} \left. \frac{d}{dh} (|B(\mathbf{p} + h\mathbf{e}, R) \cap (B_1 \cup B_2)|) \right|_{h=0} &= \int_{\partial B \cap (B_1 \cup B_2)} \mathbf{e} \cdot \frac{\mathbf{z} - \mathbf{p}}{R} d\mathcal{H}^1(\mathbf{z}) \\ &= R^{n-1} \int_{\mathbb{S}_{\mathbf{e}}^{n-2}} \int_{\theta \in (0, \pi): \mathbf{f}(\theta, \boldsymbol{\xi}) \in E} \cos \theta (\sin \theta)^{n-2} d\theta d\mathcal{H}^{n-2}(\boldsymbol{\xi}) \end{aligned}$$

We can write the integral with respect to θ as

$$\int_0^{\pi/2} \cos \theta (\sin \theta)^{n-2} (\chi_E(\mathbf{f}(\theta, \boldsymbol{\xi})) - \chi_E(\mathbf{f}(\pi - \theta, \boldsymbol{\xi}))) d\theta.$$

If we prove that

$$\mathbf{f}(\pi - \theta, \boldsymbol{\xi}) \in \overline{B_1} \cup \overline{B_2} \Rightarrow \mathbf{f}(\theta, \boldsymbol{\xi}) \in \overline{B_1} \cup \overline{B_2} \quad \text{for every } \theta \in [0, \pi/2] \quad (3.26)$$

and that

$$\chi_E(\mathbf{f}(\theta, \boldsymbol{\xi})) - \chi_E(\mathbf{f}(\pi - \theta, \boldsymbol{\xi})) = 1 \quad \text{for all } (\theta, \boldsymbol{\xi}) \text{ in a set of positive measure,} \quad (3.27)$$

we will obtain that $\frac{d}{dh} (|B(\mathbf{p} + h\mathbf{e}, R) \cap (B_1 \cup B_2)|) > 0$ at $h = 0$. The contradiction will follow by noting that if (B, B_1, B_2) solves (3.22), then $D_{\mathbf{x}} |B(\mathbf{x}, R) \cap (B_1 \cup B_2)|$ must be zero at $\mathbf{x} = \mathbf{p}$.

Suppose that $\mathbf{f}(\pi - \theta_0, \boldsymbol{\xi}) \in \overline{B_i}$ for some $i = 1, 2$ and some $\theta_0 \in [0, \frac{\pi}{2}]$. Since $\overline{B_i} \cap \partial B$ is connected and contains $\mathbf{f}(0, \boldsymbol{\xi}) = \mathbf{x}_0$, its projection to the plane $\mathbf{p} + \langle \mathbf{e}, \boldsymbol{\xi} \rangle$ must contain the whole of the arc $\mathbf{f}(\theta, \boldsymbol{\xi})$, $\theta \in [0, \pi - \theta_0)$. This proves (3.26). In order to prove (3.27), define $\theta_1(\boldsymbol{\xi}) := \sup\{\theta \in [0, \pi] : \mathbf{f}(\theta, \boldsymbol{\xi}) \in \overline{B_1} \cup \overline{B_2}\}$. Arguing as before, we see that

$$|\{\theta \in [0, \pi] : \chi_E(\mathbf{f}(\theta, \boldsymbol{\xi})) - \chi_E(\mathbf{f}(\pi - \theta, \boldsymbol{\xi})) = 1\}| > 0 \quad (3.28)$$

unless $\theta_1(\boldsymbol{\xi}) = 0$ or $\theta_1(\boldsymbol{\xi}) = \pi$ (by continuity, if (3.28) holds for at least one $\boldsymbol{\xi} \in \mathbb{S}_{\mathbf{e}}^{n-2}$, then (3.27) follows). Since $R_1, R_2 < R$, in fact $\theta_1 = \pi$ is not possible (in that case \mathbf{x}_0 and $\mathbf{x}_0 - 2R\mathbf{e}$ would belong to some $\overline{B_i}$, but $\text{diam } \overline{B_i} = 2R_i < 2R$). It remains to rule out the possibility that $\theta_1(\boldsymbol{\xi}) = 0$

for all $\boldsymbol{\xi}$, that is, that $\overline{B} \cap (\overline{B_1} \cup \overline{B_2}) = \{\mathbf{x}_0\}$. If that were the case then B and B_1 would be tangent, so for all $h < R_1$ we would have that

$$|B(\mathbf{p} + h\mathbf{e}, R) \cap (B_1 \cup B_2)| \geq |B(\mathbf{p} + h\mathbf{e}, R) \cap B_1| > 0 = |B \cap (B_1 \cup B_2)|$$

and (B, B_1, B_2) would not be a solution to (3.22). This completes the proof.

Step 2: the centres of B, B_1, B_2 lie on a same line. In all the three cases considered in (3.25), $|B \cap B_1 \cap B_2| = |(B + \mathbf{h}) \cap B_1 \cap B_2|$ for every \mathbf{h} sufficiently small. Also, for given R, R_1, R_2 , the expression $|B(\mathbf{y}_i, R_i) \cap B(\mathbf{y}, R)|$ is a decreasing function of $|\mathbf{y} - \mathbf{y}_i|$, $i = 1, 2$. If \mathbf{y} were not in the line containing \mathbf{y}_1 and \mathbf{y}_2 , both $|\mathbf{y} - \mathbf{y}_1|$ and $|\mathbf{y} - \mathbf{y}_2|$ could be reduced by displacing \mathbf{y} towards that line. By (3.18), this would increase $|B \cap (B_1 \cup B_2)|$, contradicting the choice of (B, B_1, B_2) as a solution to (3.22).

Step 3: (B, B_1, B_2) satisfies ii)-iii). Moreover, these conditions uniquely determine the distances and relative positions between the centres (that is, the solution to (3.22) is unique up to isometries).

Let h, h_1 , and h_2 denote, respectively, the radii of $\partial B_1 \cap \partial B_2$, $\partial B \cap \partial B_1$, and $\partial B \cap \partial B_2$ (or the semi-lengths of the common chords between B_1 and B_2 , B and B_1 , and B and B_2 if $n = 2$) defining these radii (or lengths) as zero in case of empty intersection. By virtue of i), both $\mathbf{p}_1 - \mathbf{p}$ and $\mathbf{p}_2 - \mathbf{p}$ are parallel to $\mathbf{e} := \frac{\mathbf{p}_2 - \mathbf{p}_1}{|\mathbf{p}_2 - \mathbf{p}_1|}$, where $\mathbf{p}, \mathbf{p}_1, \mathbf{p}_2$ are the centres of B, B_1, B_2 , respectively. Setting $q_i := (\mathbf{p}_i - \mathbf{p}) \cdot \mathbf{e}$, $i = 1, 2$, and using Cartesian coordinates (y_1, \dots, y_n) with \mathbf{p} as the origin and \mathbf{e} in the direction of the y_1 -axis, we have that $B = B((0, 0, \dots, 0), R)$, $B_1 = B((q_1, 0, \dots, 0), R_1)$, $B_2 = B((q_2, 0, \dots, 0), R_2)$. By (3.18) and⁸ Lemma 3.10,

$$\begin{aligned} \frac{\partial}{\partial q_1} |B \cap (B_1 \cup B_2)| &= \frac{\partial}{\partial q_1} |B \cap B_1| - \frac{\partial}{\partial q_1} |(B \cap B_2) \cap B_1| \\ &= \int_{\partial B_1 \cap B} \frac{z_1 - q_1}{R_1} d\mathcal{H}^{n-1}(z_1, \dots, z_n) - \int_{\partial B_1 \cap (B \cap B_2)} \frac{z_1 - q_1}{R_1} d\mathcal{H}^{n-1}(z_1, \dots, z_n). \end{aligned}$$

In the first of the possibilities considered in (3.25), B cannot intersect both B_1 and B_2 , hence (B, B_1, B_2) is not optimal (for example, it would be better if B contained completely either B_1 or B_2). In the other two cases we have $\partial B_1 \cap (B \cap B_2) = \partial B_1 \cap B_2$. Parametrize ∂B_1 by

$$\mathbf{z} \in \partial B_1 \quad \Leftrightarrow \quad \mathbf{z} - \mathbf{p}_1 = (R_1 \cos \theta)\mathbf{e} + (R_1 \sin \theta)\boldsymbol{\xi}, \quad \theta \in [0, \pi], \quad \boldsymbol{\xi} \in \mathbb{S}_{\mathbf{e}}^{n-2} := \mathbb{S}^{n-1} \cap \langle \mathbf{e} \rangle^\perp.$$

By definition of \mathbf{e} , $q_1 < q_2$. Therefore, $\mathbf{z} \in \partial B_1 \cap B_2$ if and only if $\theta \in [0, \theta_2)$, where θ_2 is one of the two angles in $[0, \pi]$ such that by $h = R_1 \sin \theta_2$ (when $h = 0$, we choose $\theta_2 = 0$ or $\theta_2 = \pi$ according to whether $B_2 \cap B_1 = \emptyset$ or $B_2 \subset B_1$). Thus,

$$\frac{\partial}{\partial q_1} |(B \cap B_2) \cap B_1| = \mathcal{H}^{n-2}(\mathbb{S}_{\mathbf{e}}^{n-2}) \int_0^{\theta_2} R^{n-1} \cos \theta (\sin \theta)^{n-2} d\theta = \omega_{n-1} h^{n-1}.$$

As for the integral on $\partial B_1 \cap B$, the same argument shows that it equals $-(\operatorname{sgn} q_1)\omega_{n-1}h_1^{n-1}$. After obtaining the corresponding expression for $\frac{\partial}{\partial q_2} |B \cap B_2|$, and by virtue of the optimality of (B, B_1, B_2) , we obtain

$$\operatorname{sgn}(q_1)h_1^{n-1} + h^{n-1} = h^{n-1} - \operatorname{sgn}(q_2)h_2^{n-1} = 0.$$

⁸There is exactly one situation not covered by Lemma 3.10, namely when $R_1 = R_2$ and $B_1 = B_2 \subset\subset B$, but it is easy to see that this does not give a maximum of $|B \cap (B_1 \cup B_2)|$.




Figure 9: Relationship between h and the distance between the centres.

The case $h = h_1 = h_2 = 0$ is not optimal (due to the assumption $R < R_1 + R_2$), hence $q_1 < 0 < q_2$ and $h = h_1 = h_2 > 0$. This proves ii)-iii). It remains to show that q_1 , q_2 and h are uniquely determined by these conditions. Denoting the hyperplane containing the intersection of the boundaries of two (intersecting) balls B', B'' by $\Pi(B', B'')$, we have that the hyperplanes $\Pi(B_1, B)$, $\Pi(B_1, B_2)$, and $\Pi(B_2, B)$ are given by $\{y_1 = a_1\}$, $\{y_1 = a\}$, and $\{y_1 = a_2\}$, for some $a_1, a, a_2 \in \mathbb{R}$. Clearly, the following must be satisfied

$$\begin{aligned} (a_1 - q_1)^2 + h^2 &= R_1^2 & (a - q_1)^2 + h^2 &= R_1^2 & a_2^2 + h^2 &= R^2 \\ a_1^2 + h^2 &= R^2 & (a - q_2)^2 + h^2 &= R_2^2 & (a_2 - q_2)^2 + h^2 &= R_2^2. \end{aligned}$$

In particular, $|a_1| = |a_2| = \sqrt{R^2 - h^2}$, $|a_1 - q_1| = |a - q_1| = \sqrt{R_1^2 - h^2}$, and $|a - q_2| = |a_2 - q_2| = \sqrt{R_2^2 - h^2}$. Conditions ii)-iii) imply that $a_1 < q_1 < a < q_2 < a_2$ and $a_1 < 0 < a_2$. Therefore

$$q_1 = \sqrt{R_1^2 - h^2} - \sqrt{R^2 - h^2}, \quad q_2 = \sqrt{R^2 - h^2} - \sqrt{R_2^2 - h^2}, \quad (3.29)$$

which shows that q_1 and q_2 are determined by h . We also find that

$$a - q_1 = \sqrt{R_1^2 - h^2}, \quad q_2 - a = \sqrt{R_2^2 - h^2}. \quad (3.30)$$

Adding the equations in (3.30) and subtracting the equations in (3.29) yields (see Figure 9)

$$q_2 - q_1 = \sqrt{R^2 - h^2} = \sqrt{R_1^2 - h^2} + \sqrt{R_2^2 - h^2}. \quad (3.31)$$

We may assume, without loss of generality, that $R_2 < R_1$. Rewrite (3.31) as

$$\frac{R^2 - R_1^2}{\sqrt{R^2 - h^2} + \sqrt{R_1^2 - h^2}} - \sqrt{R_2^2 - h^2} = 0.$$

The expression at the left-hand side is increasing in h , and equals $R - (R_1 + R_2) < 0$ at $h = 0$, and $\frac{R^2 - R_1^2}{\sqrt{R^2 - R_2^2} + \sqrt{R_1^2 - R_2^2}} > 0$ at $h = R_2$. This shows that h is uniquely determined by R, R_1, R_2 , and hence the balls B_1, B_2 too.

Step 4: proof of (3.23). For each $k \in \{2, \dots, n\}$ denote by P_k the k -dimensional polyhedron with vertices (the convex hull of)

$$\{(q_2 - R_2)\mathbf{e}, (q_1 + R_1)\mathbf{e}\} \cup \{a\mathbf{e} \pm h\mathbf{e}_i : i = 2, \dots, k\}, \quad \mathbf{e}_i := \underbrace{(0, \dots, 1, \dots, 0)}_{i\text{-th position}}.$$

It is easy to see that $\mathcal{H}^2(P_2) = h\gamma$, where $\gamma := |(q_1 + R_1) - (q_2 - R_2)|$, and that $\mathcal{H}^k(P_k) = 2h\mathcal{H}^{k-1}(P_{k-1})/k$, for $k \in \{3, \dots, n\}$. Thus, $|P_n| = (2^{n-1}h^{n-1}\gamma)/n!$.

From the previous analysis, we have that $B_1 \cap B_2$ contains P_n . From this we obtain (3.23), since, by virtue of (3.31),

$$\gamma = R_1 + R_2 - \sqrt{R^2 - h^2} > R_1 + R_2 - R, \quad (3.32)$$

$$\text{and} \quad \gamma = \frac{h^2}{R_1 + \sqrt{R_1^2 - h^2}} + \frac{h^2}{R_2 + \sqrt{R_2^2 - h^2}} < \frac{(R_1 + R_2)h^2}{R_1 R_2}. \quad (3.33)$$

□

We finally prove the main result.

Proof of Proposition 1.2. We can assume that $|E_1|^{\frac{1}{n}} + |E_2|^{\frac{1}{n}} > |E|^{\frac{1}{n}}$ (otherwise the estimate is trivially true). By (3.16) and (3.23) we have that

$$\min(\|\chi_B - \chi_{B_1} - \chi_{B_2}\|_{L^1} - (|B| - |B_1| - |B_2|)) \geq \frac{2^n}{n!} (R_1 + R_2 - R)^{\frac{n+1}{2}} \left(\frac{R_1 R_2}{R_1 + R_2} \right)^{\frac{n-1}{2}},$$

where the minimum is taken over all balls B, B_1, B_2 with $|B| = |E|$, $|B_1| = |E_1|$, $|B_2| = |E_2|$, and R, R_1, R_2 are such that $|E| = \omega_n R^n$, $|E_1| = \omega_n R_1^n$, $|E_2| = \omega_n R_2^n$. Thus, by Lemma 3.6,

$$\frac{|E|D(E)^{\frac{n}{n-1}} + |E_1|D(E_1)^{\frac{n}{n-1}} + |E_2|D(E_2)^{\frac{n}{n-1}}}{|E| + |E_1 \cup E_2|} \geq C \frac{(R_1 + R_2 - R)^{\frac{n+1}{2} \frac{n}{n-1}}}{(R^n + R_1^n + R_2^n)^{\frac{n}{n-1}}} \left(\frac{R_1 R_2}{R_1 + R_2} \right)^{\frac{n}{2}}$$

The quantities $R^n + R_1^n + R_2^n$, $R_1^n + R_2^n$, and $(R_1 + R_2)^n$ are comparable, since we are assuming that $R < R_1 + R_2$ and by virtue of the identity $a^n + b^n \leq (a + b)^n \leq 2^{n-1}(a^n + b^n)$. Hence

$$(R^n + R_1^n + R_2^n)^{\frac{n}{n-1}} \leq C(R_1 + R_2)^{\frac{n^2}{n-1}} = C(R_1 + R_2)^{\frac{n(n+1)}{2(n-1)}} (R_1 + R_2)^{\frac{n}{2}},$$

which implies that

$$\frac{|E|D(E)^{\frac{n}{n-1}} + |E_1|D(E_1)^{\frac{n}{n-1}} + |E_2|D(E_2)^{\frac{n}{n-1}}}{|E| + |E_1 \cup E_2|} \geq C \left(\frac{R_1 + R_2 - R}{R_1 + R_2} \right)^{\frac{n(n+1)}{2(n-1)}} \frac{R_1^{\frac{n}{2}} R_2^{\frac{n}{2}}}{(R_1 + R_2)^n}. \quad (3.34)$$

By the mean value theorem, there exists ξ between R and $R_1 + R_2$ such that

$$R_1 + R_2 - R = \frac{(R_1 + R_2)^n - R_1^n - R_2^n}{n\xi^{n-1}} \left(\frac{(R_1 + R_2)^n - R^n}{(R_1 + R_2)^n - R_1^n - R_2^n} \right)$$

Since we are assuming that $R < R_1 + R_2$, then $\xi \leq R_1 + R_2$ and

$$\frac{R_1 + R_2 - R}{R_1 + R_2} \geq \frac{1}{n} \frac{(R_1 + R_2)^n - R_1^n - R_2^n}{(R_1 + R_2)^n} \left(\frac{(|E_1|^{\frac{1}{n}} + |E_2|^{\frac{1}{n}})^n - |E|}{(|E_1|^{\frac{1}{n}} + |E_2|^{\frac{1}{n}})^n - |E_1 \cup E_2|} \right). \quad (3.35)$$

Suppose now that $|E_1| \geq |E_2|$, so that $\frac{R_1}{R_1 + R_2} \geq \frac{1}{2}$. By the binomial theorem,

$$\frac{(R_1 + R_2)^n - R_1^n - R_2^n}{(R_1 + R_2)^n} = \sum_{k=1}^{n-1} \binom{n}{k} \left(\frac{R_1}{R_1 + R_2} \right)^{n-k} \left(\frac{R_2}{R_1 + R_2} \right)^k \geq \frac{n}{2^{n-1}} \frac{R_2}{R_1 + R_2} \quad (3.36)$$

(we have considered only the term corresponding to $k = 1$). Combining (3.35) with (3.36) we obtain

$$\left(\frac{R_1 + R_2 - R}{R_1 + R_2} \right)^{\frac{n(n+1)}{2(n-1)}} \geq C \left(\frac{R_2}{R_1 + R_2} \right)^{\frac{n(n+1)}{2(n-1)}} \left(\frac{(|E_1|^{\frac{1}{n}} + |E_2|^{\frac{1}{n}})^n - |E|}{(|E_1|^{\frac{1}{n}} + |E_2|^{\frac{1}{n}})^n - |E_1 \cup E_2|} \right)^{\frac{n(n+1)}{2(n-1)}}.$$

The conclusion follows from (3.34) and the above equation, considering that $\frac{R_1}{R_1 + R_2} \geq \frac{1}{2}$. \square

4 Upper bounds

As explained in the Introduction, we obtain the upper bounds of Theorem 2 and Corollary 1 by finding suitable test functions opening cavities of different shapes and sizes, the main difficulties being to satisfy the incompressibility constraint and the Dirichlet condition at the boundary. We split the problem into two: in Section 4.1 we define a family of incompressible, angle-preserving maps whose energy has the right singular behaviour as $\varepsilon \rightarrow 0$, with leading order $(v_1 + v_2)|\log \varepsilon|$, and serves to define the test maps close to the singularities. In Section 4.2 we extend those maps, using the existence results of Rivière & Ye [64], in order to match the boundary conditions.

4.1 Proof of Theorem 2

In order to compute the energy of the test functions, we will need the following auxiliary lemmas, whose proof is postponed to Section 4.3.

Lemma 4.1. *Let Ω be a domain in \mathbb{R}^n , star-shaped with respect to a point $\mathbf{a} \in \mathbb{R}^n$, with Lipschitz boundary parametrized by $\zeta \mapsto \mathbf{a} + q(\zeta)\zeta$, $\zeta \in \mathbb{S}^{n-1}$. Let $v \geq 0$ and define $\mathbf{u} : \mathbb{R}^n \setminus \{\mathbf{a}\} \rightarrow \mathbb{R}^n$ by*

$$\mathbf{u}(\mathbf{a} + r\zeta) := \lambda \mathbf{a} + f(r, \zeta)\zeta, \quad f(r, \zeta)^n := r^n + (\lambda^n - 1)q(\zeta)^n, \quad r \in (0, \infty), \quad \zeta \in \mathbb{S}^{n-1}, \quad (4.1)$$

with $\lambda^n := 1 + \frac{v}{|\Omega|}$. Then \mathbf{u} is a Lipschitz homeomorphism, $\det D\mathbf{u} \equiv 1$, $\mathbf{u}(\mathbf{x}) = \lambda \mathbf{x}$ for all $\mathbf{x} \in \partial\Omega$, $\mathbf{u}(\overline{\Omega} \setminus \{\mathbf{a}\}) = \lambda \overline{\Omega} \setminus \text{im}_\Gamma(\mathbf{u}, \mathbf{a})$, $\mathbf{u}(\mathbb{R}^n \setminus \Omega) = \mathbb{R}^n \setminus \lambda\Omega$, $|\text{im}_\Gamma(\mathbf{u}, \mathbf{a})| = v$, and for all r, ζ ,

$$r^{n-1} \left| \frac{D\mathbf{u}(\mathbf{a} + r\zeta)}{\sqrt{n-1}} \right|^n \leq C \left(r + |v|^{\frac{1}{n}} \frac{\max\{q, |Dq|\}}{|\Omega|^{\frac{1}{n}}} \right)^{n-1} + \left(\frac{q(\zeta)^n}{|\Omega|} + C \frac{\max\{q^{n-1}, |Dq|^{n-1}\} |Dq|}{|\Omega|} \right) \frac{v}{r}.$$

Lemma 4.2. *Suppose that $\tilde{\mathbf{a}} \in \mathbb{R}^n$, $0 \leq d \leq \rho$, and $\mathbf{a} = \tilde{\mathbf{a}} + d\mathbf{e}$ for some $\mathbf{e} \in \mathbb{S}^{n-1}$. Let $\zeta \mapsto \mathbf{a} + q(\zeta)\zeta$, $\zeta \in \mathbb{S}^{n-1}$ be the polar parametrization of $\partial B(\tilde{\mathbf{a}}, \rho)$ taking \mathbf{a} as the origin. Then*

i) for all $\zeta \in \mathbb{S}^{n-1}$, $|q(\zeta)| \leq 2\rho$, $|Dq(\zeta)| \leq 2d|\zeta \wedge \mathbf{e}|$, and $|Dq(\zeta)| \leq 2d \left| \frac{q(\zeta)}{\sqrt{\rho(\rho-d)}} \right|^2 |\zeta \wedge \mathbf{e}|$

ii) if $\zeta \cdot (\mathbf{a} - \tilde{\mathbf{a}}) < 0$ then $q(\zeta) \geq \rho |\zeta \cdot \mathbf{e}|$ and $1 \leq \frac{q(\zeta)}{d|\zeta \cdot \mathbf{e}| + \sqrt{\rho(\rho-d)}} \leq 2$

iii) if $\zeta \cdot (\mathbf{a} - \tilde{\mathbf{a}}) > 0$ then $\frac{q(\zeta)}{\sqrt{\rho(\rho-d)}} \leq \frac{\sqrt{8}}{1 + \frac{d\zeta \cdot \mathbf{e}}{\sqrt{\rho(\rho-d)}}}$.

Lemma 4.3. Let $0 \leq d \leq \rho$, $\tilde{\mathbf{a}} \in \mathbb{R}^n$, $\mathbf{e} \in \mathbb{S}^{n-1}$, and $\Omega := \{\mathbf{x} \in B(\tilde{\mathbf{a}}, \rho) : (\mathbf{x} - \tilde{\mathbf{a}}) \cdot \mathbf{e} > \rho - 2d\}$. Then

$$n|\Omega| > \omega_{n-1} d^{\frac{n+1}{2}} (2\rho - d)^{\frac{n-1}{2}}.$$

Proof of Theorem 2. - Step 1: Construction of the domain.

Given $\mathbf{a}_1, \mathbf{a}_2 \in \mathbb{R}^n$ we claim that it is possible to find a slab of width $2d$ (where $d = |\mathbf{a}_2 - \mathbf{a}_1|$) and domains Ω_1 and Ω_2 , as in Figure 5, such that $\frac{|\Omega_2|}{|\Omega_1|} = \frac{v_2}{v_1}$. For ease of exposition, however, let us first fix $d > 0$, $\mathbf{e} \in \mathbb{S}^{n-1}$, and the slab $S = \{\mathbf{x} \in \mathbb{R}^n : |\mathbf{x} \cdot \mathbf{e}| < d\}$, and suppose that \mathbf{a}_1 and \mathbf{a}_2 (with $|\mathbf{a}_2 - \mathbf{a}_1| = d$) are still free to be chosen. Given ρ_1 and ρ_2 in $(0, d)$ define

$$B_1 = B((-d + \rho_1)\mathbf{e}, \rho_1) \quad \text{and} \quad B_2 = B((d - \rho_2)\mathbf{e}, \rho_2)$$

(the balls of radii ρ_1, ρ_2 contained in S and tangent to ∂S from the right and from the left). For future reference, note that if $\rho_i < \rho'_i$ then $B_i(\rho_i) \subset B_i(\rho'_i)$, $i = 1, 2$. If the balls intersect, let $\hat{a} \in (-d, d)$ be such that $\mathbf{x} \cdot \mathbf{e} = \hat{a}$ for all $\mathbf{x} \in \overline{B_1} \cap \overline{B_2}$ and define

$$\Omega_1 := \{\mathbf{x} \in B_1 : \mathbf{x} \cdot \mathbf{e} < \hat{a}\}, \quad \Omega_2 := \{\mathbf{x} \in B_2 : \mathbf{x} \cdot \mathbf{e} > \hat{a}\}, \quad \delta := \frac{2\rho_1 + 2\rho_2 - 2d}{2d} \quad (4.2)$$

(δ is the ratio between the width of $B_1 \cap B_2$ and that of $B_1 \cup B_2$). Set

$$d_1 := \frac{\hat{a} + d}{2} \quad \text{and} \quad d_2 := \frac{d - \hat{a}}{2}. \quad (4.3)$$

It is clear that \hat{a} , Ω_1 , Ω_2 , d_1 , d_2 , and δ , thus defined, are determined by ρ_1 and ρ_2 . Let

$$\rho_{\min} := \frac{v_1^{\frac{1}{n}} d}{v_1^{\frac{1}{n}} + v_2^{\frac{1}{n}}}, \quad B'_1 := ((-d + \rho_{\min})\mathbf{e}, \rho_{\min}), \quad B'_2 := (\rho_{\min}\mathbf{e}, d - \rho_{\min})$$

(definition analogous to that of B_1 and B_2 but with radii ρ_{\min} and $d - \rho_{\min}$). The reference radius ρ_{\min} is such that B'_1 and B'_2 are tangent, such that they fit precisely in the slab S , and such that $\frac{|B'_2|}{|B'_1|} = \frac{v_2}{v_1}$. If $0 < \rho_1 < \rho_{\min}$ and $\overline{B_1} \cap \overline{B_2} \neq \emptyset$ then $\Omega_1 \subset B'_1$ and $\Omega_2 \supset B'_2$, hence $\frac{|\Omega_2|}{|\Omega_1|} > \frac{v_2}{v_1}$. Therefore, if $0 < \rho_1 < \rho_{\min}$ there exists no $\rho_2 \in [0, d]$ such that $\overline{B_1}$ and $\overline{B_2}$ intersect and $\frac{|\Omega_2|}{|\Omega_1|} = \frac{v_2}{v_1}$. We are going to show that for every ρ_1 in the remaining interval $[\rho_{\min}, d)$ there exists one, and only one, $\rho_2 \in (0, d)$ such that $\frac{|\Omega_2|}{|\Omega_1|} = \frac{v_2}{v_1}$. In particular, we may regard δ as a function of ρ_1 , and we will see that δ increases, from 0 to 1, as ρ_1 increases from ρ_{\min} to d . From this we will conclude that for every $\delta \in [0, 1]$ there exist unique ρ_1 and ρ_2 for which Ω_1 and Ω_2 have the desired volume ratio.

Fix $\rho_1 \in [\rho_{\min}, d)$. In order for B_2 to intersect B_1 we must have that $\rho_2 \geq d - \rho_1$. Let $\rho'_2 < \rho''_2$ be any two such values of ρ_2 . Define Ω'_1, Ω'_2 as the domains obtained when $\rho_2 = \rho'_2$; analogous definition for Ω''_1, Ω''_2 . It is easy to see that $\Omega'_1 \supset \Omega''_1$ and $\Omega'_2 \subset \Omega''_2$ (the intersection plane moves to the left). Consequently, $\frac{|\Omega_2|}{|\Omega_1|}$ is strictly increasing with respect to ρ_2 . When $\rho_2 = d - \rho_1$ the ratio is $\frac{(d-\rho_1)^2}{\rho_1^2} \leq \frac{(d-\rho_{\min})^n}{\rho_{\min}^n} = \frac{v_2}{v_1} \leq 1$; when $\rho_2 = \rho_1$ the ratio is 1. This proves that there is exactly one ρ_2 for which the ratio is $\frac{v_2}{v_1}$. Moreover, the solution is such that $d - \rho_{\min} \leq \rho_2 < d$ and $\rho_1 + \rho_2 \geq d$.

Let us now prove that δ is increasing in ρ_1 and that it goes from 0 to 1. Let $\rho_1 \in [\rho_{\min}, d)$, $\rho_2 = \rho_2(\rho_1)$, and let Ω_1 and Ω_2 be the domains associated to ρ_1 and ρ_2 . Suppose that $\rho_1 < \rho'_1 < d$, and let Ω'_1 and Ω'_2 be the domains associated to the pair (ρ'_1, ρ_2) . It is easy to see that $\Omega'_1 \supset \Omega_1$ and $\Omega'_2 \subset \Omega_2$ (the intersection plane moves to the right). Hence $\frac{|\Omega'_2|}{|\Omega'_1|} < \frac{|\Omega_2|}{|\Omega_1|} = \frac{v_2}{v_1}$. In the above paragraph we showed that for every fixed ρ_1 , the ratio $\frac{|\Omega_2|}{|\Omega_1|}$ is increasing in ρ_2 . Applying this to ρ'_1 , and since the desired volume ratio for ρ'_1 is larger than $\frac{|\Omega'_2|}{|\Omega'_1|}$, the value ρ'_2 associated to ρ'_1 must be larger than ρ_2 . We conclude that ρ_2 (and, hence, δ , by virtue of (4.2)) is increasing as a function of ρ_1 . When $\rho_1 = \rho_{\min}$, it is clear that $\rho_2 = d - \rho_{\min}$ and $\delta = 0$. It can be shown that as $\rho_1 \rightarrow d$, also $\rho_2 \rightarrow d$, and, therefore, $\delta \rightarrow 1$. To prove this, note first that $B_2 \subset B'_2 := B(\mathbf{0}, d)$ and that $|B'_2 \setminus B_1| \rightarrow 0$ as $\rho_1 \rightarrow d$ (in the limit, B_1 coincides with B'_2). Note also that $B_1 \subset \Omega_1 \cup \Omega_2$ (see Figure 5). Then

$$\lim_{\rho_1 \rightarrow d} \frac{|\Omega_1|}{|B_1|} = \lim_{\rho_1 \rightarrow d} \frac{|\Omega_1|}{|\Omega_1 \cup \Omega_2|} \left(1 + \frac{|(\Omega_1 \cup \Omega_2) \setminus B_1|}{|B_1|} \right) = \frac{v_1}{v_1 + v_2} \left(1 + \frac{\lim_{\rho_1 \rightarrow d} |B_2 \setminus B_1|}{\omega_n d^n} \right) = \frac{v_1}{v_1 + v_2}.$$

For every $\rho_1 < d$, the intersection $B_1 \cap B_2$ is a set of the form

$$A(\rho_1) := \{\hat{a}(\rho_1)\mathbf{e} + r\mathbf{e}' : \mathbf{e}' \in \mathbb{S}^{n-1}, \mathbf{e}' \perp \mathbf{e}, r < \sqrt{\rho_1^2 - \hat{a}(\rho_1)^2}\}.$$

Since $\hat{a}(\rho_1)$ is determined by $\frac{|\Omega_1|}{|B_1|}$, it has a well-defined limit $\hat{a}(d)$ as $\rho_1 \rightarrow d$. For every $\rho_1 < d$, the sphere ∂B_2 can be characterized as the one containing $d\mathbf{e}$ (the right-most point of the ball) and the set $A(\rho_1)$. In the limit, it will be the sphere containing $d\mathbf{e}$ and $A(d)$ (unless $v_2 = 0$, $A(d)$ cannot consist only of $d\mathbf{e}$). But $A(\rho_1) \subset B_1(\rho_1)$ for all $\rho_1 < d$, and in the limit B_1 coincides with $B(\mathbf{0}, d)$ (which also contains $d\mathbf{e}$). Hence B_2 tends to coincide with B_1 , and $\rho_2 \rightarrow d$, as desired. The limiting domains are those given by (4.2), with \hat{a} given by the limiting value $\hat{a}(d)$.

Going back to the original statement, suppose that \mathbf{a}_1 and \mathbf{a}_2 are given, and let $d := |\mathbf{a}_2 - \mathbf{a}_1|$ and $\mathbf{e} := \frac{\mathbf{a}_2 - \mathbf{a}_1}{|\mathbf{a}_2 - \mathbf{a}_1|}$. For every $\delta \in [0, 1]$ define $\rho_1(\delta)$, $\rho_2(\delta)$, $d_1(\delta)$, and $d_2(\delta)$ as in the previous discussion (d_1 and d_2 are completely determined by ρ_1 and ρ_2 : they are the semi-distances from the intersection plane to the walls of the slab). The domains of Figure 5 are given by

$$\Omega_1 := \{\mathbf{x} \in B(\tilde{\mathbf{a}}_1, \rho_1) : (\mathbf{x} - \mathbf{a}_1) \cdot \mathbf{e} < d_1\}, \quad \Omega_2 := \{\mathbf{x} \in B(\tilde{\mathbf{a}}_2, \rho_2) : (\mathbf{x} - \mathbf{a}_2) \cdot \mathbf{e} > -d_2\}, \quad (4.4)$$

with

$$\tilde{\mathbf{a}}_1 := \mathbf{a}_1 + (\rho_1 - d_1)\mathbf{e}, \quad \tilde{\mathbf{a}}_2 := \mathbf{a}_2 - (\rho_2 - d_2)\mathbf{e}, \quad B_1 := B(\tilde{\mathbf{a}}_1, \rho_1), \quad B_2 := B(\tilde{\mathbf{a}}_2, \rho_2).$$

For future reference recall that $d_1 + d_2 = d$, $\rho_2 \leq \rho_1 \leq d$, and $\rho_1 + \rho_2 \geq d$. Note also that $2d_1$, the distance from the intersection plane to the left wall of the slab, is smaller than the diameter $2\rho_1$ of B_1 , that is, $d_1 \leq \rho_1$ and $d_2 \leq \rho_2$.

- *Step 2*: Definition of the map.

We define $\mathbf{u} : \mathbb{R}^n \setminus \{\mathbf{a}_1, \mathbf{a}_2\}$ piecewise, based on Lemma 4.1, in the following manner. Inside Ω_1 we apply Lemma 4.1 to $\Omega = \Omega_1$ and $\mathbf{a} = \mathbf{a}_1$; inside Ω_2 we apply Lemma 4.1 to $\Omega = \Omega_2$ and $\mathbf{a} = \mathbf{a}_2$. Finally, in order to define \mathbf{u} in $\mathbb{R}^n \setminus \Omega_1 \cup \Omega_2$ we define

$$\mathbf{a}^* = \frac{(\tilde{\mathbf{a}}_1 + \rho_1 \mathbf{e}) + (\tilde{\mathbf{a}}_2 - \rho_2 \mathbf{e})}{2} = \tilde{\mathbf{a}}_1 + (d - \rho_2) \mathbf{e} = \tilde{\mathbf{a}}_2 - (d - \rho_1) \mathbf{e}$$

(when $\delta = 0$, \mathbf{a}^* is the intersection point; when $\delta = 1$, \mathbf{a}^* is the center of the ball) and use Lemma 4.1 with $\Omega = \Omega_1 \cup \Omega_2$, $\mathbf{a} = \mathbf{a}^*$. Let $\zeta \mapsto \mathbf{a}_1 + q_1(\zeta)\zeta$, $\zeta \mapsto \mathbf{a}_2 + q_2(\zeta)\zeta$, and $\zeta \mapsto \mathbf{a}^* + q(\zeta)\zeta$ be, respectively, the polar parametrizations of $\partial\Omega_1$, $\partial\Omega_2$, and $\partial(\overline{\Omega_1} \cup \overline{\Omega_2})$ (with $\zeta \in \mathbb{S}^{n-1}$ in all cases). To be precise,

$$\mathbf{u}(\mathbf{x}) := \begin{cases} \lambda \mathbf{a}_1 + \left(|\mathbf{x} - \mathbf{a}_1|^n + \frac{v_1}{|\Omega_1|} q_1 \left(\frac{\mathbf{x} - \mathbf{a}_1}{|\mathbf{x} - \mathbf{a}_1|} \right)^n \right)^{\frac{1}{n}} \frac{\mathbf{x} - \mathbf{a}_1}{|\mathbf{x} - \mathbf{a}_1|} & \mathbf{x} \in \overline{\Omega_1} \setminus \{\mathbf{a}_1\} \\ \lambda \mathbf{a}_2 + \left(|\mathbf{x} - \mathbf{a}_2|^n + \frac{v_2}{|\Omega_2|} q_2 \left(\frac{\mathbf{x} - \mathbf{a}_2}{|\mathbf{x} - \mathbf{a}_2|} \right)^n \right)^{\frac{1}{n}} \frac{\mathbf{x} - \mathbf{a}_2}{|\mathbf{x} - \mathbf{a}_2|} & \mathbf{x} \in \overline{\Omega_2} \setminus \{\mathbf{a}_2\} \\ \lambda \mathbf{a}^* + \left(|\mathbf{x} - \mathbf{a}^*|^n + \frac{v_1 + v_2}{|\Omega_1 \cup \Omega_2|} q \left(\frac{\mathbf{x} - \mathbf{a}^*}{|\mathbf{x} - \mathbf{a}^*|} \right)^n \right)^{\frac{1}{n}} \frac{\mathbf{x} - \mathbf{a}^*}{|\mathbf{x} - \mathbf{a}^*|} & \mathbf{x} \in \mathbb{R}^n \setminus \overline{\Omega_1 \cup \Omega_2}, \end{cases}$$

with

$$\lambda^n - 1 := \frac{v_1}{|\Omega_1|} = \frac{v_2}{|\Omega_2|} = \frac{v_1 + v_2}{|\Omega_1 \cup \Omega_2|}.$$

Since $\frac{|\Omega_1|}{|\Omega_2|} = \frac{v_1}{v_2}$, the construction is well defined and $\mathbf{u}(\mathbf{x}) = \lambda \mathbf{x}$ for all $\mathbf{x} \in \partial\Omega_1 \cup \partial\Omega_2$. The resulting map is an incompressible homeomorphism, creates cavities at the desired locations with the desired volumes and is smooth except across $\partial\Omega_1 \cup \partial\Omega_2$ (where it is still continuous). It only remains to estimate its elastic energy.

- *Step 3* : Evaluation of the energy in $\mathbb{R}^n \setminus (\Omega_1 \cup \Omega_2)$.

By Lemma 4.2i), $\max\{q, |Dq|\} \leq 2d$, then, by Lemma 4.1

$$r^{n-1} \left| \frac{D\mathbf{u}(r\zeta)}{\sqrt{n-1}} \right|^n \leq C \left(r + \frac{d(v_1 + v_2)^{\frac{1}{n}}}{|\Omega_1 \cup \Omega_2|^{\frac{1}{n}}} \right)^{n-1} + \left(\frac{q^n}{|\Omega_1 \cup \Omega_2|} + \frac{Cd^{n-1}|Dq|}{|\Omega_1 \cup \Omega_2|} \right) \frac{v_1 + v_2}{r}.$$

Since ρ_i , $i = 1, 2$ increases with δ and assumes the value $\frac{v_i^{\frac{1}{n}} d}{v_1^{\frac{1}{n}} + v_2^{\frac{1}{n}}}$ when $\delta = 0$, it follows that

$$2\omega_n d^n > \omega_n(\rho_1^n + \rho_2^n) > |\Omega_1 \cup \Omega_2| > \frac{1}{2} \omega_n(\rho_1^n + \rho_2^n) > 2^{-n} \omega_n d^n. \quad (4.5)$$

Consequently, for any $R > 0$

$$\begin{aligned} \frac{1}{n} \int_{B(\mathbf{a}^*, R) \setminus \overline{\Omega_1 \cup \Omega_2}} \left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|^n d\mathbf{x} &\leq C(v_1 + v_2 + \omega_n R^n) + (v_1 + v_2) \frac{\int_{\mathbb{S}^{n-1}} \omega_n q^n \left(\log \frac{R}{q} \right)_+ d\mathcal{H}^{n-1}}{|\Omega_1 \cup \Omega_2|} \\ &\quad + C(v_1 + v_2) \int_{\mathbb{S}^{n-1}} \frac{|Dq|}{d} \left(\log \frac{R}{q} \right)_+ d\mathcal{H}^{n-1}, \end{aligned}$$

where $(\log x)_+$ denotes $\max\{0, \log x\}$. Note that $\int_{\mathbb{S}^{n-1}} \omega_n q(\zeta)^n d\mathcal{H}^{n-1}(\zeta) = |\Omega_1 \cup \Omega_2|$. Also, since $|\mathbf{a}^* - \tilde{\mathbf{a}}_1| + |\mathbf{a}^* - \tilde{\mathbf{a}}_2| = d(1 - \delta)$, Lemma 4.2i) implies that $|Dq| \leq 2d(1 - \delta)$. Therefore,

$$\begin{aligned} \frac{1}{n} \int_{B(\mathbf{a}^*, R) \setminus \overline{\Omega_1 \cup \Omega_2}} \left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|^n d\mathbf{x} &\leq C(v_1 + v_2 + \omega_n R^n) + (v_1 + v_2) \left(1 + C(1 - \delta)\right) \left(\log \frac{R}{d}\right)_+ \\ &+ C(v_1 + v_2) \int_{\mathbb{S}^{n-1}} \left(\frac{q^n}{d^n} + \frac{|Dq|}{d}\right) \left(\log \frac{d}{q(\zeta)}\right)_+ d\mathcal{H}^{n-1}(\zeta). \end{aligned}$$




Figure 10: Angle $\theta_0 > \frac{\pi}{2}$ and choice of spherical coordinates for $\delta = 0$.

The main problems at this point are that if $\delta \rightarrow 0$ then ρ_2 is of the order of $\frac{v_2 d}{v_1 + v_2}$ (so $\frac{d}{q} \rightarrow \infty$ on $\partial B_2 \cap \partial \Omega_2$ if $\frac{v_2}{v_1} \rightarrow 0$) and $q(\zeta)$ tends to vanish on $\partial B_1 \cap \partial B_2$ (see Figure 10). Parametrize \mathbb{S}^{n-1} by $\zeta = -\cos \theta \mathbf{e} + \sin \theta \boldsymbol{\xi}$ with $\theta \in (0, \pi)$ and $\boldsymbol{\xi} \in S := \mathbb{S}^{n-1} \cap \langle \mathbf{e} \rangle^\perp$. Since $\frac{q^n}{d^n} \left| \log \frac{d}{q} \right|$ is bounded we only study the term with $|Dq|$, that is, we are to prove that

$$\mathcal{H}^{n-2}(S) \left(\int_0^{\frac{\pi}{2}} + \int_{\frac{\pi}{2}}^\pi \right) (\sin \theta)^{n-2} \frac{|Dq(\zeta(\theta, \boldsymbol{\xi}))|}{d} \left(\log \frac{d}{q(\zeta(\theta, \boldsymbol{\xi}))} \right)_+ d\theta$$

is bounded independently of d , δ , v_1 , and v_2 . It can be shown that $\mathbf{a}^* + q(\theta, \boldsymbol{\xi})\zeta(\theta, \boldsymbol{\xi}) \in \partial B_1$ for all $\theta \in (0, \frac{\pi}{2})$ (due to the fact that $\rho_1 \geq \rho_2$, see Figure 10), and clearly $\zeta \cdot (\mathbf{a}^* - \tilde{\mathbf{a}}_1) = -\cos \theta (d - \rho_2) < 0$. Lemma 4.2ii) can thus be used to estimate the first integral by

$$2 \int_0^{\frac{\pi}{2}} \frac{\rho_1}{d} \log \frac{d}{\rho_1 \cos \theta} d\theta \leq 2 \left(\max_{t \in [0, 1]} |t \log t| \right) \int_0^{\frac{\pi}{2}} \left| \log \frac{1}{2} \left(\frac{\pi}{2} - \theta \right) \right| d\theta = \frac{\pi}{e} \left(1 + \log \frac{4}{\pi} \right).$$

As for the second integral we divide $(\frac{\pi}{2}, \pi)$ into $(\frac{\pi}{2}, \theta_0] \cup [\theta_0, \pi)$, according to whether $\mathbf{a}^* + q(\theta, \boldsymbol{\xi})\zeta(\theta, \boldsymbol{\xi})$ belongs to ∂B_1 or to ∂B_2 . For $\theta > \theta_0$ we can still use Lemma 4.2ii) (this time with $\tilde{\mathbf{a}} = \tilde{\mathbf{a}}_2$ and $\rho = \rho_2$) to obtain exactly the same upper bound as before. For $\theta \in (\frac{\pi}{2}, \theta_0)$, use parts i) and iii) of Lemma 4.2 together with $\rho_1 - |\mathbf{a}^* - \tilde{\mathbf{a}}_1| = d\delta$ to obtain

$$\frac{|Dq|}{d} \leq \frac{2(d - \rho_2)}{\delta \rho_1} \frac{q^2}{d^2} \sin \theta \quad \text{and} \quad |Dq| \leq 16(d - \rho_2) \left(1 + \frac{(d - \rho_2) |\cos \theta|}{\sqrt{\delta \rho_1 d}} \right)^{-2} \sin \theta.$$

Then, for any $\alpha \in (0, \frac{1}{2})$, using that $t^{2\alpha} |\log t| \leq (2\alpha e)^{-1}$ for every $t \in (0, 1)$,

$$\begin{aligned} \int_{\frac{\pi}{2}}^{\theta_0} \frac{|Dq|}{d} \left(\log \frac{d}{q} \right)_+ d\theta &\leq \int_{\frac{\pi}{2}}^{\theta_0} \left| \frac{Dq}{d} \right|^{1-\alpha} \left| \frac{Dq}{d} \right|^\alpha \left(\log \frac{d}{q} \right)_+ d\theta \\ &\leq \frac{2^\alpha 16^{1-\alpha}}{2\alpha e} \left(\frac{d-\rho_2}{d} \right)^{1-\alpha} \left(\frac{d-\rho_2}{\delta\rho_1} \right)^\alpha \int_{\frac{\pi}{2}}^\pi \left(1 + \frac{(d-\rho_2)|\cos\theta|}{\sqrt{\delta\rho_1 d}} \right)^{2(\alpha-1)} \sin\theta d\theta \\ &\leq \frac{8^{1-\alpha}}{\alpha e} \left(\frac{\delta\rho_1}{d} \right)^{\frac{1}{2}-\alpha} \int_0^{\frac{d-\rho_2}{\sqrt{\delta\rho_1 d}}} (1+t)^{2\alpha-2} dt \end{aligned}$$

The last integral can be bounded by means of the relation

$$(1-2\alpha) \int_0^x (1+t)^{2\alpha-2} dt = 1 - \frac{1}{(1+x)^{1-2\alpha}} < 1 - \frac{1}{1+x} = \frac{x}{1+x}.$$

Using that $\gamma + \sqrt{1-\gamma} > 1$ for all $\gamma \in (0, 1)$ (applied to $\gamma = \frac{d-\rho_2}{\rho_1} = \frac{|\mathbf{a}^* - \mathbf{a}_1|}{\rho_1}$),

$$\begin{aligned} \int_{\frac{\pi}{2}}^{\theta_0} \frac{|Dq|}{d} \left(\log \frac{d}{q} \right)_+ d\theta &\leq \frac{8^{1-\alpha}}{\alpha(1-2\alpha)e} \left(\frac{\delta\rho_1}{d} \right)^{\frac{1}{2}-\alpha} \frac{d-\rho_2}{\rho_1} \frac{1}{\gamma + \sqrt{1-\gamma}} \\ &\leq \frac{8^{1-\alpha}}{\alpha(1-2\alpha)e} \delta^{\frac{1}{2}-\alpha} \left(\frac{d-\rho_2}{d} \right)^{\frac{1}{2}-\alpha} \left(\frac{d-\rho_2}{\rho_1} \right)^{\frac{1}{2}-\alpha} \\ &\leq \frac{8^{1-\alpha}}{\alpha(1-2\alpha)e} \delta^{\frac{1}{2}-\alpha} (1-\delta)^{\frac{1}{2}-\alpha}. \end{aligned}$$

We conclude that for all $R > 0$

$$\frac{1}{n} \int_{B(\mathbf{a}^*, R) \setminus \overline{\Omega_1 \cup \Omega_2}} \left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|^n d\mathbf{x} \leq C(v_1 + v_2 + \omega_n R^n) + (v_1 + v_2) \left(1 + C(1-\delta) \right) \left(\log \frac{R}{d} \right)_+.$$

- *Step 4:* Estimating the energy in Ω_i .

Near the cavitation points we still have that $f \omega_n q_i^n d\mathcal{H}^{n-1} = |\Omega_i|$, $i = 1, 2$, so by Lemma 4.1

$$\begin{aligned} \frac{1}{n} \int_{\Omega_i \setminus B_{\varepsilon_i}(\mathbf{a}_i)} \left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|^n d\mathbf{x} &\leq C(v_i + \omega_n \rho_i^n) + v_i \left(\log \frac{2\rho_i}{\varepsilon_i} \right)_+ \\ &\quad + C \frac{v_1 + v_2}{|\Omega_1 \cup \Omega_2|} \left(\int_{\mathbb{S}^{n-1}} \max\{q_i, |Dq_i|\}^{n-1} |Dq_i| d\mathcal{H}^{n-1} \right) \log \frac{2d}{\varepsilon_i} \\ &\leq C(v_i + \omega_n \rho_i^n) + v_i \log \frac{2d}{\varepsilon_i} + C(v_1 + v_2) \frac{\rho_i^{n-1}}{d^{n-1}} \left(\int_{\mathbb{S}^{n-1}} \frac{|Dq_i|}{d} \right) \log \frac{2d}{\varepsilon_i}. \end{aligned}$$

For Ω_1 set $\zeta = -\cos\theta \mathbf{e} + \sin\theta \boldsymbol{\xi}$. If $\theta \in (0, \frac{\pi}{2})$ then, by Lemma 4.2, using that $|\mathbf{a}_1 - \tilde{\mathbf{a}}_1| = \rho_1 - d_1$,

$$\begin{aligned} \int_0^{\frac{\pi}{2}} |Dq_1| \sin^{n-2} \theta d\theta &\leq 16(\rho_1 - d_1) \int_0^{\frac{\pi}{2}} \left(1 + \frac{\rho_1 - d_1}{\sqrt{d_1 \rho_1}} \cos\theta \right)^{-2} \sin\theta d\theta \\ &= 16\sqrt{d_1 \rho_1} \int_0^{\frac{\rho_1 - d_1}{\sqrt{d_1 \rho_1}}} (1+t)^{-2} dt = \sqrt{\frac{d_1}{\rho_1}} \frac{\rho_1 - d_1}{\gamma + \sqrt{1-\gamma}}, \end{aligned}$$

with $\gamma = 1 - \frac{d_1}{\rho_1}$. Since $\gamma + \sqrt{1 - \gamma} \leq 1$ for $\gamma \in [0, 1]$,

$$\rho_1^{n-1} \int_0^{\frac{\pi}{2}} |Dq_1| \sin^{n-2} \theta \, d\theta \leq \rho_1^{n-2} \sqrt{d_1 \rho_1} (\rho_1 - d_1).$$

Define θ_1 as in Figure 5. By Lemma 4.2, $|Dq_1| \leq 2(\rho_1 - d_1) \sin \theta$ and $q_1 \geq \sqrt{d_1 \rho_1}$, hence

$$\rho_1 \int_{\frac{\pi}{2}}^{\theta_1} |Dq_1| \sin^{n-2} \theta \, d\theta \leq \rho_1 (\rho_1 - d_1) |\cos \theta_1| \leq (\rho_1 - d_1) \frac{d_1 \rho_1}{q_1(\theta_1)} \leq \sqrt{d_1 \rho_1} (\rho_1 - d_1).$$

For $\theta \in (\theta_1, \pi)$, $q_1(\boldsymbol{\zeta})$ is given by $q_1 \boldsymbol{\zeta} \cdot \mathbf{e} = d_1$ hence

$$q_1(\theta) = \frac{d_1}{\cos(\pi - \theta)} \quad \text{and} \quad |Dq_1(\boldsymbol{\zeta}(\theta, \boldsymbol{\xi}))| = \left| \frac{q_1(1 - \boldsymbol{\zeta} \otimes \boldsymbol{\zeta}) \mathbf{e}}{-\boldsymbol{\zeta} \cdot \mathbf{e}} \right| = \frac{d_1 \sin \theta}{\cos^2(\pi - \theta)}.$$

Using that $1 - |\cos \theta_1| = \frac{\sin^2 \theta_1}{1 + |\cos \theta_1|} \leq \sin^2 \theta_1$ and that $q_1(\theta_1) \geq (\rho_1 - d_1) \cos \theta_1 + \sqrt{d_1 \rho_1} \geq \sqrt{d_1 \rho_1}$,

$$\rho_1 \int_{\theta_1}^{\pi} |Dq_1| \, d\theta \leq d_1 \rho_1 \int_{|\cos \theta_1|}^1 \frac{dt}{t^2} \leq \rho_1 \frac{d_1 \sin^2 \theta_1}{\cos(\pi - \theta_1)} = \frac{\rho_1 (q_1(\theta_1) \sin \theta_1)^2}{q_1(\theta_1)} \leq 4\sqrt{d_1 \rho_1} (\rho_1 - d_1),$$

the last equality being due to the fact that $q_1(\theta_1) \cos \theta_1 = d_1$ and $\mathbf{a}_1 + q_1(\theta_1) \boldsymbol{\zeta}(\theta_1, \boldsymbol{\xi}) \in \partial B(\tilde{\mathbf{a}}_1, \rho_1)$. Now we show that $\max\{q_1, |Dq_1|\} \leq 8\rho_1$. The fact that $q_1(\theta_1) \geq \sqrt{d_1 \rho_1}$ implies that $\rho_1 |\cos \theta_1| \leq \sqrt{d_1 \rho_1}$. Clearly $q(\theta)$ is decreasing, therefore

$$q(\theta) \leq q(\theta_1) \leq 2((\rho_1 - d_1) |\cos \theta_1| + \sqrt{d_1 \rho_1}) \leq 4\sqrt{d_1 \rho_1} \leq 4\rho_1.$$

As for $|Dq_1|$, we have that $q_1(\theta) \sin \theta$ is decreasing and $q_1(\theta_1) \sin \theta_1 = 2\sqrt{d_1(\rho_1 - d_1)}$, then

$$|Dq_1| = \frac{q_1(q_1 \sin \theta)}{q_1 \cos(\pi - \theta)} \leq \frac{2q_1(\theta_1) \sqrt{d_1(\rho_1 - d_1)}}{d_1} \leq 8\sqrt{\rho_1(\rho_1 - d_1)} \leq 8\rho_1.$$

The study of \mathbf{u} in Ω_2 being completely analogous, the conclusion is that for all $R > 0$

$$\begin{aligned} \frac{1}{n} \int_{B(\mathbf{a}^*, R) \setminus (B_{\varepsilon_1}(\mathbf{a}_1) \cup B_{\varepsilon_2}(\mathbf{a}_2))} \left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|^n \, d\mathbf{x} &\leq C(v_1 + v_2 + \omega_n R^n) + v_1 \log \frac{R}{\varepsilon_1} + v_2 \log \frac{R}{\varepsilon_2} \\ &+ C(v_1 + v_2) \left((1 - \delta) \left(\log \frac{R}{d} \right)_+ + \sqrt{\frac{d_1}{d}} \frac{\rho_1 - d_1}{d} \log \frac{d}{\varepsilon_1} + \sqrt{\frac{d_2}{d}} \frac{\rho_2 - d_2}{d} \log \frac{d}{\varepsilon_2} \right) \end{aligned}$$

In the case of \mathbf{a}_1 it is $\rho_1 - d_1$ that has an interesting behaviour, whereas for \mathbf{a}_2 it is d_2 . This follows from our final ingredient: the ‘height’ of $B(\mathbf{a}_1, \rho_1) \cap B(\mathbf{a}_2, \rho_2)$, whether we measure it from the first ball or from the second, is the same. The corresponding expression is $d_1(\rho_1 - d_1) = d_2(\rho_2 - d_2)$. As a consequence,

$$\frac{\rho_1 - d_1}{d} = \frac{\delta(\rho_1 - d_1)}{(\rho_1 - d_1) + (\rho_2 - d_2)} = \frac{\delta d_2}{d_1 + d_2} = \delta \frac{d_2}{d}.$$

The theorem is thus proved since, by Lemma 4.3,

$$\left(\frac{d_2}{d} \right)^{\frac{n+1}{2}} \leq C \frac{|\Omega_2|}{\rho_2^{\frac{n-1}{2}} d^{\frac{n+1}{2}}} \leq C \frac{\frac{v_2 |\Omega_1 \cup \Omega_2|}{v_1 + v_2}}{\left(\frac{\frac{1}{v_1^n} + \frac{1}{v_2^n}}{d} \right)^{\frac{n-1}{2}} d^{\frac{n+1}{2}}} \leq C \left(\left(\frac{v_2}{v_1 + v_2} \right)^{\frac{1}{n}} \right)^{\frac{n+1}{2}}.$$

□

4.2 Transition to radial symmetry

Our proof of Theorem 3 is based on the following result (see [53, 20, 82, 51, 19, 8] for related work):

Proposition 4.4 (Rivière-Ye, [64], Thm. 8). *Let D be a smooth domain, $k = 0, 1, 2, \dots$ and suppose that $g \in C^{k,1}(\overline{D}) = W^{k+1,\infty}(D)$ with $\inf_D g > 0$ and $\int_D g = 1$. Then, there exists a diffeomorphism $\phi : \overline{D} \rightarrow \overline{D}$, satisfying $\det D\phi = g$ in D and $\phi = \mathbf{id}$ on ∂D , such that, for any $\alpha < 1$, ϕ is in $C^{k+1,\alpha}(\overline{D})$ and*

$$\|\phi - \mathbf{id}\|_{C^{k+1,\alpha}(\overline{D})} \leq C\|\mathbf{g} - 1\|_{C^{k,1}(\overline{D})}$$

for any $0 < \delta < 1$, where C depends only on $\alpha, k, D, \inf_D g, \delta$, and $\|g\|_{0,\delta}$.

Lemma 4.5. *Let $\zeta \in \mathbb{S}^{n-1} \mapsto \mathbf{a}^* + q(\zeta)\zeta$ be the polar parametrization of $\partial(\overline{\Omega_1 \cup \Omega_2})$ and define*

$$\rho(\zeta)^n := R_1^n + (v_1 + v_2) \frac{q(\zeta)^n}{|\Omega_1 \cup \Omega_2|}, \quad \zeta \in \mathbb{S}^{n-1}, \quad (4.6)$$

R_1 being fixed and such that $\Omega_1 \cup \Omega_2 \subset B(\mathbf{a}^*, R_1)$. Suppose that \mathbf{u} is a one-to-one incompressible map from $\{R_1 < |\mathbf{x} - \mathbf{a}^*| < R_2\}$ onto $\{r\zeta : \rho(\zeta) < r < R_3\}$, for some $R_2, R_3 \geq 0$. Then

$$\omega_n(R_2^n - R_1^n) > \frac{\frac{\pi}{3} - \frac{1}{2}}{2^{n-2}3\pi} (v_1 + v_2)(1 - \delta).$$

Proof. Denote $\max_{\mathbb{S}^{n-1}} q = 2\rho_1 - \delta d$ by q_{\max} . By incompressibility,

$$\omega_n R_3^n = v_1 + v_2 + \omega_n R_2^n, \quad (4.7)$$

hence the requirement that $R_3 \geq \rho(\zeta)$ for all $\zeta \in \mathbb{S}^{n-1}$ is equivalent to

$$\omega_n(R_2^n - R_1^n) > (v_1 + v_2) \frac{\omega_n \int_{\mathbb{S}^{n-1}} (q_{\max}^n - q^n) d\mathcal{H}^{n-1}}{|\Omega_1 \cup \Omega_2|}.$$

Write $\zeta := -\cos\theta \mathbf{e} + \sin\theta \boldsymbol{\xi}$ with $\theta \in [0, \pi]$, $\boldsymbol{\xi} \in S := \mathbb{S}^{n-1} \cap \langle \mathbf{e} \rangle^\perp$. For all $\theta \in (0, \frac{\pi}{2})$

$$\begin{aligned} q_{\max} - q(\theta) &= 2\rho_1 - \delta d - (\rho_1 - \delta d) \cos\theta - \sqrt{\delta d(2\rho_1 - \delta d) + (\rho_1 - \delta d)^2 \cos^2\theta} \\ &= \frac{(\rho_1 + (\rho_1 - \delta d)(1 - \cos\theta))^2 - (\delta d(2\rho_1 - \delta d) + (\rho_1 - \delta d)^2 \cos^2\theta)}{\rho_1 + (\rho_1 - \delta d)(1 - \cos\theta) + \sqrt{\delta d(2\rho_1 - \delta d) + (\rho_1 - \delta d)^2 \cos^2\theta}} \\ &> \frac{(\rho_1 - \delta d)^2(\sin^2\theta + (1 - \cos\theta)^2) + 2\rho_1(\rho_1 - \delta d)(1 - \cos\theta)}{(2\rho_1 - \delta d) + (2\rho_1 - \delta d) + \rho_1 - \delta d} \\ &= \frac{2(\rho_1 - \delta d)(2\rho_1 - \delta d)(1 - \cos\theta)}{5\rho_1 - 3\delta d} > \frac{2}{3}(d - \rho_2)(1 - \cos\theta) > \frac{2d}{3}(1 - \delta)(1 - \cos\theta), \end{aligned}$$

where we have used that $\rho_1 - \delta d = d - \rho_2$ and $\rho_2 \leq d$. Therefore,

$$\frac{\omega_n \int_{\mathbb{S}^{n-1}} (q_{\max}^n - q^n) d\mathcal{H}^{n-1}}{|\Omega_1 \cup \Omega_2|} > \frac{\mathcal{H}^{n-2}(S) \int_0^{\frac{\pi}{2}} (q_{\max} - q) q_{\max}^{n-1} (\sin\theta)^{n-2} d\theta}{n\omega_n 2d^n} > \frac{\frac{\pi}{3} - \frac{1}{2}}{2^{n-2}3\pi} (1 - \delta). \quad (4.8)$$

□

Proof of Theorem 3. We prove the theorem in the following stronger version (see the remark after the proof of Corollary 1): “Let R_1, R_2 be such that

$$R_1 \geq 2d \quad \text{and} \quad \omega_n(R_2^n - R_1^n) > 4^n n(v_1 + v_2)(1 - \delta) \quad (4.9)$$

($\delta, v_1, v_2, \mathbf{a}_1, \mathbf{a}_2, d, \varepsilon_1$, and ε_2 being as in the original statement). Then there exists \mathbf{a}^* , C_1, C_2 , and $\mathbf{u} : \mathbb{R}^n \setminus \{\mathbf{a}_1, \mathbf{a}_2\} \rightarrow \mathbb{R}^n$ such that $\mathbf{u}|_{\mathbb{R}^n \setminus B(\mathbf{a}^*, R_2)}$ is radially symmetric and for all $R \geq R_1$

$$\begin{aligned} \frac{1}{n} \int_{B(\mathbf{a}^*, R) \setminus (B_{\varepsilon_1}(\mathbf{a}_1) \cup B_{\varepsilon_2}(\mathbf{a}_2))} \left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|^n d\mathbf{x} &\leq C_1(v_1 + v_2 + \omega_n R^n) + v_1 \log \frac{R}{\varepsilon_1} + v_2 \log \frac{R}{\varepsilon_2} \\ &+ C_2(v_1 + v_2) \left((1 - \delta) \log \frac{R_1}{d} + \delta \left(\sqrt[n]{\frac{v_2}{v_1}} \log \frac{d}{\varepsilon_1} + \sqrt[2n]{\frac{v_2}{v_1}} \log \frac{d}{\varepsilon_2} \right) \right) \\ &+ \Sigma \left(\frac{(v_1 + v_2)(1 - \delta)}{\omega_n(R_2^n - R_1^n)} \right) (v_1 + v_2 + \omega_n R_2^n) \left(\frac{\min\{R, R_2\}}{R_1} - 1 \right)^n, \end{aligned}$$

the function Σ being such that $\Sigma(t) < \infty$ in for $t \in [0, \frac{1}{4^n n})$ and $\Sigma(t) = O(t^{n(n-1)})$ as $t \rightarrow 0$. The Theorem follows by choosing R_1 and R_2 as in (1.15).

Since the constant in Proposition 4.4 depends on the reference domain, we work on the annulus $D := \{\mathbf{z} \in \mathbb{R}^n : 1 \leq |\mathbf{z}| \leq \sqrt[n]{2}\}$ (we choose $\sqrt[n]{2}$ so that $|D| = \omega_n$). Our strategy is to define $\mathbf{u} : B(\mathbf{a}^*, R_1) \setminus \{\mathbf{a}_1, \mathbf{a}_2\} \rightarrow \mathbb{R}^n$ as in Theorem 2 and to look for a map

$$\mathbf{u} : \{\mathbf{x} \in \mathbb{R}^n : R_1 \leq |\mathbf{x} - \mathbf{a}^*| \leq R_2\} \rightarrow \{\mathbf{y} = \lambda \mathbf{a}^* + r\boldsymbol{\zeta} : \rho(\boldsymbol{\zeta}) \leq r \leq R_3, \boldsymbol{\zeta} \in \mathbb{S}^{n-1}\}$$

(where ρ is defined in (4.6)) of the form $\mathbf{u} = \mathbf{v} \circ \boldsymbol{\phi}^{-1} \circ \mathbf{w}^{-1}$, with $\boldsymbol{\phi} : \overline{D} \rightarrow \overline{D}$ a diffeomorphism and

$$\begin{aligned} \mathbf{w}(r\boldsymbol{\zeta}) &:= \mathbf{a}^* + ((2 - r^n)R_1^n + (r^n - 1)R_2^n)^{\frac{1}{n}} \boldsymbol{\zeta}, \\ \mathbf{v}(r\boldsymbol{\zeta}) &:= \lambda \mathbf{a}^* + ((2 - r^n)\rho(\boldsymbol{\zeta})^n + (r^n - 1)R_3^n)^{\frac{1}{n}} \boldsymbol{\zeta}. \end{aligned} \quad (4.10)$$

The maps \mathbf{w} and \mathbf{v} are parametrizations of the reference and target domains, and are defined so that $\det D\mathbf{w}$ is constant and $\mathbf{v} \circ \mathbf{w}^{-1}$ sends $\partial B(\mathbf{a}^*, R)$, $R_1 \leq R \leq R_2$ onto a curve enclosing a volume of exactly $v_1 + v_2 + \omega_n R^n$ (as can be seen by writing

$$\mathbf{v} \circ \mathbf{w}^{-1}(\mathbf{a}^* + R\boldsymbol{\zeta}) = \lambda \mathbf{a}^* + \left(R^n + \frac{v_1 + v_2}{\omega_n} \left(1 + \frac{R_2^n - R_1^n \omega_n (q^n - f q^n)}{R_2^n - R_1^n |\Omega_1 \cup \Omega_2|} \right) \right)^{\frac{1}{n}} \boldsymbol{\zeta}, \quad (4.11)$$

and by considering that $|\{\lambda \mathbf{a}^* + r\boldsymbol{\zeta} : \boldsymbol{\zeta} \in \mathbb{S}^{n-1}, 0 < r < \rho(\boldsymbol{\zeta})\}| = \int_{\mathbb{S}^{n-1}} \omega_n \rho^n d\mathcal{H}^{n-1}$). The problem for $\boldsymbol{\phi}$ is $\boldsymbol{\phi} = \mathbf{id}$ on ∂D , $\det D\boldsymbol{\phi} = g := \frac{\det D\mathbf{v}}{\det D\mathbf{w}}$ in D . To use Proposition 4.4 we need to bound

$$g(r\boldsymbol{\zeta}) - 1 = \frac{v_1 + v_2}{\omega_n(R_2^n - R_1^n)} \left(1 - \frac{\omega_n q(\boldsymbol{\zeta})^n}{|\Omega_1 \cup \Omega_2|} \right) \quad \text{and} \quad Dg(r\boldsymbol{\zeta}) = -\frac{v_1 + v_2}{R_2^n - R_1^n} \frac{nq^{n-1} Dq(\boldsymbol{\zeta})}{r|\Omega_1 \cup \Omega_2|}$$

for all $\boldsymbol{\zeta} \in \mathbb{S}^{n-1}$, $r \in [1, \sqrt[n]{2}]$. Using (4.5) and the fact that $\rho_1(\delta) \leq d$ and $q(\boldsymbol{\zeta}) \geq \delta d$ for all $\delta, \boldsymbol{\zeta}$,

$$\frac{\omega_n \int_{\mathbb{S}^{n-1}} (q_{\max}^n - q^n) d\mathcal{H}^{n-1}}{|\Omega_1 \cup \Omega_2|} \leq n(2d)^{n-1} \frac{(2\rho_1 - \delta d) - \delta d}{2^{-n} d^n} \leq 4^n n(1 - \delta). \quad (4.12)$$

By Lemma 4.2i),

$$\sup |Dg| \leq \frac{(v_1 + v_2) 2n(2d)^{n-1}(1 - \delta)d}{R_2^n - R_1^n} \leq 4^n n \frac{(v_1 + v_2)(1 - \delta)}{\omega_n(R_2^n - R_1^n)}.$$

This and Proposition 4.4 imply the existence of a (piecewise smooth) solution ϕ such that

$$\|\phi - \mathbf{id}\|_{C^1(\bar{D})} \leq \Sigma \left(\frac{(v_1 + v_2)(1 - \delta)}{\omega_n(R_2^n - R_1^n)} \right) \quad (4.13)$$

for some function Σ satisfying $\Sigma(t) < \infty$ for $t \in [0, \frac{1}{4^n n})$ and $\Sigma(t) = O(t)$ as $t \rightarrow 0$.

Define $\mathbf{u} = \mathbf{v} \circ \phi^{-1} \circ \mathbf{w}$. Writing $\mathbf{x} = \mathbf{w}(\phi(\mathbf{z}))$ we obtain

$$|D\mathbf{u}(\mathbf{x})|^n = \left| \frac{D\mathbf{v}(\mathbf{z}) \operatorname{adj} D\phi(\mathbf{z}) D\mathbf{w}^{-1}(\mathbf{x})}{\det D\phi(\mathbf{z})} \right|^n \leq C_n \frac{R_3^n}{R_1^n} \Sigma^{n(n-1)} \left(1 - 4^n n \frac{(v_1 + v_2)(1 - \delta)}{\omega_n(R_2^n - R_1^n)} \right)^{-n}.$$

The conclusion follows from (4.7). \square

Remark 2. For Dirichlet boundary conditions that are not necessarily radially symmetric, the above method can still be used provided there is an initial diffeomorphism \mathbf{v} , from the reference domain $D = \{\mathbf{z} : 1 < |\mathbf{z}| < \sqrt[n]{2}\}$ used above onto the desired target domain, for which $g := \frac{\det D\mathbf{v}}{\det D\mathbf{w}}$ is bounded away from zero. The final energy estimate will depend on $\inf_D g$, $\|D\mathbf{v}\|_\infty \|D\mathbf{w}^{-1}\|_\infty$, and $\|g\|_\infty + \|Dg\|_\infty$.

4.3 Proof of the preliminary lemmas

In this Section, we give the proofs of Lemmas 4.1, 4.2 and 4.3.

Proof of Lemma 4.1. First we show that for any map of the form $\mathbf{u}(\mathbf{x}) := \lambda \mathbf{a} + f(\mathbf{x}) \frac{\mathbf{x} - \mathbf{a}}{|\mathbf{x} - \mathbf{a}|}$ the incompressibility equation reduces to an ODE of the form $f^{n-1}(r, \zeta) \frac{\partial f}{\partial r}(r, \zeta) = r^{n-1}$. In order to see this, we consider any local parametrization $(s_1, \dots, s_{n-1}) \mapsto \zeta(s_1, \dots, s_{n-1})$ of \mathbb{S}^{n-1} and introduce a polar coordinate system of the form

$$\mathbf{x} = \mathbf{x}(r, s_1, \dots, s_{n-1}) = \mathbf{a} + r\zeta(s_1, \dots, s_{n-1}), \quad r > 0, \quad (s_1, \dots, s_{n-1}) \in D \subset \mathbb{R}^{n-1},$$

D being some parameter space. The claim follows by observing that

$$\frac{\partial \mathbf{u}}{\partial r} \wedge \frac{\partial \mathbf{u}}{\partial s_1} \wedge \dots \wedge \frac{\partial \mathbf{u}}{\partial s_{n-1}} = \det D\mathbf{u}(\mathbf{x}) \left(\frac{\partial \mathbf{x}}{\partial r} \wedge \frac{\partial \mathbf{x}}{\partial s_1} \wedge \dots \wedge \frac{\partial \mathbf{x}}{\partial s_{n-1}} \right) = \det D\mathbf{u}(\mathbf{x}) \left(\zeta \wedge \bigwedge_{k=1}^{n-1} r \frac{\partial \zeta}{\partial s_k} \right)$$

and

$$\frac{\partial \mathbf{u}}{\partial r} \wedge \frac{\partial \mathbf{u}}{\partial s_1} \wedge \dots \wedge \frac{\partial \mathbf{u}}{\partial s_{n-1}} = \frac{\partial f}{\partial r} \zeta \wedge \bigwedge_{k=1}^{n-1} \left(\frac{\partial f}{\partial s_k} \zeta + f \frac{\partial \zeta}{\partial s_k} \right) = f^{n-1} \frac{\partial f}{\partial r} \left(\zeta \wedge \bigwedge_{k=1}^{n-1} \frac{\partial \zeta}{\partial s_k} \right).$$

From the above we find that $\mathbf{u}(\mathbf{x}) := \lambda \mathbf{a} + f(\mathbf{x})\zeta$ is incompressible provided $f(r, \zeta)^n \equiv r^n + A(\zeta)^n$, for some $A : \mathbb{S}^{n-1} \rightarrow \mathbb{R}$. The definition in (4.1) is obtained by imposing the boundary condition $\mathbf{u}(\mathbf{x}) = \lambda \mathbf{x}$ on $\partial\Omega$. Differentiating f in (4.1) with respect to ζ yields

$$f^{n-1}(r, \zeta)D_\zeta f(r, \zeta) = (\lambda^n - 1)q^{n-1}(\zeta)Dq(\zeta), \quad D_\zeta f(r, \zeta), Dq(\zeta) : T_\zeta(\mathbb{S}^{n-1}) \rightarrow \mathbb{R}$$

($T_\zeta(\mathbb{S}^{n-1})$ being the tangent plane to \mathbb{S}^{n-1} at ζ). Writing $r(\mathbf{x}) = |\mathbf{x} - \mathbf{a}|$, $\zeta(\mathbf{x}) = \frac{\mathbf{x} - \mathbf{a}}{|\mathbf{x} - \mathbf{a}|}$ we obtain $Dr = \zeta$, $D\zeta = \frac{1 - \zeta \otimes \zeta}{r}$, $|D\zeta|^2 = \frac{n-1}{r^2}$. Identifying $D_\zeta f(r, \zeta) = \frac{v}{|\Omega|} \frac{q^{n-1}(\zeta)}{f^{n-1}(r, \zeta)} Dq(\zeta) \in (T_\zeta(\mathbb{S}^{n-1}))^*$ with a vector in $\langle \zeta \rangle^\perp \subset \mathbb{R}^n$ in the usual manner, from $f(x) = f(r(\mathbf{x}), \zeta(\mathbf{x}))$ we find that

$$Df(\mathbf{x}) = \frac{\partial f}{\partial r} Dr + (D\zeta)^T D_\zeta f = \frac{\partial f}{\partial r} \zeta + \frac{D_\zeta f}{r}, \quad |Df|^2 = \left| \frac{\partial f}{\partial r} \right|^2 + \left(\frac{v}{|\Omega|} \frac{q^{n-1}}{f^{n-1}} \frac{|Dq|}{r} \right)^2. \quad (4.14)$$

Since $D\mathbf{u} = \zeta \otimes Df + fD\zeta$, $(D\zeta) \cdot (\zeta \otimes Df) = \zeta \cdot ((D\zeta)Df) = 0$, and $\frac{\partial f}{\partial r} = \frac{r^{n-1}}{f^{n-1}} < 1$, then

$$|D\mathbf{u}|^2 = |Df|^2 + f^2 |D\zeta|^2 = (n-1) \frac{f^2}{r^2} + \left| \frac{\partial f}{\partial r} \right|^2 + \left| \frac{D_\zeta f}{r} \right|^2 \leq (n-1) \frac{f^2}{r^2} + 1 + \left| \frac{D_\zeta f}{r} \right|^2. \quad (4.15)$$

The leading order term $(v_1 + v_2)|\log \varepsilon|$ in the energy estimates will come from $(n-1)\frac{f^2}{r^2}$, hence we need to write $\left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|^n$ as $\frac{f^n}{r^n}$ plus a remainder term (for which we do not require an exact expression, only an upper bound). To this end we will write $a^n - b^n$, with $a = \left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|$ and $b = \sqrt{\frac{1}{n-1} + \frac{f^2}{r^2}}$, as

$$\begin{aligned} \left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|^n - \left(\frac{1}{n-1} + \frac{f^2}{r^2} \right)^{\frac{n}{2}} &\leq (a-b) |a^{n-1} + \dots + b^{n-1}| \leq n \frac{|a^2 - b^2|}{a+b} \max\{a, b\}^{n-1} \\ &\leq \frac{n}{n-1} \frac{|(D_\zeta f)/r|^2}{a} \left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|^{n-1} \leq C \frac{v^{\frac{1}{n}}}{|\Omega|^{\frac{1}{n}}} \frac{|Dq|}{r} \left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|^{n-1}, \end{aligned} \quad (4.16)$$

where we have used that $(n-1)a^2 \geq \frac{|D_\zeta f|^2}{r^2}$, and finally that $f^{n-1} \geq \left(\frac{v}{|\Omega|} q^n \right)^{\frac{n-1}{n}}$. Proceeding analogously, writing $c = \frac{f}{r}$ and $b^n - c^n \leq n \frac{b^2 - c^2}{b+c} b^{n-1}$, we obtain

$$\left(\frac{1}{n-1} + \frac{f^2}{r^2} \right)^{\frac{n}{2}} \leq \underbrace{1 + \frac{v}{|\Omega|} \frac{q^n}{r^n}}_{f^n/r^n} + C \left(1 + \frac{|v|^{\frac{1}{n}} q}{|\Omega|^{\frac{1}{n}} r} \right)^{n-1}. \quad (4.17)$$

Based on (4.15) and (4.14), we bound $\left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|^{n-1}$ by

$$\left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|^{n-1} \leq C \left(1 + \frac{v^{\frac{1}{n}}}{|\Omega|^{\frac{1}{n}}} \frac{\max\{q, |Dq|\}}{r} \right)^{n-1} \leq C \left(1 + \frac{v^{\frac{n-1}{n}}}{|\Omega|^{\frac{n-1}{n}}} \frac{\max\{q^{n-1}, |Dq|^{n-1}\}}{r^{n-1}} \right).$$

This combined with (4.16) and (4.17) yields

$$\begin{aligned} r^{n-1} \left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|^n &\leq r^{n-1} + C \left(r + \frac{|v|^{\frac{1}{n}} q}{|\Omega|^{\frac{1}{n}}} \right)^{n-1} + C \frac{v^{\frac{1}{n}}}{|\Omega|^{\frac{1}{n}}} |Dq| r^{n-2} \\ &\quad + \left(\frac{q^n}{|\Omega|} + C \frac{\max\{q^{n-1}, |Dq|^{n-1}\} |Dq|}{|\Omega|} \right) \frac{v}{r}. \end{aligned}$$

The proof is completed by substituting both r and $\frac{|v|^{\frac{1}{n}}}{|\Omega|^{\frac{1}{n}}}|Dq|$ with $r + |v|^{\frac{1}{n}} \frac{\max\{q, |Dq|\}}{|\Omega|^{\frac{1}{n}}}$. \square

Proof of Lemma 4.2. Write $\zeta = \cos \theta \mathbf{e} + \sin \theta \boldsymbol{\xi}$, $\theta \in (0, \pi)$, $\boldsymbol{\xi} \in \mathbb{S}^{n-1} \cap \langle \mathbf{e} \rangle^\perp$. By virtue of $|(\mathbf{a} + q(\zeta)\zeta) - \tilde{\mathbf{a}}|^2 \equiv \rho^2$,

$$q^2 + 2q\zeta \cdot (\mathbf{a} - \tilde{\mathbf{a}}) = \rho^2 - d^2, \quad q(\theta, \boldsymbol{\xi}) = -d \cos \theta + \sqrt{(\rho^2 - d^2) + d^2 \cos^2 \theta}.$$

Differentiate the first equation with respect to ζ and multiply by q

$$|Dq(\theta, \boldsymbol{\xi})| = \left| \frac{-2dq^2(1 - \zeta \otimes \zeta)\mathbf{e}}{q^2 + (q^2 + 2q\zeta \cdot (\mathbf{a} - \tilde{\mathbf{a}}))} \right| \leq \frac{2dq^2 \sin \theta}{(\rho - d)(\rho + d)} \leq \frac{2dq^2 \sin \theta}{\rho(\rho - d)}.$$

Part ii) is proved directly from the second equation for q , considering that $\sqrt{\rho^2 - d^2} \geq \sqrt{\rho(\rho - d)}$ and that $\gamma + \sqrt{1 - \gamma} \geq 1$ for all $\gamma \in (0, 1)$. It is clear that $q(\zeta) = \text{dist}(\mathbf{a} + q(\zeta)\zeta, \mathbf{a}) \leq \text{diam } B(\tilde{\mathbf{a}}, \rho) = 2\rho$ for all $\zeta \in \mathbb{S}^{n-1}$. The fact that $|Dq(\zeta)| \leq 2d \sin \theta$ follows from the first expression for $|Dq|$. Finally, if $\zeta \cdot \mathbf{e} = \cos \theta > 0$ then

$$\frac{2^{-1}}{1 + \frac{d \cos \theta}{\sqrt{\rho(\rho - d)}}} \leq \frac{q(\theta, \boldsymbol{\xi})}{\sqrt{\rho(\rho - d)}} = \frac{\sqrt{\rho - d} \left(\sqrt{\rho} + \frac{d}{\sqrt{\rho}} \right)}{\sqrt{(\rho^2 - d^2) + d^2 \cos^2 \theta} + d \cos \theta} \leq \frac{2\sqrt{2}}{1 + \frac{d \cos \theta}{\sqrt{\rho(\rho - d)}}}.$$

\square

Proof of Lemma 4.3. Call $\mathbf{a} := \tilde{\mathbf{a}} + (\rho - d)\mathbf{e}$. Consider the $(n - 2)$ -sphere $S := \{\mathbf{x} \in \partial B(\tilde{\mathbf{a}}, \rho) : (\mathbf{x} - \mathbf{a}) \cdot \mathbf{e} = 0\}$. It is clear that Ω contains the cone generated by $\tilde{\mathbf{a}} + \rho\mathbf{e}$ (the ‘right-most’ point on $\partial B(\tilde{\mathbf{a}}, \rho)$) and S . Since the radius of S (the ‘height’) is given by $h = \sqrt{d(2\rho - d)}$ (see Figure 11) and the base measures d , the volume of the cone is a constant times $dh^{n-1} = d^{\frac{n+1}{2}}(2\rho - d)^{\frac{n-1}{2}}$. The value of the constant is obtained from

$$|\Omega| \geq \frac{\mathcal{H}^{n-2}(\mathbb{S}^{n-2})}{n-1} \int_{\rho-d}^{\rho} \left(\frac{\rho - x_1}{d} \sqrt{\rho^2 - (\rho - d)^2} \right)^{n-1} dx_1 = \omega_{n-1} \sqrt{\rho(2\rho - d)}^{n-1} \frac{d}{n}.$$

\square




Figure 11: Cone generated by S and $\tilde{\mathbf{a}} + \rho\mathbf{e}$ (Lemma 4.3)

4.4 Numerical computations

The deformations depicted in Figure 6 are obtained by the alternative method of Dacorogna-Moser (constructive in nature and easier to implement, [20, Sect. 4]). Following the notation in Theorem 3 (and restricting now to the case $n = 2$), let $\rho(\theta) := \sqrt{R_1^2 + (v_1 + v_2) \frac{q(\theta)^2}{|\Omega_1 \cup \Omega_2|}}$, where $q(\theta)$ denotes the parametrization of $\partial(\overline{\Omega_1 \cup \Omega_2})$ using polar coordinates, taking \mathbf{a}^* to be the origin. Let also $0 < R_1 < R_2 < R_3$ be such that $B(\mathbf{a}^*, R_1) \supset \Omega_1 \cup \Omega_2$ and $\pi R_3^2 = v_1 + v_2 + \pi R_2^2$. Given parametrizations $\mathbf{w}(s, t)$ and $\mathbf{v}(s, t)$, $(s, t) \in D := [1, \sqrt{2}] \times [0, 2\pi]$ of $\{\mathbf{x} : R_1 < |\mathbf{x} - \mathbf{a}^*| < R_2\}$ and of $\{\mathbf{y} = \lambda \mathbf{a}^* + r e^{i\theta} : \rho(\theta) < r < R_3\}$, respectively, the strategy is to find an incompressible homeomorphism $\mathbf{u} : \mathbf{w}(Q) \rightarrow \mathbf{v}(Q)$ of the form

$$\mathbf{u} = \mathbf{v} \circ \phi_2 \circ \phi_1 \circ \mathbf{w}, \quad \text{with} \quad \phi_1(s, t) = (h(s, t), t), \quad \phi_2(s, t) = (s, t + \eta(s)\beta(t)).$$

Here $\eta : [1, \sqrt{2}] \rightarrow \mathbb{R}$ is any function satisfying

$$\int_1^{\sqrt{2}} \eta(s) ds = 1, \quad \eta(0) = \eta(1) = 0, \quad 0 \leq \eta \leq 1 + \varepsilon, \quad \int_1^{\sqrt{2}} |1 - \eta(s)| ds \leq \varepsilon$$

for some $\varepsilon \leq \min \left\{ \frac{\min f}{2 \max g}, \frac{\min g}{\max g} \right\}$, where $f(s, t) = \det D\mathbf{w}(s, t)$ and $g(s, t) = \det D\mathbf{v}(s, t)$. The functions β and h are found by defining $g_1(s_1, t_1) := g(\phi_2(s_1, t_1)) \det D\phi_2(s_1, t_1)$ and solving

$$\int_1^{\sqrt{2}} \int_0^{t + \eta(\sigma)\beta(t)} g(\sigma, \tau) d\tau d\sigma = \int_1^{\sqrt{2}} \int_0^t f(s, \bar{t}) d\bar{t} ds, \quad \int_1^{h(s, t)} g_1(s_1, t) ds_1 = \int_1^s f(\bar{s}, t) d\bar{s}$$

for every fixed $t \in [0, 2\pi]$. The solution is unique, and for \mathbf{v} and \mathbf{w} as in (4.10), it is such that $\int_{R_1 < |\mathbf{x} - \mathbf{a}^*| < R_2} |D\mathbf{u}|^2 \leq C$, where C is an expression that might possibly go to infinity only if the target domain is too narrow, more precisely, if $\frac{v_1 + v_2}{\pi(R_2^2 - R_1^2)} \left(\frac{\pi q_{\max}^2}{|\Omega_1 \cup \Omega_2|} - 1 \right) \nearrow 1$, (recall that $\frac{\pi q_{\max}^2}{|\Omega_1 \cup \Omega_2|} - 1$ is of the order of $1 - \delta$, equations (4.8) and (4.12)). In our computations we choose $R_1 = q_{\max} = 2\rho_1 - d\delta$ and R_2 such that $\pi(R_2^2 - R_1^2) = 2(v_1 + v_2) \left(\frac{\pi q_{\max}^2}{|\Omega_1 \cup \Omega_2|} - 1 \right)$.

5 Proof of the convergence result, Theorem 4

We follow the strategy of Struwe [77] to prove that $\sup_\varepsilon \|\mathbf{u}_\varepsilon\|_{W^{1,p}(\Omega_\varepsilon)} < \infty$ for all $p < n$. Fix $\varepsilon > 0$, call $\mathcal{B}_0 := \bigcup_{i=1}^m \overline{B}_\varepsilon(\mathbf{a}_{i,\varepsilon})$, $t_0 := r(\mathcal{B}_0) = m\varepsilon$, and let $\{\mathcal{B}(t) : t \geq t_0\}$ be the family obtained by applying Proposition 3.2 to \mathcal{B}_0 . Define $\rho = \sup\{t \geq t_0 : \bigcup \mathcal{B}(t) \subset \Omega\}$ and write $\mathcal{C}_k := \bigcup \mathcal{B}(r_k) \setminus \bigcup \mathcal{B}(r_{k+1})$, $r_k := 2^{-k}\rho$. By using Hölder's inequality, then comparing the lower bound of Proposition 3.4, to the upper bound, we find that for every $p < n$

$$\begin{aligned} \int_{\mathcal{C}_k} |D\mathbf{u}_\varepsilon|^p d\mathbf{x} &\leq C(n, p) \rho^{n-p} 2^{-(n-p)k} \left(\frac{1}{n} \int_{\Omega_\varepsilon} \left| \frac{D\mathbf{u}_\varepsilon}{\sqrt{n-1}} \right|^n d\mathbf{x} - \sum_{i=1}^m v_{i,\varepsilon} \log \frac{r_{k+1}}{t_0} \right)^{\frac{p}{n}} \\ &\leq C \rho^{n-p} 2^{-(n-p)k} \left(|\Omega| + \sum_{i=1}^m v_{i,\varepsilon} \right)^{\frac{p}{n}} \left(C + \log \frac{\text{diam } \Omega}{\rho/m} + (k+1) \log 2 \right)^{\frac{p}{n}}. \end{aligned}$$

Adding over k we find that

$$\begin{aligned}
\int_{\Omega_\varepsilon} |D\mathbf{u}_\varepsilon|^p \, d\mathbf{x} &\leq C\rho^{n-p} \left(|\Omega| + \sum_{i=1}^m v_{i,\varepsilon} \right)^{\frac{p}{n}} \left(\sum_{k=1}^{\infty} \frac{(C+k \log 2)^{\frac{p}{n}}}{2^{(n-p)k}} + \frac{\left(\log \frac{\text{diam } \Omega}{\rho/m} \right)^{\frac{p}{n}}}{2^{n-p} - 1} \right) \\
&\quad + n^{\frac{p}{n}} (n-1)^{\frac{p}{2}} |\Omega|^{1-\frac{p}{n}} \left(\frac{1}{n} \int_{\Omega_\varepsilon} \left| \frac{D\mathbf{u}_\varepsilon}{\sqrt{n-1}} \right|^n \, d\mathbf{x} - \sum_{i=1}^m v_{i,\varepsilon} \log \frac{\rho}{m\varepsilon} \right)^{\frac{p}{n}} \\
&\leq C \left(\rho^{n-p} + |\Omega|^{\frac{n-p}{n}} \right) \left(|\Omega| + \sum_{i=1}^m v_{i,\varepsilon} \right)^{\frac{p}{n}} \left(C + \log \frac{\text{diam } \Omega}{\rho/m} \right)^{\frac{p}{n}}.
\end{aligned}$$

It can be seen (as in the proof of Proposition 1.1) that $\rho \geq \frac{1}{2} \text{dist}(\{\mathbf{a}_{1,\varepsilon}, \dots, \mathbf{a}_{m,\varepsilon}\}, \partial\Omega)$. Hence, in order to prove that $\sup_\varepsilon \|D\mathbf{u}_\varepsilon\|_{L^p} < \infty$ it only remains to show that $\sum_{i=1}^m v_{i,\varepsilon}$ is uniformly bounded. Choose $r > \varepsilon$ such that the balls $\overline{B}(\mathbf{a}_{i,\varepsilon}, r)$ are disjoint and $r \in R_{\mathbf{a}_{i,\varepsilon}}$ for all $i = 1, \dots, m$. By Proposition 2.1, the topological images $E(\mathbf{a}_{i,\varepsilon}, r; \mathbf{u}_\varepsilon)$ are disjoint, contained in $B(\mathbf{0}, \|\mathbf{u}_\varepsilon\|_{L^\infty(\Omega_\varepsilon)})$ (because $E(\mathbf{a}_{i,\varepsilon}, r; \mathbf{u}_\varepsilon)$ is the region enclosed by $\mathbf{u}(\partial B(\mathbf{a}_{i,\varepsilon}, r))$), and such that $E(\mathbf{a}_{i,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon) \subset E(\mathbf{a}_{i,\varepsilon}, r; \mathbf{u}_\varepsilon)$. Therefore

$$\sum_{i=1}^m (v_{i,\varepsilon} + \omega_n \varepsilon^n) = \sum_{i=1}^m |E(\mathbf{a}_{i,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon)| \leq \left| \bigcup_{i=1}^m E(\mathbf{a}_{i,\varepsilon}, r; \mathbf{u}_\varepsilon) \right| \leq \omega_n \|\mathbf{u}_\varepsilon\|_{L^\infty(\Omega_\varepsilon)}^n.$$

Since we are assuming that $\sup_\varepsilon \|\mathbf{u}_\varepsilon\|_{L^\infty(\Omega_\varepsilon)} < \infty$, we obtain that $\sup_\varepsilon \|\mathbf{u}_\varepsilon\|_{W^{1,p}(\Omega_\varepsilon)} < \infty$, as desired.

For the existence of a limit map and for the convergence in $W_{\text{loc}}^{1,n}(\Omega \setminus \{\mathbf{a}_1, \dots, \mathbf{a}_m\}, \mathbb{R}^n)$, let $\delta > 0$ be small, assume that $|\mathbf{a}_{i,\varepsilon} - \mathbf{a}_i| < \delta/2$ for all $i = 1, \dots, m$, and consider the following energy bound, obtained again by comparing (1.16) with the lower bound of Proposition 3.4 (applied to $s = \delta/2$)

$$\frac{1}{n} \int_{\Omega \setminus \bigcup B(\delta/2)} \left| \frac{D\mathbf{u}}{\sqrt{n-1}} \right|^n \, d\mathbf{x} \leq \sum_{i=1}^m v_{i,\varepsilon} \log \frac{\text{diam } \Omega}{\delta/2m} + C \left(|\Omega| + \sum_{i=1}^m v_{i,\varepsilon} \right).$$

Since $r(B(\delta/2)) = \delta/2$, it follows that $\{\mathbf{u}_{\varepsilon_j}\}_{j \in \mathbb{N}}$ is bounded in $W^{1,n}(\Omega \setminus \bigcup_{i=1}^m \overline{B}_{\delta}(\mathbf{a}_i), \mathbb{R}^n)$. From this, and since $\delta > 0$ is arbitrary, the existence of \mathbf{u} and of a convergent subsequence follows by standard arguments (see, e.g., [74] or [37]): inductively take successive subsequences of $\{\mathbf{u}_{\varepsilon_j}\}_{j \in \mathbb{N}}$ (for some sequence $\delta_k \rightarrow 0$) converging weakly in $W^{1,n}(\Omega \setminus \bigcup_{i=1}^m \overline{B}_{\delta_k}(\mathbf{a}_i), \mathbb{R}^n)$. Choose then a diagonal sequence $\{\mathbf{u}_{\varepsilon_k}\}_{k \in \mathbb{N}}$ converging weakly in $W^{1,n}(\Omega \setminus \bigcup_{i=1}^m \overline{B}_{\delta}(\mathbf{a}_i), \mathbb{R}^n)$ for every $\delta > 0$, to some $\mathbf{u} \in W_{\text{loc}}^{1,n}(\Omega \setminus \{\mathbf{a}_1, \dots, \mathbf{a}_m\}, \mathbb{R}^n)$.

Since $\sup_\varepsilon \|\mathbf{u}_\varepsilon\|_{W^{1,p}(\Omega_\varepsilon)} < \infty$ for all $p < n$, the maps \mathbf{u}_ε can be extended, by multiplying them by suitable cut-off functions ψ_ε , inside the holes $\overline{B}(\mathbf{a}_{i,\varepsilon}, \varepsilon)$, in such a way that $\sup_\varepsilon \|\psi_\varepsilon \mathbf{u}_\varepsilon\|_{W^{1,p}(\Omega)} < \infty$. It is easy to see that any weakly convergent subsequence of $\{\psi_{\varepsilon_k} \mathbf{u}_{\varepsilon_k}\}_{k \in \mathbb{N}}$ must converge to the limit map \mathbf{u} defined above; this proves that $\mathbf{u} \in W^{1,p}(\Omega, \mathbb{R}^n)$ for all $p < n$.

By the classical result of Reshetnyak [61, Thm. 4] and Ball [2, Cor. 6.2.2], $\text{cof } D\mathbf{u}_{\varepsilon_k} \rightharpoonup \text{cof } D\mathbf{u}$ in $L_{\text{loc}}^{\frac{n}{n-1}}(\Omega \setminus \{\mathbf{a}_1, \dots, \mathbf{a}_m\}, \mathbb{R}^{n \times n})$. By the definition of $\text{Det } D\mathbf{u}$ in (2.4), and since $\{\text{Det } D\mathbf{u}_\varepsilon\}_{\varepsilon > 0}$ is bounded as a sequence in the space of measures ($\text{Det } D\mathbf{u}_\varepsilon = \mathcal{L}^n \llcorner \Omega_\varepsilon$, by hypothesis), it follows that $\text{Det } D\mathbf{u}$ coincides with \mathcal{L}^n in $\Omega \setminus \{\mathbf{a}_1, \dots, \mathbf{a}_m\}$, and that $\text{Det } D\mathbf{u}_\varepsilon \xrightarrow{*} \text{Det } D\mathbf{u}$ in $\Omega \setminus \{\mathbf{a}_1, \dots, \mathbf{a}_m\}$

in the sense of measures. Moreover, by [70, Lemma 3.2] (applied to $\Omega \setminus \bigcup_{i=1}^m \overline{B}(\mathbf{a}_i, \delta)$ instead of Ω), we obtain that $\det D\mathbf{u}(\mathbf{x}) = 1$ for a.e. $\mathbf{x} \in \Omega \setminus \{\mathbf{a}_1, \dots, \mathbf{a}_m\}$.

From Definition 5 and from the proof of [37, Lemma 4.2] it follows that the limit map \mathbf{u} satisfies condition INV. Proposition 2.2 then implies that $\text{Det } D\mathbf{u} = \mathcal{L}^n + \sum_{i=1}^m c_i \delta_{\mathbf{a}_i}$ for some coefficients $c_i \in \mathbb{R}$, and the proof of the same proposition also shows that

$$\begin{aligned} \frac{1}{n} \int_{\partial B(\mathbf{a}_i, r)} \mathbf{u}_\varepsilon \cdot (\text{cof } D\mathbf{u}_\varepsilon) \boldsymbol{\nu} \, d\mathcal{H}^{n-1} &= \omega_n r^n + \sum_{j: \mathbf{a}_j \in B(\mathbf{a}_i, r)} v_{j, \varepsilon} \\ \frac{1}{n} \int_{\partial B(\mathbf{a}_i, r)} \mathbf{u} \cdot (\text{cof } D\mathbf{u}) \boldsymbol{\nu} \, d\mathcal{H}^{n-1} &= \omega_n r^n + \sum_{j: \mathbf{a}_j \in B(\mathbf{a}_i, r)} c_j \end{aligned}$$

for a.e. $r > 0$ such that $\partial B(\mathbf{a}_i, r) \subset \Omega$ (note that if $\mathbf{a}_i = \mathbf{a}_j$ for some $i \neq j$, then the choice of the coefficients c_i is not unique). By standard arguments, for every $\delta > 0$ there exists $r < \delta$ such that $\mathbf{u}_{\varepsilon_k} \rightarrow \mathbf{u}$ uniformly on $\partial B(\mathbf{a}_i, r)$ and $\text{cof } D\mathbf{u}_{\varepsilon_k} \rightarrow \text{cof } D\mathbf{u}$ in $L^{\frac{n}{n-1}}(\partial B(\mathbf{a}_i, r))$ (passing, if necessary, to a subsequence that may depend on r). Taking, first, the limit as $\varepsilon \rightarrow 0$, then the limit as $r \rightarrow 0$, we obtain that $\text{Det } D\mathbf{u} = \mathcal{L}^n + \sum_{i=1}^m v_i \delta_{\mathbf{a}_i}$.

Consider now the case of two cavities. Set $\mathbf{a}_\varepsilon := \frac{\mathbf{a}_{1,\varepsilon} + \mathbf{a}_{2,\varepsilon}}{2}$, $d_\varepsilon := |\mathbf{a}_{2,\varepsilon} - \mathbf{a}_{1,\varepsilon}|$.

i) Suppose that $v_1 \geq v_2 > 0$ and $d = |\mathbf{a}_2 - \mathbf{a}_1| > 0$. By Lemma 3.5 we have that for all $r > \varepsilon$

$$\left| |E(\mathbf{a}_{i,\varepsilon}, r; \mathbf{u}_\varepsilon)| D(E(\mathbf{a}_{i,\varepsilon}, r; \mathbf{u}_\varepsilon))^{\frac{n}{n-1}} - |E(\mathbf{a}_{i,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon)| D(E(\mathbf{a}_{i,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon))^{\frac{n}{n-1}} \right| \leq 2^{\frac{n}{n-1}} \frac{n+1}{n-1} \omega_n r^n,$$

hence, by (3.5), for all $\alpha \in (0, 1)$ and all $R < \min\{\frac{d}{2}, \text{dist}(\{\mathbf{a}_1, \mathbf{a}_2\}, \partial\Omega)\}$ we have that

$$\begin{aligned} \frac{\int_{\Omega_\varepsilon} \frac{1}{n} \left| \frac{D\mathbf{u}(\mathbf{x})}{\sqrt{n-1}} \right|^n}{|\log \varepsilon|} &\geq \frac{\sum_{i=1}^2 \left(\int_\varepsilon^{\varepsilon^\alpha} + \int_{\varepsilon^\alpha}^R \right) \int_{\partial B(\mathbf{a}_{i,\varepsilon}, r)} \frac{1}{n} \left| \frac{D\mathbf{u}(\mathbf{x})}{\sqrt{n-1}} \right|^n \, d\mathcal{H}^{n-1} \, dr}{|\log \varepsilon|} \\ &\geq \sum_{i=1}^2 \left(v_{i,\varepsilon} \frac{\log(R/\varepsilon)}{|\log \varepsilon|} + (1-\alpha) C \left(|E(\mathbf{a}_{i,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon)| D(E(\mathbf{a}_{i,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon))^{\frac{n}{n-1}} - \varepsilon^{\alpha n} \right) \right). \end{aligned}$$

Combining this with (1.16) we obtain

$$\sum_{i=1}^2 v_{i,\varepsilon} D(E(\mathbf{a}_{i,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon))^{\frac{n}{n-1}} \leq \frac{(|\Omega| + v_{1,j} + v_{2,j}) (C_2 + \log \frac{\text{diam } \Omega}{R})}{C_1 |\log \varepsilon^{1-\alpha}|} + C \varepsilon^{\alpha n}.$$

Therefore, as $\varepsilon \rightarrow 0$, $D(E(\mathbf{a}_{i,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon)) \rightarrow 0$ (i.e., \mathbf{u}_ε tends to create spherical cavities).

As mentioned before, for every $\delta > 0$ there exists $r < \delta$ such that $\mathbf{u}_\varepsilon|_{\partial B(\mathbf{a}_i, r)}$ converges uniformly, for each $i = 1, 2$, to $\mathbf{u}|_{\partial B(\mathbf{a}_i, r)}$ (passing to a subsequence, if necessary). By continuity of the degree, this implies that $\text{im}_\Gamma(\mathbf{u}, \mathbf{a}_i)$ is contained in $E(\mathbf{a}_i, r; \mathbf{u}_\varepsilon)$ for sufficiently small ε . In particular, by definition of $v_{i,\varepsilon}$ and Proposition 2.2,

$$|E(\mathbf{a}_i, r; \mathbf{u}_\varepsilon) \Delta \text{im}_\Gamma(\mathbf{u}, \mathbf{a}_i)| = |E(\mathbf{a}_i, r; \mathbf{u}_\varepsilon)| - |\text{im}_\Gamma(\mathbf{u}, \mathbf{a}_i)| = (v_{i,\varepsilon} + \omega_n r^n) - v_i.$$

On the other hand, $B(\mathbf{a}_{i,\varepsilon}, \varepsilon) \subset B(\mathbf{a}_i, r)$ for sufficiently small ε . By Proposition 2.1 this implies that $E(\mathbf{a}_{i,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon) \subset E(\mathbf{a}_i, r; \mathbf{u}_\varepsilon)$, so, proceeding as in the proof of Proposition 2.2, we obtain

$$|E(\mathbf{a}_{i,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon) \Delta E(\mathbf{a}_i, r; \mathbf{u}_\varepsilon)| = \text{Det } D\mathbf{u}(B(\mathbf{a}_i, r) \setminus B(\mathbf{a}_{i,\varepsilon}, \varepsilon)) = |B(\mathbf{a}_i, r) \setminus B(\mathbf{a}_{i,\varepsilon}, \varepsilon)| < \omega_n \delta^n.$$

Thus,

$$\begin{aligned} & \limsup_{\varepsilon \rightarrow 0} |E(\mathbf{a}_{i,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon) \Delta \text{im}_T(\mathbf{u}, \mathbf{a}_i)| \\ & \leq \limsup_{\varepsilon \rightarrow 0} (|E(\mathbf{a}_{i,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon) \Delta E(\mathbf{a}_i, r; \mathbf{u}_\varepsilon)| + |E(\mathbf{a}_i, r; \mathbf{u}_\varepsilon) \Delta \text{im}_T(\mathbf{u}, \mathbf{a}_i)|) \leq 2\omega_n \delta^n \end{aligned} \quad (5.1)$$

for all $\delta > 0$, that is, the cavities formed by \mathbf{u}_ε converge to the cavities formed by \mathbf{u} .

Suppose, finally, that $v_{1,\varepsilon} + v_{2,\varepsilon} < \omega_n (2R)^n$ and that $B(\mathbf{a}_\varepsilon, R) \subset \Omega$ for some fixed $R > 0$ and all ε sufficiently small. If $\omega_n d^n \geq 2^{-n}(v_1 + v_2)$ there is nothing to prove, so assume, further, that $v_{1,\varepsilon} + v_{2,\varepsilon} > \omega_n (2d_\varepsilon)^n$. Then $\frac{R}{d_\varepsilon} > 1$ and $n\sqrt{\frac{v_{1,\varepsilon} + v_{2,\varepsilon}}{2^n \omega_n d_\varepsilon}} < \sqrt[n]{\frac{R}{d_\varepsilon}} < \frac{R}{d_\varepsilon}$. By Theorem 1 and (1.16), for sufficiently small $\varepsilon > 0$

$$\left(\left(\frac{v_{2,\varepsilon}}{v_{1,\varepsilon} + v_{2,\varepsilon}} \right)^{\frac{n}{n-1}} - \frac{\omega_n d_\varepsilon^n}{v_{1,\varepsilon} + v_{2,\varepsilon}} \right) \log \frac{v_{1,\varepsilon} + v_{2,\varepsilon}}{2^n \omega_n d_\varepsilon^n} \leq C \left(1 + \frac{|\Omega|}{v_{1,\varepsilon} + v_{2,\varepsilon}} + \log \frac{\omega_n (\text{diam } \Omega)^n}{\omega_n R^n} \right).$$

As $\varepsilon \rightarrow 0$, we obtain

$$\frac{\omega_n d^n}{v_1 + v_2} \geq \min \left\{ 2^{-n}, \frac{1}{2} \left(\frac{v_2}{v_1 + v_2} \right)^{\frac{n}{n-1}}, 2^{-n} F(\Omega, v_1, v_2) \right\},$$

with

$$F(\Omega, v_1, v_2) := \exp \left(- \frac{1 + \frac{|\Omega|}{v_1 + v_2} + \log \frac{\omega_n (2 \text{diam } \Omega)^n}{v_1 + v_2}}{C \left(\frac{v_2}{v_1 + v_2} \right)^{\frac{n}{n-1}}} \right). \quad (5.2)$$

Since $\exp(\frac{1}{h}) \gg \frac{1}{h}$ as $h \rightarrow 0^+$, there exists $C(n)$ such that $\frac{\omega_n d^n}{v_1 + v_2} \geq CF(\Omega, v_1, v_2)$.

ii) Suppose that $v_1 > v_2 = 0$. Applying Proposition 3.2 to $\mathcal{B}_0 := \{\overline{B}_\varepsilon(\mathbf{a}_{1,\varepsilon}), \overline{B}_\varepsilon(\mathbf{a}_{2,\varepsilon})\}$ we obtain $\mathcal{B}(t) = \{B(\mathbf{a}_{1,\varepsilon}, t/2), B(\mathbf{a}_{2,\varepsilon}, t/2)\}$ for $t \in (2\varepsilon, d_\varepsilon)$, and $\mathcal{B}(t) = \{B(\mathbf{a}_\varepsilon, t)\}$ for $t \geq d_\varepsilon$. We claim that if $R < \frac{2}{3} \text{dist}(\{\mathbf{a}_{1,\varepsilon}, \mathbf{a}_{2,\varepsilon}\}, \partial\Omega)$ then $\bigcup \mathcal{B}(R) \subset \Omega$. Indeed, if $R < d_\varepsilon$, this holds automatically. If $R \geq d_\varepsilon$, then

$$\frac{3R}{2} < \text{dist}(\mathbf{a}_{1,\varepsilon}, \partial\Omega) \leq \frac{d_\varepsilon}{2} + \text{dist}(\mathbf{a}_\varepsilon, \partial\Omega) \leq \frac{R}{2} + \text{dist}(\mathbf{a}_\varepsilon, \partial\Omega) \Rightarrow B(\mathbf{a}_\varepsilon, R) \subset \Omega.$$

Therefore, by Proposition 3.4 and Lemma 3.5, for every $\alpha \in (0, 1)$

$$\begin{aligned} & |E(\mathbf{a}_{1,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon)| D(E(\mathbf{a}_{1,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon))^{\frac{n}{n-1}} \log \frac{\varepsilon^\alpha}{2\varepsilon} \\ & \leq \int_{\Omega_\varepsilon} \frac{1}{n} \left| \frac{D\mathbf{u}_\varepsilon}{\sqrt{n-1}} \right|^n \mathbf{d}\mathbf{x} - (v_{1,\varepsilon} + v_{2,\varepsilon}) \log \frac{R}{2\varepsilon} + 2^{\frac{n}{n-1}} \frac{n+1}{n-1} (v_{2,\varepsilon} + \omega_n \varepsilon^{\alpha n}) \log \frac{\varepsilon^\alpha}{2\varepsilon}. \end{aligned}$$

By virtue of (1.16) and again Lemma 3.5,

$$v_1 D(\text{im}_T(\mathbf{u}, \mathbf{a}_1))^{\frac{n}{n-1}} \leq 2^{\frac{n}{n-1}} \frac{n+1}{n-1} \lim_{\varepsilon \rightarrow 0} (v_{2,\varepsilon} + \omega_n \varepsilon^{\alpha n} + |E(\mathbf{a}_{1,\varepsilon}, \varepsilon; \mathbf{u}_j) \Delta \text{im}_T(\mathbf{u}, \mathbf{a}_1)|).$$

Proceeding as in (5.1) we find that

$$\limsup_{\varepsilon \rightarrow 0} |E(\mathbf{a}_{1,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon) \Delta \text{im}_T(\mathbf{u}, \mathbf{a}_1)| \leq 2(v_2 + \omega_n r^n)$$

for arbitrarily small values of $r > 0$, proving that $\text{im}_T(\mathbf{u}, \mathbf{a}_1)$ is a ball.

iii) Suppose that $v_1 \geq v_2 > 0$ and $\mathbf{a}_1 = \mathbf{a}_2$. Let $R > 0$ be such that $B(\mathbf{a}_\varepsilon, R) \subset \Omega$ for all $j \in \mathbb{N}$. Since $\lim d_\varepsilon = |\mathbf{a}_2 - \mathbf{a}_1| = 0$, (3.6) and (1.16) imply that

$$\limsup_{\varepsilon \rightarrow 0} \frac{\int_{d_\varepsilon}^R |E(\mathbf{a}_\varepsilon, r; \mathbf{u}_\varepsilon)| D(E(\mathbf{a}_\varepsilon, r; \mathbf{u}_\varepsilon))^{\frac{n}{n-1}} \frac{dr}{r}}{\log d_\varepsilon} \leq C \frac{(|\Omega| + v_1 + v_2) \left(1 + \log \frac{\text{diam } \Omega}{R/2}\right)}{\lim_{\varepsilon \rightarrow 0} \log d_\varepsilon} = 0.$$

For $\alpha \in (0, 1)$ fixed and ε small $B(\mathbf{a}_\varepsilon, d_\varepsilon) \subset B(\mathbf{a}_\varepsilon, d_\varepsilon^\alpha) \subset \Omega$. By Lemma 3.5, for all $r \in (d_\varepsilon, d_\varepsilon^\alpha)$

$$\left| |E(\mathbf{a}_\varepsilon, r; \mathbf{u}_\varepsilon)| D(E(\mathbf{a}_\varepsilon, r; \mathbf{u}_\varepsilon))^{\frac{n}{n-1}} - |E(\mathbf{a}_\varepsilon, d_\varepsilon; \mathbf{u}_\varepsilon)| D(E(\mathbf{a}_\varepsilon, d_\varepsilon; \mathbf{u}_\varepsilon))^{\frac{n}{n-1}} \right| \leq 2^{\frac{n}{n-1}} \frac{n+1}{n-1} \omega_n d_\varepsilon^{\alpha n}.$$

Dividing $\int_{d_\varepsilon}^{d_\varepsilon^\alpha} |E(\mathbf{a}_\varepsilon, d_\varepsilon; \mathbf{u}_\varepsilon)| D(E(\mathbf{a}_\varepsilon, d_\varepsilon; \mathbf{u}_\varepsilon))^{\frac{n}{n-1}} \frac{dr}{r}$ by $\log d_\varepsilon^{\alpha-1}$ we obtain

$$\limsup_{\varepsilon \rightarrow 0} |E(\mathbf{a}_\varepsilon, d_\varepsilon; \mathbf{u}_\varepsilon)| D(E(\mathbf{a}_\varepsilon, d_\varepsilon; \mathbf{u}_\varepsilon))^{\frac{n}{n-1}} \leq \limsup_{\varepsilon \rightarrow 0} 2^{\frac{n}{n-1}} \frac{n+1}{n-1} \omega_n d_\varepsilon^{\alpha n} = 0. \quad (5.3)$$

Because of the continuity of the distributional determinant, $|\text{im}_T(\mathbf{u}, \mathbf{a})| = v_1 + v_2$. Using this, and proceeding as in (5.1), it can be proved that $\lim_{\varepsilon \rightarrow 0} |\text{im}_T(\mathbf{u}, \mathbf{a}_1) \Delta E(\mathbf{a}_\varepsilon, d_\varepsilon; \mathbf{u}_\varepsilon)| = 0$, which in turn implies that $D(\text{im}_T(\mathbf{u}, \mathbf{a})) = 0$.

In order to prove that at least one of the limit cavities must be distorted, we proceed as in the proof of Theorem 1 by applying Proposition 1.2 to $E_1 = E(\mathbf{a}_{1,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon)$, $E_2 = E(\mathbf{a}_{2,\varepsilon}, \varepsilon; \mathbf{u}_\varepsilon)$, and $E = E(\mathbf{a}_\varepsilon, d_\varepsilon; \mathbf{u}_\varepsilon)$. Again we define $g(\beta_1, \beta_2) := (\beta_1^{\frac{1}{n}} + \beta_2^{\frac{1}{n}})^n - (\beta_1 + \beta_2)$ and note that it is increasing in its two variables. It is easy to see that

$$\frac{(|E_1|^{\frac{1}{n}} + |E_2|^{\frac{1}{n}})^n - |E|}{(|E_1|^{\frac{1}{n}} + |E_2|^{\frac{1}{n}})^n - |E_1 \cup E_2|} \geq 1 - \frac{\omega_n d_\varepsilon^n}{g(v_{1,\varepsilon}, v_{2,\varepsilon})} \xrightarrow{\varepsilon \rightarrow 0} 1.$$

Therefore,

$$\liminf_{\varepsilon \rightarrow 0} \frac{|E| D(E)^{\frac{n}{n-1}} + |E_1| D(E_1)^{\frac{n}{n-1}} + |E_2| D(E_2)^{\frac{n}{n-1}}}{|E| + |E_1 \cup E_2|} \geq C \left(\frac{v_2}{v_1 + v_2} \right)^{\frac{n}{n-1}}.$$

Property (1.17) follows from (5.3). On the other hand, (3.6), (1.16), and Lemma 3.5 imply that

$$\begin{aligned} \sum_{i=1}^2 \int_{\varepsilon}^{\min\{\frac{d_\varepsilon}{2}, \varepsilon^\alpha\}} C \left(v_{i,\varepsilon} D(E_i)^{\frac{n}{n-1}} - 2^{\frac{n}{n-1}} \frac{n+1}{n-1} \omega_n \min\{\frac{d_\varepsilon^n}{2^n}, \varepsilon^{\alpha n}\} \right) \frac{dr}{r} \\ \leq (v_{1,\varepsilon} + v_{2,\varepsilon}) \log \frac{\text{diam } \Omega}{R/2} + C(v_{1,\varepsilon} + v_{2,\varepsilon} + |\Omega|). \end{aligned}$$

for every fixed $\alpha \in (0, 1)$. Hence,

$$\limsup_{\varepsilon \rightarrow 0} \left(\min \left\{ \log \frac{d_\varepsilon}{2\varepsilon}, \log \varepsilon^{\alpha-1} \right\} \right) \leq \frac{C \left(\log \frac{\text{diam } \Omega}{R/2} + 1 + \frac{|\Omega|}{v_1 + v_2} \right)}{\liminf_{\varepsilon \rightarrow 0} \left(\frac{v_{1,\varepsilon} D(E_1)^{\frac{n}{n-1}} + v_{2,\varepsilon} D(E_2)^{\frac{n}{n-1}}}{v_{1,\varepsilon} + v_{2,\varepsilon}} - \varepsilon^{\alpha n} \right)}.$$

By virtue of (1.17), and since $|\log \varepsilon| \rightarrow \infty$, we conclude that $\limsup_{\varepsilon \rightarrow 0} d_\varepsilon/\varepsilon$ is finite.

References

- [1] AMBROSIO, L., FUSCO, N., AND PALLARA, D. *Functions of bounded variation and free discontinuity problems*. Oxford University Press, New York, 2000.
- [2] BALL, J. M. Convexity conditions and existence theorems in nonlinear elasticity. *Arch. Rational Mech. Anal.* 63 (1976/77), 337–403.
- [3] BALL, J. M. Global invertibility of Sobolev functions and the interpenetration of matter. *Proc. Roy. Soc. Edinb. Sect. A* 88, 3-4 (1981), 315–328.
- [4] BALL, J. M. Discontinuous equilibrium solutions and cavitation in nonlinear elasticity. *Philos. Trans. R. Soc. Lond. Ser. A* 306 (1982), 557–611.
- [5] BALL, J. M. Minimizers and the Euler-Lagrange equations. In *Trends and applications of pure mathematics to mechanics (Palaiseau, 1983)*, vol. 195 of *Lecture Notes in Phys.* Springer, Berlin, 1984, pp. 1–4.
- [6] BALL, J. M. Some recent developments in nonlinear elasticity and its applications to materials science. In *Nonlinear mathematics and its applications (Guildford, 1995)*. Cambridge Univ. Press, Cambridge, 1996, pp. 93–119.
- [7] BALL, J. M., AND MURAT, F. $W^{1,p}$ -quasiconvexity and variational problems for multiple integrals. *J. Funct. Anal.* 58, 3 (1984), 225–253.
- [8] BANDYOPADHYAY, S., AND DACOROGNA, B. On the pullback equation $\phi^*(g) = f$. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 26, 5 (2009), 1717–1741.
- [9] BAYRATKAR, E., BESSRI, K., AND BATHIAS, C. Deformation behaviour of elastomeric matrix composites under static loading conditions. *Eng. Fract. Mech.* 75 (2008), 2695–2706.
- [10] BÉTHUEL, F., BREZIS, H., AND HÉLEIN, F. *Ginzburg-Landau vortices*. Progress in Nonlinear Differential Equations and their Applications, 13. Birkhäuser Boston Inc., Boston, MA, 1994.
- [11] BREZIS, H., AND NGUYEN, H.-M. Jacobian determinant revisited. *Invent. Math.* (2011). In press.

- [12] BREZIS, H., AND NIRENBERG, L. Degree theory and BMO. I. Compact manifolds without boundaries. *Selecta Math. (N.S.)* 1, 2 (1995), 197–263.
- [13] CERMELLI, P., AND LEONI, G. Renormalized energy and forces on dislocations. *SIAM J. Math. Anal.* 37, 4 (2005), 1131–1160.
- [14] CHENG, C., HILTNER, A., BAER, E., SOSKEY, P., AND MYLONAKIS, S. Cooperative cavitation in rubber-toughened polycarbonate. *Journal of Materials Science* 30, 3 (1995), 587–595.
- [15] CHO, K., GENT, A., AND LAM, P. Internal fracture in an elastomer containing a rigid inclusion. *Journal of Materials Science* 22 (1987), 2899–2905.
- [16] COIFMAN, R., LIONS, P., MEYER, Y., AND SEMMES, S. Compensated compactness and Hardy spaces. *Journal de Mathématiques Pures et Appliquées* 72, 3, 247–286.
- [17] CONTI, S., AND DE LELLIS, C. Some remarks on the theory of elasticity for compressible Neoohookean materials. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* 2, 3 (2003), 521–549.
- [18] CRISTIANO, A., MARCELLAN, A., LONG, R., HUI, C., STOLK, J., AND CRETON, C. An experimental investigation of fracture by cavitation of model elastomeric networks. *J. Polym. Sci. Pol. Phys.* 48 (2010), 1409–1422.
- [19] CUPINI, G., DACOROGNA, B., AND KNEUSS, O. On the equation $\det \nabla u = f$ with no sign hypothesis. *Calc. Var. Partial Differential Equations* 36, 2 (2009), 251–283.
- [20] DACOROGNA, B., AND MOSER, J. On a partial differential equation involving the Jacobian determinant. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 7, 1 (1990), 1–26.
- [21] DE LELLIS, C., AND GHIRALDIN, F. An extension of the identity $Det = det$. *C. R. Math. Acad. Sci. Paris* 348, 17-18 (2010), 973–976.
- [22] DORFMANN, A. Stress softening of elastomers in hydrostatic tension. *Acta Mechanica* 165 (2003), 117–137.
- [23] EVANS, L. C., AND GARIEPY, R. F. *Measure theory and fine properties of functions*. CRC Press, Boca Raton, FL, 1992.
- [24] FEDERER, H. *Geometric measure theory*. Springer, New York, 1969.
- [25] FOND, C. Cavitation criterion for rubber materials: A review of void-growth models. *Journal of Polymer Science Part B: Polymer Physics* 39, 17 (2001), 2081–2096.
- [26] FONSECA, I., AND GANGBO, W. *Degree theory in analysis and applications*. Oxford University Press, New York, 1995.
- [27] FUSCO, N., MAGGI, F., AND PRATELLI, A. The sharp quantitative isoperimetric inequality. *Ann. Math. (2)* 168 (2008), 941–980.

- [28] GARRONI, A., LEONI, G., AND PONSIGLIONE, M. Gradient theory for plasticity via homogenization of discrete dislocations. *J. Eur. Math. Soc. (JEMS)* 12, 5 (2010).
- [29] GENT, A. Cavitation in rubber: a cautionary tale. *Rubber Chem. Tech.* 63 (1991), G49–G53.
- [30] GENT, A., AND LINDLEY, P. Internal rupture of bonded rubber cylinders in tension. *Proc. Roy. Soc. London Ser. A* 249 (1959), 195 – 205.
- [31] GENT, A., AND PARK, B. Failure processes in elastomers at or near a rigid spherical inclusion. *J. Mater. Sci.* 19 (1984), 1947–1956.
- [32] GENT, A., AND WANG, C. Fracture mechanics and cavitation in rubber-like solids. *J. Mater. Sci.* 26, 12 (1991), 3392–3395.
- [33] GIAQUINTA, M., MODICA, G., AND SOUČEK, J. *Cartesian currents in the calculus of variations. I.* Springer-Verlag, Berlin, 1998.
- [34] GOODS, S., AND BROWN, L. The nucleation of cavities by plastic deformation. *Acta Metall.* 27 (1979), 1–15.
- [35] GURSON, A. Continuum theory of ductile rupture by void nucleation and growth: Part I—Yield criteria and flow rules for porous ductile media. *Journal of Engineering Materials and Technology* 99, 1 (1977), 2–15.
- [36] HAN, Z.-C., AND SHAFRIR, I. Lower bounds for the energy of S^1 -valued maps in perforated domains. *J. Anal. Math.* 66 (1995), 295–305.
- [37] HENAO, D. Cavitation, invertibility, and convergence of regularized minimizers in nonlinear elasticity. *J. Elasticity* 94 (2009), 55–68.
- [38] HENAO, D., AND MORA-CORRAL, C. Invertibility and weak continuity of the determinant for the modelling of cavitation and fracture in nonlinear elasticity. *Arch. Rational Mech. Anal.* 197 (2010), 619–655.
- [39] HENAO, D., AND MORA-CORRAL, C. Fracture surfaces and the regularity of inverses for BV deformations. *Arch. Rational Mech. Anal.* 203 (2011). doi:10.1007/s00205-010-0395-6.
- [40] HENAO, D., AND MORA-CORRAL, C. Lusin’s condition and the distributional determinant for deformations with finite energy. *Adv. Calc. Var.* (2012). <http://www.ann.jussieu.fr/publications/2009.php3>, preprint R09062.
- [41] HORGAN, C., AND POLIGNONE, D. Cavitation in nonlinearly elastic solids: A review. *Appl. Mech. Rev.* 48 (1995), 471–485.
- [42] HORGAN, C. O. Void nucleation and growth for compressible non-linearly elastic materials: an example. *Int. J. Solids Structures* 29, 3 (1992), 279–291.

- [43] HORGAN, C. O., AND ABEYARATNE, R. A bifurcation problem for a compressible nonlinearly elastic medium: growth of a microvoid. *J. Elast.* 16 (1986), 189–200.
- [44] JERRARD, R. L. Lower bounds for generalized Ginzburg-Landau functionals. *SIAM J. Math. Anal.* 30, 4 (1999), 721–746.
- [45] KUNDU, S., AND CROSBY, A. Cavitation and fracture behavior of polyacrylamide hydrogels. *Soft Matter* 5, 20 (2009), 3963–3968.
- [46] LAZZERI, A., AND BUCKNALL, C. Applications of a dilatational yielding model to rubber-toughened polymers. *Polymer* 36 (1995), 2895–2902.
- [47] LIAN, Y., AND LI, Z. A numerical study on cavitations in nonlinear elasticity - defects and configurational forces. *Math. Models Methods Appl. Sci.* (2011). In press.
- [48] LIANG, J., AND LI, R. Rubber toughening in polypropylene: A review. *J. Appl. Polym. Sci.* 77 (2000), 409–417.
- [49] LOPEZ-PAMIES, O., IDIART, M., AND NAKAMURA, T. Cavitation in elastomeric solids: I a defect-growth theory. *J. Mech. Phys. Solids* 59 (2011), 1464–1487.
- [50] LOPEZ-PAMIES, O., NAKAMURA, T., AND IDIART, M. Cavitation in elastomeric solids: II onset-of-cavitation surfaces for Neo-Hookean materials. *J. Mech. Phys. Solids* 59 (2011), 1488–1505.
- [51] MCMULLEN, C. T. Lipschitz maps and nets in Euclidean space. *Geom. Funct. Anal.* 8, 2 (1998), 304–314.
- [52] MICHEL, J., LOPEZ-PAMIES, O., PONTE CASTAÑEDA, P., AND TRIANTAFYLIDIS, N. Microscopic and macroscopic instabilities in finitely strained fiber-reinforced elastomers. *J. Mech. Phys. Solids* 58 (2010), 1776–1803.
- [53] MOSER, J. On the volume elements on a manifold. *Trans. Amer. Math. Soc.* 120 (1965), 286–294.
- [54] MÜLLER, S. Det = det. A remark on the distributional determinant. *C. R. Acad. Sci. Paris Série I* 311, 1 (1990), 13–17.
- [55] MÜLLER, S., AND SPECTOR, S. J. An existence theory for nonlinear elasticity that allows for cavitation. *Arch. Rational Mech. Anal.* 131, 1 (1995), 1–66.
- [56] MÜLLER, S., SPECTOR, S. J., AND TANG, Q. Invertibility and a topological property of Sobolev maps. *SIAM J. Math. Anal.* 27, 4 (1996), 959–976.
- [57] OBERTH, A., AND BRUENNER, R. Tear phenomena around solid inclusions in castable elastomers. *Trans. Soc. Rheol.* 9 (1965), 165–185.

- [58] PETRINIC, N., CURIEL SOSA, J. L., SIVIOUR, C. R., AND ELLIOTT, B. C. F. Improved predictive modelling of strain localisation and ductile fracture in a Ti-6Al-4V alloy subjected to impact loading. *Journal de Physique IV 134* (Aug. 2006), 147–155.
- [59] PONSIGLIONE, M. Elastic energy stored in a crystal induced by screw dislocations: from discrete to continuous. *SIAM J. Math. Anal.* 39, 2 (2007), 449–469.
- [60] REINA, C. *Multiscale Modeling and simulation of damage by void nucleation and growth*. PhD thesis, California Institute of Technology, 2010. <http://thesis.library.caltech.edu/6165/>.
- [61] RESHETNYAK, Y. G. On the stability of conformal mappings in multidimensional spaces. *Sibirsk. Mat. Ž.* 8 (1967), 91–114.
- [62] RICE, J., AND TRACEY, D. On the ductile enlargement of voids in triaxial stress fields. *Journal of the Mechanics and Physics of Solids* 17, 3 (1969), 201–217.
- [63] RIVIÈRE, T., AND YE, D. Une résolution de l'équation à forme volume prescrite. *C. R. Acad. Sci. Paris Sér. I Math.* 319, 1 (1994), 25–28.
- [64] RIVIÈRE, T., AND YE, D. Resolutions of the prescribed volume form equation. *NoDEA Nonlinear Differential Equations Appl.* 3, 3 (1996), 323–369.
- [65] SANDIER, E. Lower bounds for the energy of unit vector fields and applications. *J. Funct. Anal.* 152, 2 (1998), 379–403.
- [66] SANDIER, E. Erratum: “Lower bounds for the energy of unit vector fields and applications”. *J. Funct. Anal.* 171, 1 (2000), 233.
- [67] SANDIER, E., AND SERFATY, S. *Vortices in the magnetic Ginzburg-Landau model*. Progress in Nonlinear Differential Equations and their Applications, 70. Birkhäuser, Boston, 2007.
- [68] SCHWARTZ, J. T. *Nonlinear functional analysis*. Gordon and Breach, New York, 1969.
- [69] SIVALOGANATHAN, J. Uniqueness of regular and singular equilibria for spherically symmetric problems of nonlinear elasticity. *Arch. Rational Mech. Anal.* 96, 2 (1986), 97–136.
- [70] SIVALOGANATHAN, J., AND SPECTOR, S. J. On the existence of minimizers with prescribed singular points in nonlinear elasticity. *J. Elast.* 59 (2000).
- [71] SIVALOGANATHAN, J., AND SPECTOR, S. J. On cavitation, configurational forces and implications for fracture in a nonlinearly elastic material. *J. Elast.* 67 (2002), 25–49.
- [72] SIVALOGANATHAN, J., AND SPECTOR, S. J. On the symmetry of energy-minimising deformations in nonlinear elasticity I: incompressible materials. *Arch. Rational Mech. Anal.* 196 (2010), 363–394.

- [73] SIVALOGANATHAN, J., AND SPECTOR, S. J. On the symmetry of energy-minimising deformations in nonlinear elasticity II: compressible materials. *Arch. Rational Mech. Anal.* 196 (2010), 395–431.
- [74] SIVALOGANATHAN, J., SPECTOR, S. J., AND TILAKRAJ, V. The convergence of regularized minimizers for cavitation problems in nonlinear elasticity. *SIAM J. Appl. Math.* 66 (2006), 736–757.
- [75] SPIVAK, M. *Calculus on manifolds*. W. A. Benjamin, Inc., New York-Amsterdam, 1965.
- [76] STEENBRINK, A., AND VAN DER GIESSEN, E. On cavitation, post-cavitation and yield in amorphous polymer-rubber blends. *J. Mech. Phys. Solids* 47 (1999), 732–765.
- [77] STRUWE, M. On the asymptotic behavior of minimizers of the Ginzburg-Landau model in 2 dimensions. *Differential Integral Equations* 7, 5-6 (1994), 1613–1624.
- [78] TVERGAARD, V. Material failure by void growth and coalescence. In *Advances in Applied Mechanics*, vol. 27. Academic Press, San Diego, 1990, pp. 83–151.
- [79] ŠVERÁK, V. Regularity properties of deformations with finite energy. *Arch. Rational Mech. Anal.* 100, 2 (1988), 105–127.
- [80] WILLIAMS, M., AND SCHAPERLY, R. Spherical flaw instability in hydrostatic tension. *Int. J. Fract. Mech.* 1, 1 (1965), 64–71.
- [81] XU, X., AND HENAO, D. An efficient numerical method for cavitation in nonlinear elasticity. *Math. Models Methods Appl. Sci.* (2011). In press.
- [82] YE, D. Prescribing the Jacobian determinant in Sobolev spaces. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 11, 3 (1994), 275–296.
- [83] ZIEMER, W. P. *Weakly differentiable functions*. Springer, New York, 1989.

SYLVIA SERFATY

UPMC Univ. Paris 06, UMR 7598 Laboratoire Jacques-Louis Lions,
 Paris, F-75005 France;
 CNRS, UMR 7598 LJLL, Paris, F-75005 France
 & Courant Institute, New York University
 251 Mercer St., New York, NY 10012, USA
 serfaty@ann.jussieu.fr

DUVAN HENAO

SYLVIA SERFATY

UPMC Univ. Paris 06, UMR 7598 Laboratoire Jacques-Louis Lions,
 Paris, F-75005 France;
 CNRS, UMR 7598 LJLL, Paris, F-75005 France.