



Every plane graph of maximum degree 8 has an edge-face 9-colouring

Ross Kang, Jean-Sébastien Sereni, Matej Stehlík

► To cite this version:

Ross Kang, Jean-Sébastien Sereni, Matej Stehlík. Every plane graph of maximum degree 8 has an edge-face 9-colouring. *Siam Journal on Discrete Mathematics, Society for Industrial and Applied Mathematics*, 2011, 25 (2), pp.514–533. <10.1137/090781206>. <hal-00604185>

HAL Id: hal-00604185

<https://hal.archives-ouvertes.fr/hal-00604185>

Submitted on 28 Jun 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EVERY PLANE GRAPH OF MAXIMUM DEGREE 8 HAS AN EDGE-FACE 9-COLOURING

ROSS J. KANG*, JEAN-SÉBASTIEN SERENI[†], AND MATĚJ STEHLÍK[‡]

Abstract. An edge-face colouring of a plane graph with edge set E and face set F is a colouring of the elements of $E \cup F$ such that adjacent or incident elements receive different colours. Borodin proved that every plane graph of maximum degree $\Delta \geq 10$ can be edge-face coloured with $\Delta + 1$ colours. Borodin's bound was recently extended to the case where $\Delta = 9$. In this paper, we extend it to the case $\Delta = 8$.

Key words. graph colouring, edge-face colouring, plane graphs

AMS subject classifications. 05C15

1. Introduction. Let G be a plane graph with vertex set V , edge set E and face set F . Given a positive integer k , an *edge-face k -colouring* of G is a mapping $\lambda : E \cup F \rightarrow \{1, 2, \dots, k\}$ such that

- (i) $\lambda(e) \neq \lambda(e')$ for every pair (e, e') of adjacent edges;
- (ii) $\lambda(e) \neq \lambda(f)$ for edge e and every face f incident to e ;
- (iii) $\lambda(f) \neq \lambda(f')$ for every pair (f, f') of adjacent faces with $f \neq f'$.

The requirement in (iii) that f and f' be distinct is only relevant for graphs containing a cut-edge; such graphs would not have an edge-face colouring otherwise. Let $\chi_{ef}(G)$ be the value of the smallest integer k such that there exists an edge-face k -colouring of G . Although this problem is well-defined for graphs with loops or multiple edges, we shall throughout the paper only consider graphs that are simple (and this requirement is necessary, for instance, in Lemma 5.5). We comment here that the multigraph formed by replacing each edge in a triangle by $\Delta/2$ parallel edges is planar with maximum degree Δ and requires at least $3\Delta/2$ colours in an edge colouring (let alone an edge-face colouring).

Edge-face colourings were first studied by Jucovič [4] and Fiamčík [3], who considered 3- and 4-regular graphs. A conjecture of Mel'nikov [5] spurred research into upper bounds on $\chi_{ef}(G)$ for plane graphs G with $\Delta(G) \leq \Delta$. For small values of Δ , the best bounds known are $\Delta + 3$ for $\Delta \in \{2, \dots, 6\}$ [1, 6, 9] and $\Delta + 2$ for $\Delta = 7$ [7]. For $\Delta \geq 10$, Borodin [2] proved the bound of $\Delta + 1$. This is tight, as can be seen by considering trees. Recently, the second and third authors [8] extended the $\Delta + 1$ bound to the case $\Delta = 9$ by proving that every plane graph of maximum degree 9 has an edge-face 10-colouring. Here, we settle the case $\Delta = 8$.

THEOREM 1.1. *Every plane graph of maximum degree 8 has an edge-face 9-colouring.*

*School of Engineering and Computing Sciences, Durham University, UK (ross.kang@gmail.com). This research was conducted while this author was a Postdoctoral Fellow at McGill University, supported by the *National Sciences and Engineering Research Council of Canada* (NSERC). He is currently supported by the *Engineering and Physical Sciences Research Council* (EPSRC), grant EP/G066604/1.

[†]CNRS (LIAFA, Université Denis Diderot), Paris, France and Department of Applied Mathematics (KAM), Faculty of Mathematics and Physics, Charles University, Prague, Czech Republic (sereni@kam.mff.cuni.cz). This author's work was partially supported by the French *Agence Nationale de la Recherche* under reference ANR 10 JCJC 0204 01.

[‡]UJF-Grenoble 1 / CNRS / Grenoble-INP, G-SCOP UMR5272 Grenoble, F-38031, France (matej.stehlik@g-scop.inpg.fr).

Our result is a strengthening of the $\Delta = 7$ result of Sanders and Zhao [7]; it can also be viewed as an extension of the recent work on $\Delta = 9$ [8]. The problem of finding the provably optimal upper bounds on $\chi_{ef}(G)$ for plane graphs G with $\Delta(G) \leq \Delta$ remains open for $\Delta \in \{4, 5, 6, 7\}$.

We prove Theorem 1.1 by contradiction. From now on, we let $G = (V, E, F)$ be a counter-example to the statement of Theorem 1.1 with as few edges as possible. That is, G is a plane graph of maximum degree 8 and no edge-face 9-colouring, but every plane graph of maximum degree at most 8 with less than $|E|$ edges has an edge-face 9-colouring. In particular, for every edge $e \in E$ the plane subgraph $G - e$ of G has an edge-face 9-colouring. First, we describe various structural properties of G in Section 2; the proofs of these properties are given at the end of this paper in Section 5. In Section 3 we describe the discharging rules. In Section 4 we use the discharging rules and the structural properties of G to obtain a contradiction, and thus a proof of Theorem 1.1.

Our discharging procedure was developed through several rounds, with corrective adjustments and optimisations included in each round, starting from a naïve scheme in which only the vertices of degree at least 7 compensated for the deficit of charge on triangles. A breakthrough in the design of our strategy was the realisation that Lemma 2.1 below could allow us to conserve considerable charge at degree 7 or 8 vertices incident to faces of a particular type. We could then balance these savings against the loss of charge to incident triangles with the development of further reducible configurations. As will become apparent, the analysis of the final charge of vertices of degree 7 or 8 is particularly involved.

In the sequel, a vertex of degree d is called a d -*vertex*. A vertex is an $(\leq d)$ -*vertex* if its degree is at most d ; it is an $(\geq d)$ -*vertex* if its degree is at least d . The notions of d -*face*, $(\leq d)$ -*face* and $(\geq d)$ -*face* are defined analogously as for the vertices, where the *degree* of a face is the number of edges incident to it. A face of length 3 is called a *triangle*. For integers a, b, c , an $(\leq a, \leq b, \leq c)$ -*triangle* is a triangle xyz of G with $\deg(x) \leq a$, $\deg(y) \leq b$ and $\deg(z) \leq c$. The notions of $(a, \leq b, \leq c)$ -triangles, $(a, b, \geq c)$ -triangles, $(a, \leq b, c, d)$ -*faces*, and so on, are defined analogously. A vertex is *triangulated* if all its incident faces are triangles.

2. Reducible configurations. For our proof of Theorem 1.1, we identify that some plane graphs are *reducible configurations*, i.e. configurations that cannot be part of the chosen embedding of G . Their reducibility follows from Lemmas 2.1–5.11; these lemmas are proved in Section 5. In this section, we give an explicit description of the reducible configurations as well as the statement of Lemma 2.1.

For convenience, we depict these configurations in Figure 2.1. We use the following notational conventions for vertices: 2-, 3- and 4-vertices are depicted by black bullets, black triangles and black squares, respectively; a white bullet containing a number represents a vertex of degree that quantity; an empty white bullet represents a vertex of arbitrary degree (but at least that shown in the figure). For faces, we use the following conventions: a straight line indicates a single edge; a curved line indicates a portion of the face with an unspecified number of edges; a curved face that is shaded grey represents an (≤ 4) -face.

The following configurations are reducible. Note that, for any of the below, if an edge can be removed without affecting the prescribed incidence or facial structure, then the configuration remains reducible; for example, B6 modified by replacing the 6 by a 5 or 4 is reducible.

A0 A 1-vertex.

Configurations with faces incident to a 2-vertex.

- A1** A triangle incident to a 2-vertex.
- A2** A 4-face incident to a 2-vertex and an (≤ 3) -vertex.
- A3** A 2-vertex adjacent to a 3-vertex and an (≤ 5) -vertex.

Configurations with an edge incident to a (≤ 4) -face.

- B1** An edge uv that is incident to an (≤ 4) -face, with $\deg(u) + \deg(v) \leq 9$.
- B2** A triangle uvw with $\deg(u) + \deg(v) \leq 10$ and $\deg(w) = 6$.
- B3** A triangle uvw with uv incident to two (≤ 4) -faces, and $\deg(u) + \deg(v) \leq 10$ and $\deg(w) = 7$.
- B4** A triangle uvw with uv adjacent to two (≤ 4) -faces, vw incident to two (≤ 4) -faces, and $\deg(u) + \deg(v) \leq 10$.

Configurations with an edge incident to two (≤ 4) -faces.

- C1** An edge uv that is incident to two (≤ 4) -faces, with $\deg(u) + \deg(v) \leq 10$.
- C2** A triangle uvw with uv incident to two (≤ 4) -faces, and $\deg(u) + \deg(v) \leq 11$ and $\deg(w) = 6$.
- C3** A triangle uvw with uv and uw each incident to two (≤ 4) -faces, and $\deg(u) + \deg(v) \leq 11$ and $\deg(w) = 7$.
- C4** A triangle uvw with vw incident to the triangle vwx and wx incident to two (≤ 4) -faces, and $\deg(u) = \deg(x) = 3$.
- C5** A triangle uvw with vw incident to the triangle vwx and wx incident to two (≤ 4) -faces, and $\deg(u) + \deg(v) \leq 10$ and $\deg(v) + \deg(x) \leq 11$.

Configurations along a 2-path.

- D1** A 2-path uvw such that vwx is a triangle, with uv incident to an (≤ 4) -face, vw and vx each incident to two (≤ 4) -faces, and $\deg(u) + \deg(v) \leq 10$ and $\deg(v) + \deg(w) \leq 11$.
- D2** A 2-path uvw such that vwx is a triangle, with uv , vw and vx each incident to two (≤ 4) -faces, and $\deg(u) + \deg(v) \leq 11$ and $\deg(v) + \deg(w) \leq 11$.
- D3** A 2-path uvw such that vwx is a triangle, with vx incident to two (≤ 4) -faces, and $\deg(u) = 2$, $\deg(v) = 7$ and $\deg(w) = 3$.
- D4** A 2-path uvw such that vwx is a triangle, with vw and vx each incident to two (≤ 4) -faces, and $\deg(u) = 2$, $\deg(v) = 7$ and $\deg(w) = 4$.

Note on configurations D1 and D2. An (≤ 4) -face incident to uv is not ruled out from also being an (≤ 4) -face (distinct from vwx) incident to vw or vx . In this sense, the figures representing configurations D1 and D2 in Figure 2.1 belie the configurations' fuller forms.

Exceptional configurations.

- E1** A 4-path $uvwxy$, such that uvz , vwz , wxz and xyz are triangles, with yz incident to two (≤ 4) -faces, and $\deg(v) = 3$, $\deg(x) = 4$.
- E2** A 4-path $uvwxy$, such that uvz , vwz , wxz and xyz are triangles, and $\deg(v) = 3$, $\deg(x) = 4$ and $\deg(y) = 6$.
- E3** A triangulated 8-vertex that is adjacent to both a 3-vertex and a 4-vertex.
- E4** A 3-path uvw , with uv incident to an (≤ 4) -face, and $\deg(u) \leq 5$, $\deg(v) = 6$, $\deg(w) = 2$ and $\deg(x) = 3$.

Special lemmas for (≥ 5) -faces. An edge uv is *loose* if $\deg(u) + \deg(v) \leq 8$. The following lemma implies a general set of reducible configurations for (≥ 5) -faces. These configurations are not depicted in Figure 2.1.

LEMMA 2.1. *Let f be a d -face of G incident to x loose edges and q vertices of degree 2. If $d \geq 5$ and $x \geq 1$, then $2d - q - x \geq 9$. For (≥ 6) -faces, we require one more configuration not depicted in Figure 2.1.*

LEMMA 2.2. *Let u and v be two adjacent 2-vertices in G . If $u' \neq v$ and $v' \neq u$ are neighbours of u and v , respectively, then $u' = v'$ and $\deg(u') = 8$.*

3. Discharging rules. Recall that $G = (V, E, F)$ is a plane graph that is a minimum counter-example to the statement of Theorem 1.1, in the sense that $|E|$ is minimum. (In particular, a planar embedding of G is fixed.) We obtain a contradiction by using the Discharging Method. Each vertex and face of G is assigned an initial charge; the total sum of the charge is negative by Euler's Formula. Then vertices and faces send or receive charge according to certain redistribution rules. The total sum of the charge remains unchanged, but ultimately (by using all of the reducible configurations in Section 2) we deduce that the charge of each face and vertex is non-negative, a contradiction.

3.1. Initial charge. We assign a charge to each vertex and face. For every vertex $v \in V$, we define the initial charge $\text{ch}(v)$ to be $2 \cdot \deg(v) - 6$, while for every face $f \in F$, we define the initial charge $\text{ch}(f)$ to be $\deg(f) - 6$. The total sum is

$$\sum_{v \in V} \text{ch}(v) + \sum_{f \in F} \text{ch}(f) = -12.$$

Indeed, by Euler's formula $|E| - |V| - |F| = -2$. Thus, $6|E| - 6|V| - 6|F| = -12$. Since $\sum_{v \in V} \deg(v) = 2|E| = \sum_{f \in F} \deg(f)$, it follows that

$$\begin{aligned} -12 &= 4 \cdot |E| - 6 \cdot |V| + \sum_{f \in F} (\deg(f) - 6) \\ &= \sum_{v \in V} (2 \deg(v) - 6) + \sum_{f \in F} (\deg(f) - 6). \end{aligned}$$

3.2. Rules. We need the following definitions to state the discharging rules. Given an (≥ 7) -vertex v , a face is *special* (for v) if it is an (≥ 5) -face that is incident to a degree 2 neighbour of v (and so, in particular, such a face is incident to v). Given a 6-vertex v , a face f is *exceptional* (for v) if f is a 6-face $vv_1v_2 \dots v_5$ where v_1 is a 2-vertex and v_2 is a 3-vertex.

Since G may have cut-vertices (of a type not forbidden by Lemma 5.1), some vertices may be incident to the same face several times. Thus, in the rules below, when we say that a vertex or a face sends charge to an incident face or vertex, we mean that the charge is sent as many times as these elements are incident to each other.

The following describe how the charge is redistributed among the edges and faces in G .

R0 An (≥ 4) -face sends 1 to each incident 2-vertex.

R1 An (≥ 7) -vertex sends

R1a $3/2$ to incident $(3, \geq 7, \geq 7)$ -triangles and $(4, 6, \geq 7)$ -triangles;

R1b $7/5$ to incident $(5, 5, \geq 7)$ -triangles;

R1c $5/4$ to incident $(4, \geq 7, \geq 7)$ -triangles and $(2, 8, 4, 8)$ -faces;

R1d $6/5$ to incident $(5, 6, 8)$ -triangles;

R1e $11/10$ to incident $(5, 6, 7)$ - and $(5, \geq 7, \geq 7)$ -triangles, and incident $(2, 8, 5, 8)$ -faces;




Fig. 2.1: The reducible configurations.

- R1f** 1 to incident $(\geq 6, \geq 6, \geq 6)$ -triangles, to every incident 4-face that is not a $(2, 8, \leq 5, 8)$ -face, and to each of its incident special faces;
- R1g** $1/2$ to each of its incident non-special 5-faces.
- R2** A 6-vertex sends
- R2a** $11/10$ to incident $(5, 6, 6)$ - and $(5, 6, 7)$ -triangles;
- R2b** 1 to every other incident triangle, to each incident 4-face, and to each of its incident exceptional faces;
- R2c** $1/2$ to each of its incident 5-faces and to each of its incident unexceptional 6-faces.
- R3** A 5-vertex sends $4/5$ to each incident face.
- R4** A 4-vertex sends $1/2$ to each incident face.

Note on rules R1 and R2. Since configurations A1, B1 and B2 are reducible, it follows from rule R1 that an (≥ 7) -vertex sends positive charge to every incident triangle. We conclude that an (≥ 7) -vertex sends zero charge only to incident (≥ 6) -faces that are not special. Similarly, a 6-vertex sends zero charge only to incident (≥ 7) -faces.

4. Proof of Theorem 1.1. In this section, we prove that the final charge $\text{ch}^*(x)$ of every $x \in V \cup F$ is non-negative. Hence, we obtain

$$-12 = \sum_{x \in V \cup F} \text{ch}(x) = \sum_{x \in V \cup F} \text{ch}^*(x) \geq 0,$$

a contradiction. This contradiction establishes Theorem 1.1.

4.1. Final charge of faces. Let f be a d -face. Our goal is to show that $\text{ch}^*(f) \geq 0$. Recall that the initial charge of f is $\text{ch}(f) = d - 6$.

First suppose that $d \geq 6$. Let p be the number of occurrences of an (≥ 7) -vertex having f as an incident special face, and q the number of 2-vertices incident to f . In particular, $\text{ch}^*(f) \geq d - 6 - q + p$ by rules R0 and R1f. We define x to be the number of edges of f between a 2-vertex and an (≤ 6) -vertex, y the number of edges of f between a 2-vertex and an (≥ 7) -vertex, and z the number of edges of f between two 2-vertices. We have $2q = x + y + z$ and $2p \geq y$. If $x = 0$, then $p \geq q$, and hence $\text{ch}^*(f) \geq 0$. Assume now that $x \geq 1$. Then Lemma 2.1 implies that $2d - q - x \geq 9$, that is $d - x/2 \geq (q + 9)/2$. Now, $\text{ch}^*(f) \geq d - 6 - q + p \geq \lceil d - 6 - (x + z)/2 \rceil$ since $p \geq y/2$, $q = (x + y + z)/2$ and $d - 6 - q + p$ is integral. Hence,

$$\text{ch}^*(f) \geq \left\lceil \frac{q - 3 - z}{2} \right\rceil,$$

which is non-negative if $q - z \geq 2$. It remains to deal with the case where $q - z \leq 1$. Note that $q \geq 2z$ due to Lemma 2.2. It therefore follows that $z \leq 1$. Let us first consider the case $z = 1$, hence $q = 2$. In this case, it follows by Lemma 2.2 that $p \geq 2$ and therefore $\text{ch}^*(f) \geq d - 6 - 2 + 2 \geq 0$. We just need to check the case $z = 0$ and $q = 1$ (since $x \geq 1$). Then we may assume that $d = 6$, for if $d \geq 7$ then $\text{ch}^*(f) \geq d - 6 - 1 \geq 0$. Moreover, if $y \geq 1$ then $\text{ch}^*(f) \geq 0$ by rule R1f. Hence, $x = 2q - y - z = 2$. Let v and v' be the two (≤ 6) -neighbours of the 2-vertex of f . First, if both v and v' have degree more than 3, then each of them sends at least $1/2$ to f by rules R2, R3 and R4, and hence $\text{ch}^*(f) \geq 0$. So, as $q = 1$, we may assume that v has degree 3. Now, by the reducibility of configuration A3, the degree of v' is at least 6 and hence exactly 6. Consequently, f is exceptional for v' and f receives 1 from v' by rule R2b, which concludes the analysis for (≥ 6) -faces.

Suppose that $d = 5$, and let q be the number of 2-vertices incident to f . Lemma 2.1 implies that f is incident to at most one loose edge. Thus, if $q = 0$ (so that f sends no charge to vertices) then f is incident to at least two (≥ 4)-vertices, and hence $\text{ch}^*(f) \geq 5 - 6 + 2 \cdot 1/2 = 0$ by rules R1g, R2c, R3 and R4. Moreover, if $q \geq 1$, then Lemma 2.1 implies that f is not incident to a loose edge and f is thus incident to at least $q + 1$ vertices of degree at least 7. herefore, f receives at least $q + 1$ from its incident (≥ 7)-vertices by rule R1f, and so $\text{ch}^*(f) \geq 5 - 6 - q + q + 1 = 0$.

Next suppose that $d = 4$. Let the four vertices incident to f be v_0, \dots, v_3 in clockwise order and suppose without loss of generality that v_0 has the least degree among v_0, \dots, v_3 . First, if $\deg(v_0) \geq 4$, then by rules R1f, R2b, R3 and R4, the charge sent to f by each incident vertex is at least $1/2$, so that $\text{ch}^*(f) \geq -2 + 4 \cdot 1/2 = 0$. If $\deg(v_0) = 3$, then since configuration B1 is reducible $\deg(v_1) \geq 7$ and $\deg(v_3) \geq 7$. Thus, by rule R1f, $\text{ch}^*(f) \geq -2 + 2 = 0$. Last, assume that $\deg(v_0) = 2$. Since configuration B1 is reducible, $\deg(v_1) = \deg(v_3) = 8$, and since configuration A2 is reducible, $\deg(v_2) \geq 4$. By rule R0, f sends charge 1 to v_0 . But f receives charge 3: by rules R1c and R4 if f is a $(2, 8, 4, 8)$ -face; by rules R1e and R3 if f is a $(2, 8, 5, 8)$ -face; and by rules R1f and R2b if f is a $(2, 8, \geq 6, 8)$ -face. Thus, $\text{ch}^*(f) \geq 0$.

Finally suppose that $d = 3$. Let the three vertices incident to f be v_0, v_1 and v_2 , and let us assume without loss of generality that $\deg(v_0) \leq \deg(v_1) \leq \deg(v_2)$. Since configuration A1 is reducible, $\deg(v_0) \geq 3$. Thus f sends no charge, but needs to make up for an initial charge of -3 . We analyse several cases according to the value of $\deg(v_0)$.

$\deg(v_0) = 3$. Since configuration B1 is reducible, $\deg(v_1) \geq 7$. By rule R1a, f receives charge $2 \cdot 3/2 = 3$.

$\deg(v_0) = 4$. Since configuration B1 is reducible, $\deg(v_1) \geq 6$. If $\deg(v_1) \geq 7$, then f receives charge $2 \cdot 5/4 + 1/2 = 3$ by rules R1c and R4. Otherwise, $\deg(v_1) = 6$ and hence $\deg(v_2) \geq 7$ since configuration B2 is reducible, but then f receives charge $3/2 + 1 + 1/2 = 3$ by rules R1a, R2b and R4.

$\deg(v_0) = 5$. If $\deg(v_1) = 5$, then $\deg(v_2) \geq 7$ since configuration B2 is reducible, but then f receives charge $7/5 + 2 \cdot 4/5 = 3$ by rules R1b and R3. If $\deg(v_1) = 6$, then we separately consider the cases of $\deg(v_2) \in \{6, 7, 8\}$. If $\deg(v_2) \in \{6, 7\}$, then f receives charge $2 \cdot 11/10 + 4/5 = 3$ by rules R1e, R2a and R3; if $\deg(v_2) = 8$, then f receives charge $6/5 + 1 + 4/5 = 3$ by rules R1d, R2b and R3. Last, if $\deg(v_1) \geq 7$, then f receives charge $2 \cdot 11/10 + 4/5 = 3$ by rules R1e and R3.

$\deg(v_0) \geq 6$. The face f receives charge at least 3 by rules R1f and R2b.

This concludes our analysis of the final charge of f , verifying that $\text{ch}^*(f) \geq 0$.

4.2. Final charge of (≤ 6)-vertices. Let v be an arbitrary vertex of G . Our goal is to show that $\text{ch}^*(v) \geq 0$. Recall that the initial charge of v is $\text{ch}(v) = 2 \cdot \deg(v) - 6$. Moreover, $\deg(v) \geq 2$ since configuration A0 is reducible.

If $\deg(v) = 2$, then v is incident to two (≥ 4)-faces since configuration A1 is reducible; thus, v receives charge 1 from both incident faces by rule R0 and the final charge of v is $\text{ch}^*(v) = -2 + 2 = 0$.

If $\deg(v) = 3$, then v neither sends nor receives any charge; hence, the final charge of v is $\text{ch}^*(v) = \text{ch}(v) = 0$.

If $\deg(v) \in \{4, 5\}$, then v sends charge $\text{ch}(v)/\deg(v)$ to each incident face by rules R3 and R4; the final charge of v is $\text{ch}^*(v) = 0$.

Suppose now that $\deg(v) = 6$. The initial charge of v is $\text{ch}(v) = 6$. If v is incident to a 5-face or an unexceptional (≥ 6)-face, then it sends charge at most $1/2$ to one

of the faces by rule R2c and so by rule R2 the total charge sent by v is at most $5 \cdot 11/10 + 1/2 = 6$. If v is incident to an exceptional 6-face, then, since configuration E4 is reducible, v has no incident $(5, 6, 6)$ - or $(5, 6, 7)$ -triangles and thus the total charge sent is at most $5 \cdot 1 + 1 = 6$ by rules R2b and R2c. We conclude that v is only incident to (≤ 4) -faces. Then, since configuration C2 is reducible, v has no incident $(5, 6, 6)$ -face; furthermore, since configuration C3 is reducible, v has no incident $(5, 6, 7)$ -face. Therefore, the charge sent by v is at most 6 and the final charge of v satisfies $\text{ch}^*(v) \geq 0$.

4.3. Final charge of 7-vertices. Next, suppose that $\deg(v) = 7$. For convenience, let v_0, v_1, \dots, v_6 be the neighbours of v in clockwise order, and let f_i be the face vv_iv_{i+1} for $i \in \{0, 1, \dots, 6\}$, where the index is modulo 7. The initial charge of v is $\text{ch}(v) = 8$. We partition our analysis based on the number of incident special (≥ 5) -faces. Note that since configuration B1 is reducible, if v is adjacent to a 2-vertex then both of the 2-vertex's incident faces are special for v .

4.3.1. There is an adjacent 2-vertex. We first treat the cases in which v is adjacent to some 2-vertex. In these cases, there are at least two incident special (≥ 5) -faces. Thus, we may assume that v is incident to at most one non-special (≥ 5) -face (which is sent at most $1/2$ charge by rule R1g), for otherwise the total charge sent by v is at most $3 \cdot 3/2 + 2 + 2 \cdot 1/2 < 8$. Now note that, by rules R1a and R1b, any face that is sent charge more than $5/4$ must be a $(3, 7, \geq 7)$ -, $(4, 6, 7)$ - or $(5, 5, 7)$ -triangle. And so we assert that if f_i is such a triangle, then both f_{i-1} and f_{i+1} are (≥ 5) -faces. The assertion holds if f_i is a $(3, 7, \geq 7)$ -triangle since configurations C1 and D3 are reducible, and the fact that configuration B3 is reducible implies the assertion for the two other cases.

Case 4.3.1(1). If v is incident to (exactly) one non-special (≥ 5) -face, then v is incident to only two special (≥ 5) -faces (for otherwise the total charge sent by v is $3 \cdot 3/2 + 3 + 1/2 = 8$). Thus, the remaining four incident faces are (≤ 4) -faces. Observe that there are at least three of these faces that are adjacent around v to another (≤ 4) -face. Hence by the assertion at the end of the last paragraph, each of these three faces is sent at most $5/4$ charge. So the charge sent by v is at most $3/2 + 3 \cdot 5/4 + 2 + 1/2 < 8$. Thus, v is not incident to a non-special (≥ 5) -face.

Case 4.3.1(2). If v is incident to at least five (≥ 5) -faces, then the charge sent is at most $2 \cdot 3/2 + 5 = 8$ due to rule R1f.

Case 4.3.1(3). If v is incident to exactly four (≥ 5) -faces, all of which are special, then there must be two incident (≤ 4) -faces that are adjacent. (Recall that each special face is adjacent to another special face.) By the assertion in the second paragraph of the 7-vertex analysis, both of these are sent charge at most $5/4$. Therefore, the total charge sent by v in this case is at most $3/2 + 2 \cdot 5/4 + 4 = 8$.

Case 4.3.1(4). If v is incident to exactly three (≥ 5) -faces, all of which are special, then these faces are sequentially adjacent around v . Hence, by the assertion in the second paragraph of the 7-vertex analysis, no face is sent charge more than $5/4$ and the total charge sent is at most $4 \cdot 5/4 + 3 = 8$.

Case 4.3.1(5). Suppose that v is incident to exactly two (≥ 5) -faces, say f_0 and f_1 , both special (so v_1 is a 2-vertex). Recall that all other incident faces have size at most 4. Let us analyse which incident faces can be sent charge $5/4$. By rule R1c, such a face must be a $(4, 7, \geq 7)$ -triangle. Since configuration D4 is reducible, such a face must be adjacent to a special face for v . Thus, there are at most two such faces, namely f_2 and f_6 . Consequently, the total charge sent by v is at most $2 \cdot 5/4 + 3 \cdot 11/10 + 2 < 8$ by rules R1c, R1e and R1f.

4.3.2. There is no adjacent 2-vertex. Now we may assume that v is not adjacent to a 2-vertex. In these cases, we may assume that v is incident to at most two (non-special) (≥ 5)-faces (which are sent at most $1/2$ charge by rule R1g), for otherwise the total charge sent by v is at most $4 \cdot 3/2 + 3 \cdot 1/2 < 8$.

Case 4.3.2(1). Suppose that v is incident to two 5-faces (and hence to no other (≥ 5)-faces). First suppose that there are four (≤ 4)-faces that are sequentially adjacent around v . Since configuration B3 is reducible, none of these is a $(3, 7, 7)$ -, $(4, 6, 7)$ - or $(5, 5, 7)$ -triangle. Also, since configuration C1 is reducible, at most two of these are $(3, 7, 8)$ -triangles. It follows that at most three faces are sent charge $3/2$, and the remaining two (≤ 4)-faces are sent charge at most $5/4$; so the total charge sent by v is at most $3 \cdot 3/2 + 2 \cdot 5/4 + 2 \cdot 1/2 = 8$. Next suppose that there are only three (≤ 4)-faces that are sequentially adjacent around v . As before, we deduce that none of these is a $(3, 7, 7)$ -, $(4, 6, 7)$ - or $(5, 5, 7)$ -triangle and at most two of these are $(3, 7, 8)$ -triangles. If two of these faces are $(3, 7, 8)$ -triangles, then the middle one is either a $(7, 8, 8)$ -triangle or a 4-face, and hence sent charge 1 by rule R1f. Hence, the total charge sent by v is at most $\max\{4 \cdot 3/2 + 1 + 2 \cdot 1/2, 3 \cdot 3/2 + 2 \cdot 5/4 + 2 \cdot 1/2\} = 8$.

Case 4.3.2(2). Suppose that v is incident to exactly one 5-face, say it is f_0 without loss of generality. Then v cannot be adjacent to an (≥ 6)-face, for otherwise the total charge sent by v is at most $5 \cdot 3/2 + 1/2 = 8$. As above, since configurations B3 and C1 are reducible, none of the remaining faces (all (≤ 4)-faces) is a $(3, 7, 7)$ -, $(4, 6, 7)$ - or $(5, 5, 7)$ -triangle and at most two of them are $(3, 7, 8)$ -triangles (either f_1 or f_6). Indeed, one of f_1 and f_6 , must be a $(3, 7, 8)$ -triangle, for otherwise the charge sent by v is at most $6 \cdot 5/4 + 1/2 = 8$. Assume without loss of generality that v_0 has degree 3. Therefore, since configuration D1 is reducible, for each $i \in \{2, 3, 4, 5\}$, either f_i is a 4-face or both v_i and v_{i+1} have degree at least 5. It follows that each of f_2, f_3, f_4 and f_5 is sent charge at most $11/10$; thus, v sends total charge at most $2 \cdot 3/2 + 4 \cdot 11/10 + 1/2 < 8$.

Case 4.3.2(3). It cannot be that v is incident to two (≥ 6)-faces, for then the total charge sent by v would be at most $5 \cdot 3/2 < 8$. The case in which v is incident to exactly one (≥ 6)-face is handled by an argument identical to the one used in the previous paragraph.

Case 4.3.2(4). If v is incident only to (≤ 4)-faces, then, since configurations B3 and C1 are reducible, v is not incident to a $(3, 7, \geq 7)$ -, $(4, 6, 7)$ - or $(5, 5, 7)$ -triangle. If v is incident to a $(4, 7, \geq 7)$ -triangle, then, since configuration D2 is reducible, v cannot be adjacent to any other 4-vertex. It therefore follows that the total charge sent by v in this case is at most $2 \cdot 5/4 + 5 \cdot 11/10 = 8$.

This concludes the analysis of the final charge of the 7-vertices.

4.4. Final charge of 8-vertices. Last, suppose that $\deg(v) = 8$. For convenience, let v_0, \dots, v_7 be the neighbours of v in clockwise order, and for $i \in \{0, \dots, 7\}$, let f_i be the face of G incident with vv_i and vv_{i+1} , where the index is modulo 8. The initial charge of v is $\text{ch}(v) = 10$. We partition our analysis based on the number of incident special faces. Note that since configuration C1 is reducible, if v is adjacent to a 2-vertex then at least one of the 2-vertex's incident faces is special for v . Furthermore, since configurations A2 and B1 are reducible, if one of the 2-vertex's incident faces is an (≤ 4)-face, then it must be a $(2, 8, \geq 4, 8)$ -face.

4.4.1. There is an incident special face. We start with the cases in which there is a face that is special for v . In these cases, we may assume that there is at most one incident non-special (≥ 5)-face (that is sent charge at most $1/2$ by rule R1g),

for otherwise the total charge sent by v is at most $5 \cdot 3/2 + 1 + 2 \cdot 1/2 < 10$.

Suppose that v is incident to (exactly) one non-special (≥ 5)-face. It may not be that v is incident to more than one special face, since then v would send charge at most $5 \cdot 3/2 + 2 + 1/2 = 10$. So the remaining faces are (≤ 4)-faces. Let us suppose that v_0 is the 2-vertex, f_0 is the special face and f_7 is a $(2, 8, \geq 4, 8)$ -face (sent charge at most $5/4$ by rules R1c, R1e and R1f), without loss of generality. Among the faces f_1, \dots, f_6 , (disregarding which one is the non-special (≥ 5)-face), there must be three (≤ 4)-faces that are sequentially adjacent around v . Since configurations B4 and D1 are reducible, the middle of these faces may not be a $(3, \geq 7, 8)$ -, $(4, 6, 8)$ - or $(5, 5, 8)$ -triangle and hence is sent charge at most $5/4$. Therefore, the total charge sent by v is at most $4 \cdot 3/2 + 2 \cdot 5/4 + 1 + 1/2 = 10$.

So we assume now that every (≥ 5)-face incident to v is special for v . If v has at least four incident special faces, then the charge sent is at most $4 \cdot 3/2 + 4 = 10$.

Case 4.4.1(1). Suppose that v is incident to exactly three special faces. If v is incident to at least two $(2, 8, \geq 4, 8)$ -faces, each sent charge at most $5/4$ by rule R1c, then the total charge sent by v is at most $3 \cdot 3/2 + 2 \cdot 5/4 + 3 = 10$. If v is incident to exactly one $(2, 8, \geq 4, 8)$ -face, then it must be that v is incident to three sequentially adjacent (≤ 4)-faces, say f_0, f_1 and f_2 . Since configuration B4 is reducible, f_1 is not a $(3, 7, 8)$ -, $(4, 6, 8)$ - or $(5, 5, 8)$ -face; since v is incident to a $(2, 8, \geq 4, 8)$ -face and configuration D1 is reducible, f_1 is not a $(3, 8, 8)$ -face; hence, f_1 receives charge at most $5/4$. Consequently, the total charge sent by v is at most $3 \cdot 3/2 + 2 \cdot 5/4 + 3 = 10$. If v is not incident to a $(2, 8, \geq 4, 8)$ -face, then the three incident special faces are sequentially adjacent around v . In the following we shall assume that f_5, f_6 and f_7 are the three special faces. The remaining five faces are all triangles, otherwise (by rule R1f) the total charge sent by v is at most $4 \cdot 3/2 + 4 = 10$. Note that there is no $i \in \{1, 2, 3\}$ such that all of f_{i-1}, f_i, f_{i+1} are $(3, \geq 7, 8)$ -triangles since configuration C4 is reducible. Since configuration B4 is reducible, none of f_1, f_2, f_3 is a $(3, 7, 8)$ -, $(4, 6, 8)$ - or $(5, 5, 8)$ -triangle. Furthermore, since configuration D2 is reducible, v is incident to at most one pair of adjacent $(3, \geq 7, 8)$ -triangles. Thus, at most one of v_1, \dots, v_4 is a 3-vertex.

If none of v_1, \dots, v_4 is a 3-vertex, then the only faces that can be sent charge more than $5/4$ are f_0 and f_4 . Therefore, the total charge sent by v is at most $2 \cdot 3/2 + 3 \cdot 5/4 + 3 < 10$.

Suppose that v_2 or v_3 is a 3-vertex, say v_2 by symmetry. Then v_1 and v_3 are 8-vertices. Since configuration C4 is reducible, v_0 and v_4 are (≥ 4)-vertices and hence f_0 and f_3 are each sent charge at most $5/4$. Thus, the total charge sent by v is at most $3 \cdot 3/2 + 2 \cdot 5/4 + 3 = 10$.

Suppose that v_1 or v_4 is a 3-vertex, say v_1 by symmetry. Then f_4 is the only face other than f_0 and f_1 that can be sent charge more than $5/4$. In this case, the total charge sent by v is at most $3 \cdot 3/2 + 2 \cdot 5/4 + 3 = 10$.

Case 4.4.1(2). Suppose that v is incident to exactly two special faces (and hence is incident to at most two $(2, 8, \geq 4, 8)$ -faces). First, assume that v is incident to a $(2, 8, \geq 4, 8)$ -face. Since configuration B4 is reducible, if f_i is a $(3, 7, 8)$ -, $(4, 6, 8)$ - or $(5, 5, 8)$ -triangle, then f_{i-1} or f_{i+1} is an (≥ 5)-face; also, since configuration D1 is reducible (and v is incident to a $(2, 8, \geq 4, 8)$ -face), the same conclusion holds if f_i is a $(3, 8, 8)$ -triangle. Since each incident special face is sequentially adjacent either to an incident $(2, 8, \geq 4, 8)$ -face or to the other special face, we deduce that at most two faces are sent charge more than $5/4$. Thus, the total charge sent by v is at most $2 \cdot 3/2 + 4 \cdot 5/4 + 2 = 10$.

Now we deal with the case where v is not incident to a $(2, 8, \geq 4, 8)$ -face. Suppose f_6 and f_7 are the two special faces, with v_7 being a 2-vertex. Recall that none of f_0, \dots, f_5 is an (≥ 5) -face. Note also that at most one of f_0, \dots, f_5 is a 4-face, for otherwise the charge sent by v is at most $4 \cdot 3/2 + 4 = 10$. We analyse possible $(3, \geq 7, 8)$ -triangles among these six faces. First, note that there is no $i \in \{1, 2, 3, 4\}$ such that all of f_{i-1}, f_i, f_{i+1} are $(3, \geq 7, 8)$ -triangles since configuration C4 is reducible. Since configuration B4 is reducible, none of f_1, \dots, f_4 is a $(3, 7, 8)$ -, $(4, 6, 8)$ - or $(5, 5, 8)$ -triangle. Furthermore, since configuration D2 is reducible, v is incident to at most one pair of adjacent $(3, \geq 7, 8)$ -triangles.

Case 4.4.1(2)(a). First, we suppose that all of f_0, \dots, f_5 are triangles. It follows that at most one of v_1, \dots, v_5 is a 3-vertex. We consider several cases regarding which neighbours of v are (≤ 4) -vertices.

If none of v_1, \dots, v_5 is a 3-vertex, then the only faces that can be sent charge more than $5/4$ are f_0 and f_5 . Therefore, the total charge sent by v is at most $2 \cdot 3/2 + 4 \cdot 5/4 + 2 = 10$.

Suppose that v_3 is a 3-vertex. Hence, v_2 and v_4 are 8-vertices. We show that f_0 and f_1 are sent charge at most $5/2$ altogether by v . Indeed, if v_1 is a 4-vertex, then $\deg(v_0) \geq 7$ because configurations B1 and E2 are reducible. Hence, v sends charge $5/4$ to each of f_0 and f_1 by rule R1c. If v_1 is a 5-vertex, then $\deg(v_0) \geq 5$ as configuration B1 is reducible, and hence v sends charge $11/10$ to f_1 and at most $7/5$ to f_0 by rules R1b and R1e. Last, if $\deg(v_1) \geq 6$ then v sends charge 1 to f_1 and at most $3/2$ to f_0 by rules R1a and R1f. Similarly, we deduce that v sends charge at most $5/2$ to f_4 and f_5 altogether. Therefore, the total charge sent by v is at most $2 \cdot 3/2 + 2 \cdot 5/2 + 2 = 10$.

Suppose that v_2 or v_4 is a 3-vertex, say v_2 by symmetry. Then, v_1 and v_3 are 8-vertices. We have $\deg(v_0) \geq 4$ since configuration C4 is reducible, so that f_0 receives charge at most $5/4$ from v . Thus, it suffices to show that v sends to f_3, f_4 and f_5 charge at most $15/4$ altogether: the total charge sent by v would then be at most $2 \cdot 3/2 + 5/4 + 15/4 + 2 = 10$. First, $\deg(v_4) \geq 5$ since configuration E1 is reducible. Recall that $\deg(v_4) + \deg(v_5) \geq 11$. If $\deg(v_4) + \deg(v_5) \geq 12$, then f_3 and f_4 are sent charge at most $9/4$ altogether by v . Thus, the conclusion holds since f_6 is sent charge at most $3/2$ by v . Now, if $\deg(v_4) + \deg(v_5) = 11$, then $\deg(v_5) + \deg(v_6) \geq 11$ since configuration C5 is reducible. Consequently, each of f_4 and f_5 is sent charge at most $5/4$ by v : recall that none of v_3, v_4 and v_5 is a 3-vertex, and if v_6 were a 3-vertex then v_5 would be an 8-vertex so that v_4 would have to be a 3-vertex in order for the charge sent to be more than $5/4$. Moreover, f_3 is sent charge at most $11/10$ by rule R1e, so that the conclusion holds.

Suppose that v_1 or v_5 is a 3-vertex, say v_1 by symmetry. Then, $\deg(v_0) = 8 = \deg(v_2)$, and $\deg(v_3) \geq 5$ since configuration E1 is reducible. Further, recall that v_4 and v_5 both have degree at least 4. If $\deg(v_6) \leq 4$, then $\deg(v_5) \geq 6$. Since $\deg(v_3) + \deg(v_4) \geq 11$ (because configuration B4 is reducible), at least one of f_2 and f_4 is sent charge at most 1, implying that the total charge sent by v is at most $3 \cdot 3/2 + 2 \cdot 5/4 + 3 = 10$. If $\deg(v_6) \geq 5$, then f_5 is sent charge at most $7/5$ and f_2 is sent charge at most $11/10$, so the total charge sent by v is at most $2 \cdot 3/2 + 7/5 + 2 \cdot 5/4 + 11/10 + 2 = 10$.

Case 4.4.1(2)(b). Assume now that (exactly) one of f_0, \dots, f_5 is a 4-face. (Such a 4-face is assumed to not have an incident 2-vertex.) Without loss of generality, we may suppose that it is one of f_0, f_1 and f_2 . Recall that none of f_1, \dots, f_4 is a $(3, 7, 8)$ -,

(4, 6, 8)- or (5, 5, 8)-triangle since configuration B4 is reducible. Also, at most one of v_1, \dots, v_5 is a 3-vertex since configurations C4 and D2 are reducible. Moreover, if at most one of f_1, \dots, f_4 is sent charge $3/2$ by v (i.e. is a (3, 8, 8)-triangle), then the total charge sent by v is at most $3 \cdot 3/2 + 2 \cdot 5/4 + 3 = 10$. In particular, we assume that (exactly) one of v_2, v_3, v_4 has degree 3.

Suppose that the 4-face is f_0 . At most three of f_1, \dots, f_5 are sent charge more than $5/4$, so the total charge sent by v is at most $3 \cdot 3/2 + 2 \cdot 5/4 + 3 = 10$.

Suppose that the 4-face is f_1 . By the remark above, one of v_3 and v_4 has degree 3. If $\deg(v_4) = 3$, then $\deg(v_3) = \deg(v_5) = 8$. Further, $\deg(v_6) \geq 4$ since configuration C4 is reducible. Consequently, the total charge sent by v is at most $3 \cdot 3/2 + 2 \cdot 5/4 + 3 = 10$. If $\deg(v_3) = 3$, then $\deg(v_4) = 8$ and we shall see that f_4 and f_5 are sent at most $5/2$ altogether by v . Indeed, let us check all of the subcases: if $\deg(v_6) \leq 4$, then $\deg(v_5) \geq 6$, implying that f_4 is sent charge 1 and f_5 is sent charge at most $3/2$; if $\deg(v_6) = 5$, then $\deg(v_5) \geq 5$, implying that f_4 is sent charge at most $11/10$ and f_5 is sent charge at most $7/5$; if $\deg(v_6) = 6$, then $\deg(v_5) \geq 5$ since configuration E2 is reducible, and so each of f_4 and f_5 are sent charge at most $6/5$; if $\deg(v_6) \geq 7$, then each of f_4 and f_5 are sent charge at most $5/4$. Therefore, the total charge sent by v is at most $3 \cdot 3/2 + 3 + 5/2 = 10$.

Suppose that the 4-face is f_2 . Then, $\deg(v_4) = 3$, for otherwise at most one face among f_1, \dots, f_4 is sent charge $3/2$ by v . Then, $\deg(v_5) = 8$ and $\deg(v_6) \geq 4$ since configuration C4 is reducible. Therefore, the total charge sent by v is at most $3 \cdot 3/2 + 2 \cdot 5/4 + 3 = 10$.

Case 4.4.1(3). Suppose that v is incident to exactly one special face. Then v is incident to a $(2, 8, \geq 4, 8)$ -face and, since configurations B4 and D1 are reducible, v is incident to at most one $(3, \geq 7, 8)$ -, $(4, 6, 8)$ - or $(5, 5, 8)$ -face; the total charge sent by v is at most $3/2 + 6 \cdot 5/4 + 1 = 10$.

4.4.2. There is no incident special face. Suppose that v is not incident to a special face. Any (≥ 5) -face incident to v is sent charge at most $1/2$ by rule R1g. If there are two such faces, then the total charge sent by v is at most $6 \cdot 3/2 + 2 \cdot 1/2 = 10$.

Case 4.4.2(1). Suppose there is exactly one incident (≥ 5) -face, say f_0 . None of the faces f_2, \dots, f_6 is a $(3, 7, 8)$ -, $(4, 6, 8)$ - or $(5, 5, 8)$ -triangle since configuration B4 is reducible. Since configuration D2 is reducible, among the vertices v_2, \dots, v_7 there is at most one 3-vertex that is incident to some triangle among f_2, \dots, f_6 . None of the vertices v_2, \dots, v_7 is a 2-vertex since configuration C1 is reducible (so in particular none of f_2, \dots, f_6 is a $(2, 8, \leq 5, 8)$ -face). Consequently, f_1, \dots, f_7 are all triangles, or else the total charge sent by v is at most $4 \cdot 3/2 + 2 \cdot 5/4 + 1 + 1/2 = 10$. Moreover, one of v_2, \dots, v_7 must be a 3-vertex or else the total charge sent by v is at most $3 \cdot 3/2 + 4 \cdot 5/4 + 1/2 = 10$. Suppose, without loss of generality, that the 3-vertex is v_2, v_3 or v_4 . If it is v_2 , then at most three faces are sent charge $3/2$ and the total charge sent by v is at most $3 \cdot 3/2 + 4 \cdot 5/4 + 1/2 = 10$. If it is v_3 , then v_2 has degree 8 and v sends charge at most 10 unless v_1 has degree 3; however, this contradicts the reducibility of configuration C4. If it is v_4 , then v_3 has degree 8 and there are three sub-cases. First, if v_2 has degree 6, then v sends f_2 charge 1 and charge at most $4 \cdot 3/2 + 2 \cdot 5/4 + 1 + 1/2 = 10$ in total; second, if v_2 has degree 5, then v sends f_2 charge at most $11/10$, f_1 charge at most $7/5$, and charge at most $3 \cdot 3/2 + 7/5 + 11/10 + 2 \cdot 5/4 + 1/2 = 10$ in total; third, if v_2 has degree 4, then, since configuration E2 is reducible, v_1 has degree at least 7, so v sends f_1 and f_2 each charge at most $5/4$, and charge at most $3 \cdot 3/2 + 4 \cdot 5/4 + 1/2 = 10$ in total.

Case 4.4.2(2). Finally, we are in the case that v is only incident to (≤ 4)-faces (in particular none of which are $(2, 8, \leq 5, 8)$ -faces since configuration C1 is reducible). Since configuration B4 is reducible, v is incident to no $(3, 7, 8)$ -, $(4, 6, 8)$ - or $(5, 5, 8)$ -triangle. If v is not incident to a $(3, 8, 8)$ -triangle, then no face is sent charge more than $5/4$ and hence the total charge sent by v is at most $8 \cdot 5/4 = 10$. So assume that v_7 is a 3-vertex, and that f_7 is a $(3, 8, 8)$ -triangle. Since configuration D2 is reducible, v is adjacent to no other 3-vertices. We may assume that v is incident to fewer 4-faces than the number of $(3, 8, 8)$ -triangles incident to v . (Otherwise, if x is the number of 4-faces incident to v , then the total charge sent by v is at most $x \cdot (3/2 + 1) + (8 - 2 \cdot x) \cdot 5/4 = 10$.) If f_7 were the only $(3, 8, 8)$ -triangle then f_6 would necessarily be a 4-face. We conclude, therefore, that v is incident to exactly two $(3, 8, 8)$ -triangles, namely f_6 and f_7 , and to at most one 4-face. Now, if v is incident to only triangles, then since configuration E3 is reducible, every neighbour of v other than v_7 has degree at least 5, and so the total charge sent is at most $2 \cdot 3/2 + 4 \cdot 6/5 + 2 \cdot 11/10 = 10$ (where we observe that the faces adjacent around v to the $(3, 8, 8)$ -triangles are ($\geq 5, 8, 8$)-faces and hence sent charge at most $11/10$).

Therefore, in addition to the two $(3, 8, 8)$ -triangles, v must be incident to exactly one 4-face. By symmetry, let us assume that f_0, f_1 and f_2 are triangles. Since configuration E1 is reducible, v_1 has degree at least 5. If v_1 has degree at least 6, then v sends f_0 charge 1 and total charge at most $2 \cdot 3/2 + 4 \cdot 5/4 + 2 = 10$. If v_1 has degree 5, then v sends f_0 charge $11/10$ and v_2 has degree at least 6. If v_2 has degree at least 7, then v sends f_1 charge $11/10$ and total charge at most $2 \cdot 3/2 + 3 \cdot 5/4 + 2 \cdot 11/10 + 1 < 10$. If v_2 has degree 6, then v sends f_1 and f_2 each charge at most $6/5$, and total charge at most $2 \cdot 3/2 + 2 \cdot 5/4 + 2 \cdot 6/5 + 11/10 + 1 = 10$.

We have shown that if $\deg(v) = 8$, then $\text{ch}^*(v) \geq 0$. This allows us to conclude our analysis of the final charge of v , having shown $\text{ch}^*(v) \geq 0$ in all cases. This completes the proof of Theorem 1.1. \square

5. Proofs of reducibility. In this section, we prove that the graph G cannot contain any of the configurations given in Section 2.

Let λ be a (partial) edge-face 9-colouring of G . For each element $x \in E \cup F$, we define $\mathcal{C}(x)$ to be the set of colours (with respect to λ) of the edges and faces incident or adjacent to x . If $x \in V$ we define $\mathcal{E}(x)$ to be the set of colours of the edges incident to x . Moreover, λ is *nice* if only some (≤ 4)-faces are uncoloured. Observe that every nice colouring can be greedily extended to an edge-face 9-colouring of G , since $|\mathcal{C}(f)| \leq 8$ for each (≤ 4)-face f , i.e. f has at most 8 forbidden colours. Therefore, in the rest of the paper, we shall always suppose that such faces are coloured at the very end. More precisely, every time we consider a partial colouring of G , we uncolour all (≤ 4)-faces, and implicitly colour them at the very end of the colouring procedure of G . We make the following observation about nice colourings, which we rely on frequently.

Observation. Let e be an edge of G incident to two faces f and f' . There exists a nice colouring λ of $G - e$, and hence a partial edge-face 9-colouring of G in which only e and f are uncoloured. Moreover, if f is an (≤ 4)-face, then it suffices to properly colour the edge e with a colour from $\{1, 2, \dots, 9\}$ to extend λ to a nice colouring of G .

The following lemma implies the reducibility of configuration A0. We require the stronger form as it is necessary for later arguments, in particular, for the reducibility of configurations A1–A3.

LEMMA 5.1. *Let v be a vertex of G with neighbours v_0, v_1, \dots, v_{d-1} in clockwise order. If v is a cut-vertex of G , then no component C of $G - v$ is such that the neighbourhood of v in C is contained in $\{v_i, v_{i+1}\}$ for some $i \in \{0, 1, \dots, d-1\}$, where*

the index i is taken modulo d .

Proof. Suppose on the contrary that C is a component of $G - v$ such that the neighbourhood of v in C is contained in, say, $\{v_0, v_1\}$.

First, assume that the neighbourhood of v in C is $\{v_0, v_1\}$. Then G is the edge-disjoint union of two plane graphs $G_1 = (C \cup \{v\}, E_1)$ and $G_2 = (V \setminus C, E_2)$. The outer face f_1 of G_1 corresponds to a face f_2 of G_2 . By the minimality of G , the graph G_i has an edge-face 9-colouring λ_i for $i \in \{1, 2\}$. Since both vv_0 and vv_1 are incident in G_1 to f_1 , we may assume that $\lambda_1(f_1) = 1$, $\lambda_1(vv_0) = 8$ and $\lambda_1(vv_1) = 9$. Regarding λ_2 , we may assume that $\lambda_2(f_2) = 1$. Furthermore, up to permuting the colours, we can also assume that the colours of the edges of G_2 incident to v are contained in $\{1, 2, \dots, 7\}$, since there are at most 6 such edges.

We now define an edge-face 9-colouring λ of G as follows. For every edge e of G , set $\lambda(e) := \lambda_1(e)$ if $e \in E_1$ and $\lambda(e) := \lambda_2(e)$ if $e \in E_2$. To colour the faces of G , let f be the face of G incident to both vv_0 and vv_{d-1} . (Note that there is only one such face, since otherwise v would have degree 2, which would be a contradiction.) Now observe that there is a natural one-to-one correspondence between the faces of G_1 and a subset F_1 of the face set F of G that maps f_1 to f . Similarly, there is a natural one-to-one correspondence between the faces of G_2 and a subset F_2 of F that maps f_2 to f . Note that $F_1 \cap F_2 = \{f\}$. Now, we can colour every face $f \in F_i$ using λ_i . This is well defined since $\lambda_1(f_1) = \lambda_2(f_2) = 1$.

Let us check that λ is proper. Two adjacent edges of G are assigned different colours. Indeed, if the two edges belong to E_i for some $i \in \{1, 2\}$, then it comes from the fact that λ_i is an edge-face 9-colouring of G_i . Otherwise, both edges are incident with v , and one is in G_1 and the other in G_2 . The former is coloured either 8 or 9, and the latter with a colour of $\{1, 2, \dots, 7\}$ by the choice of λ_1 and λ_2 . Two adjacent faces in G necessarily correspond to two adjacent faces in G_1 or G_2 , and hence are assigned different colours. Last, let g be a face of G and e an edge incident to g in G . If $g \neq f$, then g and e are incident in G_1 or G_2 , and hence coloured differently. Otherwise e is incident to f_i in G_i for some $i \in \{1, 2\}$, and hence $\lambda(e) = \lambda_i(e) \neq \lambda_i(f_i) = 1 = \lambda(f)$.

The case where the neighbourhood of v in C is $\{v_0\}$, i.e. vv_0 is a cut-edge, is dealt with in the very same way so we omit it. \square

The next lemma shows the reducibility of configurations B1 and C1.

LEMMA 5.2. *Let uv be an edge of G , and let $s \in \{1, 2\}$ be the number of (≤ 4)-faces incident to uv . Then $\deg(u) + \deg(v) \geq 9 + s$.*

Proof. Suppose on the contrary that $\deg(u) + \deg(v) \leq 8 + s$. Let f and f' be the two faces incident to uv .

Without loss of generality assume that f is an (≤ 4)-face. By the minimality of G , the graph $G - uv$ has a nice colouring λ . Let f'' be the face of $G - uv$ corresponding to the union of the two faces f and f' of G after having removed the edge uv . We obtain a partial edge-face 9-colouring of G in which only uv , f and the (≤ 4)-faces are uncoloured by just assigning the colour $\lambda(f'')$ to f' , and keeping all the other assignments.

Consequently, $|\mathcal{C}(uv)| \leq \deg(u) + \deg(v) - 2 + 2 - s \leq 8$. Hence, we can properly colour the edge uv , thereby obtaining a nice colouring of G ; a contradiction. \square

In light of Lemma 5.2, we make the following definition and observation. An edge uv of G is called *tight* if $\deg(u) + \deg(v) - s = 9$, where $s \in \{1, 2\}$ is the number of (≤ 4)-faces incident to uv .

Observation. Assume that c is an edge-face 9-colouring of G in which only uv and the (≤ 4)-faces are uncoloured. Let S be the (possibly empty) set of colours assigned

by c to the (≥ 5)-faces incident to uv . If uv is tight, then the sets $\mathcal{E}(u)$, $\mathcal{E}(v)$ and S are pairwise disjoint, and $\mathcal{C}(uv) = \mathcal{E}(u) \cup \mathcal{E}(v) \cup S = \{1, \dots, 9\}$.

The reducibility of configurations B2, B3, B4, C2 and C3 follows from the next lemma.

LEMMA 5.3. *Let uvw be a triangle of G such that $\deg(u) + \deg(v) = 10 + s$, where $s \in \{0, 1\}$ is the number of (≤ 4)-faces distinct from uvw incident to uv , and let $t \in \{0, 1, 2\}$ be the number of (≤ 4)-faces distinct from uvw incident to uw or vw . Then $\deg(w) \geq 7 + t$.*

Proof. As we pointed out, there exists a partial edge-face 9-colouring c of G in which only uv and the (≤ 4)-faces are left uncoloured. Let α_{uv} , α_{vw} and α_{uw} be the colours, if any, assigned to the (≥ 5)-faces incident to uv , vw and uw , respectively. Since the edge uv is tight, $\mathcal{E}(u)$, $\mathcal{E}(v)$ and $\{\alpha_{uv}\}$ form a partition of $\{1, 2, \dots, 9\}$. Thus, if there is a colour $\xi \in \mathcal{E}(u) \cup \{\alpha_{uv}\}$ that is not in $\mathcal{E}(w) \cup \{\alpha_{vw}\}$, then we can colour uv with $c(vw)$ and next recolour vw with ξ to obtain a nice colouring of G . We deduce that $\mathcal{E}(u) \cup \{\alpha_{uv}\} \subseteq \mathcal{E}(w) \cup \{\alpha_{vw}\}$. Similarly, $\mathcal{E}(v) \cup \{\alpha_{vw}\} \subseteq \mathcal{E}(w) \cup \{\alpha_{uw}\}$. Hence, $\deg(w) + 2 - t \geq 9$, so $\deg(w) \geq 7 + t$, as required. \square

The following verifies that configuration A1 is reducible. The lemma is also needed for showing the reducibility of configurations A3, C4 and C5.

LEMMA 5.4. *Let u, v, w be vertices of G with $\deg(v) = 2$. Then uvw is not a face of G .*

Proof. Suppose on the contrary that uvw is a face of G . There exists a nice colouring c of $G - uv$. Note that the face f_1 of G other than uvw that is incident to both uv and vw must be an (≥ 5)-face, or else $|\mathcal{C}(uv)| \leq 8$ and we can immediately extend c to a nice colouring of G . Note that f_1 is distinct from the face f_2 of G other than uvw that is incident to uw ; otherwise, one of u or w is a cut-vertex of a type forbidden by Lemma 5.1. Let β_{uw} be the colour, if any, assigned by c to f_2 .

Observe that $c(f_1) \notin \{\beta_{uw}, c(uw)\}$. Indeed, since uv is tight, the sets $\mathcal{E}(u)$, $\{c(vw)\}$ and $\{c(f_1)\}$ are pairwise disjoint. Since vw is tight, we deduce that $c(f_1) \notin \mathcal{E}(w)$, for otherwise we could colour uv with $c(vw)$ and next recolour vw with a colour from $\{1, \dots, 9\} \setminus \mathcal{E}(w)$. Hence $c(f_1) \notin \mathcal{E}(u) \cup \mathcal{E}(w) \cup \{\beta_{uw}\}$, so that colouring uv with $c(uw)$ and next recolouring uw with $c(f_1)$ yields a nice colouring of G ; a contradiction. \square

The next lemma will be used to show the reducibility of configurations A3 and E4.

LEMMA 5.5. *Let v be a 2-vertex of G , and let u and w be its two neighbours. If $\deg(u) \leq 6$, then u and w are adjacent in G .*

Proof. Suppose on the contrary that u and w are not adjacent in G . Then, the graph G' obtained by contracting the edge uv is planar, simple and has maximum degree at most 8. By the minimality of G , let λ be a nice colouring of G' . Let g and g' be the faces of G' corresponding to the contracted faces f and f' of G , respectively. We obtain a partial edge-face 9-colouring of G in which only uv is uncoloured by assigning the colour $\lambda(g)$ to f , the colour $\lambda(g')$ to f' , and keeping all the other assignments.

Now, $|\mathcal{C}(uv)| \leq \deg(u) + \deg(v) - 2 + 2 \leq 8$. Consequently, we can properly colour the edge uv to obtain a nice colouring of G ; a contradiction. \square

We now deduce the reducibility of configuration A3.

COROLLARY 5.6. *Let v be a 2-vertex of G and let u and w be its two neighbours. If $\deg(u) = 3$, then $\deg(w) \geq 6$.*

Proof. Suppose on the contrary that u has degree 3 and w has degree at most 5. Lemma 5.5 implies that u and w are adjacent. Note that uvw cannot be a face by Lemma 5.4. Let u' be the neighbour of u besides v and w . By Jordan's curve theorem,

the curve uvw splits the plane into two parts, \mathcal{I} and \mathcal{O} with $u' \in \mathcal{O}$. First, note that w has a neighbour in \mathcal{O} , for otherwise u would be a cut-vertex that contradicts Lemma 5.1. Moreover, Lemma 5.4 implies that w has a neighbour in \mathcal{I} . Consequently, w has either one or two neighbours in \mathcal{I} , and hence w is a cut-vertex that contradicts Lemma 5.1. \square

The following demonstrates that configuration A2 is reducible.

LEMMA 5.7. *Let u, v, w, x be vertices of G with $\deg(v) = 2$ and $\deg(x) \leq 3$. Then $uvw x$ is not a face of G .*

Proof. Suppose on the contrary that $uvw x$ is a face of G . There exists a partial edge-face 9-colouring c of G in which only uv and the (≤ 4) -faces are uncoloured. Let α be the colour, if any, assigned to the (≥ 5) -face incident to both uv and vw , and let β_{ux} and β_{wx} be the colours, if any, assigned to the (≥ 5) -faces incident to ux and wx , respectively.

By Lemma 5.1, observe that $\alpha \notin \{\beta_{ux}, \beta_{wx}, c(ux), c(wx)\}$. Since uv is tight, the sets $\mathcal{E}(u)$, $\{c(vw)\}$ and $\{\alpha\}$ are pairwise disjoint. Since vw is tight, we deduce that $\alpha \notin \mathcal{E}(w)$, for otherwise we could colour uv with $c(vw)$ and next recolour vw with a colour from $\{1, \dots, 9\} \setminus \mathcal{E}(w)$. Hence $\alpha \notin \mathcal{E}(u) \cup \mathcal{E}(w) \cup \{\beta_{ux}, \beta_{wx}\}$.

Let x' be the vertex adjacent to x distinct from u and w . We must have $c(xx') = \alpha$, otherwise we could colour uv with $c(ux)$ and next recolour ux with α . Since $\beta_{ux} \neq \beta_{wx}$, at least one of β_{ux} and β_{wx} is distinct from $c(vw)$. Observing that we can colour uv with $c(vw)$ and next uncolour vw , we may assume without loss of generality that $\beta_{wx} \neq c(vw)$. As a result, colouring uv with $c(vw)$, and next swapping the colours of vw and xw yields a nice colouring of G ; a contradiction. \square

The following verifies that configurations C4 and C5 are reducible.

LEMMA 5.8. *Let uvw and vwx be triangles of G such that wx is incident to two (≤ 4) -faces.*

- (i) *At least one of u and x has degree at least 4.*
- (ii) *If uv is tight, then $\deg(v) + \deg(x) \geq 12$.*

Proof. (i). Suppose on the contrary that both u and x have degree less than 4. Then both have degree 3 by Lemma 5.4. Let u' (respectively x') be the neighbour of u (respectively x) distinct from v and w . Let c be a partial edge-face 9-colouring of G in which only wx and the (≤ 4) -faces are uncoloured. Let α_{uv} , α_{uw} and α_{vx} be the colours, if any, assigned to the (≥ 5) -faces incident to uv , uw and vx , respectively.

Since the edge wx is tight, the sets $\mathcal{E}(w)$ and $\mathcal{E}(x)$ are disjoint. Hence $c(xx') \in \mathcal{E}(v)$, otherwise we could colour wx with $c(vw)$ and recolour vw with $c(xx')$.

We first assert that $\alpha_{vx} \neq c(vw)$. Otherwise, $|\mathcal{C}(vx)| = |\mathcal{E}(v)| \leq 8$ and there exists $\xi \in \{1, 2, \dots, 9\} \setminus \mathcal{C}(vx)$. Now, colouring wx with $c(vx)$ and recolouring vx with ξ yields a nice colouring of G ; a contradiction. Consequently, we can safely swap the colours of vw and vx , if necessary.

Our next assertion is that $\{c(uu'), \alpha_{uw}\} = \{c(vw), c(vx)\}$. For, if $c(vx) \notin \{c(uu'), \alpha_{uw}\}$, we can colour wx with $c(uw)$ and recolour uw with $c(vx)$; a contradiction. The same argument after swapping the colours of vw and vx shows that $c(vw) \in \{c(uu'), \alpha_{uw}\}$. Thus, up to swapping the colours of vw and vx , we may assume that $c(vx) = \alpha_{uw}$.

Let us recolour uv with $c(vx)$, colour wx with $c(vx)$ and uncolour vx . The obtained colouring is proper, since $\alpha_{uv} \neq \alpha_{uw} = c(vx)$ and $\mathcal{E}(w) \cap \mathcal{E}(x) = \emptyset$. Now, if vx cannot be coloured greedily, then for the obtained colouring $\mathcal{E}(v) \cup \mathcal{E}(x) \cup \{\alpha_{vx}\} = \{1, 2, \dots, 9\}$. But then, since there are at most ten in the set of edges incident to v or x , two of which (uv and wx) have the same colour and one of which is uncoloured, it follows

that $c(xx') \notin \mathcal{E}(v) \cup \mathcal{E}(w)$. Now we can colour vx with $c(vw)$ and colour vw with $c(xx')$ to obtain a nice colouring of G ; a contradiction.

(ii). Suppose on the contrary that uv is tight and $\deg(v) + \deg(x) = 11$. Let c be a partial edge-face 9-colouring of G in which only uv and the (≤ 4)-faces are uncoloured. Let α_{uv} , α_{uw} and α_{vx} be the colours, if any, assigned to the (≥ 5)-faces incident to uv , uw and vx , respectively. Since the edge uv is tight, the sets $\mathcal{E}(u)$, $\mathcal{E}(v)$ and $\{\alpha_{uv}\}$ are pairwise disjoint.

Let ξ be the colour in $\{1, \dots, 9\} \setminus \mathcal{E}(w)$ (unique since we can assume that $\deg(w) = 8$ without loss of generality). Then $\xi \in \mathcal{E}(v)$, otherwise we could colour uv with $c(vw)$ and recolour vw with ξ . It follows that $\xi \notin \mathcal{E}(u)$. Therefore, $\alpha_{uw} = \xi$, otherwise we could colour uv with $c(uw)$ and recolour uw with ξ . Thus, the colours of uw and vw may be exchanged, if necessary.

Let us show that $\mathcal{E}(u) \cup \{\alpha_{uv}, c(vw)\} \subseteq \mathcal{E}(x) \cup \{\alpha_{vx}\}$. First, if there is a colour $\gamma \in \mathcal{E}(u) \cup \{\alpha_{uv}\}$ that is not in $\mathcal{E}(x) \cup \{\alpha_{vx}\}$, then we can recolour vx with γ and then colour uv with $c(vx)$ to obtain a nice colouring of G , which is a contradiction. Similarly, by exchanging the colours of uw and vw , we conclude that $c(vw) \in \mathcal{E}(x) \cup \{\alpha_{vx}\}$.

Since uv is tight and $\deg(v) + \deg(x) = 11$, we deduce that $\mathcal{E}(x) \cup \{\alpha_{vx}\} = \mathcal{E}(u) \cup \{\alpha_{uv}, c(vw), c(vx)\}$. (Indeed, $|\mathcal{E}(x) \cup \{\alpha_{vx}\}| \leq \deg(x) + 1 = 12 - \deg(v)$, and $|\mathcal{E}(u) \cup \{\alpha_{uv}\}| = 9 - (\deg(v) - 1) = 10 - \deg(v)$.) In particular, $\alpha_{vx} \neq c(vx)$ and $\xi \notin \mathcal{E}(x) \setminus \{c(vx)\}$. Now, colour uv with $c(vx)$, and then recolour vx with $c(wx)$ and wx with ξ to obtain a nice colouring of G ; a contradiction. \square

The next lemma implies that configurations D1–D4 are reducible.

LEMMA 5.9. *Let vwx be a triangle of G and u a neighbour of v distinct from x and w . If vx is incident to two (≤ 4)-faces, then either uv or vw is not tight.*

Proof. Suppose on the contrary that both uv and vw are tight. Let c be a partial edge-face 9-colouring of G in which only vw and the (≤ 4)-faces are left uncoloured. Let α be the colour, if any, assigned to the (≥ 5)-face incident to vw . Since vw is tight, we know that the sets $\mathcal{E}(v)$, $\mathcal{E}(w)$ and $\{\alpha\}$ form a partition of $\{1, 2, \dots, 9\}$. In particular, $c(vx) \notin \mathcal{E}(w)$ and $c(wx) \notin \mathcal{E}(v)$.

If an edge e that is adjacent to vw could be properly recoloured with a colour ξ , then colouring vw with $c(e)$ and recolouring e with ξ would yield a nice colouring of G ; a contradiction. Applying this to vx yields that $\mathcal{E}(w) \cup \{\alpha\} \subseteq \mathcal{E}(x)$, since $\mathcal{C}(vx) = \mathcal{E}(x) \cup \mathcal{E}(v)$, and as we noted above $\{1, \dots, 9\} \setminus \mathcal{E}(v) = \mathcal{E}(w) \cup \{\alpha\}$. Applying the same remark to wx , we obtain $\mathcal{E}(v) \cup \{\alpha\} \subseteq \mathcal{E}(x) \cup \{\beta\}$, where β is the colour, if any, assigned to the (≥ 5)-face incident to wx .

Since $9 = |\mathcal{E}(v) \cup \mathcal{E}(w) \cup \{\alpha\}| \leq |\mathcal{E}(x) \cup \{\beta\}| \leq 9$, we deduce that $\beta \notin \mathcal{E}(x)$. Therefore, we can safely swap the colours of vx and wx if needed (recalling that $\mathcal{E}(v) \cap \mathcal{E}(w) = \emptyset$).

Let S be the set of colours of the (≥ 5)-faces incident to uv . Thus, $|S| = 2 - s$ where s is the number of (≤ 4)-faces incident to uv . Again, we apply the same arguments as above to uv : since uv cannot be recoloured, we deduce that $\mathcal{E}(u) \cup \mathcal{E}(v) \cup S = \{1, 2, \dots, 9\}$. But $|\mathcal{E}(u) \cup \mathcal{E}(v) \cup S| \leq \deg(u) - 1 + \deg(v) - 1 + 2 - s = \deg(u) + \deg(v) - s \leq 9$ since uv is tight and vw is uncoloured. Consequently, $\mathcal{E}(u)$, $\mathcal{E}(v)$ and S are pairwise disjoint. In particular, $c(vx) \notin \mathcal{E}(u) \cup S$. As a result, colouring vw with $c(uv)$, then recolouring uv with $c(vx)$ and finally swapping the colours of vx and wx yields a nice colouring of G ; a contradiction. \square

The next lemma implies that configurations E1 and E2 are reducible.

LEMMA 5.10. *Let v be an 8-vertex of G with neighbours v_0, v_1, \dots, v_7 in anti-clockwise order. Assume that $v_i v_{i+1}$ is an edge for $i \in \{0, 1, 2, 3\}$, and that v_1 an*

(≤ 4)-vertex. If v_0 is an (≤ 6)-vertex or vv_0 is adjacent to two (≤ 4)-faces, then v_3 is an (≥ 4)-vertex.

Proof. Suppose on the contrary that v_3 is a 3-vertex. By the minimality of G , the graph $G - vv_3$ has a nice colouring and hence G has a partial edge-face 9-colouring c in which only vv_3 and the (≤ 4)-faces are left uncoloured. Since vv_3 is tight, we deduce that $|\mathcal{E}(v) \cup \mathcal{E}(v_3)| = 9$ and $\mathcal{E}(v) \cap \mathcal{E}(v_3) = \emptyset$.

Let α be the colour, if any, of the (≥ 5)-face incident with both v_2v_3 and v_3v_4 . If v_2v_3 can be recoloured with a colour ξ , then colouring vv_3 with $c(v_2v_3)$ and then v_2v_3 with ξ would yield a nice colouring of G ; a contradiction. Thus, $\mathcal{E}(v) \subseteq \mathcal{E}(v_2) \cup \{\alpha\}$.

Let $j \in \{1, 2\}$. If there exists a colour $\xi \in \mathcal{E}(v_3) \setminus \mathcal{E}(v_j)$, then colouring vv_3 with $c(vv_j)$ and then vv_j with ξ yields a nice colouring of G (recalling that $\mathcal{E}(v_3)$ and $\mathcal{E}(v)$ are disjoint). Therefore, $\mathcal{E}(v_3) \subseteq \mathcal{E}(v_j)$ for $j \in \{1, 2\}$. Letting γ be the colour, if any, of the (≥ 5)-face incident to vv_0 we similarly find that $\mathcal{E}(v_3) \subseteq \mathcal{E}(v_0) \cup \{\gamma\}$.

Since $\mathcal{E}(v_2) \cup \{\alpha\} \supseteq \mathcal{E}(v) \cup \mathcal{E}(v_3) = \{1, 2, \dots, 9\}$ and $|\mathcal{E}(v_2) \cup \{\alpha\}| \leq 9$, it follows that $\alpha \neq c(vv_2)$. As $\mathcal{E}(v) \cap \mathcal{E}(v_3) = \emptyset$, this implies that the colours of vv_2 and v_2v_3 can be freely swapped. By doing so, we can conclude that $\mathcal{E}(v_3) \cup \{c(vv_2)\} \subseteq \mathcal{E}(v_j)$ for $j \in \{1, 2\}$ and $\mathcal{E}(v_3) \cup \{c(vv_2)\} \subseteq \mathcal{E}(v_0) \cup \{\gamma\}$.

Since $\deg(v_1) = 4$, we find that $\mathcal{E}(v_1) = \{c(vv_1), c(vv_2)\} \cup \mathcal{E}(v_3)$. Furthermore, by swapping the colours of vv_2 and v_2v_3 if necessary, we may assume that $c(v_0v_1) \in \mathcal{E}(v_3)$. Now, if v_0v_1 could be recoloured with a colour ξ , then colouring vv_3 with $c(vv_1)$, then vv_1 with $c(v_0v_1)$ and then v_0v_1 with ξ would yield a nice colouring of G . Thus, letting β be the colour, if any, of the (≥ 5)-face incident to v_0v_1 we obtain $\mathcal{E}(v_0) \cup \mathcal{E}(v_1) \cup \{\beta\} = \{1, 2, \dots, 9\}$.

Let us partition our analysis now based on if v_0 is an (≤ 6)-vertex or if vv_0 is adjacent to two (≤ 4)-faces.

Suppose we are in the former case. Since $\mathcal{E}(v_3) \cup \{c(vv_2)\} \subseteq (\mathcal{E}(v_0) \cup \{\gamma\}) \cap \mathcal{E}(v_1)$, we deduce that $|\mathcal{E}(v_0) \cup \mathcal{E}(v_1)| \leq \deg(v_0) + \deg(v_1) - 2 \leq 8$. Consequently, $\beta \neq c(vv_1)$ and $c(vv_1) \notin \mathcal{E}(v_0)$. In particular, the colours of vv_1 and v_0v_1 can safely be swapped if needed. As a result, colouring vv_3 with $c(vv_1)$ and then swapping the colours of vv_1 and v_0v_1 yields a nice colouring of G ; a contradiction.

Now suppose we are in the latter case. Then there is no colour γ . For $j \in \{0, 1\}$, it cannot be that $c(vv_j) \in \mathcal{E}(v_{1-j})$ (and hence $c(vv_j) \in \mathcal{E}(v_0) \cap \mathcal{E}(v_1)$). Otherwise, we would have, using $\mathcal{E}(v_3) \cup \{c(vv_2)\} \subseteq \mathcal{E}(v_0) \cap \mathcal{E}(v_1)$, that $|\mathcal{E}(v_0) \cup \mathcal{E}(v_1)| \leq \deg(v_0) + \deg(v_1) - 4 \leq 8$, in which case, recolouring as we did in the last paragraph, we would reach a contradiction. However, for some $j \in \{0, 1\}$, we must have $\beta \neq c(vv_j)$, and so the colours of vv_j and v_0v_1 can be swapped safely. Thus, colouring vv_3 with $c(vv_j)$ and then swapping the colours of vv_j and v_0v_1 yields a nice colouring of G ; a contradiction. \square

In the following lemma, we show that configuration E3 is reducible.

LEMMA 5.11. *Let v be a triangulated 8-vertex of G with neighbours v_0, v_1, \dots, v_7 in anti-clockwise order. If v_0 is a 3-vertex, then every vertex v_i with $i \neq 0$ has degree at least 5.*

Proof. Suppose on the contrary that v_j is an (≤ 4)-vertex with $j \in \{1, \dots, 7\}$. First, note that $j \notin \{1, 7\}$ by Lemma 5.3 (the reducibility of configuration B2, in particular). By the minimality of G , the graph $G - vv_0$ has a nice colouring, and hence the graph G has a partial edge-face 9-colouring in which only vv_0 and the (≤ 4)-faces are left uncoloured. Since vv_0 is tight and incident to two triangles, we infer that $|\mathcal{E}(v) \cup \mathcal{E}(v_0)| = 9$ and $\mathcal{E}(v) \cap \mathcal{E}(v_0) = \emptyset$.

Note that $\mathcal{E}(v_0) \subset \mathcal{E}(v_i)$ for $i \neq 0$, for otherwise we could colour vv_0 with $c(vv_i)$

and then recolour vv_i with a colour in $\mathcal{E}(v_0) \setminus \mathcal{E}(v_i)$ to obtain a nice colouring of G (recalling that $\mathcal{E}(v) \cap \mathcal{E}(v_0) = \emptyset$). Since $\{c(vv_j)\} \cup \mathcal{E}(v_0) \subseteq \mathcal{E}(v_j)$ and $\deg(v_j) \leq 4$, we deduce that one of $c(vv_1)$ and $c(vv_7)$ does not belong to $\mathcal{E}(v_j)$, say $c(vv_7)$.

Let α be the colour of the face incident to both v_0v_1 and v_0v_7 . We prove that $\alpha \neq c(vv_7)$. Indeed, suppose on the contrary that $\alpha = c(vv_7)$. Then, there exists a colour ξ that does not belong to $\mathcal{E}(v_7) \cup \{\alpha\} = \mathcal{E}(v_7)$, since $\deg(v_7) \leq 8$. As $\mathcal{E}(v_0) \subset \mathcal{E}(v_7)$, we deduce that $\xi \notin \mathcal{E}(v_0) \cup \mathcal{E}(v_7) \cup \{\alpha\}$. Therefore, colouring vv_0 with $c(v_0v_7)$ and then v_0v_7 with ξ yields a nice colouring of G ; a contradiction. Hence, $\alpha \neq c(vv_7)$. Consequently, we can freely swap the colours of vv_7 and v_0v_7 . Now, colouring vv_0 with $c(vv_j)$, then recolouring vv_j with $c(vv_7)$ and last swapping the colours of vv_7 and v_0v_7 yields a nice colouring of G ; a contradiction. \square

The next lemma implies that configuration E4 is reducible.

LEMMA 5.12. *Let v be a 6-vertex of G with neighbours u and w , and suppose $x \neq v$ is a neighbour of w . Suppose w is a 2-vertex and x is a 3-vertex. Assume that uw is adjacent to an (≤ 4)-face. Then u is an (≥ 6)-vertex.*

Proof. First of all note that, due to Lemma 5.5, v and x are adjacent in G . Suppose on the contrary that u is an (≤ 5)-vertex. By the minimality of G , the graph $G - uv$ has a nice colouring and hence G has a partial edge-face 9-colouring in which only uv and the (≤ 4)-faces are left uncoloured. Let us further uncolour vw and vx . Now, $|\mathcal{C}(uv)| \leq \deg(u) - 1 + \deg(v) - 1 - 2 + 1 \leq 8$, so we may properly colour uv . It remains to colour vw and vx . Next, since vw was uncoloured, we see that $|\mathcal{C}(vx)| \leq \deg(v) - 1 + \deg(x) - 1 - 1 + 2 = 8$, so we may properly colour vx . Finally, we consider vw and notice that $|\mathcal{C}(vw)| \leq \deg(v) - 1 + \deg(w) - 1 + 2 = 8$, which does not prevent us from properly colouring vw . We have thereby obtained a nice colouring of G ; a contradiction. \square

It remains to prove Lemmas 2.1 and 2.2.

Proof of Lemma 2.1. By the minimality of G , the (proper) subgraph G' formed from G by deleting all loose edges incident to f has a nice colouring. To extend this to a nice colouring of G , it would suffice to properly colour f , as every loose edge on f can then be greedily coloured. Indeed, a loose edge uv on f is incident to at most $\deg(u) - 1 + \deg(v) - 1 \leq 6$ other edges. Consequently, since G cannot have a nice colouring, we conclude that f is incident or adjacent to elements of all nine colours. Now, f is adjacent to at most $d - q$ other faces, and incident to $d - x$ coloured edges. Therefore, $d - q + d - x \geq 9$, as asserted. \square

Proof of Lemma 2.2. That $u' = v'$ follows directly from Lemma 5.5. Note that by Lemma 5.4, uvu' is not a face. By Jordan's curve theorem, the curve uvu' splits the plane into two parts, \mathcal{I} and \mathcal{O} . Then u' must have three neighbours in \mathcal{I} and three neighbours in \mathcal{O} , or else it would be a cut vertex of a type forbidden by Lemma 5.1. This implies that u' has degree 8. \square

Acknowledgements. We thank Artem Pyatkin for pointing out an error in a previous version of this work. We are grateful to the anonymous referees for their thorough reading and helpful suggestions.

This work was initiated during a visit by Ross Kang to Centre Émile Borel, Institut Henri Poincaré in Paris for the thematic term “Statistical physics, combinatorics and probability: from discrete to continuous models”, supported by the *European Research Council* (ERC), grant ERC StG 208471 ExploreMaps.

- [1] O. V. Borodin. Coupled colorings of graphs on a plane (in Russian). *Metody Diskret. Anal.*, 45:21–27, 95, 1987.
- [2] O. V. Borodin. Simultaneous coloring of edges and faces of plane graphs. *Discrete Math.*, 128(1-3):21–33, 1994.
- [3] J. Fiamčík. Simultaneous colouring of 4-valent maps. *Mat. Časopis Sloven. Akad. Vied*, 21:9–13, 1971.
- [4] E. Jucovič. On a problem in map coloring. *Mat. Časopis Sloven. Akad. Vied*, 19:225–227, 1969. Errata *ibid.* 20:224, 1970.
- [5] L. S. Mel'nikov. Problem 9. In M. Fiedler, editor, *Recent Advances in Graph Theory, Proc. International Symposium, Prague, 1974*, page 543. Academia Praha, 1975.
- [6] D. P. Sanders and Y. Zhao. On simultaneous edge-face colorings of plane graphs. *Combinatorica*, 17(3):441–445, 1997.
- [7] D. P. Sanders and Y. Zhao. On improving the edge-face colorings theorem. *Graphs Combin.*, 17(2):329–341, 2001.
- [8] J.-S. Sereni and M. Stehlík. Edge-face colouring of plane graphs with maximum degree nine. *J. Graph Theory*, 66(4):332–346, 2011.
- [9] A. O. Waller. Simultaneously coloring the edges and faces of plane graphs. *J. Combin. Theory Ser. B*, 69(2):219–221, 1997.