



Asymptotic behavior of structures made of straight rods

Dominique Blanchard, Georges Griso

► To cite this version:

Dominique Blanchard, Georges Griso. Asymptotic behavior of structures made of straight rods. 2010. <hal-00526722>

HAL Id: hal-00526722

<https://hal.archives-ouvertes.fr/hal-00526722>

Submitted on 18 Oct 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ASYMPTOTIC BEHAVIOR OF STRUCTURES MADE OF STRAIGHT RODS.

D. Blanchard¹, G. Griso²

¹ Université de Rouen, UMR 6085, 76801 Saint Etienne du Rouvray Cedex, France,
E-mail: dominique.blanchard@univ-rouen.fr

² Laboratoire J.-L. Lions–CNRS, Boîte courrier 187, Université Pierre et Marie Curie,
4 place Jussieu, 75005 Paris, France, Email: griso@ann.jussieu.fr

Abstract

This paper is devoted to describe the deformations and the elastic energy for structures made of straight rods of thickness 2δ when δ tends to 0. This analysis relies on the decomposition of the large deformation of a single rod introduced in [7] and on the extension of this technique to a multi-structure. We characterize the asymptotic behavior of the infimum of the total elastic energy as the minimum of a limit functional for an energy of order δ^β ($2 < \beta \leq 4$).

KEY WORDS: nonlinear elasticity, junctions, rods.

Mathematics Subject Classification (2000): 74B20, 74K10, 74K30.

1 Introduction

This paper concerns the modeling of a structure \mathcal{S}_δ made of elastic straight rods whose cross sections are discs of radius δ . The centerlines of the rods form the skeleton structure \mathcal{S} . We introduce a notion of elementary deformations of this structure based on the decomposition of a large deformation of a rod introduced in [6]. A special care is devoted in the junctions of the rods where these elementary deformations are translation-rotations. An elementary deformation is characterized by two fields defined on the skeleton \mathcal{S} . The first one \mathcal{V} stands for the centerlines deformation while the second one \mathbf{R} represents the rotations of the cross sections. For linearized deformations in plates or rods structures such decompositions have been considered in [14] and [16].

Then to an arbitrary deformation v of \mathcal{S}_δ , we associate an elementary deformation v_e such that the residual part $\bar{v} = v - v_e$ is controlled by $\|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)}$ and δ . In order to construct the deformation v_e we first apply the rigidity theorem of [11] -in the form given in [6]- in a neighborhood of each junction to obtain a constant translation-rotation in each junction. Then we match the decomposition derived in [6]

in each rod with this constant translation-rotation. Doing such, we obtain estimates on \mathcal{V} and \mathbf{R} for the whole structure \mathcal{S}_δ (see Theorem 3.3). Upon the assumption that the structure is fixed on some extremities, these estimates allow us to establish a nonlinear Korn's type inequality for the admissible deformations. Moreover we are in a position to analyze the asymptotic behavior of the Green-St Venant's strain tensor $E(v_\delta) = 1/2((\nabla v_\delta)^T \nabla v_\delta - \mathbf{I}_3)$ for a sequence v_δ of admissible deformations such that $\|\text{dist}(\nabla_x v_\delta, SO(3))\|_{L^2(\mathcal{S}_\delta)} = O(\delta^K)$ for $1 < K \leq 2$.

We then consider an elastic structure, whose energy is denoted W , submitted to applied body forces $f_{\kappa,\delta}$ whose order with respect to the parameter δ is related a constant $\kappa \geq 1$ (see below the order of $f_{\kappa,\delta}$). The total energy is given by $J_{\kappa,\delta}(v) = \int_{\mathcal{S}_\delta} W(E(v)) - \int_{\mathcal{S}_\delta} f_{\kappa,\delta} \cdot (v - I_d)$ if $\det(\nabla v) > 0$ where I_d is the identity map. We adopt a few usual assumptions on W (see (7.1)). We set

$$m_{\kappa,\delta} = \inf_{v \in \mathbb{D}_\delta} J_{\kappa,\delta}(v),$$

where \mathbb{D}_δ is the set of admissible deformations.

We assume that the order of $f_{\kappa,\delta}$ is equal to $\delta^{2\kappa-2}$ if $1 \leq \kappa \leq 2$ (or δ^κ if $\kappa \geq 2$) outside the junctions and that $f_{\kappa,\delta}$ is constant in each junction with order $\delta^{2\kappa-3}$ if $1 \leq \kappa \leq 2$ (or $\delta^{\kappa-1}$ if $\kappa \geq 2$). Then using the Korn's type inequality mentioned above allows us to show that the order of $m_{\kappa,\delta}$ is $\delta^{2\kappa}$. The aim of this paper is to prove that the sequence $\frac{m_{\kappa,\delta}}{\delta^{2\kappa}}$ converges (when δ tends to 0) and to characterize its limit m_κ as the minimum of a functional defined on \mathcal{S} for $1 < \kappa \leq 2$ (we will analyze the case $\kappa > 2$ in a forthcoming paper). Indeed the derivation of this functional relies on the asymptotic behavior of the Green-St Venant's strain tensor for minimizing sequences. The limit centerline deformation \mathcal{V} is linked to the limit \mathbf{R} of the rotation fields via its derivatives and the centerline directions. For $1 < \kappa < 2$, the limit energy depends linearly on the two fields $(\mathcal{V}, \mathbf{R})$ where in particular \mathbf{R} takes its value in the convex hull of $SO(3)$. In the case $\kappa = 2$, the limit of the rotation \mathbf{R} takes its value in $SO(3)$ and the functional is quadratic with respect to its derivatives.

As general references for the theory of elasticity we refer to [1], [9]. The theory of rods is developed in e.g. [2] and [22]. In the framework of linear elasticity, we refer to [10] for the junction of a three dimensional domain and a two dimensional one, to [19] for the junction of two rods and to [20] for the junction of two plates. The junction between a rod and a plate in the linear case is investigated in [17] and [12] and in nonlinear elasticity in [18]. The decomposition of the displacements in thin structures has been introduced in [13] and in [14] and then used in [3], [4] and [5] for the homogenization of the junction of rods and a plate.

This paper is organized as follows. In Section 2 we specify the geometry of the structure. Section 3 is devoted to introduce the elementary deformation associated to a deformation and to establish estimates. A Korn's type inequality for the structure is

derived in Section 4. In Section 5 the usual rescaling of each rod is recalled. The limit behavior of the Green-St Venant's strain tensor is analyzed in Section 6 for sequences (v_δ) such that $\|\text{dist}(\nabla_x v_\delta, SO(3))\|_{L^2(\mathcal{S}_\delta)} = O(\delta^K)$ for $1 < K \leq 2$. The assumptions on the elastic energy are introduced in Section 7 together with the scaling on the applied forces. The characterization of m_κ is performed in Section 8 for $1 < \kappa < 2$ and in Section 9 for $\kappa = 2$. At the end, an appendix contains a few technical results. The results of the present paper have been announced in [8].

2 Geometry and notations.

2.1 The rod structure.

The Euclidian space \mathbb{R}^3 is related to the frame $(O; \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3)$. We denote by $\|\cdot\|_2$ the euclidian norm and by \cdot the scalar product in \mathbb{R}^3 .

For an integer $N \geq 1$ and $i \in \{1, \dots, N\}$, let γ_i be a segment parametrized by s_i , with direction the unit vector \mathbf{t}_i , origin the point $P_i \in \mathbb{R}^3$ and length L_i . We have $\gamma_i = \varphi_i([0, L_i])$ where

$$\varphi_i(s_i) = P_i + s_i \mathbf{t}_i, \quad s_i \in \mathbb{R}.$$

So, the running point of γ_i is $\varphi_i(s_i)$, $0 \leq s_i \leq L_i$. The extremities of the segments make up a set denoted Γ .

For any $i \in \{1, \dots, N\}$, we choose a unit vector \mathbf{n}_i normal to \mathbf{t}_i and we set

$$\mathbf{b}_i = \mathbf{t}_i \wedge \mathbf{n}_i.$$

The structure-skeleton \mathcal{S} is the set $\bigcup_{i=1}^N \gamma_i$. The common points to two segments are called knots and the set of knots is denoted \mathcal{K} . For any γ_i , $i \in \{1, \dots, N\}$, we denote by a_i^j the arc-length of the knots belonging to γ_i ,

$$0 \leq a_i^1 < a_i^2 < \dots < a_i^{K_i} \leq L_i.$$

Among all the knots, $\Gamma_{\mathcal{K}}$ is the set of those which are extremities of all segments containing them.

Geometrical hypothesis. *We assume the following hypotheses on \mathcal{S} :*

- \mathcal{S} is connected,
- $\forall (i, j) \in \{1, \dots, N\}^2$

$$i \neq j \quad \implies \quad \gamma_i \cap \gamma_j = \emptyset \quad \text{or} \quad \gamma_i \cap \gamma_j \text{ is reduced to one knot.}$$

We denote by ω the unit disc of center the origin and we set $\omega_\delta = \delta\omega$ for $\delta > 0$. The reference cylinder of length L_i and cross-section ω_δ is denoted $\Omega_{i,\delta} =]0, L_i[\times \omega_\delta$.

Definition 2.1. For all $i \in \{1, \dots, N\}$, the straight rod $\mathcal{P}_{i,\delta}$ is the cylinder of center line γ_i and reference cross-section ω_δ . We have $\mathcal{P}_{i,\delta} = \Phi_i(\Omega_{i,\delta})$ where

$$\Phi_i(s) = \varphi_i(s_i) + y_2 \mathbf{n}_i + y_3 \mathbf{b}_i = P_i + s_i \mathbf{t}_i + y_2 \mathbf{n}_i + y_3 \mathbf{b}_i, \quad s = (s_i, y_2, y_3) \in \mathbb{R}^3.$$

The whole structure \mathcal{S}_δ is

$$\mathcal{S}_\delta = \left(\bigcup_{i=1}^N \mathcal{P}_{i,\delta} \right) \cup \left(\bigcup_{A \in \Gamma_{\mathcal{K}}} B(A; \delta) \right).$$

Loosely speaking, in the definition of \mathcal{S}_δ a ball of radius δ is added at each knot which belongs to $\Gamma_{\mathcal{K}}$. Remark that for a knot $A \notin \Gamma_{\mathcal{K}}$, the ball $B(A; \delta)$ is at least included in one $\mathcal{P}_{i,\delta}$.

There exists $\delta_0 > 0$ such that for $0 < \delta \leq \delta_0$ and for all $(i, j) \in \{1, \dots, N\}^2$, we have

$$\mathcal{P}_{i,\delta} \cap \mathcal{P}_{j,\delta} = \emptyset \quad \text{if and only if} \quad \gamma_i \cap \gamma_j = \emptyset.$$

The reference domain associated to the straight rod $\mathcal{P}_{i,\delta}$ is the open set $\Omega_i =]0, L_i[\times \omega$ (recall that ω is the disc of center the origin and radius 1). The running point of Ω_i (resp. $\Omega_{i,\delta}$, \mathcal{S}_δ) is denoted (s_i, Y_3, Y_3) (resp. (s_i, y_2, y_3) , x).

2.2 The junctions.

In what follows we deal with portions of the rod $\mathcal{P}_{i,\delta}$. For any $h > 0$ we set

$$\mathcal{P}_{i,\delta}^{a,h} = \Phi_i(\Omega_{i,\delta}^{a,h}) \cap \mathcal{S}_\delta \quad \text{where} \quad \Omega_{i,\delta}^{a,h} =]a - h\delta, a + h\delta[\times \omega_\delta, \quad 0 \leq a \leq L_i.$$

Let A be a knot, for $h > 0$ we consider the open set

$$\mathcal{J}_{A,h\delta} = \bigcup_{i \in \{1, \dots, N\}, A \in \gamma_i} \mathcal{P}_{i,\delta}^{a_i^k, h} \quad A = \varphi_i(a_i^k) \quad \text{if} \quad A \in \gamma_i. \quad (2.1)$$

Up to choosing δ_0 smaller, it is clear that there exists a real number

$$1 \leq \rho_0 \leq \frac{1}{4\delta_0} \min_{(A,B) \in \mathcal{K}^2, A \neq B} \|\overrightarrow{AB}\|_2$$

depending on \mathcal{S} (via the angles between the segments of the skeleton \mathcal{S}) such that for all $0 < \delta \leq \delta_0$

- $\mathcal{J}_{A,(\rho_0+1)\delta} \cap \mathcal{J}_{B,(\rho_0+1)\delta} = \emptyset$ for all distinct knots A and B ,
- the set $\mathcal{S}_\delta \setminus \bigcup_{A \in \mathcal{K}} \mathcal{J}_{A,\rho_0\delta}$ is made by disjointed cylinders.

The junction in the neighborhood of A is the domain $\mathcal{J}_{A,\rho_0\delta}$.

Notice that for any knot A and for any $0 < \delta \leq \delta_0$

$$\begin{aligned} & \text{the domain } \mathcal{J}_{A,(\rho_0+1)\delta} \text{ is star-shaped with respect to the ball } B(A; \delta) \\ & \text{and has a diameter less than } (2\rho_0 + 5)\delta. \end{aligned} \quad (2.2)$$

2.3 The functional spaces.

For $q \in [1, +\infty]$, the L^q -class fields of \mathcal{S} is the product space

$$L^q(\mathcal{S}; \mathbb{R}^p) = \prod_{i=1}^N L^q(0, L_i; \mathbb{R}^p)$$

equipped with the norm

$$\begin{aligned} \|V\|_{L^q(\mathcal{S}; \mathbb{R}^p)} &= \left(\sum_{i=1}^N \|V_i\|_{L^q(0, L_i; \mathbb{R}^p)}^q \right)^{1/q}, \quad V = (V_1, \dots, V_N), \quad q \in [1, +\infty[, \\ \|V\|_{L^\infty(\mathcal{S}; \mathbb{R}^p)} &= \max_{i=1, \dots, N} \|V_i\|_{L^\infty(0, L_i; \mathbb{R}^p)}. \end{aligned}$$

The $W^{1,q}$ -class fields of \mathcal{S} make up a space denoted

$$\begin{aligned} W^{1,q}(\mathcal{S}; \mathbb{R}^p) &= \left\{ V \in \prod_{i=1}^N W^{1,q}(0, L_i; \mathbb{R}^p) \mid V = (V_1, \dots, V_N), \text{ such that} \right. \\ &\quad \text{for any } A \in \mathcal{K}, \text{ if } A \in \gamma_i \cap \gamma_j, \text{ with } A = \varphi_i(a_i^k) = \varphi_j(a_j^l), \\ &\quad \left. \text{then one has } V_i(a_i^k) = V_j(a_j^l) \right\}. \end{aligned}$$

The common value $V_i(a_i^k)$ is denoted $V(A)$. We equip $W^{1,q}(\mathcal{S}; \mathbb{R}^p)$ with the norm

$$\begin{aligned} \|V\|_{W^{1,q}(\mathcal{S}; \mathbb{R}^p)} &= \left(\sum_{i=1}^N \|V_i\|_{W^{1,q}(0, L_i; \mathbb{R}^p)}^q \right)^{1/q}, \quad q \in [1, +\infty[, \\ \|V\|_{W^{1,\infty}(\mathcal{S}; \mathbb{R}^p)} &= \max_{i=1, \dots, N} \|V_i\|_{W^{1,\infty}(0, L_i; \mathbb{R}^p)}. \end{aligned}$$

Indeed, the parametrization $\varphi = (\varphi_1, \dots, \varphi_N)$ of \mathcal{S} belongs to $W^{1,\infty}(\mathcal{S}; \mathbb{R}^3)$.

We also denote by $L^q(\mathcal{S}; SO(3))$ (respectively $W^{1,q}(\mathcal{S}; SO(3))$) the set of matrix fields \mathbf{R} in $L^q(\mathcal{S}; \mathbb{R}^{3 \times 3})$ (resp. $W^{1,q}(\mathcal{S}; \mathbb{R}^{3 \times 3})$) satisfying $\mathbf{R}(s_i) \in SO(3)$ for almost any $s_i \in]0, L_i[$, $i \in \{1, \dots, N\}$.

3 An approximation theorem.

3.1 Definition of elementary rod-structure deformations.

We recall that for $x \in \mathcal{P}_{i,\delta}$ we have $x = \Phi_i(s)$ where $s \in \Omega_{i,\delta}$.

Definition 3.1. *An elementary rod-structure deformation is a deformation v_e verifying in each rod $\mathcal{P}_{i,\delta}$ ($i \in \{1, \dots, N\}$) and each junction $\mathcal{J}_{A,\rho_0\delta}$*

$$\begin{aligned} v_e(s) &= \mathcal{V}_i(s_i) + \mathbf{R}_i(s_i)(y_2 \mathbf{n}_i + y_3 \mathbf{b}_i) \quad s = (s_i, y_2, y_3) \in \Omega_{i,\delta} \\ v_e(x) &= \mathcal{V}(A) + \mathbf{R}(A)(x - A) \quad x \in \mathcal{J}_{A,\rho_0\delta} \end{aligned} \tag{3.1}$$

where $\mathcal{V} \in H^1(\mathcal{S}; \mathbb{R}^3)$ and $\mathbf{R} \in H^1(\mathcal{S}; SO(3))$ are such that $v_e \in H^1(\mathcal{S}_\delta; \mathbb{R}^3)$.

Recall that in the above definition $\mathcal{V}(A)$ (respectively $\mathbf{R}(A)$) denote the common value of \mathcal{V}_i (resp. \mathbf{R}_i) at the knot A . Let us notice that in view of Definition 3.1 and of $\mathcal{J}_{A,\rho_0\delta}$ one has

$$\begin{aligned} v_e(s) &= \mathcal{V}(A) + (s_i - a_i^k)\mathbf{R}(A)\mathbf{t}_i + \mathbf{R}(A)(y_2\mathbf{n}_i + y_3\mathbf{b}_i), \\ s &= (s_i, y_2, y_3) \in \mathcal{J}_{A,\rho_0\delta} \cap \Omega_{i,\delta}, \quad A = \varphi_i(a_i^k), \\ \mathcal{V}_i(s_i) &= \mathcal{V}(A) + (s_i - a_i^k)\mathbf{R}(A)\mathbf{t}_i, \quad \mathbf{R}_i(s_i) = \mathbf{R}(A), \\ s_i &\in]a_i^k - \rho_0\delta, a_i^k + \rho_0\delta[\cap]0, L_i]. \end{aligned} \tag{3.2}$$

The field \mathcal{V}_i stands for the deformation of the line γ_i while $\mathbf{R}_i(s_i)$ represents the rotation of the cross section with arc length s_i on γ_i . Then Definition 3.1 impose to an elementary deformation to be a translation-rotation in each junction.

3.2 Decomposition of the deformation in each rod $\mathcal{P}_{i,\delta}$.

According to [6], we first recall that any deformation $v \in H^1(\mathcal{P}_{i,\delta}; \mathbb{R}^3)$ can be decomposed as

$$v(s) = \mathcal{V}'_i(s_i) + \mathbf{R}'_i(s_i)(y_2\mathbf{n}_i + y_3\mathbf{b}_i) + \bar{v}'_i(s), \quad s = (s_i, y_2, y_3) \in \Omega_{i,\delta}, \tag{3.3}$$

where \mathcal{V}'_i belongs to $H^1(0, L_i; \mathbb{R}^3)$, \mathbf{R}'_i belongs to $H^1(0, L_i; \mathbb{R}^{3 \times 3})$ and satisfies for any $s_i \in [0, L_i]$: $\mathbf{R}'_i(s_i) \in SO(3)$ and \bar{v}'_i belongs to $H^1(\mathcal{P}_{i,\delta}; \mathbb{R}^3)$ (or $H^1(\Omega_{i,\delta}; \mathbb{R}^3)$ using again the same convention as for v). The term \mathcal{V}'_i gives the deformation of the center line of the rod. The second term $\mathbf{R}'_i(s_i)(y_2\mathbf{n}_i + y_3\mathbf{b}_i)$ describes the rotation of the cross section (of the rod) which contains the point $\varphi_i(s_i)$. The part $\mathcal{V}'_i(s_i) + \mathbf{R}'_i(s_i)(y_2\mathbf{n}_i + y_3\mathbf{b}_i)$ of the decomposition of v is an elementary deformation of the rod $\mathcal{P}_{i,\delta}$. Let us notice that there is no reason for $\mathcal{V}' = (\mathcal{V}'_1, \dots, \mathcal{V}'_N)$ to belong to $H^1(\mathcal{S}; \mathbb{R}^3)$ or for $\mathbf{R}' = (\mathbf{R}'_1, \dots, \mathbf{R}'_N)$ to belong to $H^1(\mathcal{S}; \mathbb{R}^{3 \times 3})$. This is why, in order to define an elementary deformation of the whole structure \mathcal{S}_δ , we will modify the fields \mathcal{V}'_i and \mathbf{R}'_i near to the junction $\mathcal{J}_{A,(\rho_0+1)\delta}$ at each knot A . In view of (3.5), we choose the field $\mathbf{R}_A(x - A) + \mathbf{a}_A$ as the elementary deformation in $\mathcal{J}_{A,\rho_0\delta}$.

The following theorem is proved in [6]:

Theorem 3.2. *Let $v \in H^1(\mathcal{P}_{i,\delta}; \mathbb{R}^3)$, there exists a decomposition (3.3) such that*

$$\begin{aligned} \|\bar{v}'_i\|_{L^2(\Omega_{i,\delta}; \mathbb{R}^3)} &\leq C\delta \| \text{dist}(\nabla_x v, SO(3)) \|_{L^2(\mathcal{P}_{i,\delta})}, \\ \|\nabla_s \bar{v}'_i\|_{L^2(\Omega_{i,\delta}; \mathbb{R}^{3 \times 3})} &\leq C \| \text{dist}(\nabla_x v, SO(3)) \|_{L^2(\mathcal{P}_{i,\delta})}, \\ \left\| \frac{d\mathbf{R}'_i}{ds_i} \right\|_{L^2(0, L_i; \mathbb{R}^{3 \times 3})} &\leq \frac{C}{\delta^2} \| \text{dist}(\nabla_x v, SO(3)) \|_{L^2(\mathcal{P}_{i,\delta})}, \\ \left\| \frac{d\mathcal{V}'_i}{ds_i} - \mathbf{R}'_i \mathbf{t}_i \right\|_{L^2(0, L_i; \mathbb{R}^3)} &\leq \frac{C}{\delta} \| \text{dist}(\nabla_x v, SO(3)) \|_{L^2(\mathcal{P}_{i,\delta})}, \\ \|\nabla_x v - \mathbf{R}'_i\|_{L^2(\Omega_{i,\delta}; \mathbb{R}^{3 \times 3})} &\leq C \| \text{dist}(\nabla_x v, SO(3)) \|_{L^2(\mathcal{P}_{i,\delta})}, \end{aligned} \tag{3.4}$$

where the constant C does not depend on δ and L_i .

3.3 Decomposition of the deformation in each junction $\mathcal{J}_{A,\rho_0\delta}$.

Let v be a deformation belonging to $H^1(\mathcal{S}_\delta; \mathbb{R}^3)$ and let A be a knot. We apply the rigidity theorem of [11], formulated in the version of Theorem II.1.1 in [6] which is licit because of (2.2), to the domain $\mathcal{J}_{A,(\rho_0+1)\delta}$. Hence there exist $\mathbf{R}_A \in SO(3)$ and $\mathbf{a}_A \in \mathbb{R}^3$ such that

$$\begin{aligned} \|\nabla_x v - \mathbf{R}_A\|_{L^2(\mathcal{J}_{A,(\rho_0+1)\delta}; \mathbb{R}^{3 \times 3})} &\leq C \|\text{dist}(\nabla v, SO(3))\|_{L^2(\mathcal{J}_{A,(\rho_0+1)\delta})}, \\ \|v - \mathbf{a}_A - \mathbf{R}_A(x - A)\|_{L^2(\mathcal{J}_{A,(\rho_0+1)\delta}; \mathbb{R}^3)} &\leq C\delta \|\text{dist}(\nabla v, SO(3))\|_{L^2(\mathcal{J}_{A,(\rho_0+1)\delta})}, \end{aligned} \quad (3.5)$$

with a constant C which does not depend on δ .

3.4 The approximation theorem.

In Theorem 3.3 we show that any deformation in $H^1(\mathcal{S}_\delta; \mathbb{R}^3)$ can be approximated by an elementary rod-structure deformation $v_e \in H^1(\mathcal{S}_\delta; \mathbb{R}^3)$ of the type given in Definition 3.1

Theorem 3.3. *Let v be a deformation in $H^1(\mathcal{S}_\delta; \mathbb{R}^3)$. There exists an elementary rod-structure deformation $v_e \in H^1(\mathcal{S}_\delta; \mathbb{R}^3)$ in the sense of Definition 3.1 such that if we set $\bar{v} = v - v_e$*

$$\begin{aligned} \|\bar{v}\|_{L^2(\mathcal{S}_\delta; \mathbb{R}^3)} &\leq C\delta \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)}, \\ \|\nabla_x \bar{v}\|_{L^2(\mathcal{S}_\delta; \mathbb{R}^{3 \times 3})} &\leq C \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)}. \end{aligned} \quad (3.6)$$

Moreover the fields \mathcal{V} and \mathbf{R} associated to v_e satisfy

$$\begin{aligned} \sum_{i=1}^N \left\| \frac{d\mathbf{R}_i}{ds_i} \right\|_{L^2(0, L_i; \mathbb{R}^{3 \times 3})} &\leq \frac{C}{\delta^2} \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)}, \\ \sum_{i=1}^N \left\| \frac{d\mathcal{V}_i}{ds_i} - \mathbf{R}_i \mathbf{t}_i \right\|_{L^2(0, L_i; \mathbb{R}^3)} &\leq \frac{C}{\delta} \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)}, \\ \sum_{i=1}^N \|\nabla_x v - \mathbf{R}_i\|_{L^2(\Omega_{i,\delta}; \mathbb{R}^{3 \times 3})} &\leq C \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)}. \end{aligned} \quad (3.7)$$

In all the above estimates, the constant C is independent on δ and on the lengths L_i ($i \in \{1, \dots, N\}$).

Proof. Step 1. In this step we compare the two decompositions of v given in Subsections 3.1 and 3.2 in $\mathcal{J}_{A,(\rho_0+1)\delta} \cap \mathcal{P}_{i,\delta}$.

Let $A = \varphi_i(a_i^k)$ be a knot in $\mathcal{P}_{i,\delta}$. Using the last estimate of (3.4) and the first one in (3.5) we obtain

$$\begin{aligned} \|\mathbf{R}_A - \mathbf{R}'_i\|_{L^2([a_i^k - (\rho_0+1)\delta, a_i^k + (\rho_0+1)\delta] \cap]0, L_i[; \mathbb{R}^{3 \times 3})} &\leq \frac{C}{\delta} (\|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{P}_{i,\delta})} \\ &\quad + \|\text{dist}(\nabla v, SO(3))\|_{L^2(\mathcal{J}_{A,(\rho_0+1)\delta})}). \end{aligned} \quad (3.8)$$

Now the decomposition (3.3), the first estimate in (3.4) and the last estimate (3.5) lead to

$$\begin{aligned} & \|\mathcal{V}'_i + \mathbf{R}'_i(y_2\mathbf{n}_i + y_3\mathbf{b}_i) - \mathbf{R}_A((s_i - a_i^k)\mathbf{t}_i + y_2\mathbf{n}_i + y_3\mathbf{b}_i) - \mathbf{a}_A\|_{L^2(\Omega_{i,a_i^k,(\rho_0+1)\delta} \cap \Omega_{i,\delta}; \mathbb{R}^3)} \\ & \leq C\delta(\|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{P}_{i,\delta})} + \|\text{dist}(\nabla v, SO(3))\|_{L^2(\mathcal{J}_{A,(\rho_0+1)\delta})}) \end{aligned}$$

which in turn with (3.8) gives

$$\begin{aligned} & \|\mathcal{V}'_i - \mathbf{a}_A - (s_i - a_i^k)\mathbf{R}_A\mathbf{t}_i\|_{L^2([a_i^k - (\rho_0+1)\delta, a_i^k + (\rho_0+1)\delta] \cap]0, L_i[; \mathbb{R}^3)} \\ & \leq C(\|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{P}_{i,\delta})} + \|\text{dist}(\nabla v, SO(3))\|_{L^2(\mathcal{J}_{A,(\rho_0+1)\delta})}). \end{aligned} \quad (3.9)$$

Step 2. Here we construct v_e .

We first define a rotation field $\mathbf{R} \in H^1(\mathcal{S}; SO(3))$ which is constant and equal to \mathbf{R}_A given by (3.5) in the adequate neighborhood of the knot A and which is close to \mathbf{R}'_i on each segment γ_i . Let $i \in \{1, \dots, N\}$ be fixed. For all $k \in \{1, \dots, K_i\}$ (i.e. all the knots of the line γ_i) we first set

$$\mathbf{R}_i(s_i) = \mathbf{R}_A \quad A = \varphi_i(a_i^k) \in \mathcal{K} \cap \gamma_i, \quad s_i \in]a_i^k - \rho_0\delta, a_i^k + \rho_0\delta[, \quad (3.10)$$

and then "far from the knots"

$$\mathbf{R}_i(s_i) = \mathbf{R}'_i(s_i) \quad s_i \in]0, L_i[\setminus \bigcup_{k=1}^{K_i}]a_i^k - (\rho_0 + 1)\delta, a_i^k + (\rho_0 + 1)\delta[. \quad (3.11)$$

It remains to define \mathbf{R}_i in the intervals

$$I_+^k =]0, L_i[\cap]a_i^k + \rho_0\delta, a_i^k + (\rho_0 + 1)\delta[\quad \text{and} \quad I_-^k =]0, L_i[\cap]a_i^k - (\rho_0 + 1)\delta, a_i^k - \rho_0\delta[, \quad k \in \{1, \dots, K_i\}.$$

In these intervals, we proceed as in Appendix of [6] in order to construct the field \mathbf{R}_i by a sort of interpolation in $SO(3)$, from \mathbf{R}_A to $\mathbf{R}'_i(a_i^k + (\rho_0 + 1)\delta)$ and from \mathbf{R}_A to $\mathbf{R}'_i(a_i^k - (\rho_0 + 1)\delta)$. This construction satisfies

$$\begin{aligned} \left\| \frac{d\mathbf{R}_i}{ds_i} \right\|_{L^2(I_+^k)^{3 \times 3}}^2 & \leq \frac{C}{\delta} \|\mathbf{R}'_i(a_i^k + (\rho_0 + 1)\delta) - \mathbf{R}_A\|^2, \\ \left\| \frac{d\mathbf{R}_i}{ds_i} \right\|_{L^2(I_-^k)^{3 \times 3}}^2 & \leq \frac{C}{\delta} \|\mathbf{R}_A - \mathbf{R}'_i(a_i^k - (\rho_0 + 1)\delta)\|^2. \end{aligned} \quad (3.12)$$

The constant does not depend on δ . From the above construction we obtain a field $\mathbf{R} = (\mathbf{R}_1, \dots, \mathbf{R}_N)$ defined on \mathcal{S} and which belongs to $H^1(\mathcal{S}; SO(3))$.

Let us now define the field $\mathcal{V} \in H^1(\mathcal{S}; \mathbb{R}^3)$ of v_e . We proceed as for \mathbf{R} . Let $i \in \{1, \dots, N\}$ be fixed. For all $k \in \{1, \dots, K_i\}$ we first set

$$\begin{aligned} \mathcal{V}_i(s_i) = \mathbf{a}_A + (s_i - a_i^k)\mathbf{R}_A\mathbf{t}_i \quad & A = \varphi_i(a_i^k) \in \mathcal{K} \cap \gamma_i, \\ \text{for } s_i \in]a_i^k - \rho_0\delta, a_i^k + \rho_0\delta[, & \end{aligned} \quad (3.13)$$

while "far from the knots" we set

$$\mathcal{V}_i(s_i) = \mathcal{V}'_i(s_i) \quad s_i \in]0, L_i[\setminus \bigcup_{k=1}^{K_i}]a_i^k - (\rho_0 + 1)\delta, a_i^k + (\rho_0 + 1)\delta[. \quad (3.14)$$

At least in the remaining intervals I_+^k and I_-^k , we just perform a linear interpolation

$$\begin{aligned} \mathcal{V}_i(s_i) &= \frac{a_i^k + \rho_0\delta - s_i}{\delta} \mathcal{V}'_i(s_i) + \left(1 - \frac{a_i^k + \rho_0\delta - s_i}{\delta}\right) \left(\mathbf{a}_A + (s_i - a_i^k) \mathbf{R}_A \mathbf{t}_i\right) \\ &\quad \text{for } s_i \in I_+^k, \\ \mathcal{V}_i(s_i) &= \frac{a_i^k - \rho_0\delta - s_i}{\delta} \mathcal{V}'_i(s_i) + \left(1 - \frac{a_i^k - \rho_0\delta - s_i}{\delta}\right) \left(\mathbf{a}_A + (s_i - a_i^k) \mathbf{R}_A \mathbf{t}_i\right) \\ &\quad \text{for } s_i \in I_-^k, \quad k \in \{1, \dots, K_i\}. \end{aligned} \quad (3.15)$$

Gathering (3.13), (3.14) and (3.15), we obtain a field $\mathcal{V} = (\mathcal{V}_1, \dots, \mathcal{V}_N)$ defined on \mathcal{S} which belongs to $H^1(\mathcal{S}; \mathbb{R}^3)$. It worth noting that $(\mathcal{V}, \mathbf{R})$ verify the condition (3.2). As a consequence we can define the elementary deformation v_e associated to \mathcal{V} and \mathbf{R} through Definition 3.1 and it belongs to $H^1(\mathcal{S}_\delta; \mathbb{R}^3)$.

Step 3. Comparison between $(\mathcal{V}_i, \mathbf{R}_i)$ and $(\mathcal{V}'_i, \mathbf{R}'_i)$.

Using the third estimate in (3.4), (3.8) and (3.12) we obtain

$$\begin{aligned} &\|\mathbf{R}_i - \mathbf{R}'_i\|_{L^2(I_+^k \cup I_-^k; \mathbb{R}^{3 \times 3})} + \delta \left\| \frac{d\mathbf{R}_i}{ds_i} \right\|_{L^2(I_+^k \cup I_-^k; \mathbb{R}^{3 \times 3})} \\ &\leq \frac{C}{\delta} (\|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{P}_{i,\delta})} + \|\text{dist}(\nabla v, SO(3))\|_{L^2(\mathcal{J}_{A,(\rho_0+1)\delta})}). \end{aligned}$$

Taking into account the definition of \mathbf{R}_i , for all $i \in \{1, \dots, N\}$ we finally get

$$\begin{aligned} &\|\mathbf{R}_i - \mathbf{R}'_i\|_{L^2(0, L_i; \mathbb{R}^{3 \times 3})} + \delta \left\| \frac{d\mathbf{R}_i}{ds_i} \right\|_{L^2(0, L_i; \mathbb{R}^{3 \times 3})} \\ &\leq \frac{C}{\delta} (\|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{P}_{i,\delta})} + \sum_{A \in \gamma_i} \|\text{dist}(\nabla v, SO(3))\|_{L^2(\mathcal{J}_{A,(\rho_0+1)\delta})}). \end{aligned} \quad (3.16)$$

The constant does not depend on δ and L_i .

From (3.9) and the definition of \mathcal{V}_i (see (3.13), (3.14) and (3.15)) we deduce that

$$\begin{aligned} \|\mathcal{V}'_i - \mathcal{V}_i\|_{L^2(0, L_i; \mathbb{R}^3)} &\leq C (\|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{P}_{i,\delta})} \\ &\quad + \sum_{A \in \gamma_i} \|\text{dist}(\nabla v, SO(3))\|_{L^2(\mathcal{J}_{A,(\rho_0+1)\delta})}). \end{aligned} \quad (3.17)$$

Now, taking into account the fourth estimate in (3.4), estimates (3.8)-(3.9) and again the definition of \mathcal{V}_i we get

$$\begin{aligned} \left\| \frac{d\mathcal{V}_i}{ds_i} - \frac{d\mathcal{V}'_i}{ds_i} \right\|_{L^2(0, L_i; \mathbb{R}^3)} &\leq \frac{C}{\delta} (\|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{P}_{i,\delta})} \\ &\quad + \sum_{A \in \gamma_i} \|\text{dist}(\nabla v, SO(3))\|_{L^2(\mathcal{J}_{A,(\rho_0+1)\delta})}). \end{aligned} \quad (3.18)$$

where the constant C does not depend on δ and L_i .

Step 4. First of all, we prove the estimates (3.7). The first one in (3.7) is a direct consequence of (3.16) and the fact that the matrices \mathbf{R}_i are constants in the neighborhood of the knots (intervals I_{\pm}^k). The second one comes from the fourth estimate in (3.4), again (3.16) and (3.18). Then, from (3.4) and (3.16) we deduce the last estimate in (3.7).

Now, let us set $\bar{v} = v - v_e$ where v_e is defined in Step 2. From the decomposition (3.3) and the expression of v_e in Definition 3.1, we have for all $i \in \{1, \dots, N\}$

$$\begin{aligned} v(s) &= \mathcal{V}'_i(s_i) + \mathbf{R}'_i(s_i)(y_2 \mathbf{n}_i + y_3 \mathbf{b}_i) + \bar{v}'_i(s), \\ v(s) &= \mathcal{V}_i(s_i) + \mathbf{R}_i(s_i)(y_2 \mathbf{n}_i + y_3 \mathbf{b}_i) + \bar{v}_i(s) = v_{e,i}(s) + \bar{v}_i(s), \end{aligned} \quad s = (s_i, y_2, y_3) \in \Omega_{i,\delta}.$$

Then, using (3.16) and (3.17) it leads to

$$\begin{aligned} \|\bar{v}_i - \bar{v}'_i\|_{L^2(\Omega_{i,\delta}; \mathbb{R}^3)} &\leq C\delta (\|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{P}_{i,\delta})} \\ &\quad + \sum_{A \in \gamma_i} \|\text{dist}(\nabla v, SO(3))\|_{L^2(\mathcal{J}_{A,(\rho_0+1)\delta})}). \end{aligned}$$

Hence, due to the first estimate in (3.4) and the above inequalities we get

$$\sum_{i=1}^N \|\bar{v}_i\|_{L^2(\Omega_{i,\delta}; \mathbb{R}^3)} \leq C\delta (\|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_{i\delta})}).$$

Now, in order to take into account the knots in $\Gamma_{\mathcal{K}}$, we use (3.5) and the definition of v_e in the junction $\mathcal{J}_{A,\rho_0\delta}$ ((3.10) and (3.13)) to obtain

$$\|v - v_e\|_{L^2(B(A;\delta); \mathbb{R}^3)} \leq \|v - v_e\|_{L^2(\mathcal{J}_{A,\rho_0\delta}; \mathbb{R}^3)} \leq C\delta \|\text{dist}(\nabla v, SO(3))\|_{L^2(\mathcal{J}_{A,(\rho_0+1)\delta})}.$$

Finally due to the definition of \mathcal{S}_{δ} we deduce the first estimate in (3.6).

An easy calculation gives in $\Omega_{i,\delta}$, $i = 1, \dots, N$

$$\nabla_x v_e \mathbf{t}_i = \frac{d\mathcal{V}_i}{ds_i} + \frac{d\mathbf{R}_i}{ds_i}(y_2 \mathbf{n}_i + y_3 \mathbf{b}_i), \quad \nabla_x v_e \mathbf{n}_i = \mathbf{R}_i \mathbf{n}_i, \quad \nabla_x v_e \mathbf{b}_i = \mathbf{R}_i \mathbf{b}_i. \quad (3.19)$$

Then, from the first estimate in (3.5) and all those in (3.7) we obtain the estimate of $\nabla \bar{v}$. \square

4 A Korn's type inequality for the rod-structure.

This section is devoted to derive a nonlinear Korn's type inequality for \mathcal{S}_{δ} . We assume that the structure \mathcal{S}_{δ} is clamped on a few extremities whose set is denoted by Γ_0^{δ} , corresponding to a set Γ_0 of extremities of \mathcal{S} . Then we set

$$\mathbb{D}_{\delta} = \{v \in H^1(\mathcal{S}_{\delta}; \mathbb{R}^3) \mid v = I_d \text{ on } \Gamma_0^{\delta}\}.$$

Let v be in \mathbb{D}_δ . Proceeding as in [6] for each rod, we can choose v_e in Theorem 3.3 such that

$$v_e = I_d \quad \text{on} \quad \Gamma_0^\delta. \quad (4.1)$$

Notice that (4.1) implies that

$$\bar{v} = 0 \quad \text{on} \quad \Gamma_0^\delta, \quad \mathcal{V} = \phi, \quad \mathbf{R}_i = \mathbf{I}_3 \quad \text{on} \quad \Gamma_0 \quad \text{for} \quad i = 1, \dots, N. \quad (4.2)$$

Theorem 4.1. *There exists a constant C which depends only on \mathcal{S} such that, for all deformation $v \in \mathbb{D}_\delta$*

$$\|v - I_d\|_{H^1(\mathcal{S}_\delta; \mathbb{R}^3)} \leq \frac{C}{\delta} \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)}. \quad (4.3)$$

Moreover

$$\begin{aligned} \sum_{A \in \mathcal{K}} \|\mathcal{V}(A) - A\|_2 &\leq \frac{C}{\delta^2} \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)}, \\ \sum_{A \in \mathcal{K}} \|\mathbf{R}(A) - \mathbf{I}_3\| &\leq \frac{C}{\delta^2} \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)}. \end{aligned} \quad (4.4)$$

Proof. Taking into account the connexity of \mathcal{S} and the continuous character of \mathbf{R} , the boundary condition 4.2 implies that

$$\sum_{i=1}^N \|\mathbf{R}_i - \mathbf{I}_3\|_{L^2(0, L_i; \mathbb{R}^{3 \times 3})} \leq C \sum_{i=1}^N \left\| \frac{d\mathbf{R}_i}{ds_i} \right\|_{L^2(0, L_i; \mathbb{R}^{3 \times 3})}.$$

Then, the first estimate in (3.7) gives

$$\sum_{i=1}^N \|\mathbf{R}_i - \mathbf{I}_3\|_{L^2(0, L_i; \mathbb{R}^{3 \times 3})} \leq \frac{C}{\delta^2} \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)}. \quad (4.5)$$

From (4.5), the last estimate in (3.7), the boundary condition on v together with Poincaré inequality we deduce that 4.3 holds true. Now estimate (4.5) and the second one in (3.7) imply that

$$\sum_{i=1}^N \left\| \frac{d\mathcal{V}_i}{ds_i} - \mathbf{t}_i \right\|_{L^2(0, L_i; \mathbb{R}^3)} \leq \frac{C}{\delta^2} \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)}. \quad (4.6)$$

Hence the boundary condition (4.2) leads to

$$\|\mathcal{V} - \varphi\|_{H^1(\mathcal{S}; \mathbb{R}^3)} \leq \frac{C}{\delta^2} \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)} \quad (4.7)$$

from which the first estimate in (4.4) follows. At last the last estimate in (4.4) comes from (3.7) and the boundary condition on \mathbf{R} . The constants depend only on the skeleton \mathcal{S} . \square

Remark 4.2. Since \mathbf{R} is bounded, we also have (for $i = 1, \dots, N$)

$$\|\mathbf{R}_i - \mathbf{I}_3\|_{L^2(0, L_i; \mathbb{R}^{3 \times 3})} \leq C. \quad (4.8)$$

From the above estimate and proceeding as in the proof of Theorem 4.1 we obtain

$$\|v - I_d\|_{H^1(\mathcal{S}_\delta; \mathbb{R}^3)} \leq C \left\{ \delta + \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)} \right\}, \quad (4.9)$$

and

$$\begin{aligned} \sum_{A \in \mathcal{K}} \|\mathcal{V}(A) - A\|_2 &\leq C \left\{ 1 + \frac{1}{\delta} \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)} \right\}, \\ \sum_{A \in \mathcal{K}} \|\mathbf{R}(A) - \mathbf{I}_3\| &\leq C. \end{aligned} \quad (4.10)$$

Remark 4.3. From (3.17) and estimate (II.3.9) in [6] and the fact that $(v - I_d) - (\mathcal{V}_i - \varphi_i) = (\mathbf{R}_i - \mathbf{I}_3)(y_2 \mathbf{n}_i + y_3 \mathbf{b}_i) + \bar{v}_i$ in each rod $\mathcal{P}_{i, \delta}$, we obtain

$$\|(v - I_d) - (\mathcal{V}_i - \varphi_i)\|_{L^2(\mathcal{P}_{i, \delta}; \mathbb{R}^3)} \leq C \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{P}_{i, \delta})}.$$

Due to (4.5) and the estimate (II.3.10) in [6], we also have

$$\begin{aligned} \|\nabla_x v + (\nabla_x v)^T - 2\mathbf{I}_3\|_{L^2(\mathcal{S}_\delta; \mathbb{R}^{3 \times 3})} &\leq C \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)} \\ &\quad + \frac{C}{\delta^3} \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)}^2, \end{aligned} \quad (4.11)$$

which in turn with (3.7) give

$$\begin{aligned} \sum_{i=1}^N \|\mathbf{R}_i^T + \mathbf{R}_i - 2\mathbf{I}_3\|_{L^2(0, L_i; \mathbb{R}^{3 \times 3})} &\leq \frac{C}{\delta} \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)} \\ &\quad + \frac{C}{\delta^4} \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)}^2. \end{aligned} \quad (4.12)$$

5 Rescaling of $\Omega_{i, \delta}$ for $i = 1, \dots, N$.

We recall that $\Omega_i =]0, L_i[\times \omega$. We rescale $\Omega_{i, \delta}$ using the operator

$$\Pi_{i, \delta}(\psi)(s_i, Y_2, Y_3) = \psi(s_i, \delta Y_2, \delta Y_3) \quad \text{for a.e. } (s_i, Y_2, Y_3) \in \Omega_i$$

defined for any measurable function ψ over $\Omega_{i, \delta}$. Indeed, if $\psi \in L^2(\Omega_{i, \delta})$ then $\Pi_{i, \delta}(\psi) \in L^2(\Omega_i)$. The estimates (3.6) of \bar{v} transposed over Ω_i are

$$\begin{aligned} \|\Pi_{i, \delta}(\bar{v})\|_{L^2(\Omega_i; \mathbb{R}^3)} &\leq C \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)}, \\ \left\| \frac{\partial \Pi_{i, \delta}(\bar{v})}{\partial Y_2} \right\|_{L^2(\Omega_i; \mathbb{R}^3)} &\leq C \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)}, \\ \left\| \frac{\partial \Pi_{i, \delta}(\bar{v})}{\partial Y_3} \right\|_{L^2(\Omega_i; \mathbb{R}^3)} &\leq C \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)}, \\ \left\| \frac{\partial \Pi_{i, \delta}(\bar{v})}{\partial s_i} \right\|_{L^2(\Omega_i; \mathbb{R}^3)} &\leq \frac{C}{\delta} \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)}. \end{aligned} \quad (5.1)$$

6 Asymptotic behavior of the Green-St Venant's strain tensor.

We distinguish three main cases for the behavior of the quantity $\|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)}$ which plays an important role in the derivation of estimates in nonlinear elasticity

$$\|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)} = \begin{cases} O(\delta^\kappa), & 1 \leq \kappa < 2, \\ O(\delta^2), & \\ O(\delta^\kappa), & \kappa > 2. \end{cases}$$

This hierarchy of behavior for $\|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{P}_\delta)}$ has already been observed in [6].

In this section we investigate the behavior of a sequence $(v_\delta)_\delta$ of \mathbb{D}_δ for which $\|\text{dist}(\nabla_x v_\delta, SO(3))\|_{L^2(\mathcal{S}_\delta)} = O(\delta^\kappa)$ for $1 < \kappa \leq 2$. In Subsection 6.1 we analyse the case $1 < \kappa < 2$ and Subsection 6.2 deals with the case $\kappa = 2$. Let us emphasize that we explicit the limit of the Green-St Venant's tensor in each case.

6.1 Case $1 < \kappa < 2$. Limit behavior for a sequence such that $\|\mathbf{dist}(\nabla_x v_\delta, SO(3))\|_{L^2(\mathcal{S}_\delta)} \sim \delta^\kappa$.

Let us consider a sequence of deformations $(v_\delta)_\delta$ of \mathbb{D}_δ such that for $1 < \kappa < 2$

$$\|\text{dist}(\nabla_x v_\delta, SO(3))\|_{L^2(\mathcal{S}_\delta)} \leq C\delta^\kappa.$$

We denote by \mathcal{V}_δ , \mathbf{R}_δ and \bar{v}_δ the three terms of the decomposition of v_δ given by Theorem 3.3. The two first estimates of (3.7) and those in (5.1) lead to the following lemma:

Lemma 6.1. *There exists a subsequence still indexed by δ such that*

$$\begin{aligned} \mathbf{R}_\delta &\rightharpoonup \mathbf{R} \quad \text{weakly-}^* \text{ in } L^\infty(\mathcal{S}; \mathbb{R}^{3 \times 3}), \\ \frac{1}{\delta^{\kappa-2}} \mathbf{R}_\delta &\rightharpoonup 0 \quad \text{weakly in } H^1(\mathcal{S}; \mathbb{R}^{3 \times 3}), \\ \mathcal{V}_\delta &\rightharpoonup \mathcal{V} \quad \text{weakly in } H^1(\mathcal{S}; \mathbb{R}^3), \\ \frac{1}{\delta^{\kappa-1}} \mathbf{R}_{i,\delta}^T \left(\frac{d\mathcal{V}_{i,\delta}}{ds_i} - \mathbf{R}_{i,\delta} \mathbf{t}_i \right) &\rightharpoonup \mathcal{Z}_i \quad \text{weakly in } L^2(0, L_i; \mathbb{R}^3), \\ \frac{1}{\delta^\kappa} \Pi_{i,\delta} (\mathbf{R}_{i,\delta}^T \bar{v}_\delta) &\rightharpoonup \bar{w}_i \quad \text{weakly in } L^2(0, L_i; H^1(\omega; \mathbb{R}^3)). \end{aligned} \tag{6.1}$$

Moreover the matrix $\mathbf{R}_i(s_i)$ belongs to the convex hull of $SO(3)$ for almost any $s_i \in]0, L_i[$, $i \in \{1, \dots, N\}$, $\mathcal{V} \in W^{1,\infty}(\mathcal{S}; \mathbb{R}^3)$ and they satisfy

$$\mathcal{V} = \phi \quad \text{on} \quad \Gamma_0 \quad \text{and} \quad \frac{d\mathcal{V}_i}{ds_i} = \mathbf{R}_i \mathbf{t}_i \quad i \in \{1, \dots, N\}. \tag{6.2}$$

Furthermore, we also have

$$\begin{aligned}\Pi_{i,\delta}(v_\delta) &\rightharpoonup \mathcal{V}_i \quad \text{weakly in } H^1(\Omega_i; \mathbb{R}^3), \\ \Pi_{i,\delta}(\nabla_x v_\delta) &\rightharpoonup \mathbf{R}_i \quad \text{weakly in } L^2(\Omega_i; \mathbb{R}^{3 \times 3}).\end{aligned}$$

Now we proceed as in [6] to derive the limit of the Green-St Venant tensor as δ goes to 0. We first recall that for any function $\psi \in H^1(\Omega_{i,\delta})$

$$\frac{\partial \psi}{\partial s_i} = \nabla_x \psi \cdot \mathbf{t}_i \quad \frac{\partial \psi}{\partial y_2} = \nabla_x \psi \cdot \mathbf{n}_i \quad \frac{\partial \psi}{\partial y_3} = \nabla_x \psi \cdot \mathbf{b}_i.$$

Then in view of Lemma 6.1 and of the above relation, we obtain

$$\begin{aligned}\frac{1}{\delta^{\kappa-1}} \mathbf{R}_{i,\delta}^T (\Pi_{i,\delta}(\nabla_x v_\delta) - \mathbf{R}_{i,\delta}) \mathbf{t}_i &\rightharpoonup \mathcal{Z}_i \quad \text{weakly in } L^2(\Omega_i; \mathbb{R}^3), \\ \frac{1}{\delta^{\kappa-1}} \mathbf{R}_{i,\delta}^T (\Pi_{i,\delta}(\nabla_x v_\delta) - \mathbf{R}_{i,\delta}) \mathbf{n}_i &\rightharpoonup \frac{\partial \bar{w}_i}{\partial Y_2} \quad \text{weakly in } L^2(\Omega_i; \mathbb{R}^3), \\ \frac{1}{\delta^{\kappa-1}} \mathbf{R}_{i,\delta}^T (\Pi_{i,\delta}(\nabla_x v_\delta) - \mathbf{R}_{i,\delta}) \mathbf{b}_i &\rightharpoonup \frac{\partial \bar{w}_i}{\partial Y_3} \quad \text{weakly in } L^2(\Omega_i; \mathbb{R}^3),\end{aligned}\tag{6.3}$$

The weak convergences in (6.3) together with the relation $(\nabla_x v_\delta)^T \nabla_x v_\delta - \mathbf{I}_3 = (\nabla_x v_\delta - \mathbf{R}_{i,\delta})^T \mathbf{R}_{i,\delta} + (\mathbf{R}_{i,\delta})^T (\nabla_x v_\delta - \mathbf{R}_{i,\delta}) + (\nabla_x v_\delta - \mathbf{R}_{i,\delta})^T (\nabla_x v_\delta - \mathbf{R}_{i,\delta})$ permit to obtain the limit of the Green-St Venant's tensor in the rescaled domain Ω_i . We obtain

$$\frac{1}{2\delta^{\kappa-1}} \Pi_{i,\delta}((\nabla_x v_\delta)^T \nabla_x v_\delta - \mathbf{I}_3) \rightharpoonup \mathbf{E}_i \quad \text{weakly in } L^1(\Omega_i; \mathbb{R}^{3 \times 3}),\tag{6.4}$$

with

$$\mathbf{E}_i = \frac{1}{2} \left\{ (\mathbf{t}_i \mid \mathbf{n}_i \mid \mathbf{b}_i) \left(\mathcal{Z}_i \mid \frac{\partial \bar{w}_i}{\partial Y_2} \mid \frac{\partial \bar{w}_i}{\partial Y_3} \right)^T + \left(\mathcal{Z}_i \mid \frac{\partial \bar{w}_i}{\partial Y_2} \mid \frac{\partial \bar{w}_i}{\partial Y_3} \right) (\mathbf{t}_i \mid \mathbf{n}_i \mid \mathbf{b}_i)^T \right\}.\tag{6.5}$$

The term $(\mathbf{a} \mid \mathbf{b} \mid \mathbf{c})$ denotes the 3×3 matrix with columns \mathbf{a} , \mathbf{b} and \mathbf{c} . For any field $\bar{\psi} \in L^2(0, L_i; H^1(\omega; \mathbb{R}^3))$ we set

$$e_{22}(\bar{\psi}) = \frac{\partial \bar{\psi}}{\partial Y_2} \cdot \mathbf{n}_i, \quad e_{23}(\bar{\psi}) = \frac{1}{2} \left\{ \frac{\partial \bar{\psi}}{\partial Y_2} \cdot \mathbf{b}_i + \frac{\partial \bar{\psi}}{\partial Y_3} \cdot \mathbf{n}_i \right\}, \quad e_{33}(\bar{\psi}) = \frac{\partial \bar{\psi}}{\partial Y_3} \cdot \mathbf{b}_i.\tag{6.6}$$

Hence we can write \mathbf{E}_i as

$$\mathbf{E}_i = (\mathbf{t}_i \mid \mathbf{n}_i \mid \mathbf{b}_i) \widehat{\mathbf{E}}_i (\mathbf{t}_i \mid \mathbf{n}_i \mid \mathbf{b}_i)^T$$

where the symmetric matrix $\widehat{\mathbf{E}}_i$ is defined by

$$\widehat{\mathbf{E}}_i = \begin{pmatrix} \mathcal{Z}_i \cdot \mathbf{t}_i & * & * \\ \frac{1}{2} \frac{\partial \bar{w}_i}{\partial Y_2} \cdot \mathbf{t}_i & e_{22}(\bar{w}_i) & * \\ \frac{1}{2} \frac{\partial \bar{w}_i}{\partial Y_3} \cdot \mathbf{t}_i & e_{23}(\bar{w}_i) & e_{33}(\bar{w}_i) \end{pmatrix}\tag{6.7}$$

and where the field $\bar{w}_i \in L^2(0, L_i; H^1(\omega; \mathbb{R}^3))$ is defined by

$$\bar{w}_i = [Y_2(\mathcal{Z}_i \cdot \mathbf{n}_i) + Y_3(\mathcal{Z}_i \cdot \mathbf{b}_i)] \mathbf{t}_i + \bar{w}_i, \quad i \in \{1, \dots, N\}.\tag{6.8}$$

6.2 Case $\kappa = 2$. Limit behavior for a sequence such that $\|\mathbf{dist}(\nabla_x v_\delta, SO(3))\|_{L^2(\mathcal{S}_\delta)} \sim \delta^2$.

Let us consider a sequence of deformations $(v_\delta)_\delta$ of \mathbb{D}_δ such that

$$\|\mathbf{dist}(\nabla_x v_\delta, SO(3))\|_{L^2(\mathcal{S}_\delta)} \leq C\delta^2.$$

Indeed, in each rod $\mathcal{P}_{i,\delta}$ we get $\|\mathbf{dist}(\nabla_x v_\delta, SO(3))\|_{L^2(\mathcal{P}_{i,\delta})} \leq C\delta^2$. We denote by \mathcal{V}_δ , \mathbf{R}_δ and \bar{v}_δ the three terms of the decomposition of v_δ given by Theorem 3.3. The estimates (3.7), (3.6) and (4.7) lead to the following lemma:

Lemma 6.2. *There exists a subsequence still indexed by δ such that*

$$\begin{aligned} \mathbf{R}_\delta &\rightharpoonup \mathbf{R} \quad \text{weakly in } H^1(\mathcal{S}; SO(3)) \text{ and strongly in } L^\infty(\mathcal{S}; SO(3)), \\ \mathcal{V}_\delta &\longrightarrow \mathcal{V} \quad \text{strongly in } H^1(\mathcal{S}; \mathbb{R}^3), \\ \frac{1}{\delta} \left(\frac{d\mathcal{V}_{i,\delta}}{ds_i} - \mathbf{R}_{i,\delta} \mathbf{t}_i \right) &\rightharpoonup \mathcal{Z}_i \quad \text{weakly in } L^2(0, L_i; \mathbb{R}^3), \\ \frac{1}{\delta^2} \Pi_{i,\delta}(\bar{v}_\delta) &\rightharpoonup \bar{v}_i \quad \text{weakly in } L^2(0, L_i; H^1(\omega; \mathbb{R}^3)). \end{aligned} \tag{6.9}$$

Moreover $\mathcal{V}_i \in H^2(0, L_i; \mathbb{R}^3)$ for all $i \in \{1, \dots, N\}$ and we have

$$\mathcal{V} = \phi, \quad \mathbf{R}_i = \mathbf{I}_3, \quad \text{on } \Gamma_0 \quad \text{and} \quad \frac{d\mathcal{V}_i}{ds_i} = \mathbf{R}_i \mathbf{t}_i \quad i \in \{1, \dots, N\}. \tag{6.10}$$

Furthermore, we also have

$$\begin{aligned} \Pi_{i,\delta}(v_\delta) &\longrightarrow \mathcal{V}_i \quad \text{strongly in } H^1(\Omega_i; \mathbb{R}^3), \\ \Pi_{i,\delta}(\nabla_x v_\delta) &\longrightarrow \mathbf{R}_i \quad \text{strongly in } L^2(\Omega_i; \mathbb{R}^{3 \times 3}). \end{aligned} \tag{6.11}$$

As a consequence of the above lemma we obtain

$$\begin{aligned} \frac{1}{\delta} \left(\Pi_{i,\delta}(\nabla_x v_\delta) - \mathbf{R}_{i,\delta} \right) \mathbf{t}_i &\rightharpoonup \frac{d\mathbf{R}_i}{ds_i} (Y_2 \mathbf{n}_i + Y_3 \mathbf{b}_i) + \mathcal{Z}_i \quad \text{weakly in } L^2(\Omega_i; \mathbb{R}^3), \\ \frac{1}{\delta} \left(\Pi_{i,\delta}(\nabla_x v_\delta) - \mathbf{R}_{i,\delta} \right) \mathbf{n}_i &\rightharpoonup \frac{\partial \bar{v}_i}{\partial Y_2} \quad \text{weakly in } L^2(\Omega_i; \mathbb{R}^3), \\ \frac{1}{\delta} \left(\Pi_{i,\delta}(\nabla_x v_\delta) - \mathbf{R}_{i,\delta} \right) \mathbf{b}_i &\rightharpoonup \frac{\partial \bar{v}_i}{\partial Y_3} \quad \text{weakly in } L^2(\Omega_i; \mathbb{R}^3). \end{aligned} \tag{6.12}$$

Proceeding as in Subsection 6.1 and using convergences (6.9) and (6.12) permit to obtain the limit of the Green-St Venant's tensor in the rescaled domain Ω_i

$$\frac{1}{2\delta} \Pi_{i,\delta} \left((\nabla_x v_\delta)^T \nabla_x v_\delta - \mathbf{I}_3 \right) \rightharpoonup \mathbf{E}_i \quad \text{weakly in } L^1(\Omega_i; \mathbb{R}^{3 \times 3}), \tag{6.13}$$

with

$$\begin{aligned} \mathbf{E}_i &= \frac{1}{2} \left\{ (\mathbf{t}_i \mid \mathbf{n}_i \mid \mathbf{b}_i) \left(\frac{d\mathbf{R}_i}{ds_i} (Y_2 \mathbf{n}_i + Y_3 \mathbf{b}_i) + \mathcal{Z}_i \mid \frac{\partial \bar{v}_i}{\partial Y_2} \mid \frac{\partial \bar{v}_i}{\partial Y_3} \right)^T \mathbf{R}_i \right. \\ &\quad \left. + \mathbf{R}_i^T \left(\frac{d\mathbf{R}_i}{ds_i} (Y_2 \mathbf{n}_i + Y_3 \mathbf{b}_i) + \mathcal{Z}_i \mid \frac{\partial \bar{v}_i}{\partial Y_2} \mid \frac{\partial \bar{v}_i}{\partial Y_3} \right) (\mathbf{t}_i \mid \mathbf{n}_i \mid \mathbf{b}_i)^T \right\}. \end{aligned} \tag{6.14}$$

Setting

$$\bar{u}_i = [Y_2(\mathcal{Z}_i \cdot \mathbf{R}_i \mathbf{n}_i) + Y_3(\mathcal{Z}_i \cdot \mathbf{R}_i \mathbf{b}_i)] \mathbf{t}_i + \mathbf{R}_i^T \bar{v}_i, \quad i \in \{1, \dots, N\}. \quad (6.15)$$

and using the fact that the matrix $\mathbf{R}_i^T \frac{d\mathbf{R}_i}{ds_3}$ is antisymmetric, we can write \mathbf{E}_i as

$$\mathbf{E}_i = (\mathbf{t}_i | \mathbf{n}_i | \mathbf{b}_i) \widehat{\mathbf{E}}_i (\mathbf{t}_i | \mathbf{n}_i | \mathbf{b}_i)^T,$$

where the symmetric matrix $\widehat{\mathbf{E}}_i$ is defined by

$$\widehat{\mathbf{E}}_i = \begin{pmatrix} -Y_2 \Gamma_{i,3}(\mathbf{R}) + Y_3 \Gamma_{i,2}(\mathbf{R}) + \mathcal{Z}_i \cdot \mathbf{R}_i \mathbf{t}_i & * & * \\ -\frac{1}{2} Y_3 \Gamma_{i,1}(\mathbf{R}) + \frac{1}{2} \frac{\partial \bar{u}_i}{\partial Y_2} \cdot \mathbf{t}_i & e_{22}(\bar{u}_i) & * \\ \frac{1}{2} Y_2 \Gamma_{i,1}(\mathbf{R}) + \frac{1}{2} \frac{\partial \bar{u}_i}{\partial Y_3} \cdot \mathbf{t}_i & e_{23}(\bar{u}_i) & e_{33}(\bar{u}_i) \end{pmatrix} \quad (6.16)$$

where

$$\Gamma_{i,3}(\mathbf{R}) = \frac{d\mathbf{R}_i}{ds_i} \mathbf{t}_i \cdot \mathbf{R}_i \mathbf{n}_i, \quad \Gamma_{i,2}(\mathbf{R}) = \frac{d\mathbf{R}_i}{ds_i} \mathbf{b}_i \cdot \mathbf{R}_i \mathbf{t}_i, \quad \Gamma_{i,1}(\mathbf{R}) = \frac{d\mathbf{R}_i}{ds_i} \mathbf{n}_i \cdot \mathbf{R}_i \mathbf{b}_i, \quad (6.17)$$

and where the $e_{kl}(\bar{u}_i)$'s are given by (6.6).

7 Elastic structure

In this section we assume that the structure \mathcal{S}_δ is made of an elastic material. The associated local energy $W : \mathbf{S}_3 \rightarrow \mathbb{R}^+$ is a continuous function of symmetric matrices which satisfies the following assumptions

$$\begin{aligned} \exists c > 0 \quad \text{such that} \quad \forall E \in \mathbf{S}_3 \quad W(E) &\geq c \| \|E\| \|^2, \\ \forall \varepsilon > 0, \quad \exists \theta > 0, \quad \text{such that} & \\ \forall E \in \mathbf{S}_3 \quad \| \|E\| \leq \theta \implies |W(E) - Q(E)| &\leq \varepsilon \| \|E\| \|^2, \end{aligned} \quad (7.1)$$

where Q is a positive quadratic form defined on the set of 3×3 symmetric matrices (see e.g. [7]). Remark that Q satisfies the first inequality in (7.1) with the same constant c .

Following [9], for any 3×3 matrix F , we set

$$\widehat{W}(F) = \begin{cases} W\left(\frac{1}{2}(F^T F - \mathbf{I}_3)\right) & \text{if } \det(F) > 0, \\ +\infty & \text{if } \det(F) \leq 0. \end{cases} \quad (7.2)$$

Remark that due to (7.1) and to the inequality $\| \|F^T F - \mathbf{I}_3\| \| \geq \text{dist}(F, SO(3))$ if $\det(F) > 0$, we have for any 3×3 matrix F

$$\widehat{W}(F) \geq \frac{c}{4} \text{dist}(F, SO(3))^2. \quad (7.3)$$

Remark 7.1. A classical example of local elastic energy satisfying the above assumptions is given by the St Venant-Kirchhoff's density (see [9])

$$\widehat{W}(F) = \begin{cases} \frac{\lambda}{8}(\text{tr}(F^T F - \mathbf{I}_3))^2 + \frac{\mu}{4}\text{tr}((F^T F - \mathbf{I}_3)^2) & \text{if } \det(F) > 0, \\ +\infty & \text{if } \det(F) \leq 0. \end{cases} \quad (7.4)$$

The coefficients λ and μ are the Lamé's constants. In this case we have for all matrix $E \in \mathbf{S}_3$

$$Q(E) = \frac{\lambda}{2}(\text{tr}(E))^2 + \mu \text{tr}(E^2).$$

Now we assume that the structure \mathcal{S}_δ is submitted to applied body forces $f_{\kappa,\delta} \in L^2(\mathcal{S}_\delta; \mathbb{R}^3)$ and we define the total energy $J_{\kappa,\delta}(v)$ ¹ over \mathbb{D}_δ by

$$J_{\kappa,\delta}(v) = \int_{\mathcal{S}_\delta} \widehat{W}(\nabla_x v)(x) dx - \int_{\mathcal{S}_\delta} f_{\kappa,\delta}(x) \cdot (v(x) - I_d(x)) dx. \quad (7.5)$$

Assumptions on the forces. We set

$$\kappa' = \begin{cases} 2\kappa - 2 & \text{if } 1 \leq \kappa \leq 2, \\ \kappa & \text{if } \kappa \geq 2. \end{cases} \quad (7.6)$$

In what follows we define the forces applied to the structure by distinguish the forces applied to the junctions and to their complementary in \mathcal{S}_δ . Moreover in order to take into account the resultant and the moment of the forces near each knot and in each cross section of the rods we decompose the forces density into two types. The first one mainly works with the mean deformation \mathcal{V} while the second one is related to the rotation \mathbf{R} . We begin by the definition of the forces in the junctions.

For any knot A , let F_A and G_A be two fields belonging to $L^2(\mathcal{J}_{A,\rho_0}; \mathbb{R}^3)^2$, the second field G_A satisfying $\int_{\mathcal{J}_{A,\rho_0}} G_A(z) dz = 0$.

We define the applied forces in the junction $\mathcal{J}_{A,\rho_0\delta}$ ($A \in \mathcal{K}$) by

$$f_{\kappa,\delta}(x) = \delta^{\kappa'-1} F_A\left(A + \frac{x-A}{\delta}\right) + \delta^{\kappa'-2} G_A\left(A + \frac{x-A}{\delta}\right), \quad \text{for a.e. } x \in \mathcal{J}_{A,\rho_0\delta}. \quad (7.7)$$

In order to precise the forces $f_{\delta,\kappa}$ in the complementary of the junctions (i.e. in $\mathcal{S}_\delta \setminus \bigcup_{A \in \mathcal{K}} \mathcal{J}_{A,\rho_0\delta}$), we follow [6] and we assume that there exist f , $g^{(\mathbf{n})}$ and $g^{(\mathbf{b})}$ in $L^2(\mathcal{S}; \mathbb{R}^3)$ such that

$$f_{\kappa,\delta}(x) = \delta^{\kappa'} f_i(s_i) + \delta^{\kappa'-2} (y_2 g_i^{(\mathbf{n})}(s_i) + y_3 g_i^{(\mathbf{b})}(s_i)), \quad x = \Phi_i(s) \quad (7.8)$$

for a.e. $s \in \left(]0, L_i[\setminus \bigcup_{k=1}^{K_i}]a_i^k - \rho_0\delta, a_i^k + \rho_0\delta[\right) \times \omega_\delta$

¹For later convenience, we have added the term $\int_{\mathcal{S}_\delta} f_{\kappa,\delta}(x) \cdot I_d(x) dx$ to the usual standard energy, indeed this does not affect the minimizing problem for $J_{\kappa,\delta}$.

²The domain \mathcal{J}_{A,ρ_0} is obtained by transforming $\mathcal{J}_{A,\rho_0\delta}$ by a dilation of center A and ratio $1/\delta$.

Notice that $J_{\kappa,\delta}(I_d) = 0$. So, in order to minimize $J_{\kappa,\delta}$ we only need to consider deformations v of \mathbf{D}_δ such that $J_{\kappa,\delta}(v) \leq 0$. Now from the decomposition given in Theorem 3.3, and the definition (7.7)-(7.8) of the forces we first get

$$\begin{aligned} \left| \int_{\mathcal{S}_\delta} f_{\kappa,\delta}(x) \cdot (v(x) - I_d(x)) dx \right| &\leq C\delta^{\kappa'+2} \left[\|f\|_{L^2(\mathcal{S};\mathbb{R}^3)} \|\mathcal{V} - \phi\|_{L^2(\mathcal{S};\mathbb{R}^3)} \right. \\ &\quad \left. + (\|g^{(\mathbf{n})}\|_{L^2(\mathcal{S};\mathbb{R}^3)} + \|g^{(\mathbf{b})}\|_{L^2(\mathcal{S};\mathbb{R}^3)}) \left(\sum_{i=1}^N \|\mathbf{R}_i - \mathbf{I}_3\|_{L^2(0,L_i;\mathbb{R}^{3\times 3})} \right) \right] \\ &+ \sum_{A \in \mathcal{K}} \|F_A\|_{L^2(\mathcal{J}_{A,\rho_0};\mathbb{R}^3)} \|\mathcal{V}(A) - A\|_2 + \sum_{A \in \mathcal{K}} \|G_A\|_{L^2(\mathcal{J}_{A,\rho_0};\mathbb{R}^3)} \|\mathbf{R}(A) - \mathbf{I}_3\|. \end{aligned} \quad (7.9)$$

In the case $\kappa \geq 2$, using (4.4), (4.5) and (4.7) gives

$$\left| \int_{\mathcal{S}_\delta} f_{\kappa,\delta}(x) \cdot (v(x) - I_d(x)) dx \right| \leq C\delta^\kappa \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)} \quad (7.10)$$

while in the case $1 \leq \kappa \leq 2$, (4.8) and (4.10) lead to

$$\left| \int_{\mathcal{S}_\delta} f_{\kappa,\delta}(x) \cdot (v(x) - I_d(x)) dx \right| \leq C\delta^{2\kappa} \left\{ 1 + \frac{1}{\delta} \|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)} \right\} \quad (7.11)$$

where in both cases the constant C depends on the L^2 -norms of f , $g^{(\mathbf{n})}$, $g^{(\mathbf{b})}$, F_A and G_A (and are then independent upon δ). For a deformation v such that $J_{\kappa,\delta}(v) \leq 0$, the coerciveness assumption (7.3) and the estimates (7.10) and (7.11) allows us to obtain for $1 \leq \kappa$

$$\|\text{dist}(\nabla_x v, SO(3))\|_{L^2(\mathcal{S}_\delta)} \leq C\delta^\kappa. \quad (7.12)$$

Again estimates (7.10) and (7.11) and (7.12) lead to

$$c\delta^{2\kappa} \leq J_{\kappa,\delta}(v) \leq 0. \quad (7.13)$$

with a constant independent on δ .

Then, from (7.1)-(7.2)-(7.3) and the estimates (7.10), (7.11) and (7.12) we deduce

$$\frac{c}{4} \|(\nabla_x v)^T \nabla_x v - \mathbf{I}_3\|_{L^2(\mathcal{S}_\delta;\mathbb{R}^{3\times 3})}^2 \leq J_{\kappa,\delta}(v) + \int_{\mathcal{S}_\delta} f_{\kappa,\delta} \cdot (v - I_d) \leq C\delta^{2\kappa}.$$

Hence, the following estimate of the Green-St Venant's tensor hold true:

$$\left\| \frac{1}{2} \{(\nabla_x v)^T \nabla_x v - \mathbf{I}_3\} \right\|_{L^2(\mathcal{S}_\delta;\mathbb{R}^{3\times 3})} \leq C\delta^\kappa.$$

We deduce from the above inequality that $v \in (W^{1,4}(\mathcal{S}_\delta))^3$ with

$$\|\nabla_x v\|_{L^4(\mathcal{S}_\delta;\mathbb{R}^{3\times 3})} \leq C\delta^{\frac{1}{2}}.$$

We set

$$m_{\kappa,\delta} = \inf_{v \in \mathbb{D}_\delta} J_{\kappa,\delta}(v)$$

and we recall that, in general, a minimizer of $J_{\kappa,\delta}$ does not exist on \mathbb{D}_δ .

8 Asymptotic behavior of $m_{\kappa,\delta}$ for $1 < \kappa < 2$.

The goal of this section is to establish Theorem 8.1 below. Let us first introduce a few notations. We denote by \mathcal{C} the convex hull of the set $SO(3)$

$$\mathcal{C} = \overline{\text{conv}}(SO(3)). \quad (8.1)$$

We set

$$\mathbb{V}\mathbb{R}_1 = \left\{ (\mathcal{V}, \mathbf{R}) \in H^1(\mathcal{S}; \mathbb{R}^3) \times L^2(\mathcal{S}; \mathcal{C}) \mid \mathcal{V} = \phi \quad \text{on} \quad \Gamma_0, \right. \\ \left. \frac{d\mathcal{V}_i}{ds_i} = \mathbf{R}_i \mathbf{t}_i \quad i \in \{1, \dots, N\} \right\}. \quad (8.2)$$

We define the linear functional \mathcal{L} over $H^1(\mathcal{S}; \mathbb{R}^3) \times L^2(\mathcal{S}; \mathbb{R}^{3 \times 3})$ by

$$\mathcal{L}(\mathcal{V}, \mathbf{R}) = \sum_{i=1}^N \pi \int_0^{L_i} \left(f_i \cdot (\mathcal{V}_i - \phi_i) + \frac{g_i^{(\mathbf{n})}}{3} \cdot (\mathbf{R}_i - \mathbf{I}_3) \mathbf{n}_i + \frac{g_i^{(\mathbf{b})}}{3} \cdot (\mathbf{R}_i - \mathbf{I}_3) \mathbf{b}_i \right) ds_i \\ + \sum_{A \in \mathcal{K}} \left[\left(\int_{\mathcal{J}_{A, \rho_0}} F_A(y) dy \right) \cdot (\mathcal{V}(A) - \phi(A)) + \int_{\mathcal{J}_{A, \rho_0}} G_A(y) \cdot (\mathbf{R}(A) - \mathbf{I}_3)(y - A) dy \right]. \quad (8.3)$$

It is easy to prove that the infimum of $-\mathcal{L}$ on $\mathbb{V}\mathbb{R}_1$ is a minimum.

Theorem 8.1. *We have*

$$\lim_{\delta \rightarrow 0} \frac{m_{\kappa, \delta}}{\delta^{2\kappa}} = \min_{(\mathcal{V}, \mathbf{R}) \in \mathbb{V}\mathbb{R}_1} (-\mathcal{L}(\mathcal{V}, \mathbf{R})). \quad (8.4)$$

Proof. Step 1. In this step we show that

$$\min_{(\mathcal{V}, \mathbf{R}) \in \mathbb{V}\mathbb{R}_1} (-\mathcal{L}(\mathcal{V}, \mathbf{R})) \leq \liminf_{\delta \rightarrow 0} \frac{m_{\kappa, \delta}}{\delta^{2\kappa}}.$$

Let $(v_\delta)_\delta$ be a sequence of deformations belonging to \mathbb{D}_δ and such that

$$\lim_{\delta \rightarrow 0} \frac{J_{\kappa, \delta}(v_\delta)}{\delta^{2\kappa}} = \liminf_{\delta \rightarrow 0} \frac{m_{\kappa, \delta}}{\delta^{2\kappa}}. \quad (8.5)$$

We can always assume that $J_{\kappa, \delta}(v_\delta) \leq 0$. Then, from the estimates of the previous section we obtain

$$\|\text{dist}(\nabla v_\delta, SO(3))\|_{L^2(\mathcal{S}_\delta)} \leq C\delta^\kappa, \\ \frac{1}{2} \|\nabla v_\delta^T \nabla v_\delta - \mathbf{I}_3\|_{L^2(\mathcal{S}_\delta; \mathbb{R}^{3 \times 3})} \leq C\delta^\kappa, \\ \|\nabla v_\delta\|_{L^4(\mathcal{S}_\delta; \mathbb{R}^{3 \times 3})} \leq C\delta^{\frac{1}{2}}.$$

For any fixed δ , the deformation v_δ is decomposed as in Theorem 3.3 and we are in a position to apply the results of Subsection 6.1. There exists a subsequence still indexed

by δ such that

$$\begin{aligned}
\mathbf{R}_\delta &\rightharpoonup \mathbf{R}^{\{0\}} \quad \text{weakly-* in } L^\infty(\mathcal{S}; \mathbb{R}^{3 \times 3}), \\
\frac{1}{\delta^{\kappa-2}} \mathbf{R}_\delta &\rightharpoonup 0 \quad \text{weakly in } H^1(\mathcal{S}; \mathbb{R}^{3 \times 3}), \\
\mathcal{V}_\delta &\rightharpoonup \mathcal{V}^{\{0\}} \quad \text{weakly in } H^1(\mathcal{S}; \mathbb{R}^3), \\
\frac{1}{\delta^{\kappa-1}} \mathbf{R}_{i,\delta}^T \left(\frac{d\mathcal{V}_{i,\delta}}{ds_i} - \mathbf{R}_{i,\delta} \mathbf{t}_i \right) &\rightharpoonup \mathcal{Z}_i^{\{0\}} \quad \text{weakly in } L^2(0, L_i; \mathbb{R}^3), \\
\frac{1}{\delta^\kappa} \Pi_{i,\delta}(\mathbf{R}_{i,\delta}^T \bar{v}_\delta) &\rightharpoonup \bar{w}_i^{\{0\}} \quad \text{weakly in } L^2(0, L_i; H^1(\omega; \mathbb{R}^3)).
\end{aligned} \tag{8.6}$$

The couple $(\mathcal{V}^{\{0\}}, \mathbf{R}^{\{0\}})$ belongs to $\mathbb{V}\mathbb{R}_1$. Furthermore, we also have $(i \in \{1, \dots, N\})$

$$\begin{aligned}
\Pi_{i,\delta} v_\delta &\rightharpoonup \mathcal{V}_i^{\{0\}} \quad \text{weakly in } H^1(\Omega_i; \mathbb{R}^3), \\
\Pi_{i,\delta}(\nabla_x v_\delta) &\rightharpoonup \mathbf{R}_i^{\{0\}} \quad \text{weakly in } L^4(\Omega_i; \mathbb{R}^{3 \times 3}), \\
\frac{1}{2\delta^{\kappa-1}} \Pi_{i,\delta}((\nabla_x v_\delta)^T \nabla_x v_\delta - \mathbf{I}_3) &\rightharpoonup \mathbf{E}_i^{\{0\}} \quad \text{weakly in } L^2(\Omega_i; \mathbb{R}^{3 \times 3}),
\end{aligned} \tag{8.7}$$

where the symmetric matrix $\mathbf{E}_i^{\{0\}}$ is defined in (6.5) (see Subsection 6.1). Due to the decomposition of v_δ and the above convergences (8.6) we immediately get the limit of the term involving the body forces

$$\lim_{\delta \rightarrow 0} \frac{1}{\delta^{2\kappa}} \int_{\mathcal{S}_\delta} f_{\kappa,\delta} \cdot (v_\delta - I_d) = \mathcal{L}(\mathcal{V}^{\{0\}}, \mathbf{R}^{\{0\}})$$

where $\mathcal{L}(\mathcal{V}, \mathbf{R})$ is defined by (8.3).

Recall that we have $-\int_{\mathcal{S}_\delta} f_{\kappa,\delta} \cdot (v_\delta - I_d) \leq J_{\kappa,\delta}(v_\delta)$. Then, due to (8.5) and the above limit, we finally get

$$\min_{(\mathcal{V}, \mathbf{R}) \in \mathbb{V}\mathbb{R}_1} (-\mathcal{L}(\mathcal{V}, \mathbf{R})) \leq -\mathcal{L}(\mathcal{V}^{\{0\}}, \mathbf{R}^{\{0\}}) \leq \liminf_{\delta \rightarrow 0} \frac{m_{\kappa,\delta}}{\delta^{2\kappa}}. \tag{8.8}$$

Step 2. In this step we show that

$$\limsup_{\delta \rightarrow 0} \frac{m_{\kappa,\delta}}{\delta^{2\kappa}} \leq \min_{(\mathcal{V}, \mathbf{R}) \in \mathbb{V}\mathbb{R}_1} (-\mathcal{L}(\mathcal{V}, \mathbf{R})).$$

Let $(\mathcal{V}^{\{1\}}, \mathbf{R}^{\{1\}}) \in \mathbb{V}\mathbb{R}_1$ such that $-\mathcal{L}(\mathcal{V}^{\{1\}}, \mathbf{R}^{\{1\}}) = \min_{(\mathcal{V}, \mathbf{R}) \in \mathbb{V}\mathbb{R}_1} (-\mathcal{L}(\mathcal{V}, \mathbf{R}))$.

Using Proposition 10.4 in the Appendix, there exists a sequence $(\mathcal{V}^{(n)}, \mathbf{R}^{(n)})_{n \geq 0}$ in $\mathbb{V}\mathbb{R}_1$ which satisfies

- $\mathbf{R}^{(n)} \in W^{1,\infty}(\mathcal{S}; SO(3))$,
- $\mathbf{R}^{(n)}$ is equal to \mathbf{I}_3 in a neighbourhood of each knot $A \in \mathcal{K}$ and each fixed extremity belonging to Γ_0 ,

$$\begin{aligned}
\mathcal{V}^{(n)} &\rightharpoonup \mathcal{V}^{\{1\}} \quad \text{weakly in } H^1(\mathcal{S}; \mathbb{R}^3), \\
\mathbf{R}^{(n)} &\rightharpoonup \mathbf{R}^{\{1\}} \quad \text{weakly in } L^2(\mathcal{S}; \mathcal{C}).
\end{aligned} \tag{8.9}$$

Now we fix n . Since $(\mathcal{V}^{(n)}, \mathbf{R}^{(n)})$ in $\mathbb{V}\mathbb{R}_1$ and due to the second condition imposed on $\mathbf{R}^{(n)}$ above, we can consider the elementary deformation $v^{(n)}$ constructed by using $(\mathcal{V}^{(n)}, \mathbf{R}^{(n)})$ in Definition 3.1. Indeed the deformation $v^{(n)}$ belongs to $\mathbb{D}_\delta \cap W^{1,\infty}(\mathcal{S}_\delta; \mathbb{R}^3)$. Thanks to the expression of the gradient of $v^{(n)}$ (see (3.19)) and to the definition of $\mathbb{V}\mathbb{R}_1$ we have

$$\|\nabla_x v^{(n)} - \mathbf{R}_i^{(n)}\|_{L^\infty(\Omega_{i,\delta}; \mathbb{R}^{3 \times 3})} \leq C_n \delta.$$

Now using the identity $(\nabla_x v^{(n)})^T \nabla_x v^{(n)} - \mathbf{I}_3 = (\nabla_x v^{(n)} - \mathbf{R}_i^{(n)})^T \mathbf{R}_i^{(n)} + (\mathbf{R}_i^{(n)})^T (\nabla_x v^{(n)} - \mathbf{R}_i^{(n)}) + (\nabla_x v^{(n)} - \mathbf{R}_i^{(n)})^T (\nabla_x v^{(n)} - \mathbf{R}_i^{(n)})$ and the above estimate, we obtain

$$\frac{1}{2\delta^{\kappa-1}} \Pi_{i,\delta} \left((\nabla_x v^{(n)})^T \nabla_x v^{(n)} - \mathbf{I}_3 \right) \longrightarrow 0 \quad \text{strongly in } L^\infty(\Omega_i; \mathbb{R}^{3 \times 3}) \tag{8.10}$$

as $\delta \rightarrow 0$. Hence, if δ is small enough, we get $\det(\nabla_x v^{(n)}) > 0$ a.e. in \mathcal{S}_δ .

In what follows, we show that $\lim_{\delta \rightarrow 0} \frac{1}{\delta^{2\kappa}} J_{\kappa,\delta}(v^{(n)}) = -\mathcal{L}(\mathcal{V}^{(n)}, \mathbf{R}^{(n)})$. Let $\mathcal{P}_{i,\delta}$ be a rod of the structure. Using the third assumption in (7.1) (with $\varepsilon = 1$) and the estimate (8.10), for δ small enough we have

$$\frac{1}{\delta^{2\kappa}} \int_{\mathcal{P}_{i,\delta}} \widehat{W}(\nabla_x v^{(n)})(x) dx \leq \frac{1}{\delta^{2(\kappa-1)}} \int_{\Omega_i} (Q(E(v^{(n)})) + \|E(v^{(n)})\|^2) ds_i dY_2 dY_3$$

where

$$E(v^{(n)}) = \Pi_{i,\delta} \left((\nabla_x v^{(n)})^T \nabla_x v^{(n)} - \mathbf{I}_3 \right).$$

Thanks to the convergence (8.10) we obtain $\frac{1}{\delta^{2\kappa}} \int_{\mathcal{P}_{i,\delta}} \widehat{W}(\nabla_x v^{(n)})(x) dx \rightarrow 0$ as $\delta \rightarrow 0$.

Notice that since $v^{(n)}$ is an elementary deformation, its Green-St Venant's tensor is null in the neighbourhood of the knots. Finally we get

$$\frac{1}{\delta^{2\kappa}} \int_{\mathcal{S}_\delta} \widehat{W}(\nabla_x v^{(n)})(x) dx \longrightarrow 0 \quad \text{as } \delta \text{ tends to } 0.$$

Using again the fact that $v^{(n)}$ is an elementary deformation, and assumptions (7.7) and (7.8) on the forces, we immediately get the limit of the term involving the body forces

$$\lim_{\delta \rightarrow 0} \frac{1}{\delta^{2\kappa}} \int_{\mathcal{S}_\delta} f_{\kappa,\delta} \cdot (v^{(n)} - I_d) = \mathcal{L}(\mathcal{V}^{(n)}, \mathbf{R}^{(n)}).$$

Indeed, since $v^{(n)} \in \mathbb{D}_\delta$, we have

$$\frac{m_{\kappa,\delta}}{\delta^{2\kappa}} \leq \frac{J_{\kappa,\delta}(v^{(n)})}{\delta^{2\kappa}}.$$

Passing to the limit as δ tends to 0 we obtain

$$\limsup_{\delta \rightarrow 0} \frac{m_{\kappa, \delta}}{\delta^{2\kappa}} \leq -\mathcal{L}(\mathcal{V}^{(n)}, \mathbf{R}^{(n)}).$$

In view of the convergences (8.9) we are able to pass to the limit as n tends to infinity and we obtain

$$\limsup_{\delta \rightarrow 0} \frac{m_{\kappa, \delta}}{\delta^{2\kappa}} \leq -\mathcal{L}(\mathcal{V}^{\{1\}}, \mathbf{R}^{\{1\}}) = \min_{(\mathcal{V}, \mathbf{R}) \in \mathbb{V}\mathbb{R}_1} (-\mathcal{L}(\mathcal{V}, \mathbf{R})).$$

This concludes the proof of the theorem. \square

Remark 8.2. *Let us point out that Theorem 8.1 shows that for any minimizing sequence $(v_\delta)_\delta$ as in Step 1, the third convergence of the rescaled Green-St Venant's strain tensor in (8.7) is a strong convergence to 0 in $L^2(\Omega_i; \mathbb{R}^{3 \times 3})$.*

9 Asymptotic behavior of $m_{2, \delta}$.

The goal of this section is to establish Theorem 9.1 below. Let us first introduce a few notations. We set

$$\mathbb{V}\mathbb{R}_2 = \left\{ (\mathcal{V}, \mathbf{R}) \in H^1(\mathcal{S}; \mathbb{R}^3) \times H^1(\mathcal{S}; SO(3)) \mid \mathcal{V} = \phi, \mathbf{R}_i = \mathbf{I}_3 \quad \text{on} \quad \Gamma_0, \right. \\ \left. \frac{d\mathcal{V}_i}{ds_i} = \mathbf{R}_i \mathbf{t}_i \quad i \in \{1, \dots, N\} \right\}. \quad (9.1)$$

Theorem 9.1. *We have*

$$\lim_{\delta \rightarrow 0} \frac{m_{2, \delta}}{\delta^4} = \min_{(\mathcal{V}, \mathbf{R}) \in \mathbb{V}\mathbb{R}_2} \mathcal{J}_2(\mathcal{V}, \mathbf{R}), \quad (9.2)$$

where the functional \mathcal{J}_2 is defined by

$$\mathcal{J}_2(\mathcal{V}, \mathbf{R}) = \sum_{i=1}^N \int_0^{L_i} a_{lk} \Gamma_{i,k}(\mathbf{R}) \Gamma_{i,l}(\mathbf{R}) - \mathcal{L}(\mathcal{V}, \mathbf{R}). \quad (9.3)$$

In \mathcal{J}_2 , the 3×3 matrix $\mathbf{A} = (a_{ij})$ is symmetric and definite positive. This matrix depend on ω and on the quadratic form Q .

Note that the infimum of \mathcal{J}_2 on $\mathbb{V}\mathbb{R}_2$ is actually a minimum.

For a St-Venant- Kirchhoff material, whose energy is recalled in Remark 7.1, the expression of the matrix \mathbf{A} is explicitly derived at the end of the appendix (see Remark 10.7) and it leads to the following limit energy

$$\mathcal{J}_2(\mathcal{V}, \mathbf{R}) = \frac{\pi}{4} \sum_{i=1}^N \int_0^{L_i} (\mu |\Gamma_{i,1}(\mathbf{R})|^2 + E |\Gamma_{i,2}(\mathbf{R})|^2 + E |\Gamma_{i,3}(\mathbf{R})|^2) - \mathcal{L}(\mathcal{V}, \mathbf{R}) \quad (9.4)$$

where E and μ are respectively the Young and Poisson's coefficients.

Proof of Theorem 9.1. Step 1. In this step we show that

$$\min_{(\mathcal{V}, \mathbf{R}) \in \mathbb{V}\mathbb{R}_2} \mathcal{J}_2(\mathcal{V}, \mathbf{R}) \leq \liminf_{\delta \rightarrow 0} \frac{m_{2,\delta}}{\delta^4}.$$

Let $(v_\delta)_\delta$ be a sequence of deformations belonging to \mathbf{D}_δ and such that

$$\lim_{\delta \rightarrow 0} \frac{J_{2,\delta}(v_\delta)}{\delta^4} = \liminf_{\delta \rightarrow 0} \frac{m_{2,\delta}}{\delta^4}. \quad (9.5)$$

We can always assume that $J_{\kappa,\delta}(v_\delta) \leq 0$. Then, from the estimates of the Section 7 we obtain

$$\begin{aligned} \|\text{dist}(\nabla v_\delta, SO(3))\|_{L^2(\mathcal{S}_\delta)} &\leq C\delta^2, \\ \left\| \frac{1}{2} \{ \nabla v_\delta^T \nabla v_\delta - \mathbf{I}_3 \} \right\|_{L^2(\mathcal{S}_\delta; \mathbb{R}^{3 \times 3})} &\leq C\delta^2, \\ \|\nabla v_\delta\|_{L^4(\mathcal{S}_\delta; \mathbb{R}^{3 \times 3})} &\leq C\delta^{\frac{1}{2}}. \end{aligned} \quad (9.6)$$

For any fixed δ , the deformation v_δ is decomposed as in Theorem 3.3. There exists a subsequence still indexed by δ such that (see Subsection 6.2)

$$\begin{aligned} \mathbf{R}_\delta &\rightharpoonup \mathbf{R}^{\{0\}} \quad \text{weakly in } H^1(\mathcal{S}; SO(3)), \\ \mathcal{V}_\delta &\longrightarrow \mathcal{V}^{\{0\}} \quad \text{strongly in } H^1(\mathcal{S}; \mathbb{R}^3), \\ \frac{1}{\delta} \left(\frac{d\mathcal{V}_{i,\delta}}{ds_i} - \mathbf{R}_{i,\delta} \mathbf{t}_i \right) &\rightharpoonup \mathcal{Z}_i^{\{0\}} \quad \text{weakly in } L^2(0, L_i; \mathbb{R}^3), \\ \frac{1}{\delta^2} \Pi_{i,\delta}(\bar{v}_\delta) &\rightharpoonup \bar{v}_i^{\{0\}} \quad \text{weakly in } L^2(0, L_i; H^1(\omega; \mathbb{R}^3)). \end{aligned} \quad (9.7)$$

The couple $(\mathcal{V}^{\{0\}}, \mathbf{R}^{\{0\}})$ belongs to $\mathbb{V}\mathbb{R}_2$. Furthermore, we also have ($i \in \{1, \dots, N\}$)

$$\begin{aligned} \Pi_{i,\delta}(v_\delta) &\longrightarrow \mathcal{V}_i^{\{0\}} \quad \text{strongly in } H^1(\Omega_i; \mathbb{R}^3), \\ \Pi_{i,\delta}(\nabla_x v_\delta) &\rightharpoonup \mathbf{R}_i^{\{0\}} \quad \text{weakly in } L^4(\Omega_i; \mathbb{R}^{3 \times 3}), \\ \frac{1}{2\delta} \Pi_{i,\delta}((\nabla_x v_\delta)^T \nabla_x v_\delta - \mathbf{I}_3) &\rightharpoonup \mathbf{E}_i^{\{0\}} \quad \text{weakly in } L^2(\Omega_i; \mathbb{R}^{3 \times 3}), \end{aligned} \quad (9.8)$$

where the symmetric matrix $\mathbf{E}_i^{\{0\}} = (\mathbf{t}_i | \mathbf{n}_i | \mathbf{b}_i) \widehat{\mathbf{E}}_i^{\{0\}} (\mathbf{t}_i | \mathbf{n}_i | \mathbf{b}_i)^T$, (see Subsection 6.2). The matrix $\widehat{\mathbf{E}}_i^{\{0\}}$ is defined by

$$\widehat{\mathbf{E}}_i^{\{0\}} = \begin{pmatrix} -Y_2 \Gamma_{i,3}(\mathbf{R}^{\{0\}}) + Y_3 \Gamma_{i,2}(\mathbf{R}^{\{0\}}) + \mathcal{Z}_i^{\{0\}} \cdot \mathbf{R}_i^{\{0\}} \mathbf{t}_i & * & * \\ -\frac{1}{2} Y_3 \Gamma_{i,1}(\mathbf{R}^{\{0\}}) + \frac{1}{2} \frac{\partial \bar{u}_i^{\{0\}}}{\partial Y_2} \cdot \mathbf{t}_i & e_{22}(\bar{u}_i^{\{0\}}) & * \\ \frac{1}{2} Y_2 \Gamma_{i,1}(\mathbf{R}^{\{0\}}) + \frac{1}{2} \frac{\partial \bar{u}_i^{\{0\}}}{\partial Y_3} \cdot \mathbf{t}_i & e_{23}(\bar{u}_i^{\{0\}}) & e_{33}(\bar{u}_i^{\{0\}}) \end{pmatrix}$$

with

$$\bar{u}_i^{\{0\}} = [Y_2(\mathcal{Z}_i^{\{0\}} \cdot \mathbf{R}_i^{\{0\}} \mathbf{n}_i) + Y_3(\mathcal{Z}_i^{\{0\}} \cdot \mathbf{R}_i^{\{0\}} \mathbf{b}_i)] \mathbf{t}_i + (\mathbf{R}_i^{\{0\}})^T \bar{v}_i^{\{0\}}, \quad i \in \{1, \dots, N\}.$$

Due to the decomposition of v_δ , the assumptions (7.7), (7.8) and the above convergences (9.7) we immediately get the limit of the term involving the body forces

$$\lim_{\delta \rightarrow 0} \frac{1}{\delta^4} \int_{\mathcal{S}_\delta} f_{2,\delta} \cdot (v_\delta - I_d) = \mathcal{L}(\mathcal{V}^{\{0\}}, \mathbf{R}^{\{0\}})$$

where $\mathcal{L}(\mathcal{V}, \mathbf{R})$ is defined by (8.3).

We now consider a given rod $\mathcal{P}_{i,\delta}$. Let $\varepsilon > 0$ be fixed. Due to assumption (7.1), there exists $\theta > 0$ such that

$$\forall E \in \mathbf{S}_3, \quad |||E||| \leq \theta, \quad W(E) \geq Q(E) - \varepsilon |||E|||^2. \quad (9.9)$$

Let us denote by $\chi_{i,\delta}^\theta$ the characteristic function of the set

$$A_{i,\delta}^\theta = \left\{ s \in \Omega_i \setminus \bigcup_{k=1}^{K_i} \omega \times [a_i^k - \rho_0 \delta, a_i^k + \rho_0 \delta] ; \quad |||\Pi_{i,\delta}((\nabla_x v_\delta)^T \nabla_x v_\delta - \mathbf{I}_3)(s)||| \geq \theta \right\}$$

where a_i^k is the arc length of a knot belonging to the line γ_i . Due to (9.8), we have

$$\text{meas}(A_{i,\delta}^\theta) \leq C \frac{\delta^2}{\theta^2}. \quad (9.10)$$

Using the positive character of W , (7.2), (9.9) and (9.6) give

$$\begin{aligned} \frac{1}{\delta^4} \int_{\mathcal{S}_\delta} \widehat{W}(\nabla_x v_\delta) &\geq \sum_{i=1}^N \int_{\Omega_i} \frac{1}{\delta^2} W\left(\frac{1}{2} \Pi_{i,\delta}((\nabla_x v_\delta)^T \nabla_x v_\delta - \mathbf{I}_3)\right) (1 - \chi_{i,\delta}^\theta) \\ &\geq \sum_{i=1}^N \int_{\Omega_i} Q\left(\frac{1}{2\delta} \Pi_{i,\delta}((\nabla_x v_\delta)^T \nabla_x v_\delta - \mathbf{I}_3)\right) (1 - \chi_{i,\delta}^\theta) - C\varepsilon. \end{aligned}$$

In view of (9.10), the function $\chi_{i,\delta}^\theta$ converges a.e. to 0 as δ tends to 0 while the weak limit of $\frac{1}{2\delta} \Pi_{i,\delta}((\nabla_x v_\delta)^T \nabla_x v_\delta - \mathbf{I}_3)(1 - \chi_{i,\delta}^\theta)$ is given by (9.8). As a consequence we have

$$\liminf_{\delta \rightarrow 0} \frac{1}{\delta^4} \int_{\mathcal{S}_\delta} \widehat{W}(\nabla_x v_\delta) \geq \sum_{i=1}^N \int_{\Omega_i} Q(\mathbf{E}_i^{\{0\}}) - C\varepsilon.$$

As ε is arbitrary, this gives

$$\liminf_{\delta \rightarrow 0} \frac{1}{\delta^4} \int_{\mathcal{S}_\delta} \widehat{W}(\nabla_x v_\delta) \geq \sum_{i=1}^N \int_{\Omega_i} Q(\mathbf{E}_i^{\{0\}}). \quad (9.11)$$

Hence, due to the limit of the applied forces and (9.11) we obtain

$$\sum_{i=1}^N \int_{\Omega_i} Q(\mathbf{E}_i^{\{0\}}) - \mathcal{L}(\mathcal{V}^{\{0\}}, \mathbf{R}^{\{0\}}) \leq \liminf_{\delta \rightarrow 0} \frac{m_{2,\delta}}{\delta^4}$$

where $\mathcal{L}(\mathcal{V}, \mathbf{R})$ is defined by (8.3).

Finally, for a.e. $s_i \in]0, L_i[$ we minimize the quantity $\int_{\omega} Q(\mathbf{E}_i^{\{0\}})(s_i, Y_2, Y_3) dY_2 dY_3$ with respect to the $\mathcal{Z}_i^{\{0\}}(s_i) \cdot \mathbf{R}_i^{\{0\}}(s_i) \mathbf{t}_i(s_i)$'s and the $\bar{u}_i^{\{0\}}(s_i, \cdot, \cdot)$'s using Lemma 10.6 in the Appendix. Hence we obtain

$$\begin{aligned} \min_{(\mathcal{V}, \mathbf{R}) \in \mathbb{V}\mathbb{R}_2} \mathcal{J}_2(\mathcal{V}, \mathbf{R}) &\leq \mathcal{J}_2(\mathcal{V}^{\{0\}}, \mathbf{R}^{\{0\}}) \\ &= \sum_{i=1}^N \int_0^{L_i} a_{lk} \Gamma_{i,k}(\mathbf{R}^{\{0\}}) \Gamma_{i,l}(\mathbf{R}^{\{0\}}) - \mathcal{L}(\mathcal{V}^{\{0\}}, \mathbf{R}^{\{0\}}) \\ &\leq \sum_{i=1}^N \int_{\Omega_i} Q(\mathbf{E}_i^{\{0\}}) - \mathcal{L}(\mathcal{V}^{\{0\}}, \mathbf{R}^{\{0\}}) \leq \liminf_{\delta \rightarrow 0} \frac{m_{2,\delta}}{\delta^4}, \end{aligned} \quad (9.12)$$

where the definite positive symmetric 3×3 matrix $\mathbf{A} = (a_{lk})$ is defined in Lemma 10.6.

Step 2. In this step we show that

$$\limsup_{\delta \rightarrow 0} \frac{m_{2,\delta}}{\delta^4} \leq \min_{(\mathcal{V}, \mathbf{R}) \in \mathbb{V}\mathbb{R}_2} \mathcal{J}_2(\mathcal{V}, \mathbf{R}).$$

Let $(\mathcal{V}^{\{2\}}, \mathbf{R}^{\{2\}}) \in \mathbb{V}\mathbb{R}_2$ be such that $\mathcal{J}_2(\mathcal{V}^{\{2\}}, \mathbf{R}^{\{2\}}) = \min_{(\mathcal{V}, \mathbf{R}) \in \mathbb{V}\mathbb{R}_2} \mathcal{J}_2(\mathcal{V}, \mathbf{R})$. First, for $i \in \{1, \dots, N\}$ let \bar{v}_i be arbitrary in $W^{1,\infty}(\Omega_i; \mathbb{R}^3)$.

For each rod $\mathcal{P}_{i,\delta}$, we apply the Proposition 10.5 given in Appendix to the triplet $(\mathcal{V}_i^{\{2\}}, \mathbf{R}_i^{\{2\}}, \bar{v}_i)$ in each portion of $\mathcal{P}_{i,\delta}$ which is contained between an extremity and a knot or between two knots. Doing such leads to a sequence of deformations v_δ which belong to $\mathbb{D}_\delta \cap W^{1,\infty}(\mathcal{S}_\delta; \mathbb{R}^3)$ and which satisfy in each junction $\mathcal{J}_{A,\rho_0\delta}$

$$\forall A \in \mathcal{K}, \quad v_\delta(x) = \mathcal{V}^{\{2\}}(A) + \mathbf{R}^{\{2\}}(A)(x - A), \quad x \in \mathcal{J}_{A,\rho_0\delta}, \quad (9.13)$$

and

$$\begin{aligned} \det(\nabla v_\delta(x)) &> 0 \quad \text{for a.e. } x \in \mathcal{S}_\delta, \\ \Pi_{i,\delta}(v_\delta) &\longrightarrow \mathcal{V}_i^{\{2\}} \quad \text{strongly in } H^1(\Omega_i; \mathbb{R}^3), \\ \frac{1}{\delta} \Pi_{i,\delta}(v_\delta - \mathcal{V}_{i,\delta}) &\longrightarrow \mathbf{R}_i^{\{2\}}(Y_2 \mathbf{n}_i + Y_3 \mathbf{b}_i) \quad \text{strongly in } L^2(\Omega_i; \mathbb{R}^3), \\ \frac{1}{2\delta} \Pi_{i,\delta}((\nabla v_\delta)^T \nabla v_\delta - \mathbf{I}_3) &\longrightarrow \mathbf{E}_i^{\{2\}} = \left(\frac{d\mathbf{R}_i^{\{2\}}}{ds_i}(Y_2 \mathbf{e}_2 + Y_3 \mathbf{e}_3) \mid \frac{\partial \bar{v}_i}{\partial Y_2} \mid \frac{\partial \bar{v}_i}{\partial Y_3} \right)^T \mathbf{R}_i^{\{2\}} \\ &+ (\mathbf{R}_i^{\{2\}})^T \left(\frac{d\mathbf{R}_i^{\{2\}}}{ds_i}(Y_2 \mathbf{e}_2 + Y_3 \mathbf{e}_3) \mid \frac{\partial \bar{v}_i}{\partial Y_2} \mid \frac{\partial \bar{v}_i}{\partial Y_3} \right) \quad \text{strongly in } L^2(\Omega_i; \mathbb{R}^{3 \times 3}) \end{aligned} \quad (9.14)$$

where $\mathcal{V}_{i,\delta}$ is the average of v_δ on each cross-section of the rod $\mathcal{P}_{i,\delta}$. Moreover there exists a constant $C_1 \geq \theta$ which does not depend on δ such that

$$\forall i \in \{1, \dots, N\}, \quad \|\Pi_{i,\delta}((\nabla v_\delta)^T \nabla v_\delta - \mathbf{I}_3)\|_{L^\infty(\Omega_i; \mathbb{R}^{3 \times 3})} \leq C_1. \quad (9.15)$$

Let $\varepsilon > 0$ be fixed. Due to assumption (7.1), there exists $\theta > 0$ such that

$$\forall E \in \mathbf{S}_3, \quad \|||E||| \leq \theta, \quad W(E) \leq Q(E) + \varepsilon \|||E|||^2. \quad (9.16)$$

Let us denote by $\chi_{i,\delta}^\theta$ the characteristic function of the set

$$A_{i,\delta}^\theta = \left\{ s \in \Omega_i \setminus \bigcup_{k=1}^{K_i} \omega \times [a_i^k - \rho_0 \delta, a_i^k + \rho_0 \delta] ; \|||\Pi_{i,\delta}((\nabla_x v_\delta)^T \nabla_x v_\delta - \mathbf{I}_3)(s)\||| \geq \theta \right\}$$

where a_i^k is the arc length of a knot belonging to the line γ_i . Due to (9.8), we have

$$\text{meas}(A_{i,\delta}^\theta) \leq C \frac{\delta^2}{\theta^2}. \quad (9.17)$$

Using the fact that v_δ is equal to a rotation in the junctions (see (9.13)) the Saint Venant's strain tensor is equal to zero in the junctions. Hence we have

$$\begin{aligned} \frac{1}{\delta^4} \int_{\mathcal{S}_\delta} \widehat{W}(\nabla_x v_\delta) &= \sum_{i=1}^N \int_{\Omega_i} \frac{1}{\delta^2} W\left(\frac{1}{2} \Pi_{i,\delta}((\nabla_x v_\delta)^T \nabla_x v_\delta - \mathbf{I}_3)\right) (1 - \chi_{i,\delta}^\theta) \\ &+ \sum_{i=1}^N \int_{\Omega_i} \frac{1}{\delta^2} W\left(\frac{1}{2} \Pi_{i,\delta}((\nabla_x v_\delta)^T \nabla_x v_\delta - \mathbf{I}_3)\right) \chi_{i,\delta}^\theta. \end{aligned} \quad (9.18)$$

In view of (9.16) and the third strong convergence in (9.14), the first term of the right hand side is estimated as

$$\begin{aligned} &\sum_{i=1}^N \int_{\Omega_i} \frac{1}{\delta^2} W\left(\frac{1}{2} \Pi_{i,\delta}((\nabla_x v_\delta)^T \nabla_x v_\delta - \mathbf{I}_3)\right) (1 - \chi_{i,\delta}^\theta) \\ &\leq \sum_{i=1}^N \int_{\Omega_i} Q\left(\frac{1}{2\delta} \Pi_{i,\delta}((\nabla_x v_\delta)^T \nabla_x v_\delta - \mathbf{I}_3)\right) (1 - \chi_{i,\delta}^\theta) + C\varepsilon. \end{aligned}$$

Again, the third strong convergence in (9.14) and the pointwise convergence of the function $\chi_{i,\delta}^\theta$ allows to pass to the limit as δ in the above inequality and to obtain

$$\limsup_{\delta \rightarrow 0} \sum_{i=1}^N \int_{\Omega_i} \frac{1}{\delta^2} W\left(\frac{1}{2} \Pi_{i,\delta}((\nabla_x v_\delta)^T \nabla_x v_\delta - \mathbf{I}_3)\right) (1 - \chi_{i,\delta}^\theta) \leq \sum_{i=1}^N \int_{\Omega_i} Q(\mathbf{E}_i^{\{2\}}) + C\varepsilon.$$

Let us recall estimate (9.15). Due to the continuity of W and the third assumption of (7.1), there exists a constant C_2 such that

$$\|||E||| \leq C_1 \quad \implies \quad W(E) \leq C_2 \|||E|||^2.$$

It follows that the second term in (9.18) is less than

$$\sum_{i=1}^N \int_{\Omega_i} \frac{1}{\delta^2} W \left(\frac{1}{2} \Pi_{i,\delta} ((\nabla_x v_\delta)^T \nabla_x v_\delta - \mathbf{I}_3) \right) \chi_{i,\delta}^\theta \leq \sum_{i=1}^N \int_{\Omega_i} C_2 \left\| \frac{1}{2\delta} \Pi_{i,\delta} ((\nabla_x v_\delta)^T \nabla_x v_\delta - \mathbf{I}_3) \right\|^2 \chi_{i,\delta}^\theta.$$

We have $\chi_{i,\delta}^\theta$ tends to 0 weakly * in $L^\infty(\Omega_i)$ and the third strong convergence in (9.14), hence

$$\lim_{\delta \rightarrow 0} \sum_{i=1}^N \int_{\Omega_i} \frac{1}{\delta^2} W \left(\frac{1}{2} \Pi_{i,\delta} ((\nabla_x v_\delta)^T \nabla_x v_\delta - \mathbf{I}_3) \right) \chi_{i,\delta}^\theta = 0.$$

As ε is arbitrary, finally we get

$$\limsup_{\delta \rightarrow 0} \frac{1}{\delta^4} \int_{S_\delta} \widehat{W}(\nabla_x v_\delta) \leq \sum_{i=1}^N \int_{\Omega_i} Q(\mathbf{E}_i^{\{2\}}).$$

Thanks to the first and second convergences in (9.14) we obtain the limit of the term involving the forces. Then we obtain

$$\limsup_{\delta \rightarrow 0} \frac{m_{2,\delta}}{\delta^4} \leq \limsup_{\delta \rightarrow 0} \frac{J_{2,\delta}(v_\delta)}{\delta^4} \leq \sum_{i=1}^N \int_{\Omega_i} Q(\mathbf{E}_i^{\{2\}}) - \mathcal{L}(\mathcal{V}^{\{2\}}, \mathbf{R}^{\{2\}}) \quad (9.19)$$

where $\mathcal{L}(\mathcal{V}, \mathbf{R})$ is defined in (8.3). In view of the expression of the $\mathbf{E}_i^{\{2\}}$'s and a density argument the above inequality holds true for any family $\bar{v}_i \in L^2(0, L_i; H^1(\omega; \mathbb{R}^3))$ ($i \in \{1, \dots, N\}$). Then we can use again Lemma 10.6 in order to minimize the right hand side of this inequality with respect to the $(\mathbf{R}^{\{2\}})^T \bar{v}_i(s_i, \cdot, \cdot)$'s. This gives

$$\begin{aligned} \limsup_{\delta \rightarrow 0} \frac{m_{2,\delta}}{\delta^4} &\leq \min_{(\mathcal{V}, \mathbf{R}) \in \mathbb{V}\mathbb{R}_2} \mathcal{J}_2(\mathcal{V}, \mathbf{R}) = \mathcal{J}_2(\mathcal{V}^{\{2\}}, \mathbf{R}^{\{2\}}) \\ &= \sum_{i=1}^N \int_0^{L_i} a_{lk} \Gamma_{i,k}(\mathbf{R}^{\{2\}}) \Gamma_{i,l}(\mathbf{R}^{\{2\}}) - \mathcal{L}(\mathcal{V}^{\{2\}}, \mathbf{R}^{\{2\}}) \\ &\leq \sum_{i=1}^N \int_{\Omega_i} Q(\mathbf{E}_i^{\{2\}}) - \mathcal{L}(\mathcal{V}^{\{2\}}, \mathbf{R}^{\{2\}}). \end{aligned}$$

This concludes the proof of the theorem. \square

Remark 9.2. *Let us point out that Theorem 9.1 shows that for any minimizing sequence $(v_\delta)_\delta$ as in Step 1, the third convergence of the rescaled Green-St Venant's strain tensor in (9.8) is a strong convergence to $\mathbf{E}_i^{\{0\}}$ in $L^2(\Omega_i; \mathbb{R}^{3 \times 3})$.*

10 Appendix

In this Appendix, we first give the construction of a suitable sequence (Proposition 10.4) of deformations to prove the second step of the proof of Theorem 8.1. To do

that we first give three lemmas. In a second part of this appendix we also construct a suitable sequence (Proposition 10.5) of deformations to prove the second step of the proof of Theorem (9.1). At last Lemma 10.6 provides the elimination technique used in the proof of Theorem (9.1).

The proofs of the three lemmas below are left to the reader.

Lemma 10.1. *Let \mathbf{R} be an element in $L^2(0, L; \mathcal{C})$. We define the field \mathbf{R}'_n of $L^2(0, L; \mathcal{C})$ by*

$$\mathbf{R}'_n(t) = \frac{n}{L} \int_{kL/n}^{(k+1)L/n} \mathbf{R}(s) ds \quad \text{for any } t \text{ in }]kL/n, (k+1)L/n[, \quad k \in \{0, \dots, n-1\}$$

We have

$$\mathbf{R}'_n \longrightarrow \mathbf{R} \quad \text{strongly in } L^2(0, L; \mathcal{C}).$$

Lemma 10.2. *Let \mathbf{R} be in \mathcal{C} . There exist $(\lambda_1, \dots, \lambda_p) \in [0, 1]^p$ and $(\mathbf{R}_1, \dots, \mathbf{R}_p) \in (SO(3))^p$ such that*

$$\sum_{i=1}^p \lambda_i = 1, \quad \mathbf{R} = \sum_{i=1}^p \lambda_i \mathbf{R}_i.$$

We set

$$\mu_0 = \frac{1}{2n}, \quad \mu_i = \mu_{i-1} + \left(1 - \frac{1}{n}\right) \lambda_i, \quad i \in \{1, \dots, p\}.$$

We define \mathbf{R}'_n in $L^2(0, L; SO(3))$ by: for any $k \in \{0, \dots, n-1\}$ and for a.e. $y \in]0, 1[$ we set

$$\mathbf{R}'_n\left(\frac{k}{n} + \frac{y}{n}\right) = \begin{cases} \mathbf{I}_3 & y \in]0, \mu_0[, \\ \mathbf{R}_i & y \in]\mu_{i-1}, \mu_i[, \quad i \in \{1, \dots, p\}, \\ \mathbf{I}_3 & y \in]1 - 1/2n, 1[. \end{cases}$$

We have

$$\mathbf{R}'_n \rightharpoonup \mathbf{R} \quad \text{weakly in } L^2(0, L; \mathcal{C}).$$

Lemma 10.3. *Let $\theta^* \in [0, \pi]$ there exists a function $\theta \in W^{1,\infty}(0, 1)$ such that*

$$\theta(0) = 0, \quad \theta(1) = \theta^*, \quad \int_0^1 e^{i\theta(t)} dt = \frac{1 + e^{i\theta^*}}{2}.$$

Moreover there exists a positive constant which does not depend on θ^* such that

$$\|\theta'\|_{L^\infty(0,1)} \leq C\theta^*.$$

The above lemmas allow us to establish the following result.

Proposition 10.4. *Let $(\mathcal{V}, \mathbf{R})$ be in $\mathbb{V}\mathbb{R}_1$. There exists a sequence $((\mathcal{V}_n, \mathbf{R}_n))_n$ in $\mathbb{V}\mathbb{R}_1$ which satisfies $\mathbf{R}_n \in W^{1,\infty}(\mathcal{S}; SO(3))$, which is equal to \mathbf{I}_3 in a neighbourhood of each knot $A \in \mathcal{K}$ and each fixed extremity belonging to Γ_0 and moreover which satisfies*

$$\begin{aligned} \mathcal{V}_n &\rightharpoonup \mathcal{V} \quad \text{weakly in } H^1(\mathcal{S}; \mathbb{R}^3), \\ \mathbf{R}_n &\rightharpoonup \mathbf{R} \quad \text{weakly in } L^2(\mathcal{S}; \mathcal{C}). \end{aligned}$$

Proof. For any $n \in \mathbb{N}^*$, we first apply the Lemma 10.1 between two consecutive points of the set $\mathcal{K} \cup \Gamma$ belonging to a same line γ_i of the skeleton \mathcal{S} . It leads to a sequence $\mathbf{R}'_n \in L^2(\mathcal{S}; \mathcal{C})$ which is piecewise constant on \mathcal{S} . Then, we define \mathcal{V}'_n by integration, using the formula $\frac{d(\mathcal{V}'_n)_i}{ds_i} = (\mathbf{R}'_n)_i \mathbf{t}_i$ between two consecutive points in $\mathcal{K} \cup \Gamma$ imposing the values

$$\mathcal{V}'_n(A) = \mathcal{V}(A) \text{ for every point in } \mathcal{K} \cup \Gamma,$$

which is possible because the means of $(\mathbf{R}'_n)_i$ are preserved between two points in $\mathcal{K} \cup \Gamma$.

Hence, we obtain a sequence $((\mathcal{V}'_n, \mathbf{R}'_n))_n$ in $\mathbb{V}\mathbb{R}_1$ which satisfies

$$\begin{aligned} \mathcal{V}'_n &\longrightarrow \mathcal{V} \text{ strongly in } H^1(\mathcal{S}; \mathbb{R}^3), \\ \mathbf{R}'_n &\longrightarrow \mathbf{R} \text{ strongly in } L^2(\mathcal{S}; \mathcal{C}). \end{aligned}$$

Then, we consider the couple $\left((1 - 1/n)\mathcal{V}'_n + 1/n\phi, (1 - 1/n)\mathbf{R}'_n + 1/n\mathbf{I}_3 \right) \in \mathbb{V}\mathbb{R}_1$ and we apply the Lemma 10.2, again between two consecutive points of $\mathcal{K} \cup \Gamma$ on a same line γ_i of the skeleton \mathcal{S} . We obtain an element $(\mathcal{V}''_n, \mathbf{R}''_n) \in \mathbb{V}\mathbb{R}_1$ such that \mathbf{R}''_n belongs to $L^2(\mathcal{S}; SO(3))$ and is piecewise constant on \mathcal{S} and equal to \mathbf{I}_3 in a neighbourhood of each knot and each fixed extremity belonging to Γ_0 and moreover which satisfies

$$\begin{aligned} \mathcal{V}''_n &\rightharpoonup \mathcal{V} \text{ weakly in } H^1(\mathcal{S}; \mathbb{R}^3), \\ \mathbf{R}''_n &\rightharpoonup \mathbf{R} \text{ weakly in } L^2(\mathcal{S}; \mathcal{C}). \end{aligned}$$

Finally, thanks to the Lemma 10.3, we replace the element $(\mathcal{V}''_n, \mathbf{R}''_n)$ by another $(\mathcal{V}'''_n, \mathbf{R}'''_n)$ in $\mathbb{V}\mathbb{R}_1$ such that \mathbf{R}'''_n belongs to $W^{1,\infty}(\mathcal{S}; SO(3))$. Notice that we do this modification without changing the values of \mathcal{V}''_n at the points of $\mathcal{K} \cup \Gamma$. We get

$$\begin{aligned} \mathcal{V}'''_n &\rightharpoonup \mathcal{V} \text{ weakly in } H^1(\mathcal{S}; \mathbb{R}^3), \\ \mathbf{R}'''_n &\rightharpoonup \mathbf{R} \text{ weakly in } L^2(\mathcal{S}; \mathcal{C}). \end{aligned}$$

The proposition is proved. □

The proposition below gives an approximation result of a deformation in a rod by deformations which are rotations near the extremities. Recall that for a single rod $\Omega_\delta =]0, L[\times \omega_\delta$, the rescaling into $\Omega =]0, L[\times \omega$ is performed through the operator Π_δ defined by

$$\Pi_\delta(\psi)(y_1, Y_2, Y_3) = \psi(s, \delta Y_2, \delta Y_3) \quad \text{for a.e. } (y_1, Y_2, Y_3) \in \Omega.$$

Proposition 10.5. *Let $(\mathcal{V}, \mathbf{R}, \bar{v})$ be in $H^1(0, L; \mathbb{R}^3) \times H^1(0, L; SO(3)) \times W^{1,\infty}(\Omega; \mathbb{R}^3)$ such that $\frac{d\mathcal{V}}{dy_1} = \mathbf{R}\mathbf{e}_1$. Let $\rho > 0$ be given. There exists a sequence of deformations*

$v_\delta \in W^{1,\infty}(\Omega_\delta; \mathbb{R}^3)$ satisfying for δ small enough

$$\begin{aligned} v_\delta(y) &= \mathcal{V}(0) + \mathbf{R}(0)y, & y \in]0, \rho\delta[\times \omega_\delta, \\ v_\delta(y) &= \mathcal{V}(L) + \mathbf{R}(L)(y - L\mathbf{e}_1), & y \in]L - \rho\delta, L[\times \omega_\delta, \\ \det(\nabla v_\delta(y)) &> 0 & \text{for almost any } y \in \Omega_\delta, \\ \|\Pi_\delta((\nabla v_\delta)^T \nabla v_\delta - \mathbf{I}_3)\|_{L^\infty(\Omega; \mathbb{R}^{3 \times 3})} &\leq C, \end{aligned} \tag{10.1}$$

where the constant C does not depend on δ .

Moreover the following convergence holds true

$$\begin{aligned} \frac{1}{2\delta} \Pi_\delta((\nabla v_\delta)^T \nabla v_\delta - \mathbf{I}_3) &\longrightarrow \left(\frac{d\mathbf{R}}{dy_1} (Y_2 \mathbf{e}_2 + Y_3 \mathbf{e}_3) \mid \frac{\partial \bar{v}}{\partial Y_2} \mid \frac{\partial \bar{v}}{\partial Y_3} \right)^T \mathbf{R} \\ &+ \mathbf{R}^T \left(\frac{d\mathbf{R}}{dy_1} (Y_2 \mathbf{e}_2 + Y_3 \mathbf{e}_3) \mid \frac{\partial \bar{v}}{\partial Y_2} \mid \frac{\partial \bar{v}}{\partial Y_3} \right) \text{ strongly in } L^2(\Omega; \mathbb{R}^{3 \times 3}). \end{aligned} \tag{10.2}$$

Proof. Let $N \geq 4$ and $\varepsilon = L/N$. At the end of the proof, we will fix $\varepsilon \geq \delta$ in terms of δ and ρ or equivalently N in terms of δ and ρ . We define $\mathbf{R}_\varepsilon \in W^{1,\infty}(0, L; \mathbb{R}^{3 \times 3})$ as follows:

$$\begin{aligned} \mathbf{R}_\varepsilon &\text{ is piecewise linear on } [0, L], \\ \mathbf{R}_\varepsilon(y_1) &= \mathbf{R}(0) & y_1 \in [0, \varepsilon], \\ \mathbf{R}_\varepsilon(k\varepsilon) &= \mathbf{R}(k\varepsilon), & k \in \{2, \dots, N-2\} \\ \mathbf{R}_\varepsilon(y_1) &= \mathbf{R}(L) & y_1 \in [L - \varepsilon, L], \end{aligned}$$

There exists a constant C independent of ε such that

$$\begin{aligned} \left\| \frac{d\mathbf{R}_\varepsilon}{dy_1} \right\|_{L^2(0, L; \mathbb{R}^{3 \times 3})} &\leq C \left\| \frac{d\mathbf{R}}{dy_1} \right\|_{L^2(0, L; \mathbb{R}^{3 \times 3})}, & \left\| \frac{d\mathbf{R}_\varepsilon}{dy_1} \right\|_{L^\infty(0, L; \mathbb{R}^{3 \times 3})} &\leq \frac{C}{\varepsilon}, \\ \left\| \mathbf{R}_\varepsilon - \mathbf{R} \right\|_{L^2(0, L; \mathbb{R}^{3 \times 3})} &\leq C\varepsilon \left\| \frac{d\mathbf{R}}{dy_1} \right\|_{L^2(0, L; \mathbb{R}^{3 \times 3})}, & \left\| \mathbf{R}_\varepsilon - \mathbf{R} \right\|_{L^\infty(0, L; \mathbb{R}^{3 \times 3})} &\leq C\sqrt{\varepsilon} \left\| \frac{d\mathbf{R}}{dy_1} \right\|_{L^2(0, L; \mathbb{R}^{3 \times 3})}. \end{aligned}$$

Due to the specific construction of the sequence \mathbf{R}_ε , one also get

$$\begin{aligned} \mathbf{R}_\varepsilon &\longrightarrow \mathbf{R} \text{ strongly in } H^1(0, L; \mathbb{R}^{3 \times 3}), \\ \frac{1}{\varepsilon} (\mathbf{R}_\varepsilon - \mathbf{R}) &\longrightarrow 0 \text{ strongly in } L^2(0, L; \mathbb{R}^{3 \times 3}). \end{aligned} \tag{10.3}$$

Now we define \mathcal{V}_ε by

$$\begin{aligned} \mathcal{V}_\varepsilon &\text{ is piecewise linear on } [0, L], \\ \mathcal{V}_\varepsilon(y_1) &= \mathcal{V}(0) + y_1 \mathbf{R}(0) \mathbf{e}_1 & y_1 \in [0, \varepsilon], \\ \mathcal{V}_\varepsilon(k\varepsilon) &= \mathcal{V}(k\varepsilon), & k \in \{2, \dots, N-2\}, \\ \mathcal{V}_\varepsilon(y_1) &= \mathcal{V}(L) + (y_1 - L) \mathbf{R}(L) \mathbf{e}_1 & y_1 \in [L - \varepsilon, L]. \end{aligned}$$

We have

$$\begin{aligned} \left\| \frac{d\mathcal{V}_\varepsilon}{dy_1} - \frac{d\mathcal{V}}{dy_1} \right\|_{L^2(0, L; \mathbb{R}^3)} &\leq C\varepsilon \left\| \frac{d\mathbf{R}}{dy_1} \right\|_{L^2(0, L; \mathbb{R}^{3 \times 3})}, \\ \left\| \frac{d\mathcal{V}_\varepsilon}{dy_1} - \frac{d\mathcal{V}}{dy_1} \right\|_{L^\infty(0, L; \mathbb{R}^3)} &\leq C\sqrt{\varepsilon} \left\| \frac{d\mathbf{R}}{dy_1} \right\|_{L^2(0, L; \mathbb{R}^{3 \times 3})}, \end{aligned}$$

and the following convergence

$$\frac{1}{\varepsilon} \left(\frac{d\mathcal{V}_\varepsilon}{dy_1} - \frac{d\mathcal{V}}{dy_1} \right) \rightarrow 0 \quad \text{strongly in} \quad L^2(0, L; \mathbb{R}^3). \quad (10.4)$$

We set

$$d_\varepsilon(y_1) = \inf \left(1, \sup \left(0, \frac{y_1 - \varepsilon}{\varepsilon} \right), \sup \left(0, \frac{L - \varepsilon - y_1}{\varepsilon} \right) \right), \quad y_1 \in [0, L].$$

Now, we consider the deformation v_ε defined by

$$v_\varepsilon(y) = \mathcal{V}_\varepsilon(y_1) + \mathbf{R}_\varepsilon(y_1)(y_2\mathbf{e}_2 + y_3\mathbf{e}_3) + \delta^2 d_\varepsilon(y_1) \bar{v}(y_1, \frac{y_2}{\delta}, \frac{y_3}{\delta}), \quad x \in \Omega_\delta.$$

A straightforward calculation gives

$$\begin{aligned} (\nabla v_\varepsilon)^T \nabla v_\varepsilon - \mathbf{I}_3 &= (\nabla v_\varepsilon - \mathbf{R}_\varepsilon)^T \mathbf{R}_\varepsilon + \mathbf{R}_\varepsilon^T (\nabla v_\varepsilon - \mathbf{R}_\varepsilon) + (\nabla v_\varepsilon - \mathbf{R}_\varepsilon)^T (\nabla v_\varepsilon - \mathbf{R}_\varepsilon) \\ &\quad + (\mathbf{R}_\varepsilon - \mathbf{R})^T \mathbf{R}_\varepsilon + \mathbf{R}^T (\mathbf{R}_\varepsilon - \mathbf{R}), \\ \frac{\partial v_\varepsilon}{\partial y_1} - \mathbf{R}_\varepsilon \mathbf{e}_1 &= \frac{d\mathcal{V}_\varepsilon}{dy_1} - \mathbf{R}_\varepsilon \mathbf{e}_1 + \frac{d\mathbf{R}_\varepsilon}{dy_1} (y_2\mathbf{e}_2 + y_3\mathbf{e}_3) + \delta^2 \left(\frac{\partial \bar{v}}{\partial y_1} \left(\cdot, \frac{y_2}{\delta}, \frac{y_3}{\delta} \right) + \frac{dd_\varepsilon}{dy_1} \bar{v} \left(\cdot, \frac{y_2}{\delta}, \frac{y_3}{\delta} \right) \right), \\ \frac{\partial v_\varepsilon}{\partial y_\alpha} - \mathbf{R}_\varepsilon \mathbf{e}_\alpha &= \delta d_\varepsilon \frac{\partial \bar{v}}{\partial Y_\alpha} \left(\cdot, \frac{y_2}{\delta}, \frac{y_3}{\delta} \right), \quad \alpha \in \{2, 3\}. \end{aligned}$$

Hence, thanks to the above estimates we obtain

$$\|\nabla v_\varepsilon - \mathbf{R}\|_{L^\infty(\Omega_\delta; \mathbb{R}^{3 \times 3})} \leq C \left(\sqrt{\varepsilon} \left\| \frac{d\mathbf{R}}{dy_1} \right\|_{L^2(0, L; \mathbb{R}^{3 \times 3})} + \frac{\delta}{\varepsilon} + \frac{\delta^2}{\varepsilon} \|\bar{v}\|_{W^{1, \infty}(\Omega; \mathbb{R}^3)} \right).$$

Finally, we choose $\theta \geq \rho$ and large enough so that $\theta \geq 4C$. Then setting $\varepsilon = \theta\delta$ show that if δ is small enough (denoting now v_ε by v_δ)

$$\|\nabla v_\delta - \mathbf{R}\|_{L^\infty(\Omega_\delta; \mathbb{R}^{3 \times 3})} \leq \frac{1}{2},$$

which shows first that the determinant of ∇v_ε is positive. Moreover the above inequality also implies (e.g.) that $\|(\nabla v_\delta)^T \nabla v_\delta - \mathbf{I}_3\|_{L^\infty(\Omega_\delta; \mathbb{R}^{3 \times 3})} \leq 2$.

At least the above decomposition of $(\nabla v_\delta)^T \nabla v_\delta - \mathbf{I}_3$, the strong convergences (10.3) and (10.4) together with the definition of d_ε allows to obtain the strong convergence (10.2). \square

We know justify the elimination process used in Theorem 9.1. We set

$$\mathbb{W} = \left\{ \psi \in H^1(\omega; \mathbb{R}^3) \mid \int_\omega \psi(Y_2, Y_3) dY_2 dY_3 = 0, \int_\omega (Y_3 \psi_2(Y_2, Y_3) - Y_2 \psi_3(Y_2, Y_3)) dY_2 dY_3 = 0 \right\}$$

We equip \mathbb{W} with the norm

$$\|\psi\|_{\mathbb{W}} = \|\nabla \psi_1\|_{L^2(\omega; \mathbb{R}^2)} + \|e_{22}(\psi)\|_{L^2(\omega)} + \|e_{23}(\psi)\|_{L^2(\omega)} + \|e_{33}(\psi)\|_{L^2(\omega)}$$

where

$$e_{kl}(\psi) = \frac{1}{2} \left(\frac{\partial \psi_k}{\partial Y_l} + \frac{\partial \psi_l}{\partial Y_k} \right), \quad (k, l) \in \{2, 3\}^2.$$

In the space \mathbb{W} , the above norm is equivalent to the H^1 norm.

Let \mathbf{Q} be a matrix field belonging to $L^\infty(\omega; \mathbb{R}^{6 \times 6})$ satisfying for a. e. $(Y_2, Y_3) \in \omega$

$$\begin{aligned} \mathbf{Q}(Y_2, Y_3) &\text{ is a symmetric positive definite matrix,} \\ \mathbf{Q}(-Y_2, -Y_3) &= \mathbf{Q}(Y_2, Y_3), \\ \forall \xi \in \mathbb{R}^6, \quad c|\xi|^2 &\leq \mathbf{Q}(Y_2, Y_3) \xi \cdot \xi \leq C|\xi|^2 \end{aligned} \tag{10.5}$$

where c and C are positive constants which do not depend on (Y_2, Y_3) .

We denote $\chi_i \in \mathbb{W}$, $i \in \{1, 2, 3\}$ the solutions of the following variational problems:

$$\begin{aligned} \forall \psi \in \mathbb{W}, \\ \int_{\omega} \mathbf{Q} \begin{pmatrix} 0 \\ \frac{1}{2} \frac{\partial \chi_{i,1}}{\partial Y_2} \\ \frac{1}{2} \frac{\partial \chi_{i,1}}{\partial Y_3} \\ e_{22}(\chi_i) \\ e_{23}(\chi_i) \\ e_{33}(\chi_i) \end{pmatrix} \cdot \begin{pmatrix} 0 \\ \frac{1}{2} \frac{\partial \psi_1}{\partial Y_2} \\ \frac{1}{2} \frac{\partial \psi_1}{\partial Y_3} \\ e_{22}(\psi) \\ e_{23}(\psi) \\ e_{33}(\psi) \end{pmatrix} &= - \int_{\omega} \mathbf{Q} \mathbf{V}_i \cdot \begin{pmatrix} 0 \\ \frac{1}{2} \frac{\partial \psi_1}{\partial Y_2} \\ \frac{1}{2} \frac{\partial \psi_1}{\partial Y_3} \\ e_{22}(\psi) \\ e_{23}(\psi) \\ e_{33}(\psi) \end{pmatrix} \end{aligned} \tag{10.6}$$

where

$$\mathbf{V}_1 = \begin{pmatrix} 0 \\ -\frac{1}{2} Y_3 \\ \frac{1}{2} Y_2 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad \mathbf{V}_2 = \begin{pmatrix} Y_3 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad \mathbf{V}_3 = \begin{pmatrix} -Y_2 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}. \tag{10.7}$$

The fields χ_i , $i \in \{1, 2, 3\}$, satisfy

$$\chi_i(-Y_2, -Y_3) = \chi_i(Y_2, Y_3) \quad \text{for a.e. } (Y_2, Y_3) \in \Omega. \tag{10.8}$$

Lemma 10.6. *There exists a symmetric positive definite matrix \mathbf{A} such that for any*

$\Gamma_i \in \mathbb{R}$, $i \in \{1, 2, 3\}$,

$$\begin{aligned}
& \forall (Z, \psi) \in \mathbb{R} \times H^1(\omega; \mathbb{R}^3), \\
& \int_{\omega} \mathbf{Q} \begin{pmatrix} -Y_2\Gamma_3 + Y_3\Gamma_2 + Z \\ -\frac{1}{2}Y_3\Gamma_1 + \frac{1}{2}\frac{\partial\psi_1}{\partial Y_2} \\ \frac{1}{2}Y_2\Gamma_1 + \frac{1}{2}\frac{\partial\psi_1}{\partial Y_3} \\ e_{22}(\psi) \\ e_{23}(\psi) \\ e_{33}(\psi) \end{pmatrix} \cdot \begin{pmatrix} -Y_2\Gamma_3 + Y_3\Gamma_2 + Z \\ -\frac{1}{2}Y_3\Gamma_1 + \frac{1}{2}\frac{\partial\psi_1}{\partial Y_2} \\ \frac{1}{2}Y_2\Gamma_1 + \frac{1}{2}\frac{\partial\psi_1}{\partial Y_3} \\ e_{22}(\psi) \\ e_{23}(\psi) \\ e_{33}(\psi) \end{pmatrix} \\
& \geq \int_{\omega} \mathbf{Q} \begin{pmatrix} -Y_2\Gamma_3 + Y_3\Gamma_2 \\ -\frac{1}{2}Y_3\Gamma_1 + \frac{1}{2}\frac{\partial\bar{\chi}_1}{\partial Y_2} \\ \frac{1}{2}Y_2\Gamma_1 + \frac{1}{2}\frac{\partial\bar{\chi}_1}{\partial Y_3} \\ e_{22}(\bar{\chi}) \\ e_{23}(\bar{\chi}) \\ e_{33}(\bar{\chi}) \end{pmatrix} \cdot \begin{pmatrix} -Y_2\Gamma_3 + Y_3\Gamma_2 \\ -\frac{1}{2}Y_3\Gamma_1 + \frac{1}{2}\frac{\partial\bar{\chi}_1}{\partial Y_2} \\ \frac{1}{2}Y_2\Gamma_1 + \frac{1}{2}\frac{\partial\bar{\chi}_1}{\partial Y_3} \\ e_{22}(\bar{\chi}) \\ e_{23}(\bar{\chi}) \\ e_{33}(\bar{\chi}) \end{pmatrix} = \mathbf{A} \begin{pmatrix} \Gamma_1 \\ \Gamma_2 \\ \Gamma_3 \end{pmatrix} \cdot \begin{pmatrix} \Gamma_1 \\ \Gamma_2 \\ \Gamma_3 \end{pmatrix}
\end{aligned}$$

where

$$\bar{\chi} = \Gamma_1\chi_1 + \Gamma_2\chi_2 + \Gamma_3\chi_3 \in \mathbb{W}. \quad (10.9)$$

Proof. First of all, it remains the same to minimize the functional

$$(Z, \psi) \mapsto \int_{\omega} \mathbf{Q} \begin{pmatrix} -Y_2\Gamma_3 + Y_3\Gamma_2 + Z \\ -\frac{1}{2}Y_3\Gamma_1 + \frac{1}{2}\frac{\partial\psi_1}{\partial Y_2} \\ \frac{1}{2}Y_2\Gamma_1 + \frac{1}{2}\frac{\partial\psi_1}{\partial Y_3} \\ e_{22}(\psi) \\ e_{23}(\psi) \\ e_{33}(\psi) \end{pmatrix} \cdot \begin{pmatrix} -Y_2\Gamma_3 + Y_3\Gamma_2 + Z \\ -\frac{1}{2}Y_3\Gamma_1 + \frac{1}{2}\frac{\partial\psi_1}{\partial Y_2} \\ \frac{1}{2}Y_2\Gamma_1 + \frac{1}{2}\frac{\partial\psi_1}{\partial Y_3} \\ e_{22}(\psi) \\ e_{23}(\psi) \\ e_{33}(\psi) \end{pmatrix}$$

over $\mathbb{R} \times H^1(\omega; \mathbb{R}^3)$ or $\mathbb{R} \times \mathbb{W}$. The second property of \mathbf{Q} in (10.5) implies that the minimum is achieved for $Z = 0$. Then, it is easy to prove that the minimum of the functional over the space $\mathbb{R} \times \mathbb{W}$ is achieved for the element $(0, \bar{\chi})$, where $\bar{\chi}$ is given by (10.9).

It is easy to prove (e.g. by contradiction) that the norm

$$\|(\Gamma_1, \theta)\| = \sqrt{\int_{\omega} \left(\left| -\frac{1}{2}Y_3\Gamma_1 + \frac{1}{2}\frac{\partial\theta}{\partial Y_2} \right|^2 + \left| \frac{1}{2}Y_2\Gamma_1 + \frac{1}{2}\frac{\partial\theta}{\partial Y_3} \right|^2 \right)}$$

on the space $\mathbb{R} \times H^1(\omega)/\mathbb{R}$ is equivalent to the product norm of this space. Hence, there exists a positive constant C such that for any $\Gamma_1 \in \mathbb{R}$ and any function $\theta \in H^1(\omega)$ we

have

$$\int_{\omega} \left(\left| -\frac{1}{2}Y_3\Gamma_1 + \frac{1}{2}\frac{\partial\theta}{\partial Y_2} \right|^2 + \left| \frac{1}{2}Y_2\Gamma_1 + \frac{1}{2}\frac{\partial\theta}{\partial Y_3} \right|^2 \right) \geq C(\Gamma_1^2 + \|\nabla\theta\|_{L^2(\omega;\mathbb{R}^2)}^2).$$

Taking into account the third property in (10.5), this allows to prove the positivity of matrix \mathbf{A} . \square

Remark 10.7. *In the case where the matrix \mathbf{Q} corresponds to an isotropic and homogeneous material, we have*

$$\mathbf{Q} = \begin{pmatrix} \lambda + 2\mu & 0 & 0 & \lambda & 0 & \lambda \\ 0 & 2\mu & 0 & 0 & 0 & 0 \\ 0 & 0 & 2\mu & 0 & 0 & 0 \\ \lambda & 0 & 0 & \lambda + 2\mu & 0 & \lambda \\ 0 & 0 & 0 & 0 & 2\mu & 0 \\ \lambda & 0 & 0 & \lambda & 0 & \lambda + 2\mu \end{pmatrix}$$

where λ, μ are the Lamé's constants. We get

$$\chi_1(Y_2, Y_3) = 0, \quad \chi_2(Y_2, Y_3) = \begin{pmatrix} -\nu Y_2 Y_3 \\ -\nu \frac{Y_3^2 - Y_2^2}{2} \end{pmatrix}, \quad \chi_3(Y_2, Y_3) = \begin{pmatrix} \nu \frac{Y_2^2 - Y_3^2}{2} \\ \nu Y_2 Y_3 \end{pmatrix}$$

where ν is the Poisson's coefficient. The matrix \mathbf{A} is equal to

$$\mathbf{A} = \begin{pmatrix} \frac{\pi\mu}{4} & 0 & 0 \\ 0 & \frac{\pi E}{4} & 0 \\ 0 & 0 & \frac{\pi E}{4} \end{pmatrix}$$

where E is the Young's modulus.

References

- [1] S. S. Antman. The theory of rods, in Handbuch der Physik, Band VIa/2, S. Flügge & C. Truesdell eds., Springer-Verlag, Berlin, (1972), 641–703.
- [2] S. S. Antman. Nonlinear Problems of Elasticity, Applied Mathematical Sciences 107, Springer-Verlag, New York (1995).
- [3] D. Blanchard, A. Gaudiello, G. Griso. Junction of a periodic family of elastic rods with a 3d plate. I. J. Math. Pures Appl. (9) 88 (2007), no 1, 149-190.
- [4] D. Blanchard, A. Gaudiello, G. Griso. Junction of a periodic family of elastic rods with a thin plate. II. J. Math. Pures Appl. (9) 88 (2007), no 2, 1-33.

- [5] D. Blanchard, G. Griso. Microscopic effects in the homogenization of the junction of rods and a thin plate. *Asympt. Anal.* 56 (2008), no 1, 1-36.
- [6] D. Blanchard, G. Griso. Decomposition of deformations of thin rods. Application to nonlinear elasticity, *Ana. Appl.* 7 (1) (2009) 21-71.
- [7] D. Blanchard, G. Griso. Decomposition of the deformations of a thin shell. Asymptotic behavior of the Green-St Venant's strain tensor. *J. of Elasticity*, 101 (2), 179-205 (2010).
- [8] D. Blanchard, G. Griso. Modeling of rod-structures in nonlinear elasticity. *C. R. Acad. Sci. Paris, Ser. I*, doi:10.1016/j.crma.2010.09.008.
- [9] P.G. Ciarlet. *Mathematical Elasticity, Vol. I*, North-Holland, Amsterdam (1988).
- [10] P.G. Ciarlet, H. Le Dret and R. Nzingwa. Junctions between three dimensional and two dimensional linearly elastic structures. *J. Math. Pures Appl.* 68 (1989), 261–295.
- [11] G. Friesecke, R. D. James and S. Müller. A theorem on geometric rigidity and the derivation of nonlinear plate theory from the three-dimensional elasticity. *Communications on Pure and Applied Mathematics*, Vol. LV, 1461-1506 (2002).
- [12] A. Gaudiello, R. Monneau, J. Mossino, F. Murat, A. Sili. Junction of elastic plates and beams. *ESAIM Control Optim. Calc. Var.* 13 (2007), no. 3, 419–457.
- [13] G. Griso. Asymptotic behavior of rods by the unfolding method. *Math. Meth. Appl. Sci.* 2004; 27: 2081-2110.
- [14] G. Griso. Asymptotic behavior of structures made of plates. *Analysis and Applications* 3 (2005), 4, 325-356.
- [15] G. Griso. Decomposition of displacements of thin structures. *J. Math. Pures Appl.* 89 (2008), 199-233.
- [16] G. Griso. Asymptotic behavior of structures made of curved rods. *Analysis and Applications*, 6 (2008), 1, 11-22.
- [17] I. Gruais. Modélisation de la jonction entre une plaque et une poutre en élasticité linéarisée. *RAIRO: Modél. Math. Anal. Numér.* 27 (1993) 77-105.
- [18] I. Gruais. Modeling of the junction between a plate and a rod in nonlinear elasticity. *Asymptotic Anal.* 7 (1993) 179-194.
- [19] H. Le Dret. Modeling of the junction between two rods, *J. Math. Pures Appl.* 68 (1989), 365–397.
- [20] H. Le Dret. Modeling of a folded plate, *Comput. Mech.*, 5 (1990), 401–416.

- [21] H. Le Dret. Problèmes Variationnels dans les Multi-domaines: Modélisation des Jonctions et Applications. Masson, Paris (1991).
- [22] L. Trabucho and J.M. Viano. Mathematical Modelling of Rods, Handbook of Numerical Analysis 4. North-Holland, Amsterdam (1996).