



# The Boltzmann-Grad limit of the periodic Lorentz gas in two space dimensions

Emanuele Caglioti, François Golse

## ► To cite this version:

Emanuele Caglioti, François Golse. The Boltzmann-Grad limit of the periodic Lorentz gas in two space dimensions. *Comptes-Rendus de l'Académie des Sciences, Série 1, Mathématiques*, Elsevier, 2008, 346 (7-8), pp.477-482. <10.1016/j.crma.2008.01.016>. <hal-00176202v2>

**HAL Id: hal-00176202**

<https://hal.archives-ouvertes.fr/hal-00176202v2>

Submitted on 3 Oct 2007

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# The Boltzmann-Grad limit of the periodic Lorentz gas in two space dimensions

Emanuele Caglioti<sup>a</sup>, François Golse<sup>b</sup>

<sup>a</sup> *Università di Roma “La Sapienza”, Dipartimento di Matematica “Guido Castelnuovo”, P.le Aldo Moro 2, 00185 Rome*

<sup>b</sup> *Ecole polytechnique, Centre de Mathématiques Laurent Schwartz, 91128 Palaiseau Cedex*

## Abstract

The periodic Lorentz gas is the dynamical system corresponding to the free motion of a point particle in a periodic system of fixed spherical obstacles of radius  $r$  centered at the integer points, assuming all collisions of the particle with the obstacles to be elastic. In this Note, we study this motion on time intervals of order  $1/r$  as  $r \rightarrow 0^+$ .

## Résumé

**La limite de Boltzmann-Grad du gaz de Lorentz périodique en dimension deux d’espace.** Le gaz de Lorentz périodique est le système dynamique correspondant au mouvement libre dans le plan d’une particule ponctuelle rebondissant de manière élastique sur un système de disques de rayon  $r$  centrés aux points de coordonnées entières. On étudie ce mouvement pour  $r \rightarrow 0^+$  sur des temps de l’ordre de  $1/r$ .

Version française abrégée

On appelle gaz de Lorentz le système dynamique correspondant au mouvement libre d’une particule ponctuelle dans un système d’obstacles circulaires de rayon  $r$  centrés aux sommets d’un réseau de  $\mathbf{R}^2$ , supposant que les collisions entre la particule et les obstacles sont parfaitement élastiques. Les trajectoires de la particule sont alors données par les formules (2). La limite de Boltzmann-Grad pour le gaz de Lorentz consiste à supposer que le rayon des obstacles  $r \rightarrow 0^+$ , et à observer la dynamique de la particule sur des plages de temps longues, de l’ordre de  $1/r$  — voir (3) pour la loi d’échelle de Boltzmann-Grad en dimension 2.

Or les trajectoires de la particule s’expriment en fonction de l’application de transfert d’obstacle à obstacle  $T_r$  définie par (8) — où la notation  $Y$  désigne la transformation inverse de (7) — application qui associe, à tout paramètre d’impact  $h' \in [-1, 1]$  correspondant à une particule quittant la surface d’un

obstacle dans la direction  $\omega \in \mathbf{S}^1$ , le paramètre d'impact  $h$  à la collision suivante, ainsi que le temps  $s$  s'écoulant jusqu'à cette collision. (Pour une définition de la notion de paramètre d'impact, voir (6).)

On se ramène donc à étudier le comportement limite de l'application de transfert  $T_r$  pour  $r \rightarrow 0^+$ .

**Proposition 0.1** *Lorsque  $0 < \omega_2 < \omega_1$  et  $\alpha = \frac{\omega_2}{\omega_1} \notin \mathbf{Q}$ , l'application de transfert  $T_r$  est approchée à  $O(r^2)$  près par l'application  $\mathbf{T}_{A,B,Q,N}$  définie à la formule (14). Pour  $\omega \in \mathbf{S}^1$  quelconque, on se ramène au cas ci-dessus par la symétrie (15).*

Les paramètres  $A, B, Q, N \bmod 2$  intervenant dans l'application de transfert asymptotique sont définis à partir du développement en fraction continue (9) de  $\alpha$  par les formules (11) et (12).

On voit sur ces formules que les paramètres  $A, B, Q, N \bmod 2$  sont des fonctions très fortement oscillantes des variables  $\omega$  et  $r$ . Il est donc naturel de chercher le comportement limite de l'application de transfert  $T_r$  dans une topologie faible vis à vis de la dépendance en la direction  $\omega$ . On montre ainsi que, pour tout  $h' \in [-1, 1]$ , la famille d'applications  $\omega \mapsto T_r(h', \omega)$  converge au sens des mesures de Young (voir par exemple [8] p. 146–154 pour une définition de cette notion) lorsque  $r \rightarrow 0^+$  vers une mesure de probabilité  $P(s, h|h') ds dh$  indépendante de  $\omega$  :

**Théorème 0.2** *Pour tout  $\Phi \in C_c(\mathbf{R}_+^* \times ]-1, 1[)$  et tout  $h' \in ]-1, 1[$ , la limite (16) a lieu dans  $L^\infty(\mathbf{S}^1)$  faible-\* lorsque  $r \rightarrow 0^+$ , où la mesure de probabilité  $P(s, h|h') ds dh$  est l'image de la probabilité  $\mu$  définie dans (17) par l'application  $(A, B, Q, N) \mapsto \mathbf{T}_{A,B,Q,N}(h')$  de la formule (14). De plus, cette densité de probabilité de transition  $P(s, h|h')$  vérifie les propriétés (18).*

Le théorème ci-dessus est le résultat principal de cette Note : il montre que, dans la limite de Boltzmann-Grad, le transfert d'obstacle à obstacle est décrit de manière naturelle par une densité de probabilité de transition  $P(s, h|h')$ , où  $s$  est le laps de temps entre deux collisions successives avec les obstacles (dans l'échelle de temps de la limite de Boltzmann-Grad),  $h$  le paramètre d'impact lors de la collision future et  $h'$  celui correspondant à la collision passée.

Le fait que la probabilité de transition  $P(s, h|h')$  soit indépendante de la direction suggère l'hypothèse d'indépendance (H) des quantités  $A, B, Q, N \bmod 2$  correspondant à des collisions successives.

**Théorème 0.3** *Sous l'hypothèse (H), pour toute densité de probabilité  $f^{in} \in C_c(\mathbf{R}^2 \times \mathbf{S}^1)$ , la fonction de distribution  $f_r \equiv f_r(t, x, \omega)$  de la théorie cinétique, définie par (3) converge dans  $L^\infty(\mathbf{R}_+ \times \mathbf{R}^2 \times \mathbf{S}^1)$  vers la limite (22) lorsque  $r \rightarrow 0^+$ , où  $F$  est la solution du problème de Cauchy (21) posé dans l'espace des phases étendu  $(x, \omega, s, h) \in \mathbf{R}^2 \times \mathbf{S}^1 \times \mathbf{R}_+^* \times ]-1, 1[$ .*

Dans le cas d'obstacles aléatoires indépendants et poissonniens, Gallavotti a montré que la limite de Boltzmann-Grad du gaz de Lorentz obéit à l'équation cinétique de Lorentz (4). Le cas périodique est absolument différent : en se basant sur des estimations (cf. [3] et [7]) du temps de sortie du domaine  $Z_r$  défini dans (1), on démontre que la limite de Boltzmann-Grad du gaz de Lorentz périodique ne peut pas être décrite par l'équation de Lorentz (4) sur l'espace des phases  $\mathbf{R}^2 \times \mathbf{S}^1$  classique de la théorie cinétique : voir [6]. Si l'hypothèse (H) ci-dessous était vérifiée, le modèle cinétique (22) dans l'espace des phases étendu fournirait donc l'équation devant remplacer l'équation cinétique classique de Lorentz (4) dans le cas périodique.

## 1. The Lorentz gas

The Lorentz gas is the dynamical system corresponding to the free motion of a single point particle in a periodic system of fixed spherical obstacles, assuming that collisions between the particle and any of the obstacles are elastic. Henceforth, we assume that the space dimension is 2 and that the obstacles are disks of radius  $r$  centered at each point of  $\mathbf{Z}^2$ . Hence the domain left free for particle motion is

$$Z_r = \{x \in \mathbf{R}^2 \mid \text{dist}(x, \mathbf{Z}^2) > r\}, \quad \text{where it is assumed that } 0 < r < \frac{1}{2}. \quad (1)$$

Assuming that the particle moves at speed 1, its trajectory starting from  $x \in Z_r$  with velocity  $\omega \in \mathbf{S}^1$  at time  $t = 0$  is  $t \mapsto (X_r, \Omega_r)(t; x, \omega) \in \mathbf{R}^2 \times \mathbf{S}^1$  given by

$$\begin{aligned} \dot{X}_r(t) &= \Omega_r(t) & \text{and } \dot{\Omega}_r(t) &= 0 & \text{whenever } X_r(t) &\in Z_r, \\ X_r(t+0) &= X_r(t-0) & \text{and } \Omega_r(t+0) &= \mathcal{R}[X_r(t)]\Omega_r(t-0) & \text{whenever } X_r(t-0) &\in \partial Z_r, \end{aligned} \quad (2)$$

denoting  $\dot{\cdot} = \frac{d}{dt}$  and  $\mathcal{R}[X_r(t)]$  the specular reflection on  $\partial Z_r$  at the point  $X_r(t) = X_r(t \pm 0)$ . Assume that the initial position  $x$  and direction  $\omega$  of the particle are distributed in  $Z_r \times \mathbf{S}^1$  with some probability density  $f^{in} \equiv f^{in}(x, \omega)$ , and define

$$f_r(t, x, \omega) := f^{in}(rX_r(-t/r; x, \omega), \Omega_r(-t/r; x, \omega)) \quad \text{whenever } x \in Z_r. \quad (3)$$

We are concerned with the limit of  $f_r$  as  $r \rightarrow 0^+$  in some appropriate sense to be explained below. In the 2-dimensional setting considered here, this is precisely the Boltzmann-Grad limit.

In the case of a random (Poisson), instead of periodic, configuration of obstacles, Gallavotti [5] proved that the expectation of  $f_r$  converges to the solution of the Lorentz kinetic equation for  $(x, \omega) \in \mathbf{R}^2 \times \mathbf{S}^1$ :

$$(\partial_t + \omega \cdot \nabla_x) f(t, x, \omega) = \int_{\mathbf{S}^1} (f(t, x, \omega - 2(\omega \cdot n)n) - f(t, x, \omega)) (\omega \cdot n)_+ dn, \quad f \Big|_{t=0} = f^{in}. \quad (4)$$

In the case of a periodic distribution of obstacles, the Boltzmann-Grad limit of the Lorentz gas cannot be described by a transport equation as above: see [6] for a complete proof, based on estimates on the free path length to be found in [3] and [7]. This limit involves instead a linear Boltzmann equation on an extended phase space with two new variables taking into account correlations between consecutive collisions with the obstacles that are an effect of periodicity: see Theorem 4.1.

## 2. The transfer map

Denote by  $n_x$  the inward unit normal to  $Z_r$  at the point  $x \in \partial Z_r$ , consider

$$\Gamma_r^\pm = \{(x, \omega) \in \partial Z_r \times \mathbf{S}^1 \mid \pm \omega \cdot n_x > 0\}, \quad (5)$$

and let  $\Gamma_r^\pm / \mathbf{Z}^2$  be the quotient of  $\Gamma_r^\pm$  under the action of  $\mathbf{Z}^2$  by translation on the  $x$  variable. For  $(x, \omega) \in \Gamma_r^+$ , let  $\tau_r(x, \omega)$  be the exit time from  $x$  in the direction  $\omega$  and  $h_r(x, \omega)$  be the impact parameter:

$$\tau_r(x, \omega) = \inf\{t > 0 \mid x + t\omega \in \partial Z_r\}, \quad \text{and } h_r(x, \omega) = \sin(\omega, n_x). \quad (6)$$

Obviously, the map

$$\Gamma_r^+ / \mathbf{Z}^2 \ni (x, \omega) \mapsto (h_r(x, \omega), \omega) \in ]-1, 1[ \times \mathbf{S}^1 \quad (7)$$

coordinatizes  $\Gamma_r^+ / \mathbf{Z}^2$ , and we henceforth denote  $Y_r$  its inverse.

For each  $r \in ]0, \frac{1}{2}[$ , consider now the transfer map  $T_r : ]-1, 1[ \times \mathbf{S}^1 \rightarrow \mathbf{R}_+^* \times ]-1, 1[$  defined by

$$T_r(h', \omega) = (r\tau_r(Y_r(h', \omega)), h_r(X_r(\tau_r(Y_r(h', \omega))); Y_r(h', \omega)), \Omega_r(\tau_r(Y_r(h', \omega))); Y_r(h', \omega))). \quad (8)$$

For a particle leaving the surface of an obstacle in the direction  $\omega$  with impact parameter  $h'$ , the transition map  $T_r(h', \omega) = (s, h)$  gives the (rescaled) distance  $s$  to the next collision, and the corresponding




Figure 1. Left: the transfer map  $(s, h) = T_r(h', v)$ , with  $h' = \sin(n_{x'}, v)$  and  $h = \sin(n_x, v)$ . Right: Particles leaving the surface of one obstacle will next collide with one of generically three obstacles. The figure explains the geometrical meaning of  $A, B, Q$ .

impact parameter  $h$ . Obviously, each trajectory (2) of the particle can be expressed in terms of the transfer map  $T_r$  and iterates thereof. The Boltzmann-Grad limit of the periodic Lorentz gas is therefore reduced to computing the limiting behavior of  $T_r$  as  $r \rightarrow 0^+$ , and this is our main purpose in this Note.

We first need some pieces of notation. Assume  $\omega = (\omega_1, \omega_2)$  with  $0 < \omega_2 < \omega_1$ , and  $\alpha = \omega_2/\omega_1 \in ]0, 1[ \setminus \mathbf{Q}$ . Consider the continued fraction expansion of  $\alpha$ :

$$\alpha = [0; a_0, a_1, a_2, \dots] = \frac{1}{a_0 + \frac{1}{a_1 + \dots}}. \quad (9)$$

Define the sequences of convergents  $(p_n, q_n)_{n \geq 0}$  and errors  $(d_n)_{n \geq 0}$  by the recursion formulas

$$\begin{aligned} p_{n+1} &= a_n p_n + p_{n-1}, & p_0 &= 1, & p_1 &= 0, & d_n &= (-1)^{n-1} (q_n \alpha - p_n), \\ q_{n+1} &= a_n q_n + q_{n-1}, & q_0 &= 0, & q_1 &= 1, \end{aligned} \quad (10)$$

and let

$$N(\alpha, r) = \inf\{n \geq 0 \mid d_n \leq 2r\sqrt{1 + \alpha^2}\}, \quad \text{and } k(\alpha, r) = - \left[ \frac{2r\sqrt{1 + \alpha^2} - d_{N(\alpha, r)-1}}{d_{N(\alpha, r)}} \right]. \quad (11)$$

**Proposition 2.1** *For each  $\omega = (\cos \theta, \sin \theta)$  with  $0 < \theta < \frac{\pi}{4}$ , set  $\alpha = \tan \theta$  and  $\epsilon = 2r\sqrt{1 + \alpha^2}$ , and*

$$A(\alpha, r) = 1 - \frac{d_{N(\alpha, r)}}{\epsilon}, \quad B(\alpha, r) = 1 - \frac{d_{N(\alpha, r)-1} - k(\alpha, r)d_{N(\alpha, r)}}{\epsilon}, \quad Q(\alpha, r) = \epsilon q_{N(\alpha, r)}. \quad (12)$$

*In the limit  $r \rightarrow 0^+$ , the transition map  $T_r$  defined in (8) is explicit in terms of  $A, B, Q, N$  up to  $O(r^2)$ :*

$$T_r(h', \omega) = \mathbf{T}_{A(\alpha, r), B(\alpha, r), Q(\alpha, r), N(\alpha, r)}(h') + (O(r^2), 0) \text{ for each } h' \in ]-1, 1[. \quad (13)$$

*In the formula above*

$$\begin{aligned} \mathbf{T}_{A, B, Q, N}(h') &= (Q, h' - 2(-1)^N(1 - A)) && \text{if } (-1)^N h' \in ]1 - 2A, 1], \\ \mathbf{T}_{A, B, Q, N}(h') &= (Q', h' + 2(-1)^N(1 - B)) && \text{if } (-1)^N h' \in [-1, -1 + 2B[, \\ \mathbf{T}_{A, B, Q, N}(h') &= (Q' + Q, h' + 2(-1)^N(A - B)) && \text{if } (-1)^N h' \in [-1 + 2B, 1 - 2A], \end{aligned} \quad (14)$$

*for each  $(A, B, Q, N) \in K := ]0, 1[^3 \times \mathbf{Z}/2\mathbf{Z}$ , with the notation  $Q' = \frac{1 - Q(1 - B)}{1 - A}$ .*

The proof uses the 3-term partition of the 2-torus defined in section 2 of [4], following the work of [1]. For  $\omega = (\cos \theta, \sin \theta)$  with arbitrary  $\theta \in \mathbf{R}$ , the map  $h' \mapsto T_r(h', \omega)$  is computed using Proposition 2.1 in the following manner. Set  $\tilde{\theta} = \theta - m\frac{\pi}{2}$  with  $m = [\frac{2}{\pi}(\theta + \frac{\pi}{4})]$  and let  $\tilde{\omega} = (\cos \tilde{\theta}, \sin \tilde{\theta})$ . Then

$$T_r(h', \omega) = (s, h), \quad \text{where } (s, \text{sign}(\tan \tilde{\theta})h) = T_r(\text{sign}(\tan \tilde{\theta})h', \tilde{\omega}). \quad (15)$$

### 3. The Boltzmann-Grad limit of the transfer map $T_r$

The formulas (11) and (12) defining  $A, B, Q, N \bmod 2$  show that these quantities are strongly oscillating functions of the variables  $\omega$  and  $r$ . In view of Proposition 2.1, one therefore expects the transfer map  $T_r$  to have a limit as  $r \rightarrow 0^+$  only in the weakest imaginable sense, i.e. in the sense of Young measures — see [8], pp. 146–154 for a definition of this notion of convergence.

The main result in the present Note is the theorem below. It says that, for each  $h' \in [-1, 1]$ , the family of maps  $\omega \mapsto T_r(h', \omega)$  converges as  $r \rightarrow 0^+$  and in the sense of Young measures to some probability measure  $P(s, h|h')$  that is moreover independent of  $\omega$ .

**Theorem 3.1** *For each  $\Phi \in C_c(\mathbf{R}_+^* \times [-1, 1])$  and each  $h' \in [-1, 1]$*

$$\Phi(T_r(h', \cdot)) \rightarrow \int_0^\infty \int_{-1}^1 \Phi(s, h) P(s, h|h') ds dh \quad \text{in } L^\infty(\mathbf{S}_\omega^1) \text{ weak-}^* \text{ as } r \rightarrow 0^+, \quad (16)$$

where the transition probability  $P(s, h|h')$  is the image of the probability measure on  $K$  given by

$$d\mu(A, B, Q, N) = \frac{6}{\pi^2} \mathbf{1}_{0 < A < 1} \mathbf{1}_{0 < B < 1-A} \mathbf{1}_{0 < Q < \frac{1}{2-A-B}} \frac{dAdBdQ}{1-A} (\delta_{N=0} + \delta_{N=1}) \quad (17)$$

under the map  $K \ni (A, B, Q, N) \mapsto \mathbf{T}_{A,B,Q,N}(h') \in \mathbf{R}_+ \times [-1, 1]$ . Moreover,  $P$  satisfies:

$$(s, h, h') \mapsto (1+s)P(s, h|h') \text{ is piecewise continuous and bounded on } \mathbf{R}_+ \times [-1, 1] \times [-1, 1], \quad (18)$$

and  $P(s, h|h') = P(s, -h|-h')$  for each  $h, h' \in [-1, 1]$  and  $s \geq 0$ .

The proof of (16-17) is based on the explicit representation of the transition map in Proposition 2.1 together with Kloosterman sums techniques as in [2]. The explicit formula for the transition probability  $P$  is very complicated and we do not report it here, however it clearly entails the properties (18).

### 4. The Boltzmann-Grad limit of the Lorentz gas dynamics

For each  $r \in ]0, \frac{1}{2}[$ , denote  $d\gamma_r^+(x, \omega)$  the probability measure on  $\Gamma_r^+$  that is proportional to  $\omega \cdot n_x dx d\omega$ . This probability measure is invariant under the billiard map

$$\mathbf{B}_r : \Gamma_r^+ \ni (x, \omega) \mapsto \mathbf{B}_r(x, \omega) = (x + \tau_r(x, \omega)\omega, \mathcal{R}[x + \tau_r(x, \omega)\omega]\omega) \in \Gamma_r^+. \quad (19)$$

For  $(x^0, \omega^0) \in \Gamma_r^+$ , set  $(x^n, \omega^n) = \mathbf{B}_r^n(x^0, \omega^0)$  and  $\alpha^n = \min(|\omega_1^n/\omega_2^n|, |\omega_2^n/\omega_1^n|)$  for each  $n \geq 0$ , and define

$$b_r^n = (A(\alpha_n, r), B(\alpha_n, r), Q(\alpha_n, r), N(\alpha_n, r) \bmod 2) \in K \quad \text{for each } n \geq 0. \quad (20)$$

We make the following asymptotic independence hypothesis: for each  $n \geq 1$  and each  $\Psi \in C([-1, 1] \times K^n)$

$$(H) \quad \lim_{r \rightarrow 0^+} \int_{\Gamma_r^+} \Psi(h_r, \omega_0, b_r^1, \dots, b_r^n) d\gamma_r^+(x_0, \omega_0) = \int_{-1}^1 \frac{dh'}{2} \int_{\mathbf{S}^1} \frac{d\omega_0}{2\pi} \int_{K^n} \Psi(h', \omega_0, \beta_1, \dots, \beta_n) d\mu(\beta_1) \dots d\mu(\beta_n)$$

Under this assumption, the Boltzmann-Grad limit of the Lorentz gas is described by a kinetic model on the extended phase space  $\mathbf{R}^2 \times \mathbf{S}^1 \times \mathbf{R}_+ \times [-1, 1]$  — unlike the Lorentz kinetic equation (4), that is set on the usual phase space  $\mathbf{R}^2 \times \mathbf{S}^1$ .

**Theorem 4.1** *Assume (H), and let  $f^{in}$  be any continuous, compactly supported probability density on  $\mathbf{R}^2 \times \mathbf{S}^1$ . Denoting by  $\tilde{R}[\theta]$  the rotation of an angle  $\theta$ , let  $F \equiv F(t, x, \omega, s, h)$  be the solution of*

$$\begin{aligned} (\partial_t + \omega \cdot \nabla_x - \partial_s)F(t, x, \omega, s, h) &= \int_{-1}^1 P(s, h|h')F(t, x, \tilde{R}[\pi - 2 \arcsin(h')]\omega, 0, h')dh' \\ F(0, x, \omega, s, h) &= f^{in}(x, \omega) \int_s^\infty \int_{-1}^1 P(\tau, h|h')dh'd\tau \end{aligned} \quad (21)$$

where  $(x, \omega, s, h)$  runs through  $\mathbf{R}^2 \times \mathbf{S}^1 \times \mathbf{R}_+ \times [-1, 1]$ . Then the family  $(f_r)_{0 < r < \frac{1}{2}}$  defined in (3) satisfies

$$f_r \rightarrow \int_0^\infty \int_{-1}^1 F(\cdot, \cdot, \cdot, s, h) ds dh \text{ in } L^\infty(\mathbf{R}_+ \times \mathbf{R}^2 \times \mathbf{S}^1) \text{ weak-* as } r \rightarrow 0^+. \quad (22)$$

For each  $(s_0, h_0) \in \mathbf{R}_+ \times [-1, 1]$ , let  $(s_n, h_n)_{n \geq 1}$  be the Markov chain defined by the induction formula

$$(s_n, h_n) = \mathbf{T}_{\beta_n}(h_{n-1}) \text{ for each } n \geq 1, \quad \text{together with } \omega_n = \tilde{R}[2 \arcsin(h_{n-1}) - \pi]\omega_{n-1}, \quad (23)$$

where  $\beta_n \in K$  are independent random variables distributed under  $\mu$ . The proof of Theorem 4.1 relies upon approximating the particle trajectory  $(X_r, \Omega_r)(t)$  starting from  $(x_0, \omega_0)$  in terms of the following jump process with values in  $\mathbf{R}^2 \times \mathbf{S}^1 \times \mathbf{R}_+ \times [-1, 1]$  with the help of Proposition 2.1

$$\begin{aligned} (X_t, \Omega_t, S_t, H_t)(x_0, \omega_0, s_0, h_0) &= (x_0 + t\omega_0, \omega_0, s_0 - t, h_0) && \text{for } 0 \leq t < s_0, \\ (X_t, \Omega_t, S_t, H_t)(x_0, \omega_0, s_0, h_0) &= (X_{\tau_n} + (t - s_n)\omega_n, \omega_n, s_{n+1} - t, h_n) && \text{for } s_n \leq t < s_{n+1}. \end{aligned} \quad (24)$$

Unlike in the case of a random (Poisson) distribution of obstacles, the successive impact parameters on each particle path are not independent and uniformly distributed in the periodic case — likewise, the successive free path lengths on each particle path are not independent with exponential distribution. The Markov chain (23) is introduced to handle precisely this difficulty.

## References

- [1] S. Blank, N. Krikorian, Thom's problem on irrational flows. *Internat. J. Math.* **4**, 721–726, (1993).
- [2] F. Boca, A. Zaharescu, The distribution of the free path lengths in the periodic two-dimensional Lorentz gas in the small-scatterer limit. *Commun. Math. Phys.* **269**, 425–471, (2007).
- [3] J. Bourgain, F. Golse, B. Wennberg On the distribution of free path lengths for the periodic Lorentz gas. *Commun. Math. Phys.* **190**, 491–508, (1998).

- [4] E. Caglioti, F. Golse, On the distribution of free path lengths for the periodic Lorentz gas III. *Commun. Math. Phys.* **236**, 199–221, (2003).
- [5] G. Gallavotti, Rigorous theory of the Boltzmann equation in the Lorentz gas. Preprint mp\_arc 93-304 ([http://www.ma.utexas.edu/mp\\_arc/](http://www.ma.utexas.edu/mp_arc/)), (1972).
- [6] F. Golse, On the Periodic Lorentz Gas and the Lorentz Kinetic Equation. Preprint arXiv:math/0703812v1 [math.AP], submitted to *Ann. Fac. Sci. Toulouse*.
- [7] F. Golse, B. Wennberg, On the distribution of free path lengths for the periodic Lorentz gas II. *M2AN Modél. Math. et Anal. Numér.* **34**, 1151–1163, (2000).
- [8] L. Tartar, Compensated compactness and applications to partial differential equations. In *Nonlinear analysis and mechanics: Heriot-Watt Symposium. Vol. IV*, R.J. Knops ed., Research Notes in Mathematics 39, Pitman, London 1979.