



CP Violation Studies at Tevatron

E. Ben-Haim

► **To cite this version:**

E. Ben-Haim. CP Violation Studies at Tevatron. P. Ratoff, G. Borissov, M. Bozzo, R.W.J. Jones, C.S. Kalman, M. Smizanska and N. Solomey. 7th International Conference on Hyperons, Charm and Beauty Hadrons - BEACH, Jul 2006, Lancaster, United Kingdom. Elsevier, 167, pp.120-123, 2007, <10.1016/j.nuclphysbps.2006.12.094>. <in2p3-00142605>

HAL Id: in2p3-00142605

<http://hal.in2p3.fr/in2p3-00142605>

Submitted on 20 Apr 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CP Violation Studies at Tevatron

E. Ben-Haim^a *

^aLPNHE - Universite Pierre et Marie Curie-Paris6, UMR7585, Paris F-75005 France; IN2P3-CNRS

We present an overview of a few recent results related to CP-violation from the Tevatron. First, we discuss a measurement of the dimuon charge asymmetry from $D\bar{O}$, that extracts the CP-violation parameter of B^0 mixing and decay. This is followed by the CDF measurement of the CP-violating asymmetry in $B^0 \rightarrow K^+\pi^-$ decays. Finally we give the CDF result on the ratio $R = \frac{BR(B \rightarrow D^0 K)}{BR(B \rightarrow D^0 \pi)}$.

1. Introduction

Heavy flavor physics at the Tevatron exploits the large $b\bar{b}$ production cross section and the fact that, unlike at the B -factories, all species of B hadrons are produced. There are also several challenges, mainly due to the huge total inelastic cross section and to the abundance of combinatorical background. The analyses described below became possible thanks to the high performances of the CDF and $D\bar{O}$ detectors. CDF has an excellent mass resolution and the unique ability to trigger events with charged particles originated in vertices displaced from the primary $p\bar{p}$ vertex (displaced tracks). The particle identification, with both the Time-of-Flight detector and dE/dx from the drift chamber, is crucial. $D\bar{O}$ uses its superb muon system: large (η, ϕ) coverage, good scintillator based triggering and cosmic ray rejection, low punch-through and precision tracking. The $D\bar{O}$ analysis presented here uses the clean muon id and the ability to reverse the toroid and solenoid magnetic fields.

2. CP-Violation Parameter of B^0 Mixing and Decay

The $D\bar{O}$ experiment has extracted the CP-violation parameter of B^0 mixing and decay from the dimuon charge asymmetry. This measurement used an integrated luminosity of 970 pb^{-1} . CP-violation in mixing, which has not yet been observed for B mesons, is sensitive to several ex-

tensions of the Standard Model [1,2]. The CP-violation parameter ϵ_{B^0} can be obtained by measuring A_{SL} , the asymmetry of the same side lepton pairs coming from direct B decays[1,3]:

$$\frac{4\Re(\epsilon_{B^0})}{1 + |\epsilon_{B^0}|^2} = A_{SL} = \frac{N(b\bar{b} \rightarrow l^+l^+X) - N(b\bar{b} \rightarrow l^-l^-X)}{N(b\bar{b} \rightarrow l^+l^+X) + N(b\bar{b} \rightarrow l^-l^-X)}. \quad (1)$$

A_{SL} is extracted from the dimuon charge asymmetry $A = \frac{N^{++} - N^{--}}{N^{++} + N^{--}}$, where N^{++} (N^{--}) is the number of events with two positive (negative) muon candidates passing selection cuts. The measured A must contain only the physics part of the charged asymmetry, separating it from detector effects. In order to relate A to A_{SL} , all processes contributing to A have to be identified.

The polarities of the toroid and solenoid magnetic fields are reversed roughly every week so that the four solenoid-toroid polarity combinations are exposed to approximately the same integrated luminosity. This allows cancellation of first order detector effects, as the possibly different reconstruction efficiencies of positive and negative tracks due to different trajectories.

The muon detector is shown schematically in Figure 1. Let $n_\alpha^{\beta\gamma}$ be the number of muons passing cuts with charge $\alpha = \pm 1$, toroid polarity $\beta = \pm 1$, and $\gamma = +1$ if $\eta > 0$ and $\gamma = -1$ if $\eta < 0$. The physics and the detector are modeled as follows:

$$n_\alpha^{\beta\gamma} \equiv \frac{1}{4} N \epsilon^\beta (1 + \alpha A) (1 + \alpha \gamma A_{fb}) (1 + \gamma A_{det})$$

*On behalf of the CDF and $D\bar{O}$ collaborations




Figure 1. Schematic drawing of the magnetized iron toroids of the DØ detector, and muon tracks related by toroid polarity reversal, CP conjugation and forward-backward reflection.

$$\times (1 + \alpha\beta\gamma A_{ro})(1 + \beta\gamma A_{\beta\gamma})(1 + \alpha\beta A_{\alpha\beta}). \quad (2)$$

N is the number of muons passing cuts, and ϵ^β is the fraction of integrated luminosity with toroid polarity β ($\epsilon^+ + \epsilon^- = 1$). Equation (2) defines six asymmetries. A is the dimuon charge asymmetry, A_{fb} is the forward-backward asymmetry (that quantifies the tendency of μ^+ to go in the proton direction and μ^- to go in the anti-proton direction), A_{det} measures the north-south asymmetry of the detector, and A_{ro} is the range-out asymmetry (that quantifies the change in acceptance and range-out of muon tracks that bend toward, or away from, the iron toroid magnet). $A_{\alpha\beta}$ is a detector asymmetry between tracks bending north and tracks bending south. A and A_{fb} are physics asymmetries that we want to measure, and A_{det} , A_{ro} and $A_{\alpha\beta}$ are detector asymmetries. The model (2) fits 8 numbers $n_\alpha^{\beta\gamma}$ with 8 parameters (N , ϵ^+ , and 6 asymmetries).

The measured value of A is

$$A = -0.0013 \pm 0.0012(\text{stat}) \pm 0.0008(\text{sys}). \quad (3)$$

The resulting CP-violation parameter, taking into account the weights of the different physics processes producing charged dimuon pairs, and assuming that the asymmetry (if any) is due to

asymmetric $B^0 \leftrightarrow \bar{B}^0$ mixing and decay is

$$\frac{\Re(\epsilon_{B^0})}{1 + |\epsilon_{B^0}|^2} = -0.0011 \pm 0.0010(\text{stat}) \pm 0.0007(\text{sys}).$$

This is the world best measurement of ϵ_{B^0} . The dominating systematic error is due to prompt $\mu + K^\pm$ -decay.

3. CP asymmetry in $B^0 \rightarrow K^+\pi^-$ decays

The CDF experiment has measured the CP-violating asymmetry in $B^0 \rightarrow K^+\pi^-$ decays using an integrated luminosity of 360 pb^{-1} approximately. The flavor-specific $B^0 \rightarrow K^+\pi^-$ decay occurs in the SM through the dominant tree and penguin diagrams. Their interfering amplitudes induce the CP asymmetry $A_{CP}(B^0 \rightarrow K^+\pi^-)$ defined as follows:

$$\frac{\mathcal{B}(\bar{B}^0 \rightarrow K^-\pi^+) - \mathcal{B}(B^0 \rightarrow K^+\pi^-)}{\mathcal{B}(\bar{B}^0 \rightarrow K^-\pi^+) + \mathcal{B}(B^0 \rightarrow K^+\pi^-)} \quad (4)$$

B -factories recently measured a $\mathcal{O}(10\%)$ asymmetry with 2% accuracy, probing for the first time direct CP violation in the b -quark sector [4,5]; however, additional experimental information is needed because theoretical predictions still suffer from large (5–10%) uncertainties [6,7,8], and the observed asymmetries in neutral and charged modes are not consistent, as the SM would suggest. A measurement from the Tevatron is therefore interesting, also for the unique possibility to combine asymmetry measurements in $B^0 \rightarrow K^+\pi^-$ and $B_s^0 \rightarrow K^-\pi^+$ decays, which provide a model-independent probe for the presence of non-SM physics [9].

The sample of pairs of oppositely-charged particles reconstructed with π mass assignment, has been used to form $B_{(s)}^0$ meson candidates. After applying a set of optimized selection cuts, the resulting $\pi\pi$ -mass distribution (Histogram in Figure 2) shows a clean signal.

Despite the excellent mass resolution, the various $B_{(s)}^0 \rightarrow h^+h^-$ modes overlapped into an unresolved mass peak, while the PID resolution was insufficient for separating them on an event-by-event basis. We achieved a statistical separation instead, with a multivariate, unbinned likelihood-fit (fit of composition) that used PID information,




Figure 2. Invariant $\pi\pi$ -mass after the optimized offline selection with individual signal components (cumulative) and backgrounds (overlapping) overlaid. Contributions to background come from random pairs of tracks which satisfy the selection requirements and partially-reconstructed $B_{(s)}^0$ decays, as resulting from the invariant-mass fit.

provided by the dE/dx in the drift chamber, and kinematics. The fit used five observables: the invariant $\pi\pi$ -mass $m_{\pi\pi}$, the signed momentum-imbalance $\alpha = (1 - p_1/p_2)q_1$, the scalar sum of particles' momenta p_{tot} , and the dE/dx of both particles. In α , the momentum (charge) p_1 (q_1) refers to the softer track. By combining kinematics and charge information, the fit separated also $K^+\pi^-$ from $K^-\pi^+$ final states.

The fit found three modes contributing to the peak: $313 \pm 34 B^0 \rightarrow \pi^+\pi^-$, $1475 \pm 60 B^0 \rightarrow K^+\pi^-$, and $523 \pm 41 B_s^0 \rightarrow K^+K^-$ decays. A not yet statistically significant contribution of $64 \pm 30 B_s^0 \rightarrow K^-\pi^+$ decays was also found. Fit projections are overlaid to data in Figure 2. From 787 ± 42 reconstructed $B^0 \rightarrow K^+\pi^-$ decays and 689 ± 41 reconstructed $\bar{B}^0 \rightarrow K^-\pi^+$ decays, we measured the following uncorrected value for the direct CP asymmetry:

$$\frac{N(\bar{B}^0 \rightarrow K^-\pi^+) - N(B^0 \rightarrow K^+\pi^-)}{N(\bar{B}^0 \rightarrow K^-\pi^+) + N(B^0 \rightarrow K^+\pi^-)} = (-6.6 \pm 3.9)\%. \quad (5)$$

Above result was then corrected for differences in trigger, reconstruction, and selection efficiencies between $B^0 \rightarrow K^+\pi^-$ and $\bar{B}^0 \rightarrow K^-\pi^+$ modes

The dominant source of the systematic error was the uncertainty on the dE/dx model for kaons, pions, and track-to-track correlation. This effect is expected to partially reduce as the size of D meson decays samples, used for calibration of the dE/dx model, increases. The second important contribution derived from the statistical uncertainty on the nominal value of B_s^0 masses. Since we use the B_s^0 masses measured by CDF, this uncertainty will reduce with the increasing statistic of fully-reconstructed B_s^0 decays.

We quote the following result for the direct CP asymmetry in $B^0 \rightarrow K^+\pi^-$ decays, where all contributions to the systematic uncertainty have been summed in quadrature:

$$A_{CP}(B^0 \rightarrow K^+\pi^-) = (-5.8 \pm 3.9 \pm 0.7)\%, \quad (6)$$

which is approximately 1.5σ different from zero, and in agreement with world best results: $A_{CP}(B^0 \rightarrow K^+\pi^-) = (-10.8 \pm 2.4 \pm 0.7)\%$ from the Babar Collaboration [4] and $A_{CP}(B^0 \rightarrow K^+\pi^-) = (-9.3 \pm 1.8 \pm 0.8)\%$, from the Belle Collaboration [5].

CDF result is still limited by the statistic uncertainty; however, its systematic uncertainty, at the same level of B -factories, is promising: with significantly more data already collected, we expect to reduce the statistical uncertainty down to 2.5%, which will make CDF result competitive with B -factories soon.

4. Measurement of the ratio $\frac{BR(B \rightarrow D^0 K)}{BR(B \rightarrow D^0 \pi)}$

This CDF measurement exploited an integrated luminosity of 360 pb^{-1} approximately to measure the ratio $R = \frac{BR(B \rightarrow D_{flav}^0 K)}{BR(B \rightarrow D_{flav}^0 \pi)}$. R is a necessary input of the GLW method [10,11] to obtain the CKM angle γ .

We reconstructed the $B^- \rightarrow D^0 \pi^-$ with $D^0 \rightarrow K^-\pi^+$ (flavour eigenstate) and the $B^- \rightarrow D_{CP+}^0 \pi^-$ with $D_{CP+}^0 \rightarrow \pi^+\pi^-$ and $D_{CP+}^0 \rightarrow K^+K^-$ (CP-even eigenstate). The π mass is assigned to the B daughter track. Selection cuts have been applied. The resulting invariant $D^0 \pi$

mass from a generic Monte Carlo sample is shown in Figure 3, where various $B \rightarrow D^{(*)}h^\pm$ are present. To reject most of the background contributions while keeping the $B^+ \rightarrow \overline{D}^0 K^+$ signal, we used the narrow mass window $5.17 < m(D\pi) < 5.6$. The resulting $D^0\pi$ mass in data is the histogram in Figure 4.




Figure 3. Invariant $D\pi$ mass obtained from generic B^+ MC sample, showing the different signal and background reconstructed components: $B^+ \rightarrow \overline{D}^0\pi^+$, $B^+ \rightarrow \overline{D}^0K^+$, $B^+ \rightarrow \overline{D}^{*0}\pi^+$, $B^+ \rightarrow \overline{D}^{*0}K^+$ and other decay modes. Right: zoom on the suppressed $B^+ \rightarrow \overline{D}^0K^+$ peak region.

We separated the yields of $B^- \rightarrow D^0K^-$ from $B^- \rightarrow D^0\pi^-$ and the different background contributions using an multivariate, unbinned likelihood-fit, exploiting information provided by the dE/dx in the drift chamber and kinematics. Fit projections are overlaid to data in Figure 4.




Figure 4. Mass Projections of the multivariate likelihood fit

The raw fit results ($3265 \pm 38 B \rightarrow D\pi$ and $224 \pm 22 B \rightarrow DK$ decays) have been corrected for detector effects and analysis efficiencies in order to obtain the final result. We quote:

$$R = \frac{BR(B \rightarrow D_{flav}^0 K)}{BR(B \rightarrow D_{flav}^0 \pi)} =$$

$$0.065 \pm 0.007(stat) \pm 0.004(sys). \quad (7)$$

The world average for R is 0.0830 ± 0.0035 . It combines the results from the Belle ($0.077 \pm 0.005 \pm 0.006$ [12]), Babar ($0.0831 \pm 0.0035 \pm 0.002$ [13]) and Cleo ($0.099^{+0.014+0.007}_{-0.012-0.006}$ [14]) collaborations.

REFERENCES

1. Lisa Randall and Shufang Su, Nucl. Phys. **B 540**, 37 (1999), hep-th/9807377.
2. Anne L. Hewett, hep-ph/9803370 (1998).
3. “Review of Particle Physics”, S. Eidelman *et al.*, Phys. Lett. **B 592**, 1 (2004).
4. B. Aubert *et al.* (BaBar Coll.), Phys. Rev. Lett. **93**:131801 (2004), [hep-ex/0407057]; ICHEP06 conference talk [hep-ex/0607106].
5. K. Abe *et al.* (Belle Coll.), [hep-ex/0507045]; Y. Chao, these proceedings; ICHEP06 conference talk.
6. Y. Y. Keum and A. I. Sanda, Phys. Rev. D **67**:054009 (2003), [hep-ph/0209014].
7. M. Beneke *et al.*, Nucl. Phys. B **606**:245 (2001), [hep-ph/0104110].
8. H. N. Li and S. Mishima, [hep-ph/0602214].
9. H. J. Lipkin, Phys. Lett. B **621**:126 (2005), [hep-ph/0503022].
10. M. Gronau and D. Wyler, Phys. Lett. B **265** (1991) 172.
11. M. Gronau and D. London., Phys. Lett. B **253** (1991) 483.
12. S. K. Swain *et al.* [Belle Collaboration], Phys. Rev. D **68**, 051101 (2003) [hep-ex/0304032].
13. B. Aubert *et al.* [BABAR Collaboration], [hep-ex/0207087].
14. A. Bornheim *et al.* [CLEO Collaboration], Phys. Rev. D **68** (2003) 052002 [hep-ex/0302026].