



On the Inverse Problem for a Size-Structured Population Model

Benoît Perthame, Jorge Zubelli

► To cite this version:

Benoît Perthame, Jorge Zubelli. On the Inverse Problem for a Size-Structured Population Model. 2006. <hal-00110904>

HAL Id: hal-00110904

<https://hal.archives-ouvertes.fr/hal-00110904>

Submitted on 2 Nov 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On the Inverse Problem for a Size-Structured Population Model

Benoît Perthame^{*†}

Jorge P. Zubelli[‡]

November 2, 2006

Abstract

We consider a size-structured model for cell division and address the question of determining the division (birth) rate from the measured stable size distribution of the population. We formulate such question as an inverse problem for an integro-differential equation posed on the half line. We develop firstly a regular dependency theory for the solution in terms of the coefficients and, secondly, a novel regularization technique for tackling this inverse problem which takes into account the specific nature of the equation. Our results rely also on generalized relative entropy estimates and related Poincaré inequalities.

Key-words Size structure models. Inverse problem. Regularization technique. Entropy methods.

AMS Class. Numbers 35B30, 35R30, 92D25

1 Introduction

For many unicellular organisms the mass of the cell, its DNA content, and the level of certain proteins concentrations are often considered as the most relevant parameters for modeling the cell division and, consequently, the population dynamics. Thus, it is natural and usual to consider the evolution of the cell density n as a function of the time t and a size parameter x . See [1, 2, 3] for applications to biology and [4] for other related cases. In many situations it is of paramount importance to determine from observed data the division (birth) rate B as a function of the size parameter x . However, one is faced with the problem that the cell distribution $n(t, x)$ is not easily measured or observed. Instead of that, the dynamics of $n(t, x)$ leads to a stable steady distribution $N = N(x)$, which after a suitable re-normalization, can be observed. In this work we face the problem of determining $B = B(x)$ from noisy and scarce observed data $N = N(x)$.

Age-structured populations also constitute basic models for the division of certain cells, [5, 6]. There has been substantial interest on the inverse problem for structured population models. See for example [7, 8, 9, 10, 11] and references therein. These models typically lead to difficult inverse problems that require a substantial amount of regularization, especially if one is interested in making use of real data. In full generality, size structured models have however a somewhat different mathematical structure (only under the assumption *the smallest daughter cell is larger than half the largest mother cell*, the

^{*}Département de Mathématiques et Applications, École Normale Supérieure, CNRS UMR8553, 45 rue d'Ulm, F 75230 Paris cedex 05, France; email: perthame@dma.ens.fr

[†]INRIA Rocquencourt, Project BANG, BP 105, F78153 Le Chesnay Cedex

[‡]IMPA, Est. D. Castorina 110, RJ 22460-320, Brazil; email: zubelli@impa.br

models lead to similar theory). For this reason, our approach, differently from that of other works we are aware of, makes use of the information given by the so-called *stable size distribution* of the model and relies on a number of recent methods based on generalized relative entropy estimates.

As a starting point, we consider the following size-structured model:

$$\begin{cases} \frac{\partial}{\partial t}n(t, x) + \frac{\partial}{\partial x}[g(x)n(t, x)] + B(x)n(t, x) = 4B(2x)n(t, 2x), & x \geq 0, t \geq 0, \\ n(t, x = 0) = 0, t > 0, \\ n(0, x) = n^0(x) \geq 0. \end{cases} \quad (1)$$

This expresses that the evolution of the cell density results from two effects: At the one hand, the term $\frac{\partial}{\partial x}[g(x)n(t, x)]$ which describes the growth of cells by nutrient uptake with the rate $g(x)$. At the other hand, the terms containing $B(x)$ and $B(2x)$, which describe the division of cells of size $2x$ into two cells of size x . The resulting function-differential equation has a mathematical interest *per se*. One way to understand such interest is that the x -derivative leads to transport to right, whereas the functional term leads to transport to the left. The overall effect is an equilibrium. Another way to understand those effects, in mathematical terms, is to consider the evolution of two macroscopic quantities, the total cell number $N(t) = \int_0^\infty n(t, x)dx$ and the total biomass $M(t) = \int_0^\infty xn(t, x)dx$. Integrating the equation (1) yields

$$\frac{d}{dt}N(t) = \int_0^\infty B(x)n(t, x)dx, \quad (2)$$

this indeed means that number of cells increase only by division, and

$$\frac{d}{dt}M(t) = \int_0^\infty g(x)n(t, x)dx \quad (3)$$

which indeed means that the biomass increases only by nutrient uptake.

For $g \equiv 1$, a case we consider here for simplicity, it was shown in [12, 13] that there exists a unique eigenvalue λ_0 and eigenfunction $N = N(x)$ such that, after a suitable time re-normalization, the solutions of (1) converge to a multiple of N thus given by the solution of the eigenvalue problem

$$\begin{cases} \frac{\partial}{\partial x}N + (\lambda_0 + B(x))N = 4B(2x)N(2x), & x \geq 0, \\ N(x = 0) = 0, \\ N(x) > 0 \text{ for } x > 0, & \int_0^\infty N(x)dx = 1. \end{cases} \quad (4)$$

Namely, we have under fairly general conditions on the coefficients

$$n(t, x)e^{-\lambda_0 t} \xrightarrow[t \rightarrow \infty]{} \rho N(x),$$

in weighted L^p topologies that are related to entropy properties and that will be described later on. Moreover exponential rates have been proved to hold for fairly general rates B ([12, 2]). Such N is therefore the above mentioned *stable size distribution*; it is the distribution observed in practice and available for measurements.

The precise question under consideration in this work is the following:

Inverse Problem 1.1 *How to recover in a stable way the birth rate $B(\cdot)$ from noisy data $N(\cdot)$ and the rate λ_0 ?*

The plan for this work is the following: In Section 2 we present some preliminary remarks on a toy model that describes and motivates our approach to the regularization of the inverse problem under consideration. In Section 3 we collect a number of results concerning the direct problem associated to the model. In particular, concerning the dependence of the solutions to the model with respect to the coefficients. In Section 4 we address the inverse problem and its regularization. The regularization we propose is novel and consists of introducing the regularization parameter α directly in the equation as a coefficient of the perturbed differential equation. We establish a result concerning the strong stability of such perturbed equation and another one demonstrating the consistency of such perturbation when α goes to zero. In Section 5 we obtain a convergence rate for this regularized inverse problem. Some of the proofs of the results in Section 3 are rather technical and we post-pone them to the Appendix A.

2 Preliminaries: A Classical Example Revised

In this section we present a simplified problem which shares some similarity with (4), and allows us to present our strategy for attacking the Inverse Problem 1.1. This problem is the regularization of a classical ill-posed inverse problem, namely recovering a function from its antiderivative, from a slightly different perspective. We believe this will make our approach and our estimates more clear by looking at a specific and simpler well known example. We refer the reader to the books [14, 15] for further information and the classical treatment of the subject.

Inverse problems are characterized by the fact that they are typically ill-posed. Perhaps one of the best well-known examples is that of differentiation. Namely, find a function u from its antiderivative v . We focus on the following:

Inverse Problem 2.1 *Find $u : I \rightarrow \mathbb{R}$ such that*

$$\int_0^x \lambda(s)u(s)ds = v(x) , \quad x \in I \quad (5)$$

where $I \subset \mathbb{R}$ is an interval (finite or infinite) of the line with 0 its left endpoint and $\lambda : I \rightarrow \mathbb{R}_{>0}$ is given weight function.

Equation (5) is formally equivalent to

$$\lambda u = \frac{\partial}{\partial x} v. \quad (6)$$

We consider the problem (5), and notice that the map that sends $v \mapsto u$ is not continuous, when we endow the space of the v 's with say the $L^2(I; dx)$ norm and the space of the u 's with $L^2(I; \lambda(x)dx)$ norm, the choice of spaces being motivated by the specific extension we have in mind. In practice, this reflects itself on the fact that small changes in the measurement of v may lead to huge changes in u . Still, one is forced to face such problem in a number of applications. We now describe some of the main ingredients from inverse problem theory that are used to handle such difficulties as they will be applied to the context of this article. They consist of the following ingredients:

1. Assume some kind of “a priori” bound on the solution u to problem (5), or equivalently on the acceptable data in a sufficiently strong norm.

2. Assume that the measured data v^ϵ satisfies $\|v - v^\epsilon\| \leq \epsilon$ in a suitable weak norm $\|\cdot\|$.
3. Approximate the problem (5) by one that is well-posed for the weak norm and solve the stabilized problem instead of the unstable one choosing the regularization parameter in an optimal way.

Let us illustrate the technique in the problem under consideration. For more information we refer the reader to [14, 15]. Let us consider

$$\begin{cases} \alpha \frac{\partial}{\partial x}(\lambda u_\alpha) + \lambda u_\alpha = \frac{\partial}{\partial x} v , \\ \lambda u_\alpha(0) = 0 \end{cases} \quad (7)$$

In what follows we shall assume that $v(0) = 0$ whenever v is defined in a neighborhood of 0 due to some extra assumptions (such as $\frac{\partial}{\partial x} v \in L^2$).

In order to implement the above program to the problem under consideration, we shall establish the following two inequalities:

Lemma 2.2 *If u_α satisfies (7), then*

$$\|u_\alpha\|_{L^2(I, \lambda^2 dx)} \leq \frac{1}{\alpha} \|v\|_{L^2(I, dx)} . \quad (8)$$

If $\frac{\partial^2}{\partial x^2} v \in L^2(I, dx)$ and u_α satisfies (7), then

$$\|u_\alpha\|_{L^2(I, \lambda^2 dx)} \leq \alpha \left\| \frac{\partial^2}{\partial x^2} v \right\|_{L^2(I, dx)} . \quad (9)$$

As a direct consequence of Equation (9) we have that if u_α is a solution of equation (7) and \bar{u} is a solution of (6), then the difference $u_\alpha - \bar{u}$ satisfies

$$\|u_\alpha - \bar{u}\|_{L^2(I, \lambda^2 dx)} \leq \alpha \left\| \frac{\partial^2}{\partial x^2} v \right\|_{L^2(I, dx)} . \quad (10)$$

We now join the above estimates with the “a priori” assumption that

$$\left\| \frac{\partial^2}{\partial x^2} v \right\|_{L^2(I, dx)} \leq E , \quad (11)$$

and consider a problem with noisy data given by v^ϵ such that

$$\|v - v^\epsilon\|_{L^2(I, dx)} \leq \epsilon . \quad (12)$$

Thus, upon solving the regularized problem (7) with data v^ϵ we are left with a solution u_α^ϵ of

$$\begin{cases} \alpha \frac{\partial}{\partial x}(\lambda u_\alpha^\epsilon) + \lambda u_\alpha^\epsilon = \frac{\partial}{\partial x} v^\epsilon , \\ \lambda u_\alpha^\epsilon(0) = 0 . \end{cases} \quad (13)$$

We further assume that $\frac{\partial}{\partial x} v^\epsilon \in L^2(I, dx)$. Now we consider the distance between u_α^ϵ and \bar{u}

$$\begin{aligned} \|u_\alpha^\epsilon - \bar{u}\|_{L^2(I, \lambda^2 dx)} &\leq \|u_\alpha - \bar{u}\|_{L^2(I, \lambda^2 dx)} + \|u_\alpha^\epsilon - u_\alpha\|_{L^2(I, \lambda^2 dx)} \\ &\leq \alpha \left\| \frac{\partial^2}{\partial x^2} v \right\|_{L^2(I, dx)} + \frac{1}{\alpha} \|v^\epsilon - v\|_{L^2(I, dx)} \\ &\leq \alpha E + \frac{\epsilon}{\alpha} . \end{aligned}$$

Now, we are free to choose α in such a way as to minimize the R.H.S. of the last inequality. Using the elementary fact that $\alpha \mapsto (\alpha E + \epsilon/\alpha)$ has a minimum at $\alpha = \sqrt{E/\epsilon}$ and choosing such optimal $\alpha = \alpha(\epsilon)$ we get that

$$\|u_\alpha^\epsilon - \bar{u}\|_{L^2(I, \lambda^2 dx)} \leq 2\epsilon^{1/2} E^{1/2}. \quad (14)$$

Summing up, if we know an “a priori” bound on the data v and v^ϵ of the form (11) and the data is measured with an accuracy ϵ then we get that the stabilized computed solution is within $\mathcal{O}(\epsilon^{1/2} E^{1/2})$ of the actual solution \bar{u} in the norm of the space $L^2(I, \lambda^2 dx)$.

The reasoning developed above can be pushed in different directions and in the next subsection we will show how it can be extended further.

We remark that Equation (7) is just one of the different possibilities of regularizing the Problem (6). Another possibility would be to solve an equation of the form

$$-\alpha \frac{\partial^2}{\partial x^2} \frac{u_\alpha}{\lambda} + \lambda u_\alpha = \frac{\partial}{\partial x} v \quad (15)$$

subject to appropriate boundary conditions. It is an easy exercise to verify that Equation (15) formally corresponds to Tikhonov regularization of the Inverse Problem 2.1. The form of Equation (15), although very attractive and general is not well adapted to our hyperbolic problem and we do not consider it here.

3 Model Properties and Regular Dependency upon the Coefficients

In this section we shall address a preliminary issue, namely the smoothness of *direct* map

$$B \mapsto (\lambda_0, N)$$

in suitable function spaces. Then, the Inverse Problem 1.1 makes sense in such spaces. This requires some assumptions and preliminary mathematical material that we introduce now.

Firstly, we need some assumptions. Here we make the choice of simplicity, keeping in mind that more realistic hypotheses are possible that would lead to more technical proofs, which we prefer to avoid. The measurable division rates B are supposed to satisfy

$$\exists B_m, B_M \text{ such that } 0 < B_m \leq B(x) \leq B_M, \quad (16)$$

Secondly, because the adjoint equation is as important as the direct eigenvalue problem, we consider both the cell division equation, to find (λ_0, N) solution to

$$\begin{cases} \frac{\partial}{\partial x} N + (\lambda_0 + B(x))N = 4B(2x)N(2x), & x \geq 0, \\ N(x=0) = 0, \\ N(x) > 0 \text{ for } x > 0, & \int_0^\infty N(x)dx = 1. \end{cases} \quad (17)$$

and the adjoint equation

$$\begin{cases} \frac{\partial}{\partial x} \varphi - (\lambda_0 + B(x))\varphi = -2B(x)\varphi(\frac{x}{2}), & x \geq 0, \\ \varphi(x) > 0, & \int_0^\infty \varphi(x)N(x)dx = 1. \end{cases} \quad (18)$$

The solution φ is used in several places because it gives natural bounds for the direct problem. See the proof of Theorem 3.2 and the entropy dissipation in Appendix A.1.

The existence and uniqueness of solutions to these eigenproblems was proved in [12], and it is known that N as well as its derivative vanish at $x = 0$ and $x = \infty$. We also recall the following results that we will use later on.

Theorem 3.1 [12, 16, 2] *Under the assumption (16), the solution N to (17) satisfies the properties*

$$B_m \leq \lambda_0 = \int B(x)N(x)dx \leq B_M, \quad (19)$$

$$\frac{1}{B_M} \leq \int xN(x)dx = \frac{1}{\lambda_0} \leq \frac{1}{B_m}, \quad (20)$$

$$0 \leq N(x) \leq 2B_M, \quad (21)$$

$$e^{ax}N \in L^1 \cap L^\infty(\mathbb{R}^+), \quad \forall a < \lambda_0 + B_m, \quad \varphi(x) \leq C(1+x). \quad (22)$$

Proof. We only prove the estimates on N and refer to [16] for the sub-linearity result on φ and for more accurate results on the behavior of N at infinity, in particular, when B has a limit B_∞ at infinity, one can obtain exponential decay with $a = \lambda_0 + B_\infty$.

The inequalities (19) and (20) are obtained by integrating Equation (17) against respectively the weights dx and $x dx$.

For (21), we use (19), and from the Equation (17), we obtain

$$\frac{\partial}{\partial x}N(x) \leq 4B(2x)N(2x) \implies N(x) \leq 4 \int_0^x B(2y)N(2y)dy \leq 2B_M.$$

For (22), we multiply (17) by e^{ax} and integrate between 0 and ∞ . In fact, a standard truncation argument is needed for complete justification. See [2] for details. We obtain

$$\begin{aligned} \int_0^\infty [\lambda_0 + B(x) - a]e^{ax}N(x)dx &= 4 \int_0^\infty B(2x)N(2x)e^{ax}dx \\ &\leq 2 \int_0^\infty B(x)N(x)e^{ax/2}dx. \end{aligned} \quad (23)$$

With the choice $a = \lambda_0$, since the right hand side is bounded by

$$2 \left(\int_0^\infty B(x)N(x)e^{ax}dx \right)^{1/2} \left(\int_0^\infty B(x)N(x) \right)^{1/2} = 2 \left(\int_0^\infty B(x)N(x)e^{ax}dx \right)^{1/2} \lambda_0^{1/2},$$

we obtain that

$$\int_0^\infty B(x)N(x)e^{\lambda_0 x}dx \leq 4\lambda_0 \leq 4B_M. \quad (24)$$

Then, we can iterate on a and choose any $a < B_m + \lambda_0 \leq 2\lambda_0$ while keeping the right hand side of (23) bounded. This proves the L^1 statement of (22).

It remains to use (17) again to deduce, thanks to the chain rule, that $\frac{\partial}{\partial x}[e^{ax}N(x)] \in L^1(\mathbb{R}^+)$, and the result follows. \square

With these results and methods of proof, we can now state the main new result of this section.

Theorem 3.2 *Under the assumption (16), the map $B \mapsto (\lambda_0, N)$, from $L^\infty(\mathbb{R}^+)$ into $[B_m, B_M] \times L^1 \cap L^\infty(\mathbb{R}^+)$ is*

- (i) *Continuous in B under the weak-* topology of $L^\infty(\mathbb{R}^+)$,*
- (ii) *Locally Lipschitz continuous in B under the strong topology of $L^2(\mathbb{R}^+)$ into $L^2(\mathbb{R}^+)$, namely (26) and (33) below hold, for m large enough,*
- (iii) *Class C^1 in the spaces of statement (ii).*

In statement (i) the weak-* topology of $L^\infty(\mathbb{R}^+)$ is natural from the assumption (16). For estimates (ii) and (iii), we have chosen the L^2 space which is used later on for the inverse problem where it plays an essential role.

Proof. First of all, we introduce some notations that we will use throughout the proof. For two functions B, \bar{B} satisfying (16), we set

$$\delta B = \bar{B} - B, \quad \Delta = \|\bar{B} - B\|_{L^2(\mathbb{R}^+)}. \quad (25)$$

For the corresponding solutions we define the differences

$$\delta N = \bar{N} - N, \quad \delta \lambda = \bar{\lambda}_0 - \lambda_0.$$

(i) This first statement follows thanks to the strong uniform estimates on N in Theorem 3.1. Indeed, consider a sequence B_n that weak-* converges to B and the corresponding solutions N_n to (17). Theorem 3.1 provides us with uniform bounds on N_n and thus with uniform bounds on $\frac{\partial N_n}{\partial x}$. This means that the sequence N_n is relatively strongly compact. Therefore, it converges to the unique N solution of Equation (17) for the limit B .

(ii) Firstly, we estimate the difference between the eigenvalues. As a consequence of (17) for \bar{N} (multiplied by φ) and (18) for φ (multiplied by \bar{N}), we obtain

$$\delta \lambda \int \bar{N} \varphi = \int \delta B \varphi \left(\frac{x}{2}\right) \bar{N} + \int \delta B \varphi \bar{N}.$$

Therefore, according to the *a priori* estimates of Theorem 3.1 on φ and on N ,

$$|\delta \lambda| \leq C(B, \bar{B}) \Delta, \quad (26)$$

with $C(B, \bar{B}) = \|\frac{\varphi}{1+x}\|_{L^\infty(\mathbb{R}^+)} \|(1+x)\bar{N}\|_{L^2(\mathbb{R}^+)}/\int \bar{N} \varphi$.

Secondly, we estimate δN . Writing the difference of the solutions to the cell-division equation, we have

$$\frac{\partial}{\partial x} \delta N + (\lambda_0 + B(x)) \delta N = 4B(2x)\delta N(2x) + \delta R(x), \quad (27)$$

with

$$\delta R(x) = 4\delta B(2x) \bar{N}(2x) + (\delta \lambda + \delta B(x)) \bar{N}(x). \quad (28)$$

We have, since N has exponential decay at infinity using again Theorem 3.2, and for all $m \in \mathbb{N}$,

$$\|\delta R(1 + x^m)\|_{L^2(\mathbb{R}^+)} \leq C(B, \bar{B}) \Delta, \quad (29)$$

for another constant C still controlled by B_m and dB_M . Also, since φ has at most linear growth at infinity, we know that

$$\int_0^\infty |\delta R(x)| \varphi(x) dx \leq C(B, \bar{B}) \Delta. \quad (30)$$

It is also useful to recall that the Equation (27) is spectral (N is the Krein-Rutman eigenvector) and one readily checks the solvability and the uniqueness conditions

$$\int_{\mathbb{R}^+} \varphi \delta R = 0, \quad \int_{\mathbb{R}^+} \delta N = 0. \quad (31)$$

Our local lipschitz regularity result then follows from the two next lemmas

Lemma 3.3 (Entropy dissipation) *Assume that Equation (16) holds and that N and φ are the solutions of (17) and (18). Then, the solution δN to (27), with the conditions (31), satisfies for all convex function $H : \mathbb{R} \rightarrow \mathbb{R}$,*

$$\begin{aligned} \int_0^\infty 4\varphi(x)B(2x)N(2x) \left[H'\left(\frac{\delta N(x)}{N(x)}\right)\left(\frac{\delta N(2x)}{N(2x)} - \frac{\delta N(x)}{N(x)}\right) - H\left(\frac{\delta N(x)}{N(x)}\right) + H\left(\frac{\delta N(2x)}{N(2x)}\right) \right] dx \\ = \int_0^\infty H'\left(\frac{\delta N(x)}{N(x)}\right) \delta R(x) \varphi(x) dx. \end{aligned} \quad (32)$$

This lemma is a variant of the generalized relative entropy inequalities introduced in [17, 13]. Its proof will be presented in the Appendix A. We point out, however, the main difficulty that arises here: Since N and \bar{N} do not have necessarily the same exponential decay at infinity, one cannot use the natural weighted quantities $\int_0^\infty \varphi N \left| \frac{\delta R}{N} \right|^p$ which are not necessarily finite. Therefore, we are sometimes forced to work in L^1 . This type of entropy inequality is related to Poincaré type inequalities which would be a way to conclude the proof at this stage. This area is still a very active field of research for so-called hypocoercive operators. See for example [18]. Here, it does not seem to follow from standard methods and we prefer to prove directly the following consequence of Lemma (3.3)

Lemma 3.4 (Spectral gap) *There is a $\nu > 0$ such that*

$$\nu \|\delta N\|_{L^2(\mathbb{R}^+)} \leq \|\delta R(1 + x^m)\|_{L^2(\mathbb{R}^+)} \leq C\Delta, \quad (33)$$

where the parameter m (large enough) is chosen so that $\lambda_0 > \frac{B_M}{2^{m-1}}$.

We again refer to the Appendix for a proof.

This Lemma expresses the Lipschitz regularity stated in point (ii).

(iii) The third claim is now a general fact on the Fréchet derivatives of mapping with quadratic nonlinearities. Whenever it is Lipschitz continuous on a subset, it is C^1 by usual algebraic manipulations. This concludes the proof of Theorem 3.2. \square

4 The Inverse Problem and its Regularization

The inverse problem consists in finding B from the knowledge of the population growth rate and the cell density (λ_0, N) . If one could guarantee that the measurement N is very smooth one could directly consider to solve the equation on B , equivalent to the cell-division Equation (17) written with $y = 2x$,

$$4B(y)N(y) = B\left(\frac{y}{2}\right)N\left(\frac{y}{2}\right) + \lambda_0 N\left(\frac{y}{2}\right) + 2\frac{\partial}{\partial y}N\left(\frac{y}{2}\right), \quad y > 0. \quad (34)$$

This is a well-posed equation on B as long as N satisfies regularity properties such as $\frac{\partial}{\partial y}N\left(\frac{y}{2}\right) \in L^p$ for some $p \geq 1$. But our interest lies on the fact that fluctuations on N make it a mere L^p function because, given a set of measurements on N , one does not have a way of controlling the precision of the measurements on $\frac{\partial}{\partial y}N$. This motivates working with a regularized problem

$$\begin{cases} \alpha \frac{\partial}{\partial y}(B_\alpha N) + 4B_\alpha(y)N(y) = B_\alpha\left(\frac{y}{2}\right)N\left(\frac{y}{2}\right) + \lambda_0 N\left(\frac{y}{2}\right) + 2\frac{\partial}{\partial y}N\left(\frac{y}{2}\right), & y > 0, \\ B_\alpha N(0) = 0, \end{cases} \quad (35)$$

where $0 < \alpha < 1$ is a small parameter adapted to the level of noise as explained in Section 2 for the toy model. Notice that this is still a well-posed problem in appropriate function spaces when written with a general source $F(y)$

$$\begin{cases} \alpha \frac{\partial}{\partial y}(B_\alpha N) + 4B_\alpha(y)N(y) = B_\alpha\left(\frac{y}{2}\right)N\left(\frac{y}{2}\right) + F(y), & y > 0, \\ B_\alpha N(0) = 0, \end{cases} \quad (36)$$

Before we analyze the issue of estimating the inverse problem, let us explain why this is indeed a well-posed problem and for simplicity we restrict ourselves to L^2 . In particular, we did not impose boundary conditions because we expect that the point $x = 0$ is characteristic since the cell density satisfies

$$N \in H^2(\mathbb{R}^+), \quad N(x) > 0 \quad \text{for } x > 0, \quad N(0) = 0 \quad \frac{\partial}{\partial x}N(0) = 0. \quad (37)$$

Theorem 4.1 (Strong stability) *Assume (37) and that $F \in L^2$. Then, there is a unique solution to (36) such that $BN \in H^1$ with*

$$\alpha |B_\alpha N(y)|^2 + \int |B_\alpha N|^2 \leq C \int |F|^2, \quad \forall y \geq 0,$$

$$\alpha^2 \int \left| \frac{\partial}{\partial y} B_\alpha N \right|^2 \leq C \int |F|^2,$$

and

$$\int \left| \frac{\partial}{\partial y} B_\alpha N \right|^2 \leq C \int \left| \frac{\partial}{\partial y} F \right|^2 \quad \text{if } F(0) = 0.$$

Proof. We drop the index α in order to simplify the notation. This model is a variant of the cell division equation and the Cauchy-Lipschitz theory applies and gives the existence for small y . The

important point is to establish the *a priori* bounds. For the first set of estimates, we multiply the Equation (36) by BN and integrate from 0 to y . This yields

$$\begin{aligned} & \frac{\alpha}{2}|BN(y)|^2 + 4 \int_0^y |BN|^2(s) ds \\ &= \int_0^y (BN)(z) (BN)\left(\frac{z}{2}\right) dz + \int_0^y (BN)(z) F(z) dz \\ &\leq \frac{1}{2} \int_0^y |BN|^2 + \frac{1}{2} \int_0^y |BN\left(\frac{z}{2}\right)|^2 dz + \frac{1}{2} \int_0^y |BN|^2 + \frac{1}{2} \int_0^y |F(z)|^2 dz. \end{aligned}$$

Altogether, this gives the two bounds claimed in the first statement above.

The second estimate, on $\alpha \int |\frac{\partial}{\partial y} BN|^2$, is then a consequence of the first two using Equation (36) once again.

For the third inequality, we differentiate the equation and write $Q = \frac{\partial}{\partial y} BN$. We find

$$\alpha \frac{\partial}{\partial y} Q + 4Q(y) = \frac{1}{2} Q\left(\frac{y}{2}\right) + \frac{\partial}{\partial y} F.$$

Therefore, multiplying by Q and integrating by parts gives

$$4 \int_0^\infty Q^2 \leq \frac{\alpha}{2} Q^2(0) + \frac{1}{2} \int_0^\infty Q(y) Q\left(\frac{y}{2}\right) dy + \int_0^\infty Q(y) \frac{\partial}{\partial y} F(y) dy,$$

and thus

$$4 \int_0^\infty Q^2 \leq \frac{\alpha}{2} Q^2(0) + \frac{1}{4} \int_0^\infty Q^2 + \frac{1}{4} \int_0^\infty Q^2\left(\frac{y}{2}\right) dy + \frac{1}{2} \int_0^\infty Q^2 + \frac{1}{2} \int_0^\infty \left|\frac{\partial}{\partial y} F\right|^2.$$

Altogether this gives

$$\frac{11}{4} \int_0^\infty Q^2 \leq \frac{\alpha}{2} Q^2(0) + \int_0^\infty \left|\frac{\partial}{\partial y} F\right|^2.$$

It remains to notice that from the Equation (36) on BN and the condition $N(0) = 0$ we have also $\alpha Q(0) = \alpha \frac{\partial}{\partial y} BN(0) = F(0)$ and the result is proved. \square

The first question we want to answer is how much this regularization on B differs from its exact value

Theorem 4.2 (Consistency) *Assume (37), then the solutions to (34) and (35) satisfy*

$$\int |B_\alpha - B|^2 N \leq C\alpha^2 \int \left[\left|\frac{\partial}{\partial y} N\right|^2 + \left|\frac{\partial^2}{\partial y^2} N\right|^2 \right].$$

Proof. We write $\delta B = B_\alpha - B$ and we have

$$\alpha \frac{\partial}{\partial y} (\delta B N) + 4\delta B(y) N(y) = \delta B\left(\frac{y}{2}\right) N\left(\frac{y}{2}\right) - \alpha \frac{\partial}{\partial y} (BN).$$

This is again an equation of the form (36) with source term

$$F = -\alpha \frac{\partial}{\partial y}(BN)$$

and thus we can apply the first estimate of Theorem 4.1 and obtain

$$\int |\delta B \ N|^2 \leq C\alpha^2 \int \left| \frac{\partial}{\partial y}(BN) \right|^2.$$

Next we apply the second estimate of Theorem 4.1 to (34) with $F = \lambda_0 N + \frac{\partial}{\partial y}N$ and obtain

$$\int \left| \frac{\partial}{\partial y}(BN) \right|^2 \leq C \int [|\frac{\partial}{\partial y}N|^2 + |\frac{\partial^2}{\partial y^2}N|^2].$$

Altogether these two estimates give the announced inequality. \square

5 Recovering Estimates for the Regularized Inverse Problem

Consider now a smooth solution (λ_0, N) to Equation (18) corresponding to division rate B , that we want to recover from a noisy measurement N_ε with

$$\int |N - N_\varepsilon|^2 \leq \varepsilon^2, \quad N_- \leq N_\varepsilon \leq N_+, \quad (38)$$

where N_\pm are two smooth functions that can serve as a filter for unrealistic data. We assume their behavior contains possible properties of the true solution and serves to assert the boundary condition is well defined with $N_+(0) = N_+(\infty) = 0$. Notice $N_- = 0$ is a possibility. The regularized method is aimed at furnishing an approximation $B_{\varepsilon, \alpha}$ of the exact coefficient B through Equation (35) with N_ε in place of N ,

$$\begin{cases} \alpha \frac{\partial}{\partial y}(B_{\varepsilon, \alpha} N_\varepsilon) + 4B_{\varepsilon, \alpha}(y)N_\varepsilon(y) = B_{\varepsilon, \alpha}\left(\frac{y}{2}\right)N_\varepsilon\left(\frac{y}{2}\right) + \lambda_0 N\left(\frac{y}{2}\right) + 2\frac{\partial}{\partial y}N_\varepsilon\left(\frac{y}{2}\right), & y > 0, \\ (B_{\varepsilon, \alpha} N_\varepsilon)(0) = 0. \end{cases} \quad (39)$$

We prove the following error estimate on the coefficient recovered by the above regularization procedure

Theorem 5.1 (Convergence rate) *Assume (37) and (38), then the solutions B to (34) and $B_{\varepsilon, \alpha}$ to (39), satisfy the error estimate,*

$$\|B_{\varepsilon, \alpha} - B\|_{L^2(N_\varepsilon^2 dx)} \leq C\alpha \|N\|_{H^2} + \frac{C + \|B\|_{L^\infty}}{\alpha} \|N_\varepsilon - N\|_{L^2}.$$

This theorem relies on a first estimate which expresses weak stability and improves Theorem 4.1 using the special structure in the RHS of Equation (39). Namely, we have

Proposition 5.2 (Weak stability) Assume (37) and (38), given two solutions B_α to (35) and (39) satisfy

$$\int |B_{\varepsilon,\alpha}N_\varepsilon - B_\alpha N|^2 \leq \frac{C}{\alpha^2} \|N_\varepsilon - N\|_{L^2}^2.$$

Proof. We set $Q = B_{\varepsilon,\alpha}N_{\varepsilon,\alpha} - BN$, $R = N_\varepsilon - N$ which satisfies

$$\alpha \frac{\partial}{\partial y} Q + 4Q(y) = Q\left(\frac{y}{2}\right) + \lambda_0 R + 2 \frac{\partial}{\partial y} R\left(\frac{y}{2}\right).$$

Then, we obtain after multiplication by Q and integration

$$\begin{aligned} \frac{\alpha}{2} [Q^2(y) - Q^2(0)] + 4 \int_0^y |Q(z)|^2 dz &= \int_0^y Q(z) Q\left(\frac{z}{2}\right) dz + \\ &\quad \lambda_0 \int_0^y QR - 2 \int_0^y R\left(\frac{z}{2}\right) \frac{\partial}{\partial y} Q + 2R\left(\frac{y}{2}\right) Q(y) - 2QR(0). \end{aligned}$$

Using again the equation on Q (and the Cauchy-Schwarz inequality) we find

$$\begin{aligned} \frac{\alpha}{2} [Q^2(y) - Q^2(0)] + 2 \int_0^y |Q(z)|^2 dz &\leq C \int_0^y |R|^2 + 2Q(y)R\left(\frac{y}{2}\right) - 2QR(0) \\ &\quad - \frac{2}{\alpha} \int_0^y R\left(\frac{z}{2}\right) [Q\left(\frac{z}{2}\right) + \lambda_0 R(z) + \\ &\quad \frac{\partial}{\partial y} R\left(\frac{z}{2}\right) - 4Q(z)] dy. \end{aligned}$$

Therefore

$$\frac{\alpha}{2} |Q(y) - \frac{2}{\alpha} R\left(\frac{y}{2}\right)|^2 + \int_0^y |Q(z)|^2 dz \leq \frac{\alpha}{2} |Q(0) - \frac{2}{\alpha} R(0)|^2 + \frac{C}{\alpha^2} \int_0^y |R|^2.$$

Notice that even though R is not defined point-wise, $Q(y) - \frac{2}{\alpha} R\left(\frac{y}{2}\right)$ is well defined (as a H^1 function) and it vanishes due to (38). \square

We are now ready to prove our main result, Theorem 5.1. We write

$$\begin{aligned} \|B_{\varepsilon,\alpha} - B\|_{L^2(N_\varepsilon^2 dx)}^2 &\leq 2 \int |B_{\varepsilon,\alpha}N_\varepsilon - BN|^2 + 2 \int |BN_\varepsilon - BN|^2 \\ &\leq 4 \int |B_{\varepsilon,\alpha}N_\varepsilon - B_\alpha N|^2 + 4 \int |B_\alpha N - BN|^2 + 2 \int |BN_\varepsilon - BN|^2. \end{aligned}$$

The first term is controlled thanks to Proposition 5.2 and gives the second error term in the estimate of Theorem 5.1. The second term is controlled because of Theorem 4.2 and gives the first error term. The third term gives the last contribution once $\|B\|_{L^\infty}$ has been factored out and the Theorem 5.1 is proved. \square

6 Conclusion and Suggestions for Further Research

We have proposed a non-standard regularization method for the Inverse Problem associated with the cell-division equations. It is based on a first order operator that leads us to solving a new equation whose structure is very close to the equation itself (but for a different unknown). This allowed us to develop a consistency and convergence analysis that can be used in practice because the outcome are

standard inequalities taking into account the noise in the measured data.

Our theoretical approach has been focused on the equal mitosis, when mother cells divide into two equal daughter cells. A first extension of the method would be to deal with more general division equations as

$$\begin{cases} \frac{\partial}{\partial t}n(t, x) + \frac{\partial}{\partial x}[g(x)n(t, x)] + B(x)n(t, x) = 2 \int_0^\infty B(y)\beta(\frac{x}{y})n(t, y)\frac{dy}{y}, & x \geq 0, t \geq 0, \\ n(t, x = 0) = 0, t > 0. \end{cases}$$

The function β represents now the repartition of daughter cell sizes and a natural inverse problem would be to recover the division rate B from measurements of the cell density N once β is known. In principle, our method can be extended but technical estimates have to be reformulated in this new context.

The numerical validation of the procedure proposed here is straightforward because it relies on a regularized equation that has a standard form and numerical methods are available. As far as validation on real data is concerned, let us point out that present experimental devices allow us to measure not only the size repartitions in a cellular culture, but also the molecular content of certain representative proteins. Mathematical models describing the division process based on molecular contents are being developed. To our knowledge, identification of coefficients in this context is still largely open.

Acknowledgments

J.P.Z. acknowledges financial support from CNPq, grants 302161/2003-1 and 474085/2003-1. This research was made possible through the Brazil-France cooperation program, which is gratefully acknowledged by both authors.

A Appendix

This appendix is devoted to the proof of two technical lemmas that were used in the Section 3 to obtain Lipschitz regularity of the solution to the eigenproblem for the cell division equation.

A.1 Generalized Relative Entropy. Proof of Lemma 3.3.

Using (27) and (17), we have

$$\frac{\partial}{\partial x} \frac{\delta N(x)}{N(x)} = 4B(2x) \frac{N(2x)}{N(x)} \left[\frac{\delta N(2x)}{N(2x)} - \frac{\delta N(x)}{N(x)} \right] + \frac{\delta R(x)}{N(x)}, \quad (40)$$

and thus, for $H(\cdot)$ a convex function,

$$\frac{\partial}{\partial x} H\left(\frac{\delta N(x)}{N(x)}\right) = 4B(2x) \frac{N(2x)}{N(x)} H'\left(\frac{\delta N(x)}{N(x)}\right) \left[\frac{\delta N(2x)}{N(2x)} - \frac{\delta N(x)}{N(x)} \right] + H'\left(\frac{\delta N(x)}{N(x)}\right) \frac{\delta R(x)}{N(x)}.$$

On the other hand, combining Equations (17) and (18), we have

$$\frac{\partial}{\partial x} \left(N(x) \varphi(x) \right) = 4\varphi(x)B(2x)N(2x) - 2N(x)B(x)\varphi\left(\frac{x}{2}\right).$$

Therefore,

$$\begin{aligned}
\frac{\partial}{\partial x} [N(x)\varphi(x)H(\frac{\delta N(x)}{N(x)})] &= 4B(2x)N(2x)\varphi(x)H'(\frac{\delta N(x)}{N(x)})[\frac{\delta N(2x)}{N(2x)} - \frac{\delta N(x)}{N(x)}] \\
&+ [4\varphi(x)B(2x)N(2x) - 2N(x)B(x)\varphi(\frac{x}{2})]H(\frac{\delta N(x)}{N(x)}) \\
&+ H'(\frac{\delta N(x)}{N(x)})\delta R(x)\varphi(x).
\end{aligned}$$

After integration in x we arrive at

$$\begin{aligned}
0 &= \int_0^\infty 4\varphi(x)B(2x)N(2x)H'(\frac{\delta N(x)}{N(x)})[\frac{\delta N(2x)}{N(2x)} - \frac{\delta N(x)}{N(x)}]dx \\
&+ \int_0^\infty 4\varphi(x)B(2x)N(2x)[H(\frac{\delta N(x)}{N(x)}) - H(\frac{\delta N(2x)}{N(2x)})]dx \\
&+ \int_0^\infty H'(\frac{\delta N(x)}{N(x)})\delta R(x)\varphi(x)dx.
\end{aligned}$$

Once the terms are reorganized, this is exactly the statement of Lemma 3.3. \square

A.2 A Poincaré-Type Inequality. Proof of Lemma 3.4.

We argue by contradiction and assume that for a sequence $\bar{B}_k \rightarrow B$, the corresponding ν_k vanishes. Then, there is a family of $\delta R_k, \delta N_k$ that satisfy

$$\nu_k \|\delta N_k\|_{L^2(\mathbb{R}^+)} \geq \|\delta R_k (1 + x^m)\|_{L^2(\mathbb{R}^+)}. \quad (41)$$

After re-normalizing (by multiplication), we can always assume that

$$\|\delta N_k\|_{L^2(\mathbb{R}^+)} = 1, \quad \|\delta R_k(1 + x^m)\|_{L^2(\mathbb{R}^+)} \xrightarrow{k \rightarrow \infty} 0.$$

Then, our proof consists in several steps: First, we prove compactness and then pass to the limit, next we use the generalized relative entropy to identify the limiting solution and finally, prove a contradiction.

Compactness. Notice that the Equation (27) automatically implies that (recall $\varphi > 0$ is smooth and sublinear at infinity), that δN_k is bounded in $H^1(\mathbb{R}^+)$. This provides us with local compactness. It remains to obtain a control at infinity; we also notice that

$$\frac{1}{2} \frac{\partial}{\partial x} (\delta N_k)^2 + (\lambda_0 + B(x)) (\delta N_k)^2 = 4B(2x)\delta N_k(2x) \delta N_k(x) + \delta R_k(x) \delta N_k(x),$$

which, after integration in x with the weight x^m gives for all $a > 0$

$$\begin{aligned}
-\frac{m}{2} \int_{\mathbb{R}^+} x^{m-1} (\delta N_k)^2 + \int_{\mathbb{R}^+} x^m (\lambda_0 + B(x)) (\delta N_k)^2 &\leq 4 \int_{\mathbb{R}^+} x^m B(2x) [a \delta N_k(2x)^2 + \frac{1}{a} \delta N_k(x)^2] \\
&+ \sqrt{\int_{\mathbb{R}^+} x^m (\delta N_k)^2 \int_{\mathbb{R}^+} x^m (\delta R_k)^2}.
\end{aligned}$$

Therefore, we also obtain

$$\begin{aligned} \int_{\mathbb{R}^+} x^m (\lambda_0 + B(x)) (\delta N_k)^2 &\leq \frac{m}{2} \int_{\mathbb{R}^+} x^{m-1} (\delta N_k)^2 + a 2^{-m+1} \int_{\mathbb{R}^+} x^m B(x) \delta N_k(x)^2 \\ &\quad + \frac{B_M}{a} \int_{\mathbb{R}^+} x^m \delta N_k(x)^2 + \sqrt{\int_{\mathbb{R}^+} x^m (\delta N_k)^2 \int_{\mathbb{R}^+} x^m (\delta R_k)^2}. \end{aligned}$$

With $a = 2^{m-1}$, we conclude that

$$\begin{aligned} \lambda_0 \int_{\mathbb{R}^+} x^m (\delta N_k)^2 &\leq \frac{m}{2} \int_{\mathbb{R}^+} x^{m-1} (\delta N_k)^2 + \frac{B_M}{a} \int_{\mathbb{R}^+} x^m (\delta N_k)^2 + \sqrt{\int_{\mathbb{R}^+} x^m (\delta N_k)^2 \int_{\mathbb{R}^+} x^m (\delta R_k)^2} \\ &\leq C \int_{\mathbb{R}^+} (\delta N_k)^2 + \left(\frac{\lambda_0}{2} + \frac{B_M}{a}\right) \int_{\mathbb{R}^+} x^m (\delta N_k)^2 + \sqrt{\int_{\mathbb{R}^+} x^m (\delta N_k)^2 \int_{\mathbb{R}^+} x^m (\delta R_k)^2}. \end{aligned}$$

It remains to choose m large enough so that $\lambda_0 > 2 \frac{B_M}{a}$ and we conclude on a uniform control

$$\int_{\mathbb{R}^+} x^m (\delta N_k)^2 \leq C \left[1 + \int_{\mathbb{R}^+} x^m (\delta R_k)^2 \right], \quad (42)$$

that concludes the global compactness in $L^2(\mathbb{R}^+)$ of the sequence δN_k .

Limit. Therefore, due to the above facts and using Lemma 3.3, we may pass to the strong limit and obtain

$$\begin{aligned} \delta N_k &\rightarrow n \in H^1(\mathbb{R}^+) \quad x^m n \in L^2(\mathbb{R}^+), \quad \forall m > 0, \\ \frac{\partial}{\partial x} n + (\lambda_0 + B(x)) n &= 4B(2x)n(2x), \quad \int_{\mathbb{R}^+} \delta n = 0. \end{aligned}$$

Notice that, thanks to the moment estimate, we also have for all $m > 0$,

$$(1 + x^m)n \in L^1(\mathbb{R}^+), \text{ and } \int_{\mathbb{R}^+} |n| \varphi < \infty.$$

This allows us to give a meaning to the vanishing integral. This follows from the Cauchy-Schwarz inequality

$$\int_{\mathbb{R}^+} (1 + x^m) |n| \leq \left(\int_{\mathbb{R}^+} (1 + x^m)^2 (1 + |x^2|) |n|^2(x) dx \int_{\mathbb{R}^+} \frac{dx}{1 + x^2} \right)^{1/2}.$$

Application of the GRE. We can now choose the family of convex functions $H(u) = (u - \xi)_+$ in the Generalized Relative Entropy of Lemma 3.3, with $\xi > 0$ a parameter. We find, because we now deal with $\delta R = 0$, after reorganizing the terms in (32), that

$$\int_0^\infty 4\varphi(x) B(2x) N(2x) \left(\frac{n(2x)}{N(2x)} - \frac{n(x)}{N(x)} \right) \left(\left(\text{sgn}_+ \left(\frac{n(x)}{N(x)} - \xi \right) - \text{sgn}_+ \left(\frac{n(2x)}{N(2x)} - \xi \right) \right) dx = 0. \right)$$

In fact, one needs to justify it by a preliminary truncation of the integral at $x = 0$ and $x \approx \infty$, we leave to the reader the corresponding analysis. Again, let us point out that in L^1 this is possible because the corresponding x -derivative term

$$[\varphi(x) N(x) \left(\frac{n(x)}{N(x)} - \xi \right)_+]$$

is integrable (but for the square entropy we arrive at $[\varphi(x)N(x)(\frac{n(x)}{N(x)})^2]$ which is not integrable!).

Conclusion. From the previous step, we conclude that, for all $x > 0$, $\xi > 0$, we have

$$\operatorname{sgn}_+(\frac{n(x)}{N(x)} - \xi) = \operatorname{sgn}_+(\frac{n(2x)}{N(2x)} - \xi),$$

which implies $\frac{n(x)}{N(x)} = \frac{n(2x)}{N(2x)}$ and thus, tells us that the limit satisfies

$$\frac{\partial}{\partial x} \frac{n(x)}{N(x)} = 0.$$

From $\frac{n(x)}{N(x)} = \frac{n(2x)}{N(2x)}$, we conclude that $n = CN$ and the vanishing total integral of n implies that $n = 0$. This gives the sought contradiction and concludes the proof of Lemma 3.4. \square

References

- [1] J. A. J. Metz and O. Diekmann. Formulating models for structured populations. In *The dynamics of physiologically structured populations (Amsterdam, 1983)*, volume 68 of *Lecture Notes in Biomath.*, pages 78–135. Springer, Berlin, 1986.
- [2] Benoît Perthame. Transport equations arising in biology. In *Frontiers in Mathematics*, Frontiers in Mathematics. Birkhauser, 2006.
- [3] Britta Basse, Bruce C. Baguley, Elaine S. Marshall, Wayne R. Joseph, Bruce van Brunt, Graeme Wake, and David J. N. Wall. A mathematical model for analysis of the cell cycle in cell lines derived from human tumors. *J. Math. Biol.*, 47(4):295–312, 2003.
- [4] Jean Bertoin. On small masses in self-similar fragmentations. *Stochastic Process. Appl.*, 109(1):13–22, 2004.
- [5] G. Chiorino, J. A. J. Metz, D. Tomasoni, and P. Ubezio. Desynchronization rate in cell populations: mathematical modeling and experimental data. *J. Theor. Biol.*, 208:185–199, 2001.
- [6] O. Arino, E. Sánchez, and G. F. Webb. Necessary and sufficient conditions for asynchronous exponential growth in age structured cell population with quiescence. *J. Math. Anal. and Appl.*, 215:499–513, 1997.
- [7] Heinz W. Engl, William Rundell, and Otmar Scherzer. A regularization scheme for an inverse problem in age-structured populations. *J. Math. Anal. Appl.*, 182(3):658–679, 1994.
- [8] Mats Gyllenberg, Andrei Osipov, and Lassi Päiväranta. The inverse problem of linear age-structured population dynamics. *J. Evol. Equ.*, 2(2):223–239, 2002.
- [9] William Rundell. Determining the birth function for an age structured population. *Math. Population Stud.*, 1(4):377–395, 397, 1989.
- [10] J. A. J. Metz and O. Diekmann, editors. *The dynamics of physiologically structured populations*, volume 68 of *Lecture Notes in Biomathematics*. Springer-Verlag, Berlin, 1986. Papers from the colloquium held in Amsterdam, 1983.

- [11] Michael Pilant and William Rundell. Determining a coefficient in a first-order hyperbolic equation. *SIAM J. Appl. Math.*, 51(2):494–506, 1991.
- [12] Benoît Perthame and Lenya Ryzhik. Exponential decay for the fragmentation or cell-division equation. *J. Differential Equations*, 210(1):155–177, 2005.
- [13] Philippe Michel, Stéphane Mischler, and Benoît Perthame. General entropy equations for structured population models and scattering. *C. R. Math. Acad. Sci. Paris*, 338(9):697–702, 2004.
- [14] Johann Baumeister and Antonio Leitão. *Topics in inverse problems*. Publicações Matemáticas do IMPA. [IMPA Mathematical Publications]. Instituto Nacional de Matemática Pura e Aplicada (IMPA), Rio de Janeiro, 2005. 25º Colóquio Brasileiro de Matemática. [25th Brazilian Mathematics Colloquium].
- [15] Heinz W. Engl, Martin Hanke, and Andreas Neubauer. *Regularization of inverse problems*, volume 375 of *Mathematics and its Applications*. Kluwer Academic Publishers Group, Dordrecht, 1996.
- [16] Philippe Michel. Existence of a solution to the cell division eigenproblem. *Model. Math. Meth. Appl. Sci.*, 16(suppl. issue 1):1125–1153, 2006.
- [17] Stéphane Mischler, Benoît Perthame, and Lenya Ryzhik. Stability in a nonlinear population maturation model. *Math. Models Methods Appl. Sci.*, 12(12):1751–1772, 2002.
- [18] Cédric Villani. Hypocoercive diffusion operators. Invited Lectures ICM, 2006.