



# The Brouwer Lecture 2005: Statistical estimation with model selection

Lucien Birgé

► **To cite this version:**

Lucien Birgé. The Brouwer Lecture 2005: Statistical estimation with model selection. 2006. <hal-00067763>

**HAL Id: hal-00067763**

**<https://hal.archives-ouvertes.fr/hal-00067763>**

Submitted on 8 May 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# The Brouwer Lecture 2005

## Statistical estimation with model selection

Lucien Birgé

Université Paris VI

Laboratoire de Probabilités et Modèles Aléatoires

U.M.R. C.N.R.S. 7599

21/03/2006

### Abstract

The purpose of this paper is to explain the interest and importance of (approximate) models and model selection in Statistics. Starting from the very elementary example of histograms we present a general notion of finite dimensional model for statistical estimation and we explain what type of risk bounds can be expected from the use of one such model. We then give the performance of suitable model selection procedures from a family of such models. We illustrate our point of view by two main examples: the choice of a partition for designing a histogram from an  $n$ -sample and the problem of variable selection in the context of Gaussian regression.

## 1 Introduction: a story of histograms

### 1.1 Histograms as graphical tools

Assume we are given a (large) set of real valued measurements or data  $x_1, \dots, x_n$ , corresponding to lifetimes of some human beings in a specific area, or lifetimes of some manufactured goods, or to the annual income of families in some country,  $\dots$ . Such measurements have a bounded range  $[a, b]$  which is often known in advance (for instance  $[0, 120]$  would do for lifetimes of human beings) or can be extrapolated from the data using the extreme values. By a proper affine transformation this range can be transformed to  $[0, 1]$ , which we shall assume here, for the simplicity of our presentation. To represent in a convenient, simplified, but suggestive way, this set of data, it is common to use what is called a *histogram*. To design a histogram, one first chooses some finite partition  $m = \{I_0, \dots, I_D\}$  ( $D \in \mathbb{N}$ ) of  $[0, 1]$  into intervals  $I_j$ , generated by an increasing sequence of endpoints  $y_0 = 0 < y_1 < \dots < y_{D+1} = 1$  so that  $I_j = [y_j, y_{j+1})$  for  $0 \leq j < D$  and  $I_D = [y_D, y_{D+1}]$ . Then, for each  $j$ , one computes the number  $n_j$  of observations falling in  $I_j$  and one represents the data set by the piecewise constant function  $\hat{s}_m$  defined on  $[0, 1]$  by

$$\hat{s}_m(x) = \sum_{j=0}^D \frac{n_j}{n|I_j|} \mathbb{1}_{I_j}(x) \quad \text{with } n_j = \sum_{i=1}^n \mathbb{1}_{I_j}(x_i) \quad \text{and} \quad |I_j| = y_{j+1} - y_j. \quad (1.1)$$

Any such histogram  $\hat{s}_m$  provides a summary of the data with three obvious properties. It is nonnegative; its integral is equal to one ( $\int_0^1 \hat{s}_m(x) dx = 1$ ) and it belongs to the  $(D + 1)$ -dimensional linear space  $V_m$  of piecewise constant functions built on the partition  $m$ , i.e.

$$V_m = \left\{ t = \sum_{j=0}^D a_j \mathbb{1}_{I_j} \mid a_0, \dots, a_D \in \mathbb{R} \right\}. \quad (1.2)$$

If the points  $y_j$  are equispaced, i.e. all intervals  $I_j$  have the same length  $(D + 1)^{-1}$ , the partition and the histogram are called *regular*. If  $D \geq 1$  and all intervals do not have the same length, the partition is called *irregular*.

Even within this very elementary framework, some questions are in order: what is a “good” partition, i.e. how can one measure the quality of the representation of the data by a histogram, and how can one choose such a good partition? One can easily figure out that a partition with too few intervals, as compared with  $n$ , will lead to an uninformative representation. Alternatively, if there are too few data per interval the histogram may be quite erratic and meaningless. But these are purely qualitative properties which cannot lead to a sound criterion of quality for a partition which could be used to choose a proper one.

## 1.2 Histograms as density estimators

### 1.2.1 The stochastic point of view

To go further with this analysis, we have to put the whole thing into a more mathematical framework and a convenient one, for this type of problem, is of statistical nature. In many situations, our data  $x_i$  can be considered as successive observations of some random phenomenon which means that  $x_i = X_i(\omega)$  is the realization of a random variable  $X_i$  from some probability space  $(\Omega, \mathcal{A}, \mathbb{P})$  with values in  $[0, 1]$  (with its Borel  $\sigma$ -algebra). If we assume that the random phenomenon was stable during the observation period and the measurements were done independently of each other, the random variables  $X_i$  can be considered as i.i.d. (independent and identically distributed) with common distribution  $Q$  so that

$$\mathbb{P}[\{\omega \in \Omega \mid X_1(\omega) \in A_1, \dots, X_n(\omega) \in A_n\}] = \prod_{i=1}^n Q(A_i),$$

for any family of Borel sets  $A_1, \dots, A_n \subset [0, 1]$ . Such assumptions are justified (at least approximately) in many practical situations and  $(X_1, \dots, X_n)$  is then called an  $n$ -sample from the distribution  $Q$ .

With this new probabilistic interpretation,  $\hat{s}_m = \hat{s}_m(x, \omega)$  becomes a random function, more precisely a random element of  $V_m$ , and (1.1) becomes

$$\hat{s}_m(x, \omega) = \sum_{j=0}^D \frac{N_j(\omega)}{n|I_j|} \mathbb{1}_{I_j}(x) \quad \text{with} \quad N_j(\omega) = \sum_{i=1}^n \mathbb{1}_{I_j}(X_i(\omega)). \quad (1.3)$$

From now on, following the probabilistic tradition, we shall, most of the time, omit the variable  $\omega$  when dealing with random elements.

It follows from (1.3) that the random variables  $N_j$  are binomial random variables with parameters  $n$  and  $p_j = Q(I_j)$  and, if we assume that  $Q$  has a density  $s$  with

respect to the Lebesgue measure on  $[0, 1]$ , then  $p_j = \int_{I_j} s(x) dx$ . If  $s$  also belongs to  $\mathbb{L}_2([0, 1], dx)$ , the piecewise constant element  $s_m = \sum_{j=0}^D p_j |I_j|^{-1} \mathbb{1}_{I_j}$  of  $\mathbb{L}_\infty([0, 1], dx)$  is the orthogonal projection of  $s$  onto  $V_m$  and

$$p_j = \int_{I_j} s_m(x) dx \quad \text{and} \quad \|s - \hat{s}_m\|^2 = \|s - s_m\|^2 + \|s_m - \hat{s}_m\|^2, \quad (1.4)$$

where  $\|t\|$  denotes the  $\mathbb{L}_2$ -norm of  $t$ .

### 1.2.2 Density estimators and their risk

From a practical point of view, even if it is reasonable to assume that the variables  $X_i$  are i.i.d. with distribution  $Q$  and density  $s = dQ/dx$ , this distribution is typically unknown and its density as well and it is often useful, in order to have an idea of the stochastic nature of the phenomenon that produced the data, to get as much information as possible about the unknown density  $s$ . For instance, comparing the shapes of lifetime densities among different populations or their evolution with time brings much more information than merely comparing the corresponding expected lifetimes. The very purpose of Statistics is to derive information about the deterministic, but unknown, parameter  $s$  from the stochastic, but observable, data  $X_i(\omega)$ . In our problem,  $\hat{s}_m$ , which is a density, can be viewed as a random approximation of  $s$  solely based on the available information provided by the sample  $X_1, \dots, X_n$ , i.e., in statistical language, an *estimator* of  $s$ . The distortion of the estimated density  $\hat{s}_m$  from the true density  $s$  can be measured by the quantity  $\|s - \hat{s}_m\|^2$ . It is clearly not the only way but this one, as seen from (1.4), has the advantage of simplicity. Note that  $\|s - \hat{s}_m\|^2$  is a random quantity depending on  $\omega$  as  $\hat{s}_m$  does. In order to average out this randomness, the statisticians often consider, as a measure of the quality of the estimator  $\hat{s}_m$ , its *risk at  $s$*  which is the expectation of the distortion  $\|s - \hat{s}_m\|^2$  given by

$$R(\hat{s}_m, s) = \mathbb{E}_s \left[ \|s - \hat{s}_m\|^2 \right] = \int \|s - \hat{s}_m(\omega)\|^2 d\mathbb{P}_s(\omega).$$

Here  $\mathbb{P}_s$  and  $\mathbb{E}_s$  respectively denote the probability and the expectation of functions of  $X_1, \dots, X_n$  when these variables are i.i.d. with density  $s$ . Of course, due to randomness,  $R(\hat{s}_m, s)$  does not provide any information on the actual distortion  $\|s - \hat{s}_m(\omega)\|^2$  in our experiment. But, by the law of large numbers, it provides a good approximation of the average distortion one would get if one iterated many times the procedure of drawing a sample  $X_1, \dots, X_n$  and building the corresponding histogram. The importance of the risk, as a measure of the quality of the estimator  $\hat{s}_m$  also derives from Markov Inequality which implies that, for any  $z > 0$ ,

$$\mathbb{P}_s \left[ \|s - \hat{s}_m\| \geq \sqrt{zR(\hat{s}_m, s)} \right] \leq z^{-1}. \quad (1.5)$$

Hence, with a guaranteed probability  $1 - z^{-1}$ , the distance between  $s$  and its estimator is bounded by  $\sqrt{zR(\hat{s}_m, s)}$ . When  $z$  is large, there are only two cases: either we were very unlucky and an event of probability not larger than  $z^{-1}$  occurred, or we were not and  $\|s - \hat{s}_m\| \leq \sqrt{zR(\hat{s}_m, s)}$ . Of course, there is no way to know which of the two cases occurred, but this is the rule in Statistics: there is always some uncertainty in our conclusions.

### 1.2.3 Risk bounds for histograms

In any case, (1.5) shows that the risk can be viewed as a good indicator of the performance of an estimator. Moreover, it follows from (1.4) that it can be written as

$$R(\hat{s}_m, s) = \|s - s_m\|^2 + \mathbb{E}_s \left[ \|s_m - \hat{s}_m\|^2 \right]. \quad (1.6)$$

With this special choice of distortion, the risk can be decomposed into the sum of two terms. The first one has nothing to do with the stochastic nature of the observations but simply measures the quality of approximation of  $s$  by the linear space  $V_m$  since it is the square of the distance from  $s$  to  $V_m$ . It only depends on the partition and the true unknown density  $s$ , not on the observations.

The second term in the risk, which is due to the stochastic nature of the observations, hence of  $\hat{s}_m$ , can be bounded in the following way, since  $N_j$  is a binomial random variable with parameters  $n$  and  $p_j$  and both  $s_m$  and  $\hat{s}_m$  are constant on each interval  $I_j$ :

$$\begin{aligned} \mathbb{E}_s \left[ \|s_m - \hat{s}_m\|^2 \right] &= \sum_{j=0}^D \mathbb{E}_s \left[ \int_{I_j} (s_m(x) - \hat{s}_m(x))^2 dx \right] \\ &= \sum_{j=0}^D \mathbb{E}_s \left[ |I_j| \left( \frac{p_j}{|I_j|} - \frac{N_j}{n|I_j|} \right)^2 \right] \\ &= \sum_{j=0}^D \frac{1}{n^2 |I_j|} \text{Var}(N_j) = \frac{1}{n} \sum_{j=0}^D \frac{p_j(1-p_j)}{|I_j|}. \end{aligned} \quad (1.7)$$

This quantity is easy to bound in the special case of a regular partition since then  $|I_j| = (D+1)^{-1}$  and we get, using the concavity of the function  $x \mapsto x(1-x)$ ,

$$\begin{aligned} \mathbb{E}_s \left[ \|s_m - \hat{s}_m\|^2 \right] &= \frac{(D+1)^2}{n} \sum_{j=0}^D \frac{p_j(1-p_j)}{D+1} \\ &\leq \frac{(D+1)^2}{n} \frac{\sum_{j=0}^D p_j}{D+1} \left( 1 - \frac{\sum_{j=0}^D p_j}{D+1} \right) = \frac{D}{n}. \end{aligned} \quad (1.8)$$

Note that  $D = 0$  corresponds to the degenerate partition  $m_0 = \{[0, 1]\}$  for which  $s_{m_0} = \mathbb{1}_{[0,1]}$  which is the density of the uniform distribution on  $[0, 1]$ , independently of  $s$ . Then  $\hat{s}_{m_0} = s_{m_0} = \mathbb{1}_{[0,1]}$  and  $R(\hat{s}_{m_0}, s) = \|s - \mathbb{1}_{[0,1]}\|^2$ .

For general irregular partitions we derive from (1.4) that  $p_j \leq |I_j| \|s_m\|_\infty$ , hence, by (1.7),

$$\mathbb{E}_s \left[ \|s_m - \hat{s}_m\|^2 \right] \leq \frac{\|s_m\|_\infty}{n} \sum_{j=0}^D (1-p_j) = \frac{D \|s_m\|_\infty}{n}. \quad (1.9)$$

There is actually little space for improvement in (1.9) as shown by the following example. Define the partition  $m$  by  $I_j = [\alpha j, \alpha(j+1))$  for  $0 \leq j < D$  and  $I_D = [\alpha D, 1]$  with  $0 < \alpha < D^{-1}$ . Set  $s = s_m = (\alpha D)^{-1} (1 - \mathbb{1}_{I_D})$ . Then  $p_j = D^{-1}$  for  $0 \leq j < D$  and, by (1.7),

$$\mathbb{E}_s \left[ \|s_m - \hat{s}_m\|^2 \right] = \frac{D-1}{\alpha D n} = \frac{(D-1) \|s_m\|_\infty}{n}.$$

If we make the extra assumption that  $s$  belongs to  $\mathbb{L}_\infty([0, 1], dx)$ , then  $\|s_m\|_\infty \leq \|s\|_\infty$  and (1.9) becomes  $\mathbb{E}_s \left[ \|s_m - \hat{s}_m\|^2 \right] \leq \|s\|_\infty n^{-1} D$ . This bound is also valid for regular partitions but always worse than (1.8) since  $\|s\|_\infty \geq 1$  for all densities with respect to Lebesgue measure on  $[0, 1]$  and strictly worse if  $s$  is not the uniform density. Finally, by (1.6),

$$R(\hat{s}_m, s) \leq \|s - s_m\|^2 + \|s\|_\infty n^{-1} D. \quad (1.10)$$

As we shall see later the rather unpleasant presence of the unknown and possibly unbounded  $\|s\|_\infty$  factor in the second term is due to the way we measure the distance between densities, i.e. through the  $\mathbb{L}_2$ -norm.

### 1.3 A first approach to model selection

#### 1.3.1 An alternative interpretation of histograms

The decomposition (1.4) suggests another interpretation for the construction of  $\hat{s}_m$ . What do we do here? Since  $s$  is possibly a complicated object, we replace it by a much simpler one  $s_m$  and estimate it by  $\hat{s}_m$ . Note that  $s_m$  is unknown, as  $s$  is, and what is available to the statistician is the partition  $m$ , the corresponding linear space  $V_m$  and, consequently, the set  $S_m$  of all densities belonging to  $V_m$ , i.e.

$$S_m = \left\{ t = \sum_{j=0}^D a_j \mathbb{1}_{I_j}(x) \mid a_0, \dots, a_D \in \mathbb{R}_+ \quad \text{and} \quad \sum_{j=0}^D a_j |I_j| = 1 \right\}. \quad (1.11)$$

It is a convex subset of some  $D$ -dimensional linear space and  $s_m$  is given by  $\|s - s_m\| = \inf_{t \in S_m} \|s - t\|$ . It is the best approximation of  $s$  in  $S_m$ . As to  $\hat{s}_m$  it only depends on the set  $S_m$  and the observations in the following way, as can easily be checked:

$$\hat{s}_m = \operatorname{argmax}_{t \in S_m} \sum_{i=1}^n \log(t(X_i)),$$

which means that it maximizes the so-called *likelihood function*  $t \mapsto \prod_{i=1}^n t(X_i)$  for  $t \in S_m$ , the likelihood at  $t$  being the joint density of the sample computed at the observations. The estimator  $\hat{s}_m$  is called the *maximum likelihood estimator* (m.l.e. for short) with respect to  $S_m$ . Note that, if  $s = s_m$  actually belongs to  $S_m$ , the m.l.e. converges in probability to  $s$  at rate at least as fast as  $n^{-1/2}$  when  $n$  goes to infinity since then, by (1.5), (1.6) and (1.9),

$$\mathbb{P}_s \left[ \|s - \hat{s}_m\| \geq n^{-1/2} \sqrt{z \|s_m\|_\infty D} \right] \leq z^{-1}.$$

The m.l.e. therefore appears to be a suitable estimator to use if the model  $S_m$  is correct, i.e. if  $s \in S_m$ . When we use the histogram estimator  $\hat{s}_m$ , we just do as if  $s$  did belong to  $S_m$ , using  $S_m$  as an *approximate model* for  $s$ . The resulting risk is then the sum of two terms, an approximation error equal to the square of the distance from  $s$  to  $S_m$  and due to the fact that  $s$  does not in general belong to the model  $S_m$ , and an estimation term  $\mathbb{E}_s \left[ \|s_m - \hat{s}_m\|^2 \right]$  which is the risk corresponding to the estimation within the model when  $s = s_m$  since  $\|s_m - \hat{s}_m\|^2$  has the same expectation when the observations are i.i.d. with density  $s$  or  $s_m$ .

### 1.3.2 Model selection and oracles

Let us denote by  $m_D$  the regular partition with  $D + 1$  pieces and set  $S_D = S_{m_D}$ ,  $\hat{s}_D = \hat{s}_{m_D}$  and  $s_D = s_{m_D}$ , for simplicity. It follows from (1.6) and (1.8) that

$$R(\hat{s}_D, s) \leq \|s - s_D\|^2 + n^{-1}D. \quad (1.12)$$

From the approximation point of view, a good partition should lead to a small value of  $\|s - s_D\|$  which typically requires a partition into many intervals, hence a large value of  $D$ , while the estimation point of view requires a model  $S_D$  defined by few parameters, hence a small value of  $D$ . Obviously, these requirements are contradictory and one should look for a compromise between them in order to minimize the right-hand side of (1.12). Unfortunately, the value  $D_{opt}$  which satisfies

$$\|s - s_{D_{opt}}\|^2 + n^{-1}D_{opt} = \inf_{D \in \mathbb{N}} \{\|s - s_D\|^2 + n^{-1}D\}$$

cannot be computed since it depends on the unknown density  $s$  via the approximation term  $\|s - s_D\|$  and is not accessible to the statistician. This is why the random variable  $\hat{s}_{D_{opt}}$  based on the partition  $m_{D_{opt}}$  is called an ‘‘oracle’’. It is not an estimator because it makes use of the number  $D_{opt}$  which is unknown to the statistician. The problem of model selection is to find a genuine estimator, solely based on the data, that mimics an oracle, i.e. to use the data  $X_1, \dots, X_n$  to select a number  $\hat{D}(X_1, \dots, X_n)$  such that the resulting histogram  $\tilde{s} = \hat{s}_{\hat{D}}$  has a performance which is comparable to that of the oracle:

$$R(\tilde{s}, s) \leq C [\|s - s_{D_{opt}}\|^2 + n^{-1}D_{opt}],$$

where  $C$  is a constant that neither depends on the unknown density  $s$  nor on  $n$ .

### 1.3.3 An illustrative example

Still working with the regular partitions  $m_D$ , let us now assume that the unknown density  $s$  satisfies some Hölderian continuity condition,

$$|s(x) - s(y)| \leq L|x - y|^\beta, \quad L > 0, \quad 0 < \beta \leq 1 \quad \text{for all } x, y \in [0, 1]. \quad (1.13)$$

If  $0 \leq j \leq D$  and  $x \in I_j$ , then  $s_D(x) = s(y)$  for some  $y \in I_j$ , hence  $|s(x) - s_D(x)| \leq L(D + 1)^{-\beta}$ , from which we derive that  $\|s - s_D\|^2 \leq \|s - s_D\|_\infty^2 \leq L^2(D + 1)^{-2\beta}$ . Therefore (1.12) implies that  $R(\hat{s}_D, s) \leq n^{-1}D + L^2(D + 1)^{-2\beta}$ . Since the minimum of the function  $x \mapsto n^{-1}x + L^2x^{-2\beta}$  is obtained for  $x = (2\beta n L^2)^{1/(2\beta+1)}$ , we choose  $D$  so that  $D + 1$  is the smallest integer  $\geq (nL^2)^{1/(2\beta+1)}$ . If  $nL^2 \leq 1$ , this leads to  $D = 0$  and  $R(\hat{s}_0, s) \leq L^2 \leq n^{-1}$ . Otherwise, and this necessarily happens for large enough  $n$ ,  $1 \leq D < (nL^2)^{1/(2\beta+1)}$ , hence  $R(\hat{s}_D, s) \leq 2(Ln^{-\beta})^{2/(2\beta+1)}$ . Finally, in any case,

$$R(\hat{s}_D, s) \leq \max \left\{ 2 \left( Ln^{-\beta} \right)^{2/(2\beta+1)} ; n^{-1} \right\}.$$

Unfortunately, we can only get a risk bound of this form if we fix  $D$  as a function of  $L$  and  $\beta$ , as indicated above. Typically,  $L$  and  $\beta$  are also unknown so that we do not know how to choose  $D$  and cannot get the right risk bound. The situation is even more complicated since, for a given  $s$ , there are many different pairs  $L, \beta$  that satisfy (1.13), leading to different values of  $D$  and risk bounds. Of course, one would like to choose the optimal one which means choosing the value of  $D$  that minimizes the right-hand side of (1.12).

## 1.4 A brief summary of this paper

The study of histograms as density estimators shows us that a convenient method to estimate a complicated object as a density  $s$  on  $[0, 1]$  works as follows: choose an approximate model  $S_m$  for  $s$  involving only a limited number of unknown parameters and then do as if the model were correct, i.e. if  $s \in S_m$ , using an estimator  $\hat{s}_m$  which is a good estimator when the model is actually correct. The resulting risk is the sum of an approximation term which measures the quality of approximation of  $s$  by the model and an estimation term which is roughly proportional to the number of parameters needed to describe an element of the model, reflecting its complexity. As a consequence, a good model should be simple (described by few parameters) and accurate (close to the true density  $s$ ). Unfortunately, because of the second requirement, a theoretical choice of a good model should be based on the knowledge of  $s$ . Given a family of possible models, a major problem is therefore to understand to what extent one can guess from the data which model in the family is appropriate.

The remainder of this paper is devoted to giving some hints to justify and understand the various steps needed to formally develop the previous arguments. The next section will present the classical parametric theory of estimation which assumes that one works with the correct model and that this model satisfies some specific regularity conditions. Under such conditions the m.l.e. enjoys some good asymptotic properties that we shall recall, but this classical theory does not handle the case of approximate models or infinite dimensional parameters. It has therefore been extended in the recent years in many directions to (partly) cover such situations. We shall present here one such generalization that attempts to solve (at least theoretically) most of the difficulties connected with the classical theory. In Section 3, we shall depart from the classical theory, assuming only an approximate model and checking on some examples that the results we got for histograms essentially extend to these cases with a risk bounded by an approximation term plus an estimation term which again leads to the problem of selecting a good model. Section 4 is devoted to a more general approach to estimation based on an approximate model with *finite dimension* for a suitably defined and purely metric notion of dimension. We show here that some specific estimators (sometimes discretized versions of the m.l.e., sometimes more complicated ones) do lead to risk bounds of the required form: an approximation term plus an estimation term which is proportional to the *dimension* (when suitably defined) of the model. In the last section, we explain how to handle many such approximate models with finite dimensions simultaneously. Ideally, we would like to choose, using only the data, the best model in the family, i.e. the one with the smallest risk. This is unfortunately not possible, but we shall explain to what extent one can approximate this ideal risk.

## 2 Some historical considerations

### 2.1 The classical parametric point of view

To be specific, let us assume again that our observations  $X_1, \dots, X_n$  are i.i.d. random variables with an unknown density  $s$  with respect to some reference measure  $\nu$  defined on the underlying measurable set  $(E, \mathcal{E})$  (not necessarily the Lebesgue measure on



$[0, 1]$ ) so that the joint distribution  $P_s$  of the observations on  $E^n$  is given by

$$\frac{dP_s}{d\nu^{\otimes n}}(x_1, \dots, x_n) = \prod_{i=1}^n s(x_i).$$

In the sequel, we shall call the problem of estimating the unknown density  $s$  from the i.i.d. sample  $X_1, \dots, X_n$  the *density estimation problem* or the *i.i.d. framework*.

The classical parametric approach to density estimation that developed after milestone papers by Fisher (1921 and 1925) up to the sixties and is still quite popular nowadays is somewhat different from what we described before. It typically assumes a *parametric model*  $\bar{S}$  for  $s$ , which means that the true unknown density  $s$  of our observations belongs to some particular set  $\bar{S} = \{t_\theta \mid \theta \in \Theta\}$  of densities parametrized by some subset  $\Theta$  of a Euclidean space  $\mathbb{R}^k$ . Then  $s = t_{\theta_0}$  for some particular  $\theta_0 \in \Theta$  which is called the *true parameter value*. One assumes moreover that the mapping  $\theta \mapsto t_\theta$  from  $\Theta$  to  $\bar{S}$  is smooth (in a suitable sense) and one-to-one, so that estimating  $s$  is equivalent to estimating the parameter  $\theta_0$ . An estimator  $\hat{\theta}_n(X_1, \dots, X_n)$  of  $\theta_0$  is then defined via a measurable mapping  $\hat{\theta}_n$  from  $E^n$  to  $\Theta$  (with its Borel  $\sigma$ -algebra) and its quadratic risk is given by

$$R(\hat{\theta}_n, \theta_0) = \mathbb{E}_s \left[ \|\hat{\theta}_n - \theta_0\|^2 \right],$$

where  $\|\cdot\|$  now denotes the Euclidian norm in  $\mathbb{R}^k$ . Typical examples of parametric models for densities on the real line are given by

i) the Gaussian densities  $\mathcal{N}(\mu, \sigma^2)$  with  $\theta = (\mu, \sigma^2)$  and  $\Theta = \mathbb{R} \times (0, +\infty)$  given by

$$t_\theta(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left[ -\frac{1}{2\sigma^2}(x - \mu)^2 \right];$$

ii) the gamma densities  $\Gamma(v, \lambda)$  with  $\theta = (v, \lambda)$  and  $\Theta = (0, +\infty)^2$  given by

$$t_\theta(x) = [\Gamma(v)]^{-1} \lambda^v x^{v-1} \exp[-\lambda x];$$

iii) the uniform density on the interval  $[\theta, \theta + 1]$  given by  $t(\theta) = \mathbb{1}_{[\theta, \theta+1]}$  with  $\theta \in \mathbb{R}$ .

## 2.2 The maximum likelihood method

### 2.2.1 Consistency and asymptotic normality of the parametric m.l.e.

Fisher's approach to parametric estimation is mainly connected with the method of maximum likelihood. We recall from Section 1.3.1 that the likelihood function on  $\Theta$  is given by  $\theta \mapsto \prod_{i=1}^n t_\theta(X_i)$  and a maximum likelihood estimator  $\hat{\theta}_n$  is any maximizer of this function or equivalently of the *log-likelihood function*

$$L(\theta) = \sum_{i=1}^n \log(t_\theta(X_i)).$$

For Gaussian densities, the maximum likelihood estimator  $\hat{\theta}_n = (\hat{\mu}_n, \hat{\sigma}_n^2)$  is unique and given by  $\hat{\mu}_n = n^{-1} \sum_{i=1}^n X_i$  and  $\hat{\sigma}_n^2 = n^{-1} \sum_{i=1}^n (X_i - \hat{\mu}_n)^2$ . Moreover  $\hat{\theta}_n$  converges in probability to the true parameter  $\theta_0$  when  $n$  goes to infinity. We say that  $\hat{\theta}_n$  is *consistent*. Unfortunately, this situation is not general. The study of our second and third examples show that explicit computation of the m.l.e. is not always possible

(gamma densities) or the m.l.e. may not be unique (uniform densities). One can also find examples of *inconsistency* of the m.l.e., but, as shown by Wald (1949), it can be proved that, under suitably strong assumptions, any sequence of maximum likelihood estimators is consistent.

If the mapping  $\theta \mapsto l_\theta(x) = \log(t_\theta(x))$  satisfies suitable differentiability assumptions, the parametric model is called *regular*. This is the case for the Gaussian and gamma densities, not for the uniform. If the model is regular and the m.l.e. is consistent we can expand the derivative of the function  $L$  in a vicinity of  $\theta_0$  when it is an inner point of  $\Theta$ . Restricting ourselves, for simplicity, to the case  $\Theta \subset \mathbb{R}$ , we get

$$L'(\theta) = L'(\theta_0) + (\theta - \theta_0)L''(\theta_0) + (1/2)(\theta - \theta_0)^2L'''(\theta')$$

and since  $\hat{\theta}_n$  is a maximizer for  $L$ ,

$$L'(\hat{\theta}_n) = 0 = L'(\theta_0) + (\hat{\theta}_n - \theta_0)L''(\theta_0) + (1/2)(\hat{\theta}_n - \theta_0)^2L'''(\theta'_n),$$

for some sequence  $(\theta'_n)$  converging to  $\theta_0$  in probability as  $\hat{\theta}_n$  does. Equivalently, setting  $\delta_n = \sqrt{n}(\hat{\theta}_n - \theta_0)$ ,

$$0 = \frac{1}{\sqrt{n}} \sum_{i=1}^n l'_{\theta_0}(X_i) + \left[ \frac{1}{n} \sum_{i=1}^n l''_{\theta_0}(X_i) \right] \delta_n + \frac{\hat{\theta}_n - \theta_0}{2} \left[ \frac{1}{n} \sum_{i=1}^n l'''_{\theta'_n}(X_i) \right] \delta_n. \quad (2.1)$$

Since  $\int t_\theta(x) d\nu(x) = 1$  for all  $\theta$ , it follows from the regularity assumptions that

$$\mathbb{E}_s [l'_{\theta_0}(X_i)] = \int [t'_{\theta_0}(x)/t_{\theta_0}(x)] t_{\theta_0}(x) d\nu(x) = \int t'_{\theta_0}(x) d\nu(x) = 0$$

and

$$\begin{aligned} \mathbb{E}_s [l''_{\theta_0}(X_i)] &= \int [t''_{\theta_0}(x)/t_{\theta_0}(x)] t_{\theta_0}(x) d\nu(x) - \int [t'_{\theta_0}(x)/t_{\theta_0}(x)]^2 t_{\theta_0}(x) d\nu(x) \\ &= 0 - \int ([t'_{\theta_0}(x)]^2 / t_{\theta_0}(x)) d\nu(x) = -I(\theta_0), \end{aligned}$$

where the last equality defines the *Fisher Information*  $I(\theta_0)$ . Moreover

$$\text{Var} [l'_{\theta_0}(X_i)] = \mathbb{E}_s [(l'_{\theta_0}(X_i))^2] = \int [t'_{\theta_0}(x)/t_{\theta_0}(x)]^2 t_{\theta_0}(x) d\nu(x) = I(\theta_0).$$

It then follows from the law of large numbers that

$$\frac{1}{n} \sum_{i=1}^n l''_{\theta_0}(X_i) \xrightarrow{P} \mathbb{E}_s [l''_{\theta_0}(X_i)] = -I(\theta_0)$$

and from the central limit theorem that

$$\frac{1}{\sqrt{n}} \sum_{i=1}^n l'_{\theta_0}(X_i) \rightsquigarrow \mathcal{N}(0, I(\theta_0)),$$

where  $\xrightarrow{P}$  and  $\rightsquigarrow$  denote respectively the convergences in probability and in distribution. The regularity assumptions also ensure that  $n^{-1} \sum_{i=1}^n l'''_{\theta'_n}(X_i)$  is asymptotically

bounded so that the third term in (2.1) is asymptotically negligible as compared to the other two. We finally deduce from (2.1) that

$$\delta_n = \sqrt{n}(\hat{\theta}_n - \theta_0) \rightsquigarrow \mathcal{N}(0, [I(\theta_0)]^{-1}). \quad (2.2)$$

This is the so-called asymptotic normality and efficiency of the maximum likelihood estimator and a formal proof of this result can be found in Cramér (1946, Section 33.3). It can also be proved that the asymptotic variance  $[I(\theta_0)]^{-1}$  of  $\delta_n$  is, in various senses, optimal, as shown by Le Cam (1953) and Hajek (1970 and 1972). Much less restrictive conditions of regularity which still imply the asymptotic normality and efficiency of the m.l.e. have been given by Le Cam (1970) — see also Theorem 12.3 in van der Vaart (2002) —. A good account of the theory can be found in Ibragimov and Has'minskii (1981). A more recent point of view on the theory of regularity and the m.l.e., based on empirical process theory, is to be found in van der Vaart (1998).

### 2.2.2 A more general point of view on the maximum likelihood method

The limitations of the classical parametric theory of maximum likelihood have been recognized for a long time. We already mentioned problems of inconsistency. Examples and further references can be found in Le Cam (1990). Moreover, although it is widely believed among non-specialists that (2.2) typically holds, this is definitely not true, even under consistency. For instance, if  $t_\theta = \theta^{-1}\mathbb{1}_{[0,\theta]}$  is the uniform density on  $[0, \theta]$  and  $\Theta = (0, +\infty)$ , the m.l.e. satisfies  $n(\theta_0 - \hat{\theta}_n) \rightsquigarrow \Gamma(1, \theta_0)$ . Additional examples can be found in Ibragimov and Has'minskii (1981, Chapters 5 and 6) showing that neither the rate  $\sqrt{n}$  nor the limiting normal distribution are general.

Another drawback of the classical point of view on maximum likelihood estimation is its purely asymptotic nature. Not only does it require specific assumptions and can fail under small departures from these assumptions but it tells us nothing about the real performances of the m.l.e. for a given (even large) number  $n_0$  of observations, just as the central limit theorem does. Suppose that our observations  $X_1, \dots, X_n$  are i.i.d. Bernoulli variables taking only the values 0 and 1 with respective probabilities  $1 - \theta_0$  and  $\theta_0$  and  $\Theta = [0, 1]$ . Then  $\hat{\theta}_n = n^{-1} \sum_{i=1}^n X_i$  and, if  $0 < \theta_0 < 1$ ,  $\delta_n = \sqrt{n}(\hat{\theta}_n - \theta_0) \rightsquigarrow \mathcal{N}(0, \theta_0[1 - \theta_0])$  as expected. But it is well-known that if  $n = 1000$  and  $0 < \theta_0 \leq 0.002$ , the distribution of  $\hat{\theta}_n$  looks rather like a Poisson distribution with parameter  $1000\theta_0$  than like a normal  $\mathcal{N}(\theta_0, n^{-1}\theta_0[1 - \theta_0])$  as predicted by the asymptotic theory. A discussion about the relevance of the asymptotic point of view for practical purposes can be found in Le Cam and Yang (2000, Section 7.1).

A further limitation of the classical m.l.e. theory is the fact that the assumed parametric model is true, i.e. the unknown distribution of the observations has a density  $s$  with respect to  $\nu$  which is of the form  $t_{\theta_0}$  for some  $\theta_0 \in \Theta$ . If this assumption is violated, even slightly, the whole theory fails as can be seen from the following example. We assume a Gaussian distribution  $P_\theta = \mathcal{N}(\theta, 1)$  with density  $t_\theta$  with respect to the Lebesgue measure and  $\Theta = \mathbb{R}$  but the observations actually follow the distribution  $Q = (99P_0 + P_{300})/100$ . It is actually rather close to the  $P_0$  distribution, which belongs to the model, in the sense that, for any measurable set  $A$ ,  $|Q(A) - P_0(A)| \leq 1/100$ . Nevertheless, the m.l.e.  $n^{-1} \sum_{i=1}^n X_i$  converges to 3 so that the estimated distribution based on the wrong model will be close to  $P_3$ , hence quite different from the true distribution which is close to  $P_0$ .

For all these reasons, the classical approach to maximum likelihood estimation has been substantially generalized in the recent years. Nonparametric and semiparametric maximum likelihood allows to deal with families of distributions  $P_s$  where  $s$  belongs to some infinite-dimensional set, while sieved m.l.e. involves situations where the true distribution does not belong to the model. Both extensions lead to truly nonasymptotic results. Among the many papers dealing with such extensions, let us mention here Grenander (1981), Silverman (1982), Wahba (1990), Groeneboom and Wellner (1992), van de Geer (1993, 1995 and 2000), Birgé and Massart (1993 and 1998), Shen and Wong (1994), Wong and Shen (1995), van der Vaart and Wellner (1996), Barron, Birgé and Massart (1999) and Massart (2006). Let us now explain what are the novelties brought by some of these extensions.

### 3 An alternative point of view

#### 3.1 Nonparametric density estimation

The assumption that the unknown density  $s$  of the observations belongs to a parametric model, i.e. a smooth image of some subset of a Euclidean space, appears to be definitely too strong and unsatisfactory in many situations. Let us give here two illustrations. If we assume that  $s$  belongs to the set  $\mathcal{S}_1$  of Lipschitz densities on  $[0, 1]$  (i.e.  $s$  satisfies  $|s(x) - s(y)| \leq |x - y|$ ), one cannot represent  $\mathcal{S}_1$  in a smooth way by a finite number of real parameters. The same holds if we simply assume that  $s \in \mathcal{S}_2$ , the set of all densities in  $\mathbb{L}_2([0, 1], dx)$ . In this case, given some orthonormal basis  $(\varphi_j)_{j \geq 1}$  of  $\mathbb{L}_2([0, 1], dx)$ , there exists a natural parametrization of  $\mathcal{S}_2$  by  $\mathbf{l}_2(\mathbb{N}^*)$  ( $\mathbb{N}^* = \mathbb{N} \setminus \{0\}$ ) via the coordinates, but it is definitely not finite-dimensional. These two problems are examples of *nonparametric density estimation problems*.

##### 3.1.1 Projection estimators

In order to solve the second estimation problem, Cencov (1962) proposed a general class of estimators called *projection estimators*. The idea is to estimate the coefficients  $s_j$  of  $s$  in the orthonormal expansion  $s = \sum_{j=1}^{+\infty} s_j \varphi_j$  using estimators  $\hat{s}_j$  chosen in such a way that  $\sum_{j=1}^{+\infty} \hat{s}_j^2 < +\infty$  a.s. so that  $\hat{s} = \sum_{j=1}^{+\infty} \hat{s}_j \varphi_j$  belongs to  $\mathbb{L}_2([0, 1], dx)$  a.s.. Since  $s_j = \int_0^1 s(x) \varphi_j(x) dx = \mathbb{E}_s[\varphi_j(X_i)]$ , a natural estimator for  $s_j$  is  $\bar{\varphi}_j = n^{-1} \sum_{i=1}^n \varphi_j(X_i)$ . Indeed

$$\mathbb{E}_s [\bar{\varphi}_j] = s_j \quad \text{and} \quad \text{Var} (\bar{\varphi}_j) = n^{-1} \text{Var}(\varphi_j(X_1)) \leq n^{-1} \int_0^1 \varphi_j^2(x) s(x) dx. \quad (3.1)$$

Assuming, for simplicity, that we take for  $(\varphi_j)_{j \geq 0}$  the trigonometric basis which is bounded by  $\sqrt{2}$ , we derive that  $\text{Var} (\bar{\varphi}_j) \leq 2/n$ . We cannot use  $\sum_{j=1}^{+\infty} \bar{\varphi}_j \varphi_j$  as an estimator of  $s$  because the series does not converge. This is actually not surprising because we are trying to estimate infinitely many parameters (the  $s_j$ ) from a finite number of observations. But, for any finite subset  $m$  of  $\mathbb{N}^*$ , the estimator  $\hat{s}_m = \sum_{j \in m} \bar{\varphi}_j \varphi_j$  does belong to  $\mathbb{L}_2([0, 1], dx)$  and

$$\|\hat{s}_m - s\|^2 = \sum_{j \in m} (\bar{\varphi}_j - s_j)^2 + \sum_{j \notin m} s_j^2.$$

If we denote by  $|m|$  the cardinality of  $m$ , we conclude from (3.1) that

$$\mathbb{E}_s [\|\hat{s}_m - s\|^2] \leq 2n^{-1}|m| + \|s_m - s\|^2 \quad \text{with} \quad s_m = \sum_{j \in m} s_j \varphi_j. \quad (3.2)$$

Note that  $\hat{s}_m$  is not necessarily a genuine estimator, i.e. a density, but this is a minor point since  $\mathcal{S}_2$  is a closed convex subset of  $\mathbb{L}_2([0, 1], dx)$  on which we may always project  $\hat{s}_m$ , getting a genuine estimator which is even closer to  $s$  than  $\hat{s}_m$ .

### 3.1.2 Approximate models for nonparametric estimation

The construction of the projection estimator  $\hat{s}_m$  can also be interpreted in terms of a model since it is actually based on the parametric model

$$S_m = \left\{ t = \sum_{j \in m} t_j \varphi_j \mid t_j \in \mathbb{R} \text{ for } j \in m \right\}.$$

To build  $\hat{s}_m$ , we proceed as if  $s$  did belong to  $S_m$ , estimating the  $|m|$  unknown parameters  $s_j$  for  $j \in m$  by their natural estimators  $\bar{\varphi}_j$ . But there are three main differences with the classical parametric approach:

- i) we do not assume that  $s \in S_m$  so that  $S_m$  is an *approximate model* for the true density;
- ii) apart from some exceptional cases, like histogram estimation, projection estimators are not maximum likelihood estimators with respect to  $S_m$ ;
- iii) there is no asymptotic point of view here and the risk bound (3.2) is valid for any value of  $n$ .

The histogram estimator can actually be viewed as a particular projection estimator. With the notations of Section 1, we set  $\varphi_j = |I_j|^{-1/2} \mathbb{1}_{I_j}$  for  $1 \leq j \leq D$ , we complete this orthonormal family into a basis of  $\mathbb{L}_2([0, 1], dx)$  and take for  $m$  the set  $\{1, \dots, D\}$ . Then, for  $j \in m$ ,

$$\bar{\varphi}_j = n^{-1} \sum_{i=1}^n |I_j|^{-1/2} \mathbb{1}_{I_j}(X_i) = n^{-1} |I_j|^{-1/2} N_j \quad \text{and} \quad \sum_{j \in m} \bar{\varphi}_j \varphi_j = \hat{s}_m.$$

## 3.2 Approximate models for parametric estimation

### 3.2.1 Gaussian linear regression

An extremely popular parametric model is *Gaussian linear regression*. In this case we observe  $n$  independent variables  $X_1, \dots, X_n$  from the Gaussian linear regression set up

$$X_i = \sum_{j=1}^p \beta_j Z_i^j + \sigma \xi_i \quad \text{for } 1 \leq i \leq n, \quad (3.3)$$

where the random variables  $\xi_i$  are i.i.d. standard normal while the numbers  $Z_i^j$ ,  $1 \leq i \leq n$  denote the respective deterministic and observable values of some *explanatory variable*  $Z^j$ . Here, “variable” is taken in its usual sense of an “economic variable” or a “physical variable”. Practically speaking,  $X_i$  corresponds to an observation in the  $i^{\text{th}}$  experiment and it is assumed that this value depends linearly on the values  $Z_i^j$  of the variables  $Z^j$ ,  $1 \leq j \leq p$  in this experiment but with some additional random

perturbation represented by the random variable  $\sigma\xi_i$ . We assume here that all  $p$  parameters  $\beta_j$  are unknown but that  $\sigma$  is known (this is not usually the case but will greatly simplify our analysis). This set-up results in a parametric model with  $p$  unknown parameters, since the distribution in  $\mathbb{R}^n$  of the vector  $\mathbf{X}$  with coordinates  $X_i$  is entirely defined by the parameters  $\beta_j$ . More precisely, the random variables  $X_1, \dots, X_n$  are independent with respective normal distributions  $\mathcal{N}(s_i, \sigma^2)$  with  $s_i = \sum_{j=1}^p \beta_j Z_i^j$ . Equivalently  $\mathbf{X}$  is a Gaussian vector with mean vector  $s = (s_i)_{1 \leq i \leq n}$  and covariance matrix  $\sigma^2 I_n$  where  $I_n$  denotes the identity matrix in  $\mathbb{R}^n$ . If we denote by  $\mathbf{Z}^j$  the vector with coordinates  $Z_i^j$  and assume that the vectors  $\mathbf{Z}^j, 1 \leq j \leq p$  span a  $p$ -dimensional linear space  $\overline{S}_p$ , which we shall do, it is equivalent to estimate the parameters  $\beta_j$  or the vector  $s \in \overline{S}_p$ .

The estimation problem can then be summarized as follows: observing the Gaussian vector  $\mathbf{X}$  with distribution  $\mathcal{N}(s, \sigma^2 I_n)$  with a known value of  $\sigma$ , estimate the parameter  $s$  which is assumed to belong to  $\overline{S}_p$ . This is a parametric problem similar to those we considered in Section 2 and it can be solved via the maximum likelihood method. The density of  $\mathbf{X}$  with respect to the Lebesgue measure on  $\mathbb{R}^n$  and the log-likelihood of  $s$  are respectively given by

$$\frac{1}{(2\pi\sigma^2)^{n/2}} \exp \left[ -\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - s_i)^2 \right] \quad \text{and} \quad -\frac{n}{2} \log(2\pi\sigma^2) - \frac{1}{2\sigma^2} \sum_{i=1}^n (X_i - s_i)^2,$$

so that the maximum likelihood estimator  $\hat{s}_p$  over  $\overline{S}_p$  is merely the orthogonal projection of  $\mathbf{X}$  onto  $\overline{S}_p$  with risk  $\mathbb{E}_s [\|s - \hat{s}_p\|^2] = \sigma^2 p$ . This estimator actually makes sense even if  $s \notin \overline{S}_p$  since, whatever the true value of  $s \in \mathbb{R}^n$ ,

$$\mathbb{E}_s [\|s - \hat{s}_p\|^2] = \sigma^2 p + \inf_{t \in \overline{S}_p} \|s - t\|^2. \quad (3.4)$$

The risk is the sum of two terms, one which is proportional to the number  $p$  of parameters to be estimated and another one which measures the accuracy of the model  $\overline{S}_p$  we use. This second term vanishes when the model is correct (contains  $s$ ).

### 3.2.2 Model choice again

In the classical regression problem, the model  $\overline{S}_p$  is assumed to be correct so that  $\mathbb{E}_s [\|s - \hat{s}_p\|^2] = \sigma^2 p$  but this approach leads to two opposite problems. In order to keep the term  $\sigma^2 p$  in (3.4) small, we may be tempted to put too few explanatory variables in the model, omitting some important ones so that not only  $s \notin \overline{S}_p$  but  $\inf_{t \in \overline{S}_p} \|s - t\|^2$  may be very large, possibly larger than  $\sigma^2 n$ . In this case, it would be wiser to use the largest possible model  $\mathbb{R}^n$  for  $s$  and the corresponding m.l.e.  $\hat{s} = \mathbf{X}$  resulting in the better risk  $\mathbb{E}_s [\|s - \hat{s}\|^2] = \sigma^2 n$ . In order to avoid this difficulty, we may alternatively introduce many explanatory variables  $Z^j$  in the model  $\overline{S}_p$ . Then even if it is correct, we shall get a large risk bound  $\sigma^2 p$ . It may then happen that only a small number  $q$  of the  $p$  explanatory variables determining the model are really influential. This means that if  $\overline{S}_q$  is the linear span of those  $q$  variables, say  $Z^1, \dots, Z^q$ ,  $\inf_{t \in \overline{S}_q} \|s - t\|^2$  is small. As a consequence, the risk bound of the m.l.e.  $\hat{s}_q$  with respect to  $\overline{S}_q$ , i.e.  $\sigma^2 q + \inf_{t \in \overline{S}_q} \|s - t\|^2$  may be much smaller than  $\sigma^2 p$ .

These examples show that, even in the parametric case, the use of an approximate model may be preferable to the use of a correct model, although a grossly wrong model may lead to terrible results. The choice of a suitable model is therefore crucial:

a large model including many explanatory variables automatically results in a large risk bound due to the component  $\sigma^2 p$  of the risk in (3.4) while the choice of a too parsimonious model including only a limited number of variables may result in a poor estimator based on a grossly wrong model if we have omitted some very influential variables.

A natural idea to solve this dilemma would be to start with some large family  $\{\bar{S}_m, m \in \mathcal{M}\}$  of linear models indexed by some set  $\mathcal{M}$  and with respective dimensions  $D_m$ . For each of them, the corresponding m.l.e.  $\hat{s}_m$  (the projection of  $\mathbf{X}$  onto  $\bar{S}_m$ ) satisfies

$$\mathbb{E}_s [\|s - \hat{s}_m\|^2] = \sigma^2 D_m + \inf_{t \in \bar{S}_m} \|s - t\|^2,$$

and an optimal model  $\bar{S}_{\bar{m}}$  is one that minimizes this quantity. But, as in the case of histograms, this optimal model depends on the unknown parameter  $s$  via  $\inf_{t \in \bar{S}_m} \|s - t\|$  so that  $\hat{s}_{\bar{m}}$  is an ‘‘oracle’’, not a genuine estimator. Since this oracle is not available to the statistician, he has to try an alternative method and use the observation  $\mathbf{X}$  to build a selection procedure  $\hat{m}(\mathbf{X})$  of one model  $\bar{S}_{\hat{m}}$ , estimating  $s$  by  $\tilde{s} = \hat{s}_{\hat{m}}$ . An ideal *model selection procedure* should have the performance of an oracle, i.e. satisfy

$$\mathbb{E}_s [\|s - \tilde{s}\|^2] = \inf_{m \in \mathcal{M}} \left\{ \sigma^2 D_m + \inf_{t \in \bar{S}_m} \|s - t\|^2 \right\}, \quad (3.5)$$

but such a procedure cannot exist and the best that one can expect is to find selection procedures satisfying a risk bound which is close to (3.5).

## 4 Model based statistical estimation

In three different contexts, namely histogram estimation for densities, projection estimation for densities and Gaussian linear regression, we have seen that the use of an approximate model associated with a convenient estimator with values in the model leads to three risk bounds, namely (1.12), (3.2) and (3.4), which share the same structure. These bounds are the sum of two terms, one is the squared distance of the unknown parameter to the model, the second is proportional to the number of parameters that are involved in the model. One can therefore wonder to what extent this situation is typical.

### 4.1 A general statistical framework

Before we proceed to the solution of the problem, let us make the statistical framework on which we work somewhat more precise. We observe a random phenomenon  $\mathbf{X}(\omega)$  (real variable, vector, sequence, process, set, ...) from the abstract probability space  $(\Omega, \mathcal{A}, \mathbb{P})$  with values in the measurable set  $(\Xi, \mathcal{X})$  and with unknown probability distribution  $P_{\mathbf{X}}$  on  $(\Xi, \mathcal{X})$  given by

$$P_{\mathbf{X}}[A] = \mathbb{P}[\mathbf{X}^{-1}(A)] = \mathbb{P}[\{\omega \in \Omega \mid \mathbf{X}(\omega) \in A\}] \quad \text{for all } A \in \mathcal{X}.$$

The purpose of statistical estimation is to get some information on this distribution from one observation  $\mathbf{X}(\omega)$  of the phenomenon. We assume that  $P_{\mathbf{X}}$  belongs to some given subset  $\mathcal{P} = \{P_t, t \in M\}$  of the set of all distributions on  $(\Xi, \mathcal{X})$ , where  $M$  denotes a one-to-one parametrization of  $\mathcal{P}$ . We moreover assume that  $M$  is a metric space with a distance  $d$ . Therefore  $P_{\mathbf{X}} = P_s$  for some  $s \in M$  and we want to *estimate*

$P_s$ , or equivalently  $s$ , in view of this one-to-one correspondence which also allows us to consider  $d$  as a distance on  $\mathcal{P}$  as well. As in Section 1.2.2, we look for an estimator of  $s$ , i.e. a measurable mapping  $\hat{s}$  from  $(\Xi, \mathcal{X})$  to  $M$  (with its Borel  $\sigma$ -algebra) such that  $\hat{s}(\mathbf{X})$  provides a good approximation of the unknown value  $s$ . Such a mapping is called an *estimator* of  $s$ . We measure the performance of the estimator  $\hat{s}(\mathbf{X})$  via its *quadratic risk*

$$R(\hat{s}, s) = \mathbb{E}_s [d^2(s, \hat{s})]. \quad (4.1)$$

There is a very large number of possibilities for the choice of  $\mathcal{P}$  depending on the structure of  $(\Xi, \mathcal{X})$  and the problem we have to solve. In this paper we focus on the two particular but typical examples that we considered earlier, namely the density estimation problem and the Gaussian regression problem which amounts to the estimation of the mean of a Gaussian vector. In both cases  $\Xi = E^n$  is a product space with a product  $\sigma$ -algebra  $\mathcal{X} = \mathcal{E}^{\otimes n}$  so that  $\mathbf{X}$  is the vector  $(X_1, \dots, X_n)$  and the  $X_i$  are random variables with values in  $(E, \mathcal{E})$ .

**Density estimation** For the *density estimation problem* we are given some reference measure  $\nu$  on  $(E, \mathcal{E})$  and we assume that the  $X_i$  are i.i.d. random variables with a density  $s$  with respect to  $\nu$ , in which case  $M$  can be chosen as the set of all densities with respect to  $\nu$ , i.e. the subset of  $\mathbb{L}_1(\nu)$  of nonnegative functions which integrate to one. Such a situation occurs when one replicates the same experiment  $n$  times under identical conditions and assumes that each experiment has no influence on the others, for instance when we observe the successive outcomes of a “roulette” game. Then, for each  $t \in M$ ,  $P_t$  has the density  $\prod_{i=1}^n t(x_i)$  with respect to  $\mu = \nu^{\otimes n}$ .

**Gaussian regression** This is the case that we considered in Section 3.2 with  $(E, \mathcal{E})$  being the real line with its Borel  $\sigma$ -algebra. Here  $\mathbf{X}$  is a Gaussian vector in  $\mathbb{R}^n$  with known covariance matrix  $\sigma^2 I_n$ . Then  $M = \mathbb{R}^n$  and  $t = (t_1, \dots, t_n) \in M$  is the unknown mean vector of the Gaussian distribution  $P_t = \mathcal{N}(t, \sigma^2 I_n)$  with density

$$g_t(x) = \frac{1}{(2\pi\sigma^2)^{n/2}} \exp \left[ -\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - t_i)^2 \right], \quad (4.2)$$

with respect to the Lebesgue measure  $\mu$  on  $\mathbb{R}^n$ .

## 4.2 Two point parameter sets

Before we come to the general situation, it will be useful to analyze a special, quite unrealistic, but very simple case. Let us make the extra assumption that  $s$  belongs to the smallest possible parameter set, i.e. a subset  $S$  of  $M$  containing only two elements  $v$  and  $u$ . Note that the statistical problem would be void if  $S$  contained only one point since  $s$  would then be known.

A solution to this estimation problem is provided by the maximum likelihood method described in Section 2.2. Let  $\mu$  be any measure dominating both  $P_v$  and  $P_u$  ( $P_v + P_u$  would do) and denote by  $g_v$  and  $g_u$  the respective densities of  $P_v$  and  $P_u$  with respect to  $\mu$ . Then define an estimator  $\hat{\varphi}(\mathbf{X})$  with values in  $S$  by

$$\hat{\varphi}(\mathbf{X}) = \begin{cases} v & \text{if } g_v(\mathbf{X}) > g_u(\mathbf{X}); \\ u & \text{if } g_u(\mathbf{X}) > g_v(\mathbf{X}). \end{cases} \quad (4.3)$$



Take any decision you like in case of equality. If  $s = v$ , we get

$$R(\hat{\varphi}, s) = d^2(v, u) \mathbb{P}_v[\hat{\varphi} = u] \leq d^2(v, u) \mathbb{P}_v[g_u(\mathbf{X}) \geq g_v(\mathbf{X})].$$

Since the distribution of  $\mathbf{X}$  is  $P_s = P_v = g_v \cdot \mu$ ,  $\mathbb{P}_v[g_v(\mathbf{X}) > 0] = 1$  and

$$\begin{aligned} \mathbb{P}_v[g_u(\mathbf{X}) \geq g_v(\mathbf{X})] &= \mathbb{P}_v \left[ \sqrt{g_u(\mathbf{X})/g_v(\mathbf{X})} \geq 1 \right] \leq \mathbb{E}_v \left[ \sqrt{g_u(\mathbf{X})/g_v(\mathbf{X})} \right] \\ &= \int_{\Xi} \sqrt{g_u(x)/g_v(x)} g_v(x) d\mu(x) = \int_{\Xi} \sqrt{g_u(x)g_v(x)} d\mu(x). \end{aligned}$$

Hence  $R(\hat{\varphi}, s) \leq d^2(v, u) \rho(P_v, P_u)$  with

$$\rho(P_v, P_u) = \rho(P_u, P_v) = \int_{\Xi} \sqrt{\frac{dP_u}{d\mu}(x) \frac{dP_v}{d\mu}(x)} d\mu(x). \quad (4.4)$$

It is easily seen that the definition of  $\rho(P_v, P_u)$  via (4.4) is independent of the choice of the dominating measure  $\mu$ . Since the same risk bound holds when  $s = u$ , we finally get

$$\sup_{s \in \{u, v\}} R(\hat{\varphi}, s) \leq d^2(v, u) \rho(P_v, P_u). \quad (4.5)$$

This bound demonstrates the importance of the so-called *Hellinger affinity*  $\rho(P, Q)$  between two probability measures  $P$  and  $Q$ . It satisfies in particular by the Cauchy-Schwarz Inequality and the Fubini Theorem

$$0 \leq \rho(P, Q) \leq 1 \quad \text{and} \quad \rho(P^{\otimes n}, Q^{\otimes n}) = \rho^n(P, Q). \quad (4.6)$$

It is, moreover, closely related to a well-known distance between probabilities, the *Hellinger distance*  $h$  defined by

$$h^2(P, Q) = \frac{1}{2} \int_{\Xi} \left( \sqrt{\frac{dP}{d\mu}(x)} - \sqrt{\frac{dQ}{d\mu}(x)} \right)^2 d\mu(x) = 1 - \rho(P, Q). \quad (4.7)$$

The Hellinger distance is merely the  $\mathbb{L}_2(\mu)$ -distance between the square roots of the densities with respect to any dominating measure  $\mu$  (and actually independent of  $\mu$ ). Here, we follow Le Cam who normalizes the integral so that the Hellinger distance has range  $[0, 1]$ . An alternative definition is without the factor  $1/2$  in (4.7). He also showed in Le Cam (1973) that

$$\rho(P, Q) \geq \int_{\Xi} \inf \left\{ \frac{dP}{d\mu}(x); \frac{dQ}{d\mu}(x) \right\} d\mu(x) \geq 1 - \sqrt{1 - \rho^2(P, Q)}. \quad (4.8)$$

It is easy to compute  $\rho(P_v, P_u)$  for our two special frameworks. In the case of Gaussian distributions  $P_u = \mathcal{N}(u, \sigma^2 I_n)$  and  $P_v = \mathcal{N}(v, \sigma^2 I_n)$ , we get

$$\rho(P_v, P_u) = \exp \left[ -\|v - u\|^2 / (8\sigma^2) \right],$$

so that  $[-\log \rho]^{1/2}$  is a multiple of the Euclidian distance between parameters, modulo the identification of  $t$  and  $P_t$ . Note that, in general,  $[-\log \rho]^{1/2}$  is not a distance since it may be infinite and does not satisfy the triangle inequality. Setting  $d(v, u) = \|v - u\|$ , (4.5) becomes

$$\sup_{s \in \{u, v\}} R(\hat{\varphi}, s) \leq \|v - u\|^2 \exp \left[ -\|v - u\|^2 / (8\sigma^2) \right] \leq 8e^{-1} \sigma^2,$$

independently of  $v$  and  $u$ . In the i.i.d. case, we use the Hellinger distance to define the risk, setting  $d(u, v) = h(u, v) = h(P_u, P_v)$  and (4.5) becomes, whatever the densities  $v$  and  $u$ ,

$$\sup_{s \in \{u, v\}} R(\hat{\varphi}, s) \leq h^2(v, u) [1 - h^2(v, u)]^n \leq n^n (n+1)^{-(n+1)} \leq (ne)^{-1}.$$

### 4.3 Two point models for the Gaussian framework

As we pointed out at the beginning of the last section, assuming that  $s$  is either  $v$  or  $u$  is definitely unrealistic. A more realistic problem would rather be as follows:  $s$  is unknown but we believe that one of two different situations can occur implying that  $s$  is close (not necessarily equal) to either  $v$  or  $u$ . Then it seems natural to use  $S = \{v, u\}$  as an approximate model for  $s$  and just proceed as before, using the estimator  $\hat{\varphi}(\mathbf{X})$  defined by (4.3). We can then try to mimic the proof which lead to (4.5), apart from the fact that the argument leading to

$$\mathbb{P}_v[g_u(\mathbf{X}) \geq g_v(\mathbf{X})] \leq \rho(P_v, P_u) = \exp[-\|v - u\|^2 / (8\sigma^2)]$$

then fails. One can instead prove the following result (Birgé, 2006).

**Proposition 1** *Let  $P_t$  denote the Gaussian distribution  $\mathcal{N}(t, \sigma^2 I_n)$  in  $\mathbb{R}^n$ . If  $\mathbf{X}$  is a Gaussian vector with distribution  $P_s$  and  $\|s - v\| \leq \|v - u\|/6$ , then*

$$\mathbb{P}_s[g_u(\mathbf{X}) \geq g_v(\mathbf{X})] \leq \exp[-\|v - u\|^2 / (24\sigma^2)].$$

We can then proceed as before and conclude that, if  $\|s - v\| \leq \|v - u\|/6$ , then

$$\begin{aligned} R(\hat{\varphi}, s) &\leq 2(\|s - v\|^2 + \mathbb{E}_s[\|\hat{\varphi} - v\|^2]) \\ &\leq 2(\|s - v\|^2 + \|v - u\|^2 \exp[-\|v - u\|^2 / (24\sigma^2)]) \\ &\leq 2\|s - v\|^2 + 48e^{-1}\sigma^2. \end{aligned}$$

A similar bound holds with  $u$  replacing  $v$  if  $\|s - u\| \leq \|v - u\|/6$ . Finally, if  $\min\{\|s - v\|, \|s - u\|\} > \|v - u\|/6$ , since  $\hat{\varphi}$  is either  $v$  or  $u$ ,

$$\begin{aligned} R(\hat{\varphi}, s) &\leq (\max\{\|s - v\|, \|s - u\|\})^2 \\ &\leq (\min\{\|s - v\|, \|s - u\|\} + \|v - u\|)^2 \\ &\leq 49(\min\{\|s - v\|, \|s - u\|\})^2. \end{aligned}$$

We finally conclude that, whatever  $s \in M$ , even if our initial assumption that  $s$  is close to  $S$  is wrong,

$$R(\hat{\varphi}, s) \leq 48e^{-1}\sigma^2 + 49 \inf_{t \in S} \|s - t\|^2,$$

which, apart from the constants, is similar to (3.4).

## 4.4 General models for the Gaussian framework

### 4.4.1 Linear models

Instead of assuming that  $s$  is close to a two-points set, let us now assume that it is close to some  $D$ -dimensional linear subspace  $V$  of  $\mathbb{R}^n$  ( $D > 0$ ). Choose some

$\lambda \geq 4\sqrt{3}\sigma$  and, identifying  $V$  to  $\mathbb{R}^D$  via some orthonormal basis, consider the lattice  $S = (2\lambda\mathbb{Z})^D \subset V$ . The maximum likelihood estimator  $\hat{s}(\mathbf{X})$  with respect to  $S$  is given by  $\hat{s}(\mathbf{X}) = \operatorname{argmax}_{t \in S} g_t(\mathbf{X})$ . Its unicity follows from the facts that  $S$  is countable and  $\mathbb{P}_s[g_t(\mathbf{X}) = g_u(\mathbf{X})] = 0$  for each pair  $(t, u) \in S^2$  such that  $t \neq u$ . As to its existence (with probability one), it is a consequence of the following result.

**Proposition 2** *For  $s$  an arbitrary point in  $M = \mathbb{R}^n$ ,  $s' \in S$ , and*

$$y \geq y_0 = \max \left\{ \lambda\sqrt{2D}, 6\|s' - s\| \right\}, \quad (4.9)$$

then

$$\mathbb{P}_s \left[ \exists t \in S \text{ with } \|s' - t\| \geq y \text{ and } g_t(\mathbf{X}) \geq g_{s'}(\mathbf{X}) \right] \leq 1.14 \exp \left[ -\frac{y^2}{48\sigma^2} \right]. \quad (4.10)$$

*Proof:* Let  $S_k = \{t \in S \mid 2^{k/2}y \leq \|s' - t\| < 2^{(k+1)/2}y\}$  with cardinality  $|S_k|$ . If we denote by  $P(y)$  the left-hand side of (4.10), we get

$$P(y) \leq \sum_{k=0}^{+\infty} \mathbb{P}_s[\exists t \in S_k \text{ with } g_t(\mathbf{X}) \geq g_{s'}(\mathbf{X})] \leq \sum_{k=0}^{+\infty} |S_k| \sup_{t \in S_k} \mathbb{P}_s[g_t(\mathbf{X}) \geq g_{s'}(\mathbf{X})]. \quad (4.11)$$

Since, for  $t \in S_k$ ,  $\|s' - t\| \geq 2^{k/2}y \geq 6\|s' - s\|$ , we may apply Proposition 1 to get

$$\sup_{t \in S_k} \mathbb{P}_s[g_t(\mathbf{X}) \geq g_{s'}(\mathbf{X})] \leq \exp \left[ -2^k y^2 / (24\sigma^2) \right]. \quad (4.12)$$

Moreover, for any ball  $\mathcal{B}(s', r)$  with center  $s'$  and radius  $r = x\lambda\sqrt{D}$  with  $x \geq 2$ ,

$$|S \cap \mathcal{B}(s', r)| < \exp \left[ x^2 D / 2 \right]. \quad (4.13)$$

To prove this, we apply the next inequality which follows from a comparison of the volumes of cubes and balls in  $\mathbb{R}^D$  as in the proof of Lemma 2 from Birgé and Massart (1998).

$$|S \cap \mathcal{B}(s', r)| \leq \frac{(\pi e/2)^{D/2}}{\sqrt{\pi D}} \left( \frac{r}{\lambda\sqrt{D}} + 1 \right)^D < \exp[D(0.73 + \log(x+1))].$$

We then get (4.13) since  $x \geq 2$ . Applying it with  $r = 2^{(k+1)/2}y \geq 2^{1+k/2}\lambda\sqrt{D}$  by (4.9), leads to  $|S_k| \leq \exp \left[ 2^k (y/\lambda)^2 \right]$ . Together with (4.11) and (4.12), this shows that

$$\begin{aligned} P(y) &\leq \sum_{k=0}^{+\infty} \exp \left[ 2^k \frac{y^2}{\lambda^2} - 2^k \frac{y^2}{24\sigma^2} \right] \leq \sum_{k=0}^{+\infty} \exp \left[ -2^k \frac{y^2}{48\sigma^2} \right] \\ &= \exp \left[ -\frac{y^2}{48\sigma^2} \right] \sum_{k=0}^{+\infty} \exp \left[ -\frac{y^2}{48\sigma^2} (2^k - 1) \right]. \end{aligned}$$

The conclusion follows from the fact that  $y^2 \geq y_0^2 \geq 2\lambda^2 D \geq 2\lambda^2 \geq 96\sigma^2$ .  $\square$

Proposition 2 implies that, for  $y \geq y_0$ , there exists a set  $\Omega_y \subset \Omega$  with  $\mathbb{P}_s(\Omega_y) \geq 1 - 1.14 \exp \left[ -y^2 / (48\sigma^2) \right]$  and such that, for  $\omega \in \Omega_y$ , the function  $t \mapsto g_t(\mathbf{X}(\omega))$  has

a maximum in the ball  $\mathcal{B}(s', y)$ . This shows that, if  $\omega \in \Omega_y$ , the m.l.e.  $\hat{s}(\mathbf{X}(\omega))$  exists and satisfies  $\|\hat{s}(\mathbf{X}) - s'\| \leq y$ . As a consequence, the m.l.e.  $\hat{s}(\mathbf{X})$  exists a.s. and

$$\begin{aligned} \mathbb{E}_s [\|\hat{s}(\mathbf{X}) - s'\|^2] &= \int_0^{+\infty} \mathbb{P}_s [\|\hat{s}(\mathbf{X}) - s'\|^2 \geq z] dz \\ &\leq y_0^2 + \int_{y_0^2}^{+\infty} \mathbb{P}_s [\|\hat{s}(\mathbf{X}) - s'\| \geq \sqrt{z}] dz \\ &\leq y_0^2 + 1.14 \int_{y_0^2}^{+\infty} \exp\left[-\frac{z}{48\sigma^2}\right] dz \\ &= y_0^2 + 1.14 \times 48\sigma^2 \exp[-y_0^2/(48\sigma^2)] \\ &\leq y_0^2 + 55e^{-2}\sigma^2. \end{aligned}$$

Then

$$\begin{aligned} \mathbb{E}_s [\|\hat{s}(\mathbf{X}) - s\|^2] &\leq 2 [\|s - s'\|^2 + \mathbb{E}_s [\|\hat{s}(\mathbf{X}) - s'\|^2]] \\ &\leq 2 [\|s - s'\|^2 + y_0^2 + 55e^{-2}\sigma^2] \\ &\leq 2 [37\|s - s'\|^2 + 2\lambda^2 D + 55e^{-2}\sigma^2]. \end{aligned}$$

Note that the construction of  $S$  as a lattice in  $V$  implies that any point in  $V$  is at a distance of some point in  $S$  not larger than  $\lambda\sqrt{D}$  which means that one can choose  $s'$  in such a way that  $\|s - s'\| \leq \inf_{t \in V} \|s - t\| + \lambda\sqrt{D}$ . With such a choice for  $s'$ , we get

$$\mathbb{E}_s [\|\hat{s}(\mathbf{X}) - s\|^2] \leq 2 \left[ 74 \inf_{t \in V} \|s - t\|^2 + 76\lambda^2 D + 55e^{-2}\sigma^2 \right].$$

Setting  $\lambda$  to its minimum value  $4\sqrt{3}\sigma$ , we conclude, since  $D \geq 1$ , that

$$\mathbb{E}_s [\|\hat{s}(\mathbf{X}) - s\|^2] \leq 148 \inf_{t \in V} \|s - t\|^2 + 7311\sigma^2 D. \quad (4.14)$$

#### 4.4.2 General models with finite metric dimension

Note that, apart from the huge constants that we actually did not try to optimize in order to keep the computations as simple as possible, (4.14) is quite similar to (3.4), although we actually used a different estimation procedure, and also a different method of proof which has an important advantage: it did not make any use of the fact that  $V$  is a linear space. What we actually used are the metric properties of the  $D$ -dimensional linear subspace  $V$  of  $M = \mathbb{R}^n$ , which can be summarized as follows.

**Property P** *Whatever  $\eta > 0$ , one can find a subset  $S$  of  $M$  such that:*

- i) for each  $t \in V$  there exists some  $t' \in S$  with  $\|t - t'\| \leq \eta$ ;*
- ii) for any ball  $\mathcal{B}(t, x\eta)$  with center  $t \in M$  and radius  $x\eta$ ,*

$$|S \cap \mathcal{B}(t, x\eta)| \leq \exp[x^2 D/2] \quad \text{for } x \geq 2.$$

In the previous example we simply defined  $S$  so that  $\eta = \lambda\sqrt{D} = 4\sigma\sqrt{3D}$ .

The fact that the previous property of  $V$  was a key argument in the proof motivates the following general definition.

**Definition 1** Let  $\bar{S}$  be a subset of some metric space  $(M, d)$  and  $\bar{D}$  be some real number  $\geq 1/2$ . We say that  $\bar{S}$  has a finite metric dimension bounded by  $\bar{D}$  if, for every  $\eta > 0$ , one can find a subset  $S_\eta$  of  $M$  such that:

- i) for each  $t \in \bar{S}$  there exists some  $t' \in S_\eta$  with  $d(t, t') \leq \eta$  (we say that  $S_\eta$  is an  $\eta$ -net for  $\bar{S}$ );
- ii) for any ball  $\mathcal{B}(t, x\eta)$  with center  $t \in M$  and radius  $x\eta$ ,

$$|S_\eta \cap \mathcal{B}(t, x\eta)| \leq \exp [x^2 \bar{D}] \quad \text{for } x \geq 2.$$

Note that any subset of  $\bar{S}$  also has a finite metric dimension bounded by  $\bar{D}$ . It follows from the Property P that a  $D$ -dimensional linear subspace of a Euclidean space has a metric dimension bounded by  $D/2$ . Note that, apart from the factor  $1/2$ , this result cannot be improved in view of the following lower bound for the metric dimension of a  $D$ -dimensional ball.

**Lemma 1** Let  $\bar{S}$  be a ball of the metric space  $(M, d)$  which is isometric to a ball in the Euclidean space  $\mathbb{R}^D$ . Then a bound  $\bar{D}$  for its metric dimension cannot be smaller than  $D/13$ .

*Proof:* Let  $\bar{S} = \mathcal{B}(t, r)$  have a finite metric dimension bounded by  $\bar{D}$  and  $\eta < r/3$ . One can find  $S_\eta$  in  $M$  which is an  $\eta$ -net for  $\bar{S}$  and such that  $N = |S_\eta \cap \mathcal{B}(t, 3\eta)| \leq \exp [9\bar{D}]$ . Moreover,  $S_\eta$  is also an  $\eta$ -net for  $\mathcal{B}(t, 2\eta)$  so that  $\mathcal{B}(t, 2\eta)$  can be covered by the  $N$  balls with radius  $\eta$  and centers in  $S_\eta \cap \mathcal{B}(t, 3\eta)$ . Since  $\mathcal{B}(t, 3\eta) \subset \bar{S}$  we can use the isometry to show, comparing the volumes of the balls, that  $N \geq 2^D$  so that  $9\bar{D} \geq D \log 2$  and the conclusion follows.  $\square$

Introducing Definition 1 in the proof of Proposition 2, we get the following result.

**Theorem 1** Let  $\mathbf{X}$  be a Gaussian vector in  $\mathbb{R}^n$  with unknown mean  $s$  and known covariance matrix  $\sigma^2 I_n$ . Let  $\bar{S}$  be a subset of the Euclidean space  $\mathbb{R}^n$  with a finite metric dimension bounded by  $\bar{D}$ . Then one can build an estimator  $\hat{s}_{\bar{S}}(\mathbf{X})$  of  $s$  such that, for some universal constant  $C$  (independent of  $s$ ,  $n$  and  $\bar{S}$ ),

$$\mathbb{E}_s [ \|\hat{s}_{\bar{S}}(\mathbf{X}) - s\|^2 ] \leq C \left[ \inf_{t \in \bar{S}} \|s - t\|^2 + \sigma^2 \bar{D} \right]. \quad (4.15)$$

This theorem implies that we can use for models non-linear sets that have a finite metric dimension. In particular, various types of manifolds could be used as models. To build the estimator  $\hat{s}_{\bar{S}}(\mathbf{X})$ , we set  $\eta = 4\sigma\sqrt{6\bar{D}}$  and choose an  $\eta$ -net  $S_\eta$  for  $\bar{S}$  satisfying the properties of Definition 1. Then we take for  $\hat{s}_{\bar{S}}(\mathbf{X})$  the m.l.e. with respect to  $S_\eta$ .

## 4.5 Density estimation

When we want to extend the results obtained for the Gaussian framework to density estimation we encounter new difficulties. The two key arguments used in the proof of Proposition 2 are that  $V$  has a finite metric dimension and Proposition 1. For i.i.d. observations  $X_1, \dots, X_n$  with density  $s$  and in view of the fact that

$$\mathbb{P}_s \left[ \prod_{i=1}^n u(X_i) \geq \prod_{i=1}^n v(X_i) \right] \leq \exp [-nh^2(u, v)] \quad \text{if } s = v,$$

an analogous result would be as follows:

**Conjecture C** Let  $X_1, \dots, X_n$  be i.i.d. random variables with an unknown density  $s$  with respect to some measure  $\nu$  on  $(E, \mathcal{E})$ . There exist two constants  $\kappa \geq 2$  and  $A > 0$  such that, whatever the densities  $u, v$  on  $(E, \mathcal{E})$  such that  $h(s, v) \leq \kappa^{-1}h(u, v)$ , then

$$\mathbb{P}_s \left[ \prod_{i=1}^n u(X_i) \geq \prod_{i=1}^n v(X_i) \right] \leq \exp [-Anh^2(u, v)].$$

If this conjecture were true one could mimic the proof for the Gaussian case, starting from a subset  $\bar{S}$  of the metric space  $(M, h)$  with finite metric dimension, choosing a suitable  $\eta$ -net  $S_\eta$  for  $\bar{S}$  and computing the m.l.e. with respect to  $S_\eta$  to get an analogue of Theorem 1. Unfortunately Conjecture C is wrong and, as a consequence, one can find situations in the i.i.d. framework where the m.l.e. with respect to  $S_\eta$  does not behave at all as expected. To get an analogue of Theorem 1 for density estimation, one cannot work with the maximum likelihood method any more. An alternative method that allows to deal with the problem of density estimation has been proposed by Le Cam (1973 and 1975) who also introduced a notion of metric dimension, and then extended by the present author in Birgé (1983 and 1984). In the sequel, we shall follow the generalized approach of Birgé (2006) from which we borrow this substitute to Conjecture C:

**Proposition 3** Let  $X_1, \dots, X_n$  be i.i.d. random variables with an unknown density  $s$  with respect to some measure  $\nu$  on  $(E, \mathcal{E})$ . Whatever the densities  $u, v$ , one can design a procedure  $\varphi_{u,v}(X_1, \dots, X_n)$  with values in  $\{u, v\}$  and such that

$$\mathbb{P}_s [\varphi_{u,v}(X_1, \dots, X_n) = u] \leq \exp [-(n/4)h^2(u, v)] \quad \text{if } h(s, v) \leq h(u, v)/4;$$

$$\mathbb{P}_s [\varphi_{u,v}(X_1, \dots, X_n) = v] \leq \exp [-(n/4)h^2(u, v)] \quad \text{if } h(s, u) \leq h(u, v)/4.$$

The main difference with Conjecture C lies in the fact that the procedure  $\varphi_{u,v}$  does not choose between  $u$  and  $v$  by merely comparing  $\prod_{i=1}^n u(X_i)$  and  $\prod_{i=1}^n v(X_i)$ . It is more complicated. This implies that, in this case, we have to design a new estimator  $\hat{s}_{\bar{S}}(\mathbf{X})$ , based on Proposition 3, to replace the m.l.e.. The construction of this estimator is more complicated than that of the m.l.e. and we shall not describe it here. The following analogue of Theorem 1 is proved in Birgé (2006).

**Theorem 2** Let  $\mathbf{X} = (X_1, \dots, X_n)$  be an i.i.d. sample with unknown density  $s$  with respect to some measure  $\nu$  on  $(E, \mathcal{E})$  and  $(M, h)$  be the metric space of all such densities with Hellinger distance. Let  $\bar{S}$  be a subset of  $(M, h)$  with a finite metric dimension bounded by  $\bar{D}$ . Then one can build an estimator  $\hat{s}_{\bar{S}}(X_1, \dots, X_n)$  of  $s$  such that, for some universal constant  $C$ ,

$$\mathbb{E}_s [h^2(\hat{s}_{\bar{S}}, s)] \leq C \left[ \inf_{t \in \bar{S}} h^2(s, t) + n^{-1}\bar{D} \right]. \quad (4.16)$$

Analogues of Proposition 3 do hold for various statistical frameworks, although not all. Additional examples are to be found in Birgé (2004 and 2006). For each such case, one can, starting from a model  $\bar{S}$  with finite metric dimension bounded by  $\bar{D}$ , design a suitable estimator  $\hat{s}_{\bar{S}}(\mathbf{X})$  and then get an analogue of Theorem 2. Within the general framework of Section 4.1, the resulting risk bound takes the following form:

$$\mathbb{E}_s [d^2(\hat{s}_{\bar{S}}, s)] \leq C_1 \inf_{t \in \bar{S}} d^2(s, t) + C_2 \bar{D} \quad \text{for all } s \in M, \quad (4.17)$$

where the constants  $C_1$  and  $C_2$  depend on the corresponding statistical framework — compare with (4.15) and (4.16) — but not on  $s$  or  $\overline{\mathcal{S}}$ . The main task is indeed to prove the proper alternative to Proposition 3. Once this has been done, (4.17) follows more or less straightforwardly.

To what extent can maximum likelihood or related estimators provide bounds of the form (4.17) has been studied in various papers among which van de Geer (1990, 1993, 1995 and 2000), Shen and Wong (1994) and Wong and Shen (1995), Birgé and Massart (1993 and 1998), Györfi, Kohler, Kryżak and Walk (2002) and Massart (2006).

## 5 Model selection

Let us consider a statistical framework for which an analogue of Proposition 3 holds so that any model  $\overline{\mathcal{S}}$  with finite metric dimension bounded by  $\overline{D}$  provides an estimator  $\hat{s}_{\overline{\mathcal{S}}}(\mathbf{X})$  with a risk bounded by (4.17). Then the quality of a given model  $\overline{\mathcal{S}}$  for estimating  $s$  can be measured by the right-hand side of (4.17). Since this quality depends on the unknown  $s$  via the approximation term  $\inf_{t \in \overline{\mathcal{S}}} d^2(s, t)$ , we cannot know it. Introducing a large family  $\{\overline{\mathcal{S}}_m, m \in \mathcal{M}\}$  of models, each one with finite metric dimension bounded by  $\overline{D}_m$ , instead of one single model, gives more chance to get an estimator  $\hat{s}_m = \hat{s}_{\overline{\mathcal{S}}_m}$  in the family with the smaller risk bound  $\inf_{m \in \mathcal{M}} \{C_1 \inf_{t \in \overline{\mathcal{S}}} d^2(s, t) + C_2 \overline{D}_m\}$ . Since we do not know which estimator reaches this bound, the challenge of *model selection* is to design a random choice  $\hat{m}(\mathbf{X})$  of  $m$  such that the corresponding estimator  $\hat{s}_{\hat{m}}$  approximately reaches this optimal risk, i.e. satisfies

$$\mathbb{E}_s [d^2(s, \hat{s}_{\hat{m}})] \leq C \inf_{m \in \mathcal{M}} \left\{ C_1 \inf_{t \in \overline{\mathcal{S}}_m} d^2(s, t) + C_2 \overline{D}_m \right\}, \quad (5.1)$$

for some constant  $C$  independent of  $s$  and the family of models.

### 5.1 Some natural limitations to the performances of model selection

Let us show here, in the context of Gaussian regression, that getting a bound like (5.1) for arbitrary families of models is definitely too optimistic. If, in this context, (5.1) were true, we would be able to design a model selection procedure  $\hat{m}$  satisfying, in view of (4.14)

$$\mathbb{E}_s [\|s - \hat{s}_{\hat{m}}\|^2] \leq C' \inf_{m \in \mathcal{M}} \left\{ \sigma^2 \overline{D}_m + \inf_{t \in \overline{\mathcal{S}}_m} \|s - t\|^2 \right\}, \quad (5.2)$$

for some universal constant  $C'$ , independent of  $s$ ,  $n$  and the family of models. It is not difficult to see that this is impossible, even if we restrict ourselves to countable families of models. Indeed, if (5.2) were true, we could choose for  $\{\overline{\mathcal{S}}_m, m \in \mathcal{M}\}$  a countable family of one-dimensional linear spaces such that each point  $s \in \mathbb{R}^n$  could be approximated by one space in the family with arbitrary accuracy. We would then get  $\overline{D}_m = 1/2$  for each  $m$  and (5.2) would imply that

$$\mathbb{E}_s [\|s - \hat{s}_{\hat{m}}\|^2] \leq C' \sigma^2 / 2 \quad \text{for all } s \in \mathbb{R}^n.$$

But it is known that the best bound one can expect for any estimator  $\hat{s}$  uniformly with respect to  $s \in \mathbb{R}^n$  is

$$\sup_{s \in \mathbb{R}^n} \mathbb{E}_s [\|s - \hat{s}\|^2] = n\sigma^2,$$

which contradicts the fact that  $C'$  should be a universal constant. One actually has to pay a price for using many models simultaneously and, as we shall see, this price depends on the *complexity* (with a suitable sense) of the chosen family of models.

## 5.2 Risk bounds for model selection

### 5.2.1 The main theorems

We shall not get here into the details of the construction of the selection procedure that we use but content ourselves to give the main results and analyze their consequences. A key idea for the construction appeared in Barron and Cover (1991). Further approaches to selection procedures have been developed in Barron, Birgé and Massart (1999), Birgé and Massart (1997 and 2001), van de Geer (2000), Györfi, Kohler, Kryżak and Walk (2002) and Massart (2006) who provides an extensive list of references. We follow here the approach based on dimension from Birgé (2006), providing hereafter two theorems corresponding to our two problems of interest, Gaussian regression and density estimation. In both cases, the construction of the estimators requires the introduction of a family of positive weights  $\{\Delta_m, m \in \mathcal{M}\}$ , to be chosen by the statistician and satisfying the condition

$$\sum_{m \in \mathcal{M}} \exp[-\Delta_m] \leq 1. \quad (5.3)$$

In case of equality in (5.3), the family  $\{q_m\}_{m \in \mathcal{M}}$  with  $q_m = \exp[-\Delta_m]$  defines a probability  $Q$  on the family of models and choosing a large value for  $\Delta_m$  means putting a small probability on the model  $\bar{S}_m$ . One can then see  $q_m$  as a probability that the statistician puts on  $\bar{S}_m$  and which influences the result of the estimation procedure, as shown by the next theorems. Such an interpretation of the weights  $\Delta_m$  corresponds to the so-called *Bayesian* point of view. A detailed analysis of this interpretation can be found in Birgé and Massart (2001, Sect. 3.4).

**Theorem 3** *Let  $\mathbf{X}$  be a Gaussian vector in  $\mathbb{R}^n$  with unknown mean  $s$  and known covariance matrix  $\sigma^2 I_n$ . Let  $\{\bar{S}_m, m \in \mathcal{M}\}$  be a finite or countable family of subsets of  $\mathbb{R}^n$  with finite metric dimensions bounded by  $\bar{D}_m$ , respectively. Let  $\{\Delta_m, m \in \mathcal{M}\}$  be a family of positive weights satisfying (5.3). One can build an estimator  $\tilde{s}(\mathbf{X})$  of  $s$  such that, for some universal constant  $C$ ,*

$$\mathbb{E}_s [\|s - \tilde{s}\|^2] \leq C \inf_{m \in \mathcal{M}} \left\{ \sigma^2 \max\{\bar{D}_m, \Delta_m\} + \inf_{t \in \bar{S}_m} \|s - t\|^2 \right\}. \quad (5.4)$$

**Theorem 4** *Let  $\mathbf{X} = (X_1, \dots, X_n)$  be an i.i.d. sample with unknown density  $s$  with respect to some measure  $\nu$  on  $(E, \mathcal{E})$  and  $(M, h)$  be the metric space of all such densities with Hellinger distance. Let  $\{\bar{S}_m, m \in \mathcal{M}\}$  be a finite or countable family of subsets of  $(M, h)$  with finite metric dimensions bounded by  $\bar{D}_m$ , respectively. Let  $\{\Delta_m, m \in \mathcal{M}\}$  be a family of positive weights satisfying (5.3). One can build an estimator  $\tilde{s}(X_1, \dots, X_n)$  of  $s$  such that, for some universal constant  $C$ ,*

$$\mathbb{E}_s [h^2(\tilde{s}, s)] \leq C \inf_{m \in \mathcal{M}} \left\{ n^{-1} \max\{\bar{D}_m, \Delta_m\} + \inf_{t \in \bar{S}_m} h^2(s, t) \right\}. \quad (5.5)$$

*Remark:* The choice of the bound 1 in (5.3) has nothing canonical and was simply made for convenience. Any small constant would do since we did not provide the actual value of  $C$  which depends on the right-hand side of (5.3).



### 5.2.2 About the complexity of families of models

The only difference between the ideal bound (5.2) and (5.4) is the replacement of  $\overline{D}_m$  by  $\max\{\overline{D}_m, \Delta_m\}$  with weights  $\Delta_m$  satisfying (5.3) and we see, comparing (4.16) and (5.5), that the same difference holds for density estimation. More generally, in a framework for which an analogue of Proposition 3 holds, leading to (4.17), we proved in Birgé (2006) that

$$\mathbb{E}_s [d^2(s, \hat{s}_{\hat{m}})] \leq C \inf_{m \in \mathcal{M}} \left\{ C_1 \inf_{t \in \overline{S}} d^2(s, t) + C_2 \max\{\overline{D}_m, \Delta_m\} \right\}, \quad (5.6)$$

holds instead of (5.1). In all situations, apart from the constant  $C$ , the loss with respect to the ideal bound is due to the replacement of  $\overline{D}_m$  by  $\max\{\overline{D}_m, \Delta_m\}$  where the weights  $\Delta_m$  satisfy (5.3). If  $\Delta_m$  is not much larger than  $\overline{D}_m$  for all  $m$ , we have almost reached the ideal risk, otherwise not and we can now explain what we mean by the complexity of a family of models.

For each positive integer  $j$ , let us denote by  $H(j)$  the cardinality of the set  $\mathcal{M}_j$  of those  $m$  such that  $j/2 \leq \overline{D}_m < (j+1)/2$ . If  $H(j)$  is finite for all  $j$ , let us choose  $\Delta_m = (j+1)/2 + \log_+(H(j))$  for  $m \in \mathcal{M}_j$  where  $\log_+(x) = \log x$  for  $x \geq 1$  and  $\log_+(0) = 0$ . Then

$$\sum_{m \in \mathcal{M}} \exp[-\Delta_m] = \sum_{j \geq 1} \sum_{m \in \mathcal{M}_j} \exp[-(j+1)/2 - \log_+(H(j))] \leq \sum_{i \geq 2} \exp[-i/2] < 1$$

and (5.3) holds. Moreover,

$$\max\{\overline{D}_m, \Delta_m\} = \Delta_m \leq 2\overline{D}_m[1 + j^{-1} \log_+(H(j))] \quad \text{for } m \in \mathcal{M}_j.$$

If  $j^{-1} \log_+[H(j)]$  is uniformly bounded and the bound is not large, then (5.6) and (5.1) are comparable and we can consider that the family of models is not complex. On the other hand, if, for some  $j$ ,  $\log[H(j)]$  is substantially larger than  $j$ ,  $\Delta_m$  is substantially larger than  $\overline{D}_m$ , at least for some  $m$ , which may result in a bound (5.6) much larger than (5.1). If  $H(j) = +\infty$  for some  $j$ , (5.3) requires that  $\Delta_m$  be unbounded for  $m \in \mathcal{M}_j$ , which is even worse. A reasonable measure of the complexity of a family of models is therefore  $\sup_{j \geq 1} j^{-1} \log_+[H(j)]$ , high complexity of the family corresponding to large values of this index.

### 5.3 Application 1: variable selection in Gaussian regression

Let us now give some concrete illustrations of more or less complex families of models corresponding to the examples that motivated our investigations about model selection. To begin with, we consider the situation of Section 3.2.1 with a large number  $p \leq n$  of potentially influential explanatory variables  $Z^j$  and set  $\Lambda = \{1; \dots; p\}$ . For any subset  $m$  of  $\Lambda$  we define  $\overline{S}_m$  as the linear span of the vectors  $Z^j$  for  $j \in m$ . According to Section 4.4.2,  $\overline{S}_m$  has a metric dimension bounded by  $|m|/2$ .

Let us assume that we have ordered the variables according to their supposed relevance,  $Z^1$  being the more relevant. In such a situation it is natural to consider the models spanned by the  $q$  more relevant variables  $Z^1, \dots, Z^q$  for  $1 \leq q \leq p$  and therefore to set  $\mathcal{M} = \mathcal{M}_1 = \{\{1; \dots; q\}, 1 \leq q \leq p\}$ . This is not a complex family of models and the choice  $\Delta_m = |m|$  ensures that (5.3) holds. It follows from Theorem 3 that one can design an estimator  $\tilde{s}(\mathbf{X})$  satisfying

$$\mathbb{E}_s [\|s - \tilde{s}\|^2] \leq C \inf_{m \in \mathcal{M}_1} \left\{ \sigma^2 |m| + \inf_{t \in \overline{S}_m} \|s - t\|^2 \right\}. \quad (5.7)$$

Comparing this with the performance of the m.l.e. with respect to each model  $\overline{\mathcal{S}}_m$  given by (3.4), we see that, apart from the constant  $C$ , we recover the performance of the best model in the family.

This simple approach has, nevertheless, some drawbacks. First, we have to order the explanatory variables which is often not easy. Then the result is really bad if we make a serious mistake in this ordering. Imagine, for instance, that  $s$  only depends on four highly influential variables so that if the variables had been ordered correctly, the best model, i.e. the one minimizing  $\sigma^2|m| + \inf_{t \in \overline{\mathcal{S}}_m} \|s - t\|^2$ , would be  $\overline{\mathcal{S}}_{\{1;2,3,4\}}$  and the corresponding risk  $4\sigma^2$ . If one of these four very influential variables has been neglected and appears in the sequence with a high index  $l$ , it may happen that, because of this wrong ordering, the best model becomes  $\overline{\mathcal{S}}_{\{1;\dots;l\}}$  leading to the much higher risk  $\sigma^2 l$ .

In order to avoid the difficulties connected with variables ordering, one may introduce many more models, defining  $\mathcal{M} = \mathcal{M}_2$  as the set of all nonvoid subsets  $m$  of  $\Lambda$ . Since the number of nonvoid subsets of  $\Lambda$  with cardinality  $q$  is  $\binom{p}{q} \leq p^q/q!$ , we may choose  $\Delta_m = 1 + |m| \log p$  to get (5.3) so that, by Theorem 3, one can find an estimator  $\tilde{s}(\mathbf{X})$  satisfying

$$\mathbb{E}_s [\|s - \tilde{s}\|^2] \leq C \inf_{m \in \mathcal{M}} \left\{ \sigma^2(1 + |m| \log p) + \inf_{t \in \overline{\mathcal{S}}_m} \|s - t\|^2 \right\}. \quad (5.8)$$

With this method, we avoid the problems connected with variables ordering and may even introduce more explanatory variables than observations ( $p > n$ ), hoping that with so many variables at disposal, one can find a small subset  $m$  of them that provides an accurate model for  $s$ . There is a price to pay for that! We now have a complex family of models when  $p$  is large resulting in values of  $\Delta_m$  which are much larger than  $|m|$  and we pay the extra factor  $\log p$  in our risk bounds.

One can actually cumulate the advantages of the two approaches by mixing the two families in the following way. We first order the  $p$  variables as we did at the beginning, giving the smallest indices to the variables we believe are more influential and set again  $\mathcal{M} = \mathcal{M}_2$ . We then fix  $\Delta_m = |m| + 1/2$  for  $m \in \mathcal{M}_1$  and  $\Delta_m = 1 + |m| \log p$  for  $m \in \mathcal{M} \setminus \mathcal{M}_1$  so that (5.3) still holds. Theorem 3 shows that

$$\begin{aligned} \mathbb{E}_s [\|s - \tilde{s}\|^2] \leq C \min & \left[ \inf_{m \in \mathcal{M} \setminus \mathcal{M}_1} \left\{ \sigma^2(1 + |m| \log p) + \inf_{t \in \overline{\mathcal{S}}_m} \|s - t\|^2 \right\} \right. \\ & \left. ; \inf_{m \in \mathcal{M}_1} \left\{ \sigma^2(|m| + 1/2) + \inf_{t \in \overline{\mathcal{S}}_m} \|s - t\|^2 \right\} \right]. \end{aligned}$$

If our ordering of the variables is right, the best  $m$  belongs to  $\mathcal{M}_1$  and we get an analogue of (5.7). If not, we lose a factor  $\log p$  from the risk of the best model as in (5.8).

## 5.4 Application 2: histograms and density estimation

### 5.4.1 Problems connected with the use of the $\mathbb{L}_2$ -distance in density estimation

Let us now come back to density estimation with histograms. In Section 1.2 we used the  $\mathbb{L}_2$ -distance to measure the distortion between  $s$  and its estimator. This is certainly the most popular and more widely studied measure of distortion for density

estimation but it actually has some serious drawbacks as shown by Devroye and Györfi (1985). For histograms it results in risk bounds (1.10) depending on  $\|s\|_\infty$  for irregular partitions, which are not of the form

$$R(\hat{s}_m, s) \leq C [\|s - s_m\|^2 + n^{-1}|m|],$$

for some universal constant  $C$ , independent of  $s$ ,  $n$  and the partition  $m$ . It is actually impossible to get an analogue of Theorem 2 where the  $\mathbb{L}_2$ -distance would replace the Hellinger distance, as shown by the following proposition motivated by Theorem 2.1 of Rigollet and Tsybakov (2005). Indeed, if such a theorem were true, we could apply it to the model  $\overline{\mathcal{S}}$  provided by this proposition and conclude that the corresponding estimator  $\hat{s}_{\overline{\mathcal{S}}}$  would satisfy the analogue of (4.16) leading to the uniform risk bound

$$\mathbb{E}_s [\|\hat{s}_{\overline{\mathcal{S}}} - s\|^2] \leq CD/(2n), \quad \text{for all } s \in \overline{\mathcal{S}}$$

and some universal constant  $C$ , therefore independent of  $L$ . This would clearly contradict (5.9) below for large enough values of  $L$ .

**Proposition 4** *For each  $L > 0$  and each integer  $D$  with  $1 \leq D \leq 3n$ , one can find a finite set  $\overline{\mathcal{S}}$  of densities with the following properties:*

- i) *it is a subset of some  $D$ -dimensional affine subspace of  $\mathbb{L}_2([0, 1], dx)$  with a metric dimension bounded by  $D/2$ ;*
- ii)  $\sup_{s \in \overline{\mathcal{S}}} \|s\|_\infty \leq L + 1$ ;
- iii) *for any estimator  $\hat{s}(X_1, \dots, X_n)$  belonging to  $\mathbb{L}_2([0, 1], dx)$  and based on an i.i.d. sample with density  $s \in \overline{\mathcal{S}}$ ,*

$$\sup_{s \in \overline{\mathcal{S}}} \mathbb{E}_s [\|\hat{s} - s\|^2] > 0.0139DLn^{-1}. \quad (5.9)$$

*Proof:* Let us set  $a = D/(4n) \leq 3/4$ , define  $\theta$  by  $(1 - \theta)/\theta = 4nL/D$  and introduce the functions  $f(x) = \mathbb{1}_{[0, 1]}(x)$  and  $g(x) = -a\mathbb{1}_{[0, (1-\theta)/D]} + a(1 - \theta)\theta^{-1}\mathbb{1}_{[(1-\theta)/D], 1/D}[\cdot]$ . Then  $\int_0^{1/D} g(x) dx = 0$ ,  $\sup_x g(x) = L$ ,  $\inf_x g(x) = -a \geq -3/4$  and

$$\|g\|^2 = \int_0^{1/D} g^2(x) dx = a^2 \frac{1 - \theta}{D} [1 + (1 - \theta)\theta^{-1}] = \frac{a^2(1 - \theta)}{\theta D} = \frac{L}{4n}. \quad (5.10)$$

It follows that  $\|f - (f + g)\|^2 = L/(4n)$ . Moreover

$$\begin{aligned} h^2(f, f + g) &= \frac{1}{2} \int_0^{1/D} [1 - \sqrt{1 + g(x)}]^2 dx \\ &= \frac{1}{2} \int_0^{1/D} [2 + g(x) - 2\sqrt{1 + g(x)}] dx = \frac{1}{D} - \int_0^{1/D} \sqrt{1 + g(x)} dx \\ &= D^{-1} [1 - (1 - \theta)\sqrt{1 - a} - \theta\sqrt{1 + a(1 - \theta)\theta^{-1}}] \\ &\leq D^{-1} [1 - \sqrt{1 - a}] \leq D^{-1}(2a/3) = (6n)^{-1}, \end{aligned} \quad (5.11)$$

since  $a \leq 3/4$ . Let us now set, for  $1 \leq j \leq D$ ,  $g_j(x) = g(x - D^{-1}(j - 1))$ , so that these  $D$  translates of  $g$  have disjoint supports and  $g_1 = g$ . Let  $\mathcal{D} = \{0; 1\}^D$  with the distance  $\Delta$  given by  $\Delta(\delta, \delta') = \sum_{j=1}^D |\delta_j - \delta'_j|$ . For each  $\delta \in \mathcal{D}$  we consider the density  $s_\delta(x) = f(x) + \sum_{j=1}^D \delta_j g_j(x)$  and set  $\overline{\mathcal{S}} = \{s_\delta, \delta \in \mathcal{D}\}$ . Clearly  $\|s_\delta\|_\infty \leq L + 1$  for all  $\delta \in \mathcal{D}$  and it follows from (5.10) that

$$\|s_\delta - s_{\delta'}\|^2 = \sum_{j=1}^D (\delta_j - \delta'_j)^2 \int_0^{1/D} g_j^2(x) dx = \frac{L}{4n} \sum_{j=1}^D (\delta_j - \delta'_j)^2 = \frac{L}{4n} \Delta(\delta, \delta'). \quad (5.12)$$

Moreover, since  $\bar{S}$  is a subset of some  $D$ -dimensional affine subspace of  $\mathbb{L}_2([0, 1], dx)$ , it follows from the arguments used in the proof of Proposition 2 that its metric dimension is bounded by  $D/2$ .

Defining  $P_\delta$  by  $dP_\delta/dx = s_\delta$ , we derive from (5.11) that  $h^2(P_\delta, P_{\delta'}) \leq (6n)^{-1}$ , hence  $\rho(P_\delta, P_{\delta'}) \geq \bar{\rho} = 1 - (6n)^{-1}$ , for each pair  $(\delta, \delta') \in \mathcal{D}^2$  such that  $\Delta(\delta, \delta') = 1$ . We may then apply Assouad's Lemma below to conclude from (5.12) that, whatever the estimator  $\hat{\delta}$  with values in  $\mathcal{D}$ ,

$$\sup_{\delta \in \mathcal{D}} \mathbb{E}_s [\|s_{\hat{\delta}} - s_\delta\|^2] = \frac{L}{4n} \sup_{\delta \in \mathcal{D}} \mathbb{E}_s [\Delta(\hat{\delta}, \delta)] \geq \frac{L}{4n} \frac{D}{2} \left[ 1 - \sqrt{1 - [1 - (6n)^{-1}]^{2n}} \right].$$

Let  $\hat{s}$  be any density estimator based on  $X_1, \dots, X_n$  and set  $\hat{\delta}(X_1, \dots, X_n)$  to satisfy  $\|\hat{s} - s_{\hat{\delta}}\| = \inf_{\delta \in \mathcal{D}} \|\hat{s} - s_\delta\|$  so that, whatever  $\delta \in \mathcal{D}$ ,  $\|s_{\hat{\delta}} - s_\delta\| \leq 2\|\hat{s} - s_\delta\|$ . We derive from our last bound that

$$\sup_{\delta \in \mathcal{D}} \mathbb{E}_s [\|\hat{s} - s_\delta\|^2] \geq \frac{1}{4} \sup_{\delta \in \mathcal{D}} \mathbb{E}_s [\|s_{\hat{\delta}} - s_\delta\|^2] \geq \frac{LD}{32n} \left[ 1 - \sqrt{1 - [1 - (6n)^{-1}]^{2n}} \right].$$

We conclude by observing that  $[1 - (6n)^{-1}]^{2n}$  is increasing with  $n$ , hence  $\geq 25/36$ .  $\square$

**Lemma 2 (Assouad, 1983)** *Let  $\{P_\delta, \delta \in \mathcal{D}\}$  be a family of distributions indexed by  $\mathcal{D} = \{0; 1\}^D$  and  $X_1, \dots, X_n$  an i.i.d. sample from a distribution in the family. Assume that  $\rho(P_\delta, P_{\delta'}) \geq \bar{\rho}$  for each pair  $(\delta, \delta') \in \mathcal{D}^2$  such that  $\Delta(\delta, \delta') = 1$ . Then for any estimator  $\hat{\delta}(X_1, \dots, X_n)$  with values in  $\mathcal{D}$ ,*

$$\sup_{\delta \in \mathcal{D}} \mathbb{E}_\delta [\Delta(\hat{\delta}(X_1, \dots, X_n), \delta)] \geq \frac{D}{2} \left[ 1 - \sqrt{1 - \bar{\rho}^{2n}} \right] \geq \frac{D\bar{\rho}^{2n}}{4}, \quad (5.13)$$

where  $\mathbb{E}_\delta$  denotes the expectation when the  $X_i$  have the distribution  $P_\delta$ .

*Proof:* Let us set  $P_\delta^n$  for the joint distribution of the  $X_i$  with individual distribution  $P_\delta$  and consider some measure  $\mu$  which dominates the probabilities  $P_\delta^n$  for  $\delta \in \mathcal{D}$ . First note that the left-hand side of (5.13) is at least as large as the average risk

$$R_B = 2^{-D} \sum_{\delta \in \mathcal{D}} \mathbb{E}_\delta [\Delta(\hat{\delta}, \delta)] = 2^{-D} \sum_{\delta \in \mathcal{D}} \int \sum_{k=1}^D |\hat{\delta}_k - \delta_k| dP_\delta^n.$$

Then, setting  $Q_k^j = 2^{-D+1} \sum_{\{\delta \in \mathcal{D} \mid \delta_k=j\}} P_\delta^n$  with  $j = 0$  or  $1$ , we get

$$\begin{aligned} R_B &= 2^{-D} \sum_{k=1}^D \left( \sum_{\{\delta \in \mathcal{D} \mid \delta_k=0\}} \int \hat{\delta}_k dP_\delta^n + \sum_{\{\delta \in \mathcal{D} \mid \delta_k=1\}} \int (1 - \hat{\delta}_k) dP_\delta^n \right) \\ &= \frac{1}{2} \sum_{k=1}^D \left( \int \hat{\delta}_k \frac{dQ_k^0}{d\mu} d\mu + \int (1 - \hat{\delta}_k) \frac{dQ_k^1}{d\mu} d\mu \right) \\ &\geq \frac{1}{2} \sum_{k=1}^D \int \inf \left\{ \frac{dQ_k^0}{d\mu}, \frac{dQ_k^1}{d\mu} \right\} d\mu. \end{aligned}$$

Since  $\inf\{x; y\}$  is a concave function of the pair  $(x, y)$ , it follows that

$$\inf \left\{ \frac{dQ_k^0}{d\mu}, \frac{dQ_k^1}{d\mu} \right\} \geq 2^{-D+1} \sum_{(\delta, \delta') \in \mathcal{D}_k} \inf \left\{ \frac{dP_\delta^n}{d\mu}, \frac{dP_{\delta'}^n}{d\mu} \right\},$$

with  $\mathcal{D}_k = \{(\delta, \delta') \mid \delta_k = 0, \delta'_k = 1, \delta_j = \delta'_j \text{ for } j \neq k\}$ , hence

$$R_B \geq \frac{1}{2} \sum_{k=1}^D 2^{-D+1} \sum_{(\delta, \delta') \in \mathcal{D}_k} \int \inf \left\{ \frac{dP_\delta^n}{d\mu}; \frac{dP_{\delta'}^n}{d\mu} \right\} d\mu.$$

We now use (4.8) to conclude that

$$R_B \geq \frac{1}{2} \sum_{k=1}^D 2^{-D+1} \sum_{(\delta, \delta') \in \mathcal{D}_k} \left[ 1 - \sqrt{1 - \rho^2(P_\delta^n, P_{\delta'}^n)} \right].$$

By assumption,  $\rho(P_\delta, P_{\delta'}) \geq \bar{\rho}$  for  $(\delta, \delta') \in \mathcal{D}_k$ , hence  $\rho^2(P_\delta^n, P_{\delta'}^n) = \rho^{2n}(P_\delta, P_{\delta'}) \geq \bar{\rho}^{2n}$ . The conclusion follows.  $\square$

#### 5.4.2 Partition selection for histograms

If we use the Hellinger distance instead of the  $\mathbb{L}_2$ -distance to evaluate the risk of histograms, we can improve (1.10), getting a universal bound which does not involve  $\|s\|_\infty$ . We recall that  $S_m$  is the set of densities which are constant on the elements of the partition  $m$  as defined in (1.11).

**Theorem 5** *Let  $s$  be some density with respect to the Lebesgue measure on  $[0, 1]$ ,  $X_1, \dots, X_n$  be an  $n$ -sample from the corresponding distribution and  $m = \{I_0, \dots, I_D\}$  be a partition of  $[0, 1]$  into intervals  $I_j$  with respective lengths  $|I_j|$ . Let  $\hat{s}_m$  be the histogram estimator based on this partition and given by*

$$\hat{s}_m(x) = \sum_{j=0}^D \left[ \frac{1}{n|I_j|} \sum_{i=1}^n \mathbb{1}_{I_j}(X_i) \right] \mathbb{1}_{I_j}(x).$$

The Hellinger risk of  $\hat{s}_m$  is bounded by

$$\mathbb{E}_s [h^2(s, \hat{s}_m)] \leq 2 \inf_{t \in S_m} h^2(s, t) + D/(2n). \quad (5.14)$$

*Proof:* It is shown in Birgé and Rozenholc (2006) that

$$\mathbb{E}_s [h^2(s, \hat{s}_m)] \leq h^2(s, s_m) + \frac{D}{2n} \quad \text{with } s_m = \sum_{j=0}^D \left[ \frac{1}{|I_j|} \int_{I_j} s(x) dx \right] \mathbb{1}_{I_j}.$$

Let  $f$  be the  $\mathbb{L}_2$ -orthogonal projection of  $\sqrt{s}$  onto the linear span  $V_m$  of  $\mathbb{1}_{I_0}, \dots, \mathbb{1}_{I_D}$ . Then

$$f = \sum_{j=0}^D \left[ \frac{1}{|I_j|} \int_{I_j} \sqrt{s(x)} dx \right] \mathbb{1}_{I_j} \quad \text{and} \quad \|f - \sqrt{s}\|^2 \leq 2h^2(s, t) \quad \text{for all } t \in V_m.$$

Setting  $s_m = \sum_{j=0}^D a_j \mathbb{1}_{I_j}$  and  $f = \sum_{j=0}^D b_j \mathbb{1}_{I_j}$ , we get from Jensen's Inequality that  $b_j \leq \sqrt{a_j}$ . It follows that

$$h^2(s, s_m) = 1 - \sum_{j=0}^D \int_{I_j} \sqrt{a_j s(x)} dx = 1 - \sum_{j=0}^D \sqrt{a_j} b_j |I_j| \leq 1 - \sum_{j=0}^D b_j^2 |I_j|,$$

while

$$\|f - \sqrt{s}\|^2 = 1 + \sum_{j=0}^D \int_{I_j} b_j^2 dx - 2 \sum_{j=0}^D \int_{I_j} b_j \sqrt{s(x)} dx = 1 - \sum_{j=0}^D b_j^2 |I_j|.$$

Hence

$$h^2(s, s_m) \leq \|f - \sqrt{s}\|^2 \leq 2 \inf_{t \in S_m} h^2(s, t). \quad \square \quad (5.15)$$

If, in particular,  $\sqrt{s}$  is Hölder continuous and satisfies (1.13), we derive as in Section 1.3.3 that one can find a regular partition  $m$ , depending on  $L$  and  $\beta$ , such that,

$$\mathbb{E}_s [h^2(s, \hat{s}_m)] \leq \max \left\{ (5/2) \left( Ln^{-\beta} \right)^{2/(2\beta+1)} ; n^{-1} \right\}. \quad (5.16)$$

Then, a useful remark is as follows. If we have at disposal a sample  $X_1, \dots, X_{2n}$  of size  $2n$  and a family  $\mathcal{M}$  of partitions of  $[0, 1]$ , one can use the first half of the sample to build the corresponding histograms  $\hat{s}_m(X_1, \dots, X_n)$  and use the second half of the sample to select one estimator in the family. For this, we merely have to apply Theorem 4 to the sample  $X_{n+1}, \dots, X_{2n}$  conditionally on  $X_1, \dots, X_n$ . Conditionally on  $X_1, \dots, X_n$ , each histogram  $\hat{s}_m$  is simply a density which can be considered as a model  $\bar{S}_m$  containing only one point, hence with a finite metric dimension bounded by  $1/2$ . Let  $\{\Delta_m, m \in \mathcal{M}\}$  be a family of weights satisfying

$$\sum_{m \in \mathcal{M}} \exp[-\Delta_m] \leq 1 \quad \text{and} \quad \Delta_m \geq 1 \quad \text{for all } m. \quad (5.17)$$

We derive from Theorem 4 applied to the models  $\bar{S}_m = \{\hat{s}_m\}$  that there exists an estimator  $\tilde{s}(X_1, \dots, X_{2n})$  such that

$$\mathbb{E}_s [h^2(\tilde{s}, s) | X_1, \dots, X_n] \leq C \inf_{m \in \mathcal{M}} \{n^{-1} \Delta_m + h^2(s, \hat{s}_m(X_1, \dots, X_n))\}.$$

Integrating with respect to  $X_1, \dots, X_n$  and using (5.14) finally leads to

$$\begin{aligned} \mathbb{E}_s [h^2(\tilde{s}, s)] &\leq C \inf_{m \in \mathcal{M}} \left\{ n^{-1} \Delta_m + 2 \inf_{t \in S_m} h^2(s, t) + (|m| - 1)/(2n) \right\} \\ &\leq C' \inf_{m \in \mathcal{M}} \left\{ n^{-1} \max\{|m|, \Delta_m\} + \inf_{t \in S_m} h^2(s, t) \right\}. \end{aligned} \quad (5.18)$$

### 5.4.3 A straightforward application of partition selection

To give a concrete application of this result, let us introduce some special classes of partitions. For any finite partition  $m = \{I_0, \dots, I_D\}$  into intervals, we denote by  $A_m$  the set  $\{y_0 < \dots < y_{D+1}\}$ ,  $y_0 = 0, y_{D+1} = 1$  of endpoints of the intervals  $I_j$ . Introducing, for  $k \geq 1$ , the set  $\mathcal{J}_k$  of dyadic numbers  $\{j2^{-k}, 0 \leq j \leq 2^k\}$ , we denote by  $\mathcal{M}_{D,k}$ , for  $1 \leq D < 2^k$ , the set of those partitions  $m$  which satisfy

$$|m| = D + 1; \quad A_m \in \mathcal{J}_k \quad \text{and} \quad A_m \notin \mathcal{J}_{k-1}.$$

Denoting by  $m_0$  the trivial partition with one element  $[0, 1]$ , we define  $\mathcal{M}$  by

$$\mathcal{M} = \{m_0\} \cup \left( \bigcup_{k \geq 1} \bigcup_{1 \leq D < 2^k} \mathcal{M}_{D,k} \right).$$

The partitions in  $\mathcal{M}$  are dense in the set of finite partitions into intervals in the following sense: given any such partition  $m$ , an element  $t$  in  $S_m$ , as defined by (1.11), and  $\varepsilon > 0$ , we can find  $m' \in \mathcal{M}$  and  $t' \in S_{m'}$  such that  $h(t, t') \leq \varepsilon$ . This means that the approximation properties of  $\bigcup_{m \in \mathcal{M}} S_m$  are the same as those of all possible histograms. Since  $|\mathcal{M}_{D,k}| \leq \binom{2^k - 1}{D} \leq 2^{kD}$ , if we set  $\Delta_m = \Delta_m^0 = [(k+1)(D+1) + 1] \log 2$  for  $m \in \mathcal{M}_{D,k}$  and  $\Delta_{m_0} = 1$ , we get

$$\sum_{k \geq 1} \sum_{1 \leq D < 2^k} \sum_{\mathcal{M}_{D,k}} e^{-\Delta_m^0} \leq \sum_{k \geq 1} \sum_{1 \leq D < 2^k} 2^{-k-D-2} \leq \frac{1}{4} \sum_{k \geq 1} 2^{-k} \sum_{D \geq 1} 2^{-D} = \frac{1}{4}.$$

It follows that (5.17) holds so that by (5.18), one can find an estimator  $\tilde{s}(X_1, \dots, X_{2n})$  which satisfies

$$\mathbb{E}_s [h^2(\tilde{s}, s)] \leq C \inf_{k \geq 1} \inf_{1 \leq D < 2^k} \inf_{m \in \mathcal{M}_{D,k}} \left\{ \frac{kD}{n} + \inf_{t \in S_m} h^2(s, t) \right\}. \quad (5.19)$$

If, in the right-hand side of (5.19), we set  $m$  to be the regular partition with  $2^k$  elements, which belongs to  $\mathcal{M}_{D,2^k-1}$ , we get a bound of the form

$$\mathbb{E}_s [h^2(\tilde{s}, s)] \leq C \left[ k2^k n^{-1} + \inf_{t \in S_m} h^2(s, t) \right].$$

For densities  $s$  with  $\sqrt{s}$  satisfying (1.13), we get

$$\mathbb{E}_s [h^2(\tilde{s}, s)] \leq C' \inf_{k \geq 1} \left\{ k2^k n^{-1} + L^2 2^{-2k\beta} \right\},$$

but an optimization with respect to  $k$  does not allow to recover the bound (5.16) because of an extra factor  $\log(nL^2)$ . This factor is connected with the complexity of the families  $\mathcal{M}_{D,k}$  which forces us to fix  $\Delta_m$  much larger than  $|m| = D + 1$  for most elements of  $\mathcal{M}_{D,k}$  when  $k$  is large. Most, but not all! It is in particular easy to modify the value of  $\Delta_m$  for the regular partitions without violating (5.17). If  $m_k$  denotes the regular partition with  $2^k$  elements and  $\mathcal{M}_R$  the set of such partitions, we may choose  $\Delta_{m_k} = |m_k|$  instead of  $\Delta_m^0$  so that

$$\sum_{m \in \mathcal{M}_R} e^{-\Delta_m} = \sum_{k \geq 0} e^{-2^k} < 0.522$$

and (5.17) still holds. It is easy to check that, with this new choice of the weights for the regular partitions, we improve the estimation for those densities such that  $\sqrt{s}$  is Hölder continuous. In particular, if  $\sqrt{s}$  satisfies (1.13) for some unknown values of  $L$  and  $\beta$ ,

$$\mathbb{E}_s [h^2(\tilde{s}, s)] \leq C \max \left\{ \left( Ln^{-\beta} \right)^{2/(2\beta+1)} ; n^{-1} \right\}, \quad (5.20)$$

which is comparable to (5.16) although  $L$  and  $\beta$  are unknown, the only loss being at the level of the constant  $C$ .

#### 5.4.4 Introducing more sophisticated Approximation Theory

The consequences of the previous modification of the weights for partitions in  $\mathcal{M}_R$  is a simple illustration of the use of elementary Approximation Theory to improve

the estimation of smooth densities. One can actually do much better with the use of more sophisticated Approximation Theory. In a milestone paper, Birman and Solomjak (1967) introduced a family  $\mathcal{M}_T$  of partitions of the cube  $[0, 1]^k$  which are such that piecewise constant (and more generally piecewise polynomials) based on the partitions in the family have excellent approximation properties with respect to functions in Sobolev spaces (and functions of bounded variation when  $k = 1$ ). Moreover, Birman and Solomjak provide a control on the number of such partitions with a given cardinality. For the case  $k = 1$  which is the one we deal with here, the number of elements  $m$  of  $\mathcal{M}_T$  with  $|m| = D$  is bounded  $4^D$  which allows us to set  $\Delta_m = 2D$  for those partitions.

The algorithm leading to the construction of the partitions in  $\mathcal{M}_T$ , which is called an “adaptive approximation algorithm”, is also described in Section 3.3 of DeVore (1998) and it works as follows. We choose a positive threshold  $\varepsilon$  and some non-negative functional  $J(f, I)$  depending on the function  $f$  to be approximated and the interval  $I$ . Roughly speaking, the functional measures the quality of approximation of  $f$  by a piecewise constant (or more generally a piecewise polynomial) function on  $I$ . At step one, the algorithm starts with the trivial partition  $m^1 = m_0$  with one single interval. At step  $j$  it provides a partition  $m^j$  into  $j$  intervals and it checks whether  $\sup_{I \in m^j} J(f, I) \leq \varepsilon$  or not. If this is the case, the algorithm stops, if not we choose one of the intervals  $I$  for which the criterion  $J(f, I) \leq \varepsilon$  is violated and divide it into two interval of equal length to derive  $m^{j+1}$ . Then we iterate the procedure. For the functions  $f$  of interest, which satisfy some smoothness condition related to the functional  $J$ , the procedure necessarily stops at some stage, leading to a final partition  $m$ . Let  $\mathcal{M}_T$  be the set of all the partitions that can be obtained in this way. Then  $\mathcal{M}_R \subset \mathcal{M}_T$ . Building a partition  $m$  in  $\mathcal{M}_T$  is actually equivalent to growing a complete binary tree for which the initial interval  $[0, 1]$  corresponds to the root of the tree, each node of the tree to an interval and each split of an interval to adding two sons to a terminal node of the tree, the partition  $m$  being in one-to-one correspondance to the set of terminal nodes of the tree. When viewed as a tree algorithm, this construction is similar to the CART algorithm of Breiman, Friedman, Olshen and Stone (1984). The analysis of CART from the model selection point of view that we explain here has been made by Gey and Nédélec (2005).

It follows from the correspondence between the partitions in  $\mathcal{M}_T$  and the complete binary trees that the number of elements  $m$  of  $\mathcal{M}_T$  such that  $|m| = j + 1$ ,  $j \in \mathbb{N}$ , is equal to the number of complete binary trees with  $j + 1$  terminal nodes which is given by the Catalan numbers  $(j + 1)^{-1} \binom{2j}{j}$ . Setting  $\Delta_m^1 = 2|m|$  for  $m \in \mathcal{M}_T$  and using  $\binom{2j}{j} \leq 4^j$  which follows from Stirling’s expansion, we derive that

$$\sum_{m \in \mathcal{M}_T} e^{-\Delta_m^1} \leq \sum_{j \geq 0} \frac{e^{-2(j+1)}}{j+1} \binom{2j}{j} \leq \sum_{j \geq 0} \frac{4^j e^{-2(j+1)}}{j+1} < \frac{1}{4}.$$

It follows that (5.17) holds if we set  $\Delta_m = \Delta_m^1$  for  $m \in \mathcal{M}_T$  and  $\Delta_m = \Delta_m^0$  for  $m \in \mathcal{M} \setminus \mathcal{M}_T$  and we then derive from (5.18) that not only (5.19) still holds but also

$$\mathbb{E}_s [h^2(\tilde{s}, s)] \leq C \inf_{m \in \mathcal{M}_T} \left\{ n^{-1}|m| + \inf_{t \in S_m} h^2(s, t) \right\},$$

which is indeed a substantial improvement over (5.19). In particular, since  $\mathcal{M}_T$  contains  $\mathcal{M}_R$ , (5.20) still holds when  $\sqrt{s}$  is Hölderian, but the introduction of the



much larger class  $\mathcal{M}_T$  leads to a much more powerful result which follows from the approximation properties of functions in  $V_m$  given by (1.2) with  $m \in \mathcal{M}_T$ . We refer the reader to the book by DeVore and Lorentz (1993) for the precise definitions of Besov spaces and semi-norms and the variation  $\text{Var}^*$  in the following theorem.

**Theorem 6** *Let  $\mathcal{M}_T$  be the set of partitions  $m$  of  $[0, 1]$  previously defined. For any  $p > 0$ ,  $\alpha$  with  $1 > \alpha > (1/p - 1/2)_+$ , any positive integer  $j$  and any function  $t$  belonging to the Besov space  $B_{p,\infty}^\alpha([0, 1])$  with Besov semi-norm  $|t|_{B_{p,\infty}^\alpha}$ , one can find some  $m \in \mathcal{M}_T$  with  $|m| = j$  and some  $t' \in V_m$  such that*

$$\|t - t'\|_2 \leq C(\alpha, p) |t|_{B_{p,\infty}^\alpha} j^{-\alpha}, \quad (5.21)$$

where  $\|\cdot\|_2$  denotes the  $\mathbb{L}_2(dx)$ -norm on  $[0, 1]$ .

If  $t$  is a function of bounded variation on  $[0, 1]$ , there exists  $m \in \mathcal{M}_T$  with  $|m| = j$  and  $t' \in V_m$  such that  $\|t - t'\|_2 \leq C' \text{Var}^*(t) j^{-1}$ .

The bound (5.21) is given in DeVore and Yu (1990). The proof for the bounded variation case has been kindly communicated to the author by Ron DeVore.

Applying the previous theorem to  $t = \sqrt{s}$ , we may always choose for  $t'$  the projection of  $\sqrt{s}$  onto  $V_m$  and it follows from (5.15) that the result still holds with  $t' = \sqrt{s_m}$ . In particular, if  $\sqrt{s} \in B_{p,\infty}^\alpha([0, 1])$ , then for a suitable  $m$  with  $|m| = j$ ,  $h^2(s, s_m) \leq C(\alpha, p) |t|_{B_{p,\infty}^\alpha}^2 j^{-2\alpha}$ . Putting this into (5.18) with  $\Delta_m = 2j$  and optimizing with respect to  $j$  shows that

$$\mathbb{E}_s [h^2(\tilde{s}, s)] \leq C \max \left\{ \left( |t|_{B_{p,\infty}^\alpha} n^{-\alpha} \right)^{2/(2\alpha+1)} ; n^{-1} \right\} \quad \text{if } \sqrt{s} \in B_{p,\infty}^\alpha([0, 1]).$$

Similarly, we can show that

$$\mathbb{E}_s [h^2(\tilde{s}, s)] \leq C \max \left\{ (\text{Var}^*(\sqrt{s})/n)^{2/3} ; n^{-1} \right\} \quad \text{if } \sqrt{s} \text{ has a bounded variation.}$$

## 5.5 Model choice and Approximation Theory

In any statistical framework for which we can prove a risk bound of the form (5.6) provided that (5.3) holds, the technical problem of model selection can be considered as being solved but the question of how to choose the family of models to which we shall apply the procedure remains. There is no general recipe to make such a choice without any ‘‘a priori’’ information on  $s$ . If we have some information about the true  $s$  or at least we suspect that it may have some specific properties, or if we wish that some particular  $s$  should be accurately estimated, we should choose our family of models in such a way that the right-hand side of (5.6) be as small as possible for the  $s$  of interest. Finding models of low dimension with good approximation properties for some specific functions  $s$  is one purpose of Approximation Theory. One should therefore base our choice of suitable families of models on Approximation Theory, which accounts for the numerous connections between modern Statistics and Approximation Theory.

We may also have the choice between several families of models with different approximation properties and complexity levels. Typically, the more complex families have better approximation properties but we have to pay a price for the complexity. A good example is the alternative regular versus irregular partitions for histograms.

As shown in the previous sections, it is possible to mix families with different approximation and complexity properties by playing with the weights  $\Delta_m$ . In particular, it is important that as many models as possible, and particularly those with good approximation properties with respect to functions of greater interest, do satisfy  $\Delta_m \leq c|m|$  for some fixed constant  $c$ . The introduction of the family of models  $\{S_m, m \in \mathcal{M}_T\}$  in Section 5.4.4 illustrates this fact. These models, which have especially good approximation properties with respect to a large class of Besov spaces, form a much richer class than those solely based on regular partitions. Nevertheless, the number of such models with dimension  $D$  remains bounded by  $\exp[c'D]$ , which allows to fix  $\Delta_m$  of the order of  $D$  for these models. By (5.6), this implies that, when we use such a family of models, the performance of the estimator based on model selection is almost (up to constants) as good as the performance of the estimator based on the best individual model.

A detailed analysis of the problems of model choice is given in Section 4.1 of Birgé and Massart (2001) which also provides additional information about the relationship between model selection and Approximation Theory. Further results in this direction are to be found in Barron, Birgé and Massart (1999). It follows from these presentations that all results in Approximation Theory that describe precisely the approximation properties of some particular classes of finite dimensional models are of special interest for the statistical applications we have in mind. Statistics has been using various approximation methods and we would like to emphasize here two main trends. One is based on approximation of functions by piecewise polynomials (or similar functions like splines), some major references here being Birman and Solomjak (1967) and the book by DeVore and Lorentz (1993). The statistical methods based on this approach to approximation lead to estimators which are generalizations of histograms, the selection procedure handling the choice of the partition (and also, possibly, the degree of the polynomials). Another trend is based on the expansion of functions on suitable bases, formerly the trigonometric basis, more recently bases derived from a multiresolution analysis (wavelet bases and the like). The related estimators are based on the estimation of the coefficients in the expansion and the selection chooses the finite set of coefficients to be kept in the expansion of the final estimator. Statistical procedures based on wavelet thresholding are of this type. Theorem 6 based on DeVore and Yu (1990) provides a set of partitions which are relevant for approximation of functions in Besov spaces. A parallel result by Birgé and Massart (2000) applies to the second approach, providing a family of subsets of coefficients to keep in order to get similar approximation properties. A good overview of nonlinear approximation based on wavelets or piecewise polynomials with many useful references is to be found in DeVore (1998).

The use of metric entropy or dimensional arguments in Statistics is not new. The first general results connecting the metric dimension of the parameter set to the performance of estimators are given by Le Cam (1973 and 1975) and statistical applications of the classical entropy results by Kolmogorov and Tikhomirov (1961) are developed in Birgé (1983). An up to date presentation with extensions to model selection following ideas by Barron and Cover (1991) is in Birgé (2006). There is also a huge amount of empirical process literature based on entropy arguments with statistical applications. Many illustrations and references are to be found in van der Vaart and Wellner (1996), van der Vaart (1998), van de Geer (2000) and Massart (2006). More generally, connexions between estimation and Approximation Theory, in particular via wavelet thresholding, have been developed in many papers. Besides

the authors' works already cited, a short selection with further references is as follows: DeVore, Kerkyacharian, Picard and Temlyakov (2004), Donoho and Johnstone (1994, 1995, 1996 and 1998), Donoho, Johnstone, Kerkyacharian and Picard (1995, 1996 and 1997), Kerkyacharian and Picard (1992 and 2000) and Johnstone (1999).

**Acknowledgements** Many thanks to Yannick Baraud, Richard Gill, Piet Groeneboom, Pascal Massart and Aad van der Vaart for their comments on earlier versions of this paper.

## References:

- ASSOUAD, P. (1983). Deux remarques sur l'estimation. *C. R. Acad. Sc. Paris Sér. I Math.* **296**, 1021-1024.
- BARRON, A.R., BIRGÉ, L. and MASSART, P. (1999). Risk bounds for model selection via penalization. *Probab. Theory Relat. Fields* **113**, 301-415.
- BARRON, A.R. and COVER, T.M. (1991). Minimum complexity density estimation. *IEEE Transactions on Information Theory* **37**, 1034-1054.
- BIRGÉ, L. (1983). Approximation dans les espaces métriques et théorie de l'estimation. *Z. Wahrscheinlichkeitstheorie Verw. Geb.* **65**, 181-237.
- BIRGÉ, L. (1984). Stabilité et instabilité du risque minimax pour des variables indépendantes équidistribuées. *Ann. Inst. H. Poincaré Sect. B* **20**, 201-223.
- BIRGÉ, L. (1986). On estimating a density using Hellinger distance and some other strange facts. *Probab. Theory Relat. Fields* **71**, 271-291.
- BIRGÉ, L. (2004). Model selection for Gaussian regression with random design. *Bernoulli* **10**, 1039 -1051.
- BIRGÉ, L. (2006). Model selection via testing : an alternative to (penalized) maximum likelihood estimators. To appear in *Ann. Inst. Henri Poincaré*.
- BIRGÉ, L. and MASSART, P. (1993). Rates of convergence for minimum contrast estimators. *Probab. Th. Rel. Fields* **97**, 113-150.
- BIRGÉ, L. and MASSART, P. (1997). From model selection to adaptive estimation. In *Festschrift for Lucien Le Cam: Research Papers in Probability and Statistics* (D. Pollard, E. Torgersen and G. Yang, eds.), 55-87. Springer-Verlag, New York.
- BIRGÉ, L. and MASSART, P. (1998). Minimum contrast estimators on sieves: exponential bounds and rates of convergence. *Bernoulli* **4**, 329-375.
- BIRGÉ, L. and MASSART, P. (2000). An adaptive compression algorithm in Besov spaces. *Constructive Approximation* **16** 1-36.
- BIRGÉ, L. and MASSART, P. (2001). Gaussian model selection. *J. Eur. Math. Soc.* **3**, 203-268.
- BIRGÉ, L. and ROZENHOLC, Y. (2006). How many bins should be put in a regular histogram. *ESAIM-PS* **10**, 24-45.
- BIRMAN, M.S. and SOLOMJAK, M.Z. (1967). Piecewise-polynomial approximation of functions of the classes  $W_p$ . *Mat. Sbornik* **73**, 295-317.
- BREIMAN, L., FRIEDMAN, J.H., OLSHEN, R.A. and STONE, C.J. (1984). *Classification and Regression Trees*. Wadsworth, Belmont.
- CENCOV, N.N. (1962). Evaluation of an unknown distribution density from observations. *Soviet Math.* **3**, 1559-1562.
- CRAMÉR, H. (1946). *Mathematical Methods of Statistics*. Princeton University Press, Princeton.
- DEVORE, R.A. (1998). Nonlinear Approximation. *Acta Numerica* **7**, 51-150.
- DEVORE, R.A., KERKYACHARIAN, G., PICARD, D. and TEMLYAKOV, V. (2004). Mathematical methods for supervised learning. Tech. report 0422, IMI, University of South Carolina, Columbia.
- DEVORE, R.A. and LORENTZ, G.G. (1993). *Constructive Approximation*. Springer-Verlag, Berlin.
- DEVORE, R.A. and YU, X.M. (1990). Degree of adaptive approximation. *Math. Comp.* **55**, 625-635.

- DEVROYE, L. and GYÖRFI, L. (1985). *Nonparametric Density Estimation: The  $L_1$  View*. John Wiley, New York.
- DONOHU, D.L. and JOHNSTONE, I.M. (1994). Ideal spatial adaptation by wavelet shrinkage. *Biometrika* **81**, 425-455.
- DONOHU, D.L. and JOHNSTONE, I.M. (1995). Adapting to unknown smoothness via wavelet shrinkage. *JASA* **90**, 1200-1224.
- DONOHU, D.L. and JOHNSTONE, I.M. (1996). Neo-classical minimax problems, thresholding and adaptive function estimation. *Bernoulli* **2**, 39-62.
- DONOHU, D.L. and JOHNSTONE, I.M. (1998). Minimax estimation via wavelet shrinkage. *Ann. Statist.* **26**, 879-921.
- DONOHU, D.L., JOHNSTONE, I.M., KERKYACHARIAN, G. and PICARD, D. (1995). Wavelet shrinkage: Asymptopia? *J. R. Statist. Soc. B* **57**, 301-369.
- DONOHU, D.L., JOHNSTONE, I.M., KERKYACHARIAN, G. and PICARD, D. (1996). Density estimation by wavelet thresholding *Ann. Statist.* **24**, 508-539.
- DONOHU, D.L., JOHNSTONE, I.M., KERKYACHARIAN, G. and PICARD, D. (1997). Universal near minimaxity of wavelet shrinkage. In *Festschrift for Lucien Le Cam: Research Papers in Probability and Statistics* (D. Pollard, E. Torgersen and G. Yang, eds.), 183-218. Springer-Verlag, New York.
- FISCHER, R.A. (1922). On the mathematical foundations of theoretical statistics. *Philos. Trans. Royal Soc. London Ser. A* **222**, 309-368.
- FISCHER, R.A. (1925). Theory of statistical estimation. *Proc. Cambridge Philos. Soc.* **22**, 700-725.
- GEY, S. and NÉDÉLEC, E. (2005). Model selection for CART regression trees. *IEEE Transactions on Information Theory* **51**, 658-670.
- GRENANDER, U. (1981). *Abstract inference*. John Wiley, New York.
- GROENEBOOM, P. and WELLNER, J.A. (1992). *Information Bounds and Nonparametric Maximum Likelihood Estimation*. Birkhäuser, Basel.
- GYÖRFI, L., KOHLER, M., KRYŽAK, A. and WALK, H. (2002). *A Distribution-Free Theory of Nonparametric Regression*. Springer, New York.
- HÁJEK, J. (1970). A characterization of limiting distributions of regular estimates. *Z. Wahrsch. Verw. Gebiete* **14**, 323-330.
- HÁJEK, J. (1972). Local asymptotic minimax and admissibility in estimation. *Proc. Sixth Berkeley Symp. Math. Statist. Probab.* **1**, 174-194. Univ. California Press, Berkeley.
- IBRAGIMOV, I.A. and HAS'MINSKII, R.Z. (1981). *Statistical Estimation: Asymptotic Theory*. Springer-Verlag, New York.
- JOHNSTONE, I. (1999). *Function Estimation and Gaussian Sequence Models*. Book in preparation.  
<http://www-stat.stanford.edu/people/faculty/johnstone/baseb.pdf>
- KERKYACHARIAN, G. and PICARD, D. (1992). Density estimation in Besov spaces. *Statist. and Probab. Lett.* **13**, 15-24.
- KERKYACHARIAN, G. and PICARD, D. (2000). Thresholding algorithms, maxisets and well-concentrated bases. *Test* **9**, 283-344.
- KOLMOGOROV, A.N. and TIKHOMIROV, V.M. (1961).  $\varepsilon$ -entropy and  $\varepsilon$ -capacity of sets in function spaces. *Amer. Math. Soc. Transl. (2)* **17**, 277-364.
- Le CAM, L.M. (1953). On some asymptotic properties of maximum likelihood estimates and related Bayes' estimates. *Univ. California Publ. Statist.* **1**, 277-329.
- Le CAM, L.M. (1970). On the assumptions used to prove asymptotic normality of maximum likelihood estimates. *Ann. Math. Stat.* **41**, 802-828.
- Le CAM, L.M. (1973). Convergence of estimates under dimensionality restrictions. *Ann. Statist.* **1**, 38-53.
- Le CAM, L.M. (1975). On local and global properties in the theory of asymptotic normality of experiments. *Stochastic Processes and Related Topics, Vol. 1* (M. Puri, ed.), 13-54. Academic Press, New York.
- Le CAM, L.M. and YANG, G.L. (2000). *Asymptotics in Statistics: Some Basic Concepts. Second Edition*. Springer-Verlag, New York.

- MASSART, P. (2006). Concentration Inequalities and Model Selection. In *Lecture on Probability Theory and Statistics, Ecole d'Eté de Probabilités de Saint-Flour XXXIII - 2003* (J. Picard, ed.). Lecture Note in Mathematics, Springer-Verlag, Berlin.
- RIGOLLET, T. and TSYBAKOV, A.B. (2005). Linear and convex aggregation of density estimators. Technical report, University Paris VI.
- SHEN, X. and WONG, W.H. (1994). Convergence rates of sieve estimates. *Ann. Statist.* **22**, 580-615.
- SILVERMAN, B.W. (1982). On the estimation of a probability density function by the maximum penalized likelihood method. *Ann. Statist.* **10**, 795-810.
- van de GEER, S. (1990). Estimating a regression function. *Ann. Statist.* **18**, 907-924.
- van de GEER, S. (1993). Hellinger-consistency of certain nonparametric maximum likelihood estimates. *Ann. Statist.* **21**, 14-44.
- van de GEER, S. (1995). The method of sieves and minimum contrast estimators. *Math. Methods Statist.* **4**, 20-38.
- van de GEER, S. (2000). *Empirical Processes in M-Estimation*. Cambridge University Press, Cambridge.
- van der VAART, A.W. (1998). *Asymptotic Statistics*. Cambridge University Press, Cambridge.
- van der VAART, A.W. (2002). The statistical work of Lucien Le Cam. *Ann. Statist.* **30**, 631-682.
- van der VAART, A.W. and WELLNER, J.A. (1996). *Weak Convergence and Empirical Processes, With Applications to Statistics*. Springer-Verlag, New York.
- WAHBA, G. (1990). *Spline Models for Observational Data*. S.I.A.M., Philadelphia.
- WALD, A. (1949). Note on the consistency of the maximum likelihood estimate. *Ann. Math. Statist.* **20**, 595-601.
- WONG, W.H. and SHEN, X. (1995). Probability inequalities for likelihood ratios and convergence rates of sieve MLEs. *Ann. Statist.* **23**, 339-362.

Lucien BIRGÉ  
 UMR 7599 "Probabilités et modèles aléatoires"  
 Laboratoire de Probabilités, boîte 188  
 Université Paris VI, 4 Place Jussieu  
 F-75252 Paris Cedex 05  
 France  
 e-mail: LB@CCR.JUSSIEU.FR