



The Ising-Sherrington-Kirpatrick model in a magnetic field at high temperature

Francis Comets, Francesco Guerra, Fabio Lucio Toninelli

► To cite this version:

Francis Comets, Francesco Guerra, Fabio Lucio Toninelli. The Ising-Sherrington-Kirpatrick model in a magnetic field at high temperature. *Journal of Statistical Physics*, Springer Verlag, 2005, 120, pp.147-165. <10.1007/s10955-005-5471-1>. <hal-00003866>

HAL Id: hal-00003866

<https://hal.archives-ouvertes.fr/hal-00003866>

Submitted on 11 Jan 2005

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Ising-Sherrington-Kirpatrick model in a magnetic field at high temperature

Francis Comets* Francesco Guerra[†]
 Fabio Lucio Toninelli[‡]

December 2004

Abstract

We study a spin system on a large box with both Ising interaction and Sherrington-Kirpatrick couplings, in the presence of an external field. Our results are: (i) existence of the pressure in the limit of an infinite box. When both Ising and Sherrington-Kirpatrick temperatures are high enough, we prove that: (ii) the value of the pressure is given by a suitable replica symmetric solution, and (iii) the fluctuations of the pressure are of order of the inverse of the square of the volume with a normal distribution in the limit. In this regime, the pressure can be expressed in terms of random field Ising models.

Key Words: Ising model, Sherrington-Kirpatrick model, spin-glass, thermodynamic limit, pressure, quadratic coupling

AMS (1991) subject classifications: 82B44; 60K35, 82D30

Short title: Ising-Sherrington-Kirpatrick model

1 Introduction

We consider a d -dimensional Ising model in a magnetic field h , perturbed by a mean field interaction of spin-glass type. The Hamiltonian contains two

* University Paris 7 - Denis Diderot. Partially supported by CNRS, UMR 7599 “Probabilités et Modèles Aléatoires.”

[†]Università di Roma “La Sapienza” and INFN, Sezione di Roma. Partially supported by MIUR and INFN

[‡]Laboratoire de Physique, Ecole Normale Supérieure de Lyon. On leave at Institut für Mathematik, Universität Zürich

parameters, β and κ , which play the role of two inverse temperatures. When $\beta = 0$ the model reduces to the Ising model at temperature $1/\kappa$, while for $\kappa = 0$ one recovers the Sherrington-Kirkpatrick (SK) model at temperature $1/\beta$. The understanding of the SK model has recently witnessed great progress (see, e.g., [10] [8] [14]). The main interest in the analysis of this model is the possibility of investigating the robustness of the phenomena typical of mean field spin glass models, in the presence of additional interactions of non-mean-field character.

The model has been previously considered in [3] under the additional assumptions that $h = 0$ and that the Ising model is ferromagnetic. Under these conditions it was proven that, if β and κ are small enough, the infinite volume pressure is given by the sum of the Ising pressure and of the SK one, at the respective temperatures. Moreover, the disorder fluctuations of the pressure were found to be of order $1/V$, V being the volume of the system, and to satisfy a central limit theorem.

In the present paper, the Ising interaction decays exponentially fast with distance, but is not necessarily ferromagnetic. It turns out that the presence of the magnetic field changes qualitatively the picture with respect to [3]. Indeed we find, still for κ and β small enough, that the limit pressure is given in terms of the pressure of an Ising model with random external field, the strength of the randomness being related to the typical value of the overlap between two replicas of the system. It is remarkable, though natural, that the random field Ising model, which has its own interest [2], plays such an important role in our model. Note also that the pressure of our model can be computed only via the thermodynamic limit of another disordered system. This is contrast with the case $h = 0$ we mentioned above, and also of course with the case $\kappa = 0$ of the standard SK model. The second difference is that the fluctuations of the pressure in presence of h satisfy a central limit theorem on the scale $1/\sqrt{V}$ rather than $1/V$. The same phenomenon is known to happen in the case of the usual SK model, see for instance [1] [4] [11].

In the region of thermodynamic parameters we consider, the system is in a “replica-symmetric” (RS) phase, the overlap between two independent replicas being a non-random value in the thermodynamic limit. Our methods fail beyond some values $\beta_0(h)$ and $\kappa_0(h)$, which we believe to be an artifact of our approach, rather than representing the true boundary of the RS region. The same inconvenient has been previously encountered in the analysis of the SK model [15], [9]. In principle one could improve these values by employing the replica symmetry breaking scheme which for instance enabled M. Talagrand [16] to control the whole replica-symmetric region of the SK model, and later the entire phase space [14]. However, in the present case (as

well as in [3]) one of the reasons why we do not reach the true critical line is due to an incomplete control of the underlying random field Ising model, and this problem would not be fixed by the methods of [16]. For this reason, we prefer to use a generalization of the technically simpler “quadratic replica coupling” technique introduced in [9].

It would be of course an interesting challenge to go beyond the present approach, and to deal with lower-temperature situations where the Ising-SK system possibly shows a RSB-like behavior.

2 Description of the model and results

The model we consider is defined on the d -dimensional hypercubic box $\Lambda_N = \{-N, \dots, N\}^d$ and its partition function is:

$$Z_N(\kappa, \beta, h; J) = \sum_{\sigma \in \{-1, +1\}^{\Lambda_N}} \exp \left(-\kappa H_N^I(\sigma) - \beta H_N^{SK}(\sigma; J) + h \sum_{i \in \Lambda_N} \sigma_i \right). \quad (1)$$

The Hamiltonian of the Sherrington-Kirkpatrick (SK) model is defined as

$$H_N^{SK}(\sigma; J) = -\frac{1}{\sqrt{2|\Lambda_N|}} \sum_{i, j \in \Lambda_N} J_{ij} \sigma_i \sigma_j$$

and the couplings J_{ij} are i.i.d. Gaussian random variables $\mathcal{N}(0, 1)$. On the other hand, the Hamiltonian of the Ising model is

$$H_N^I(\sigma) = -\frac{1}{2} \sum_{i, j \in \Lambda_N} K(i - j) \sigma_i \sigma_j,$$

where we assume that the interaction decays exponentially, i.e.,

$$|K(i)| \leq C_1 e^{-C_2|i|} \quad (2)$$

for some $C_1, C_2 > 0$. We do not require the interaction to be ferromagnetic. The finite volume (disorder-dependent) pressure is defined as usual as

$$p_N(\kappa, \beta, h; J) = \frac{1}{|\Lambda_N|} \log Z_N(\kappa, \beta, h; J).$$

Later, we will need to consider the (Gaussian) Random Field Ising Model, defined by the partition function

$$Z_N^{\text{RFIM}}(\kappa, h, \gamma; J) = \sum_{\sigma \in \{-1, +1\}^{\Lambda_N}} \exp \left(-\kappa H_N^I(\sigma) + \sum_{i \in \Lambda_N} \sigma_i (h + \gamma J_i) \right),$$

J_i being i.i.d. standard Gaussian variables, and being also independent from the J_{ij} 's in the sequel. The existence of the infinite volume pressure of the RFIM,

$$\begin{aligned} p^{\text{RFIM}}(\kappa, h, \gamma) &= \lim_{N \rightarrow \infty} \mathbb{E} p_N^{\text{RFIM}}(\kappa, h, \gamma; J) \\ &= a.s. - \lim_{N \rightarrow \infty} p_N^{\text{RFIM}}(\kappa, h, \gamma; J), \end{aligned}$$

is a well known consequence of additivity and of the ergodic theorem, e.g. [17].

Our main results can be summarized as follows:

Theorem 1. *For all $h \in \mathbb{R}$, and all $\kappa, \beta > 0$, the limit*

$$p(\kappa, \beta, h) = \lim_{N \rightarrow \infty} \mathbb{E} p_N(\kappa, \beta, h; J) \quad (3)$$

exists and

$$p_N(\kappa, \beta, h; J) \xrightarrow{N \rightarrow \infty} p(\kappa, \beta, h) \quad (4)$$

for a.e. J and in the L^p -norm ($p \in [1, \infty)$).

Theorem 2. *For all $h \in \mathbb{R}$, there exist $\kappa_0(h) > 0$ and $\beta_0(h) > 0$ such that, for $0 \leq \kappa \leq \kappa_0(h)$ and $0 \leq \beta \leq \beta_0(h)$*

$$p(\kappa, \beta, h) = \inf_{0 \leq q \leq 1} \left(p^{\text{RFIM}}(\kappa, h, \beta\sqrt{q}) + \frac{\beta^2}{4}(1-q)^2 \right) \quad (5)$$

In the case $\kappa = 0$, it is established in [7] that the infimum in (5) is achieved at a unique point. As explained below (see Section 3.2), we take small enough $\kappa_0(h)$ so that, for $0 \leq \kappa \leq \kappa_0(h)$ and $0 \leq \beta \leq \beta_0(h)$, the infimum in (5) is achieved at a unique q . In the following, we will always denote by $\bar{q} = \bar{q}(\kappa, \beta, h)$ the value that realizes the infimum in (5).

We emphasize that $\kappa_0(h)$ is taken small enough so that the RFIM is inside Dobrushin's uniqueness region for every realization of the external fields $J_i, i \in \mathbb{Z}^d$. Let $\langle \cdot \rangle_{\infty, J}$ be the unique infinite volume Gibbs measure for the RFIM with $\gamma = \beta\sqrt{\bar{q}}$, depending on $J_i, i \in \mathbb{Z}^d$. If \preceq denotes the lexicographic order in \mathbb{Z}^d , define

$$\Gamma = \Gamma(\kappa, \beta, h) = \mathbb{E}_{J_i, i \geq 0} \left(\mathbb{E}_{J'_0, J_i, i < 0} \log \left\langle e^{\beta\sqrt{\bar{q}}(J'_0 - J_0)\sigma_0} \right\rangle_{\infty, J} \right)^2,$$

where J'_0 is an independent copy of J_0 , J'_0 being independent of $(J_i, i \in \mathbb{Z}^d)$. Here, the \mathbb{E} -expectations are conditional, and the subscripts of \mathbb{E} indicate on which variables the expectation is performed. Then, we have the following central limit theorem:

Theorem 3. For $0 \leq \kappa \leq \kappa_0(h)$ and $0 \leq \beta \leq \beta_0(h)$,

$$\sqrt{|\Lambda_N|} \left(p_N(\kappa, \beta, h; J) - \mathbb{E} p_N(\kappa, \beta, h; J) \right) \xrightarrow{\text{law}} \mathcal{N} \left(0, \Gamma - \frac{\beta^2}{2} \bar{q}^2 \right). \quad (6)$$

One can check by expansion around $\kappa = \beta = 0, h \neq 0$, that the limiting variance $\Gamma - \frac{\beta^2}{2} \bar{q}^2$ is strictly positive for any fixed non-zero h and small κ, β . This proves that the fluctuations of $p_N(\kappa, \beta, h; J)$ are truly of order of the square root of the volume inverse in this region of the parameters. To match with the breakdown in the order of magnitude of fluctuations at zero external field, we observe that both \bar{q} and Γ vanish as $h \rightarrow 0$, and also that they are equal to zero when $h = 0$.

3 Proofs

3.1 For small κ the RFIM is inside the Dobrushin uniqueness region

Let $i \neq k$ be lattice points. Under any (infinite volume) RFIM Gibbs measure $\langle \cdot \rangle_{\infty, J}$ the law $\langle \cdot | \eta \rangle_{i, \infty, J}$ of σ_i given $\sigma_j = \eta_j, j \neq i$, is Bernoulli with parameter proportional to $\exp\{\sigma_i [\mathcal{H}_{i, k; \eta} + \kappa K(i - k) \eta_k]\}$ with

$$\mathcal{H}_{i, k; \sigma} = \kappa \sum_{j \neq i, k} K(i - j) \sigma_j + h + \gamma J_i.$$

Following e.g. Section 2 in [5], we define the Dobrushin's influence coefficient

$$C_{ki} = \sup \left\{ \frac{1}{2} \left\| \langle \cdot | \eta \rangle_{i, \infty, J} - \langle \cdot | \eta' \rangle_{i, \infty, J} \right\|_{\text{var}}; \eta = \eta' \text{ off } k \right\}$$

where $\| \cdot \|_{\text{var}}$ is the variation norm. With a straightforward computation,

$$\begin{aligned} \left\| \langle \cdot | \eta \rangle_{i, \infty, J} - \langle \cdot | \eta' \rangle_{i, \infty, J} \right\|_{\text{var}} &= 2 \left| \langle + | \eta \rangle_{i, \infty, J} - \langle + | \eta' \rangle_{i, \infty, J} \right| \\ &\leq \frac{2 \left| \sinh 2\kappa K(i - k) \right|}{\cosh[2\mathcal{H}_{i, k; \eta}] + \cosh[2\kappa K(i - k)]} \end{aligned}$$

for such η, η' , and so

$$C_{ki} \leq \left| \tanh 2\kappa K(i - k) \right| \leq \left| 2\kappa K(i - k) \right| \quad (7)$$

Therefore, for $\kappa < \kappa_1 = (2 \sum_{i \neq 0} |K(i)|)^{-1}$ we derive from (7) and (2) that $a = \sup_i \sum_k \rho^{|i-k|} C_{ki} < 1$ for some $\rho > 1$, which implies (see Section 2.3 in

[5]) that the Gibbs measure $\langle \cdot \rangle_{\infty, J}$ is unique and has exponentially decreasing correlations. More precisely, for all local functions f, g , there is a finite constant $C = C(a, f, g)$ such that for all $i \in \mathbb{Z}^d$

$$|\langle f; g \circ \theta_i \rangle_{\infty, J}| \leq C\rho^{-|i|}, \quad (8)$$

with θ_i the shift of vector i and $\langle f; g \rangle_{\infty, J}$ the covariance of f, g .

3.2 Uniqueness of \bar{q}

Introduce

$$F(q, \kappa) = F_{\beta, h}(q, \kappa) = p^{\text{RFIM}}(\kappa, h, \beta\sqrt{q}) + \frac{\beta^2}{4}(1 - q)^2$$

which is, in view of (8), a smooth function of all its arguments if $\kappa < \kappa_1$ (e.g., Cor. 8.37 in [6]).

Proposition 1. *Assume $h \neq 0$. For all β , there exists $\kappa_2 = \kappa_2(\beta, h) > 0$ such that $q \mapsto F(q, \kappa)$ has a unique minimizer on $[0, 1]$ for $\kappa < \kappa_2$.*

By [7], we know that the minimizer q_0 of $F(q, 0)$ is unique, strictly positive and that¹

$$\frac{\partial^2}{\partial^2 q} F(q_0, 0) > 0. \quad (9)$$

By (8), the function F is continuous in κ uniformly in $q \in (0, 1]$. Hence,

$$\forall \delta > 0 \quad \exists \tilde{\kappa} : \forall \kappa < \tilde{\kappa} \quad \operatorname{argmin}_{[0, 1]} F(\cdot, \kappa) \subset (q_0 - \delta, q_0 + \delta) \quad (10)$$

Again by (8), the function F is C^2 in a neighborhood of $(q_0, 0)$. By the implicit function theorem for the equation

$$\frac{\partial}{\partial q} F(q, \kappa) = 0$$

(which applies thanks to condition (9)) we then derive that there exist neighborhoods U of q_0 and V of $\kappa = 0$ and a function $\bar{q} : V \mapsto U$ such that, for $q \in U, \kappa \in V$, the above equation is equivalent to $q = \bar{q}(\kappa)$. With δ small enough so that $(q_0 - \delta, q_0 + \delta) \subset U$, we choose now $\kappa_2 < \tilde{\kappa}$ (with $\tilde{\kappa}$ from (10)) such that $(-\kappa_2, \kappa_2) \subset V$. Then, for $\kappa < \kappa_2$, $F(\cdot, \kappa)$ has a unique minimum at $q = \bar{q}(\kappa)$.

As for the case $h = 0$, one can prove similarly the following results, that we mention for comparison but will neither prove nor use.

¹[7], p.166

Proposition 2. *For all $\beta \neq 1$, there exists $\kappa_3 = \kappa_3(\beta) > 0$ such that $q \mapsto F_{\beta,0}(q, \kappa)$ has a unique minimizer on $[0, 1]$ for $\kappa < \kappa_3$. If $\beta < 1$, the minimizer is $q = 0$.*

Remark The restriction $\beta \neq 1$ is due to the fact that for $\beta = \beta_c = 1$ (the critical point of the Sherrington-Kirkpatrick model) one has $q_0(\beta_c, 0) = 0$ and, in contrast with (9),

$$\frac{\partial^2}{\partial^2 q} F_{\beta_c,0}(0, 0) = 0.$$

3.3 Proof of Theorem 1

Proof of (3) The proof is standard, we just sketch the main steps. Consider the box Λ_{mN} , with $m, N \in \mathbb{N}$, and partition it into sub-boxes $\Lambda_N^{(\ell)}$, $\ell = 1, \dots, m^d$, congruent to Λ_N . Moreover, let \tilde{Z}_{mN} be the partition function of the system where, with respect to (1), $H_{mN}^{SK}(\sigma; J)$ is replaced by

$$- \sum_{\ell=1}^{m^d} \frac{1}{\sqrt{2|\Lambda_N|}} \sum_{i,j \in \Lambda_N^{(\ell)}} J_{ij}^{(\ell)} \sigma_i \sigma_j,$$

the $J_{ij}^{(\ell)}$ being m^d independent families of standard Gaussian variables. Note that, in this system, the different sub-boxes interact only through the Ising potential $K(\cdot)$. Then, following the ideas of [10], it is easy to prove that

$$p_{mN}(\kappa, \beta, h) \geq \frac{1}{|\Lambda_{mN}|} \mathbb{E} \log \tilde{Z}_{mN}(\kappa, \beta, h; J^{(\ell)}).$$

Since the Ising potential is summable, the interaction among the different sub-boxes due to the potential K grows at most proportionally to κ and to the total surface, $d m^d N^{d-1}$. As a consequence, one has the approximate monotonicity

$$p_{mN}(\kappa, \beta, h) \geq p_N(\kappa, \beta, h) - \kappa \frac{C}{N},$$

for some constant C depending on the potential $K(\cdot)$. From this, it is a standard fact to deduce that the sequence $\{p_{mN}\}_N$ has a limit when $N \rightarrow \infty$, that it does not depend on m , and that it coincides with $\lim_N p_N$.

Proof of (4) The almost sure convergence is standard and follows from exponential self-averaging of the pressure (see for instance Proposition 2.18 of [12]), which in the present case reads

$$\mathbb{P} (|p_N(\kappa, \beta, h; J) - \mathbb{E} p_N(\kappa, \beta, h; J)| \geq u) \leq D_1 e^{-D_2(\beta)|\Lambda_N|u^2},$$

together with Borel-Cantelli's lemma. The L^p -convergence comes from uniform integrability, which again follows from exponential concentration.

3.4 Proof of Theorem 2

For $0 \leq t \leq 1$ and any $q \geq 0$, define the interpolating partition function

$$Z(t) = \sum_{\sigma \in \{-1, +1\}^{\Lambda_N}} \exp\left(-H^{(t)}(\sigma)\right) \quad (11)$$

$$H^{(t)}(\sigma) = \kappa H_N^I(\sigma) + \beta\sqrt{t}H_N^{SK}(\sigma; J) - \sum_{i \in \Lambda_N} \sigma_i \left(h + \beta\sqrt{q(1-t)}J_i\right)$$

with the properties

$$\begin{aligned} Z(0) &= Z_N^{\text{RFIM}}(\kappa, h, \beta\sqrt{q}; J) \\ Z(1) &= Z_N(\kappa, \beta, h; J). \end{aligned}$$

The t -derivative of the corresponding pressure

$$p_N(t) = \frac{1}{|\Lambda_N|} \mathbb{E} \log Z(t)$$

is easily computed: We denote by $\langle \cdot \rangle_t$ the Gibbs measure associated to $H^{(t)}$, by $\langle \cdot \rangle_t^{\otimes 2}$ its tensor product acting on a pair $(\sigma^1, \sigma^2) \in \{-1, +1\}^{\Lambda_N} \times \{-1, +1\}^{\Lambda_N}$, by $q_{12} = |\Lambda_N|^{-1} \sum_{i \in \Lambda_N} \sigma_i^1 \sigma_i^2$ the overlap between configurations σ^1 and σ^2 , and we get by the Gaussian integration by parts formula $\mathbb{E} J_i F(J_i) = \mathbb{E} F'(J_i)$,

$$\begin{aligned} \frac{d}{dt} p_N(t) &= \frac{\beta}{2|\Lambda_N|} \mathbb{E} \left\{ \frac{1}{\sqrt{2|\Lambda_N|t}} \sum_{i, j \in \Lambda_N} J_{ij} \langle \sigma_i \sigma_j \rangle_t - \frac{\sqrt{q}}{\sqrt{1-t}} \sum_{i \in \Lambda_N} J_i \langle \sigma_i \rangle_t \right\} \\ &\stackrel{\text{int. by parts}}{=} \frac{\beta}{2|\Lambda_N|} \mathbb{E} \left\{ \frac{\beta}{2|\Lambda_N|} \sum_{i, j \in \Lambda_N} (1 - \langle \sigma_i^1 \sigma_j^1 \sigma_i^2 \sigma_j^2 \rangle_t^{\otimes 2}) - \beta q \sum_{i \in \Lambda_N} (1 - \langle \sigma_i^1 \sigma_i^2 \rangle_t^{\otimes 2}) \right\} \\ &= \frac{\beta^2}{4} (1 - q)^2 - \frac{\beta^2}{4} \mathbb{E} \langle (q_{12} - q)^2 \rangle_t^{\otimes 2} \quad (12) \end{aligned}$$

so that, integrating in t between 0 and 1, taking the $N \rightarrow \infty$ limit and optimizing on q , we have the “first half” of Eq. (5):

$$p(\kappa, \beta, h) \leq \inf_{q \geq 0} \left(p^{\text{RFIM}}(\kappa, h, \beta\sqrt{q}) + \frac{\beta^2}{4} (1 - q)^2 \right). \quad (13)$$

Note that the inequality holds for any values of β and κ .

Next, consider a system of two coupled replicas: For $\lambda > 0$ let

$$Z^{(2)}(t, \lambda) = \sum_{\sigma^1, \sigma^2 \in \{-1, +1\}^{\Lambda_N}} \exp\left(-H^{(t)}(\sigma^1) - H^{(t)}(\sigma^2) + \frac{\beta^2}{2} |\Lambda_N| \lambda (q_{12} - q)^2\right)$$

and denote by $\langle \cdot \rangle_{t,\lambda}$ the Gibbs measure on $\{-1, +1\}^{\Lambda_N} \times \{-1, +1\}^{\Lambda_N}$ associated to this partition function. (Of course, $\langle \cdot \rangle_{t,0} = \langle \cdot \rangle_t^{\otimes 2}$ and $Z^{(2)}(t, 0) = Z(t)^2$.) With

$$p_N^{(2)}(t, \lambda) = \frac{1}{2|\Lambda_N|} \mathbb{E} \log Z^{(2)}(t, \lambda)$$

we have by a computation similar to (12) and using symmetry in σ^1, σ^2 ,

$$\begin{aligned} & \frac{d}{dt} p_N^{(2)}(t, \lambda_0 - t) = \\ &= \frac{\beta}{4|\Lambda_N|} \mathbb{E} \left\{ \frac{2}{\sqrt{2|\Lambda_N|t}} \sum_{i,j \in \Lambda_N} J_{ij} \langle \sigma_i^1 \sigma_j^1 \rangle_{t,\lambda_0-t} - \frac{2\sqrt{q}}{\sqrt{1-t}} \sum_{i \in \Lambda_N} J_i \langle \sigma_i^1 \rangle_{t,\lambda_0-t} \right. \\ & \quad \left. - \beta |\Lambda_N| \langle (q_{12} - q)^2 \rangle_{t,\lambda_0-t} \right\} \\ & \stackrel{\text{int. by parts}}{=} \frac{\beta}{4|\Lambda_N|} \mathbb{E} \left\{ \frac{\beta}{|\Lambda_N|} \sum_{i,j \in \Lambda_N} (1 + \langle \sigma_i^1 \sigma_j^1 \sigma_i^2 \sigma_j^2 \rangle_{t,\lambda_0-t} - 2 \langle \sigma_i^1 \sigma_j^1 \sigma_i^3 \sigma_j^3 \rangle_{t,\lambda_0-t}^{\otimes 2}) \right. \\ & \quad \left. - 2\beta q \sum_{i \in \Lambda_N} (1 + \langle \sigma_i^1 \sigma_i^2 \rangle_{t,\lambda_0-t} - 2 \langle \sigma_i^1 \sigma_i^3 \rangle_{t,\lambda_0-t}^{\otimes 2}) - \beta |\Lambda_N| \langle (q_{12} - q)^2 \rangle_{t,\lambda_0-t} \right\} \\ &= \frac{\beta^2}{4} (1 - q)^2 - \frac{\beta^2}{2} \mathbb{E} \langle (q_{13} - q)^2 \rangle_{t,\lambda_0-t}^{\otimes 2} \end{aligned}$$

so that, integrating,

$$p_N^{(2)}(t, \lambda) \leq \frac{\beta^2}{4} (1 - q)^2 t + \frac{1}{2|\Lambda_N|} \mathbb{E} \log Z^{(2)}(0, t + \lambda). \quad (14)$$

Now, setting $u_N(t) = p_N^{\text{RFIM}}(\kappa, h, \beta\sqrt{q}) + \frac{\beta^2}{4} (1 - q)^2 t - p_N(t)$, which is a non negative function by (12), and using convexity of the pressure with respect to λ and the identity $p_N^{(2)}(t, 0) = p_N(t)$, we obtain for any $\lambda > 0$

$$\begin{aligned} \frac{d}{dt} u_N(t) & \stackrel{(12)}{=} \frac{\partial}{\partial \lambda} p_N^{(2)}(t, 0) \\ & \stackrel{\text{convexity}}{\leq} \frac{p_N^{(2)}(t, \lambda) - p_N(t)}{\lambda} \\ & \stackrel{(14)}{\leq} \frac{1}{\lambda} \left[u_N(t) + \frac{1}{2|\Lambda_N|} \mathbb{E} \log \frac{Z^{(2)}(0, t + \lambda)}{Z^{(2)}(0, 0)} \right]. \end{aligned} \quad (15)$$

Since $Z^{(2)}$ is increasing in λ , Eq. (15) implies

$$\frac{d}{dt} \log \left[u_N(t) + \frac{1}{2|\Lambda_N|} \mathbb{E} \log \frac{Z^{(2)}(0, 1 + \lambda)}{Z^{(2)}(0, 0)} \right] \leq \frac{1}{\lambda},$$

which, recalling that $u_N(0) = 0$, can be immediately integrated to give

$$u_N(t) \leq (e^{t/\lambda} - 1) \times \frac{1}{2|\Lambda_N|} \mathbb{E} \log \frac{Z^{(2)}(0, 1 + \lambda)}{Z^{(2)}(0, 0)} \quad (16)$$

and Eq. (5) follows if we can prove that

$$\lim_{N \rightarrow \infty} \frac{1}{2|\Lambda_N|} \mathbb{E} \log \frac{Z^{(2)}(0, \lambda_0)}{Z^{(2)}(0, 0)} = 0 \quad (17)$$

for some $\lambda_0 > 1$, if q is chosen properly. (Note that $\lambda_0 = \lambda + 1 > 1$ is required so that one can take t up to 1 and still have $\lambda > 0$, which is needed in (15).)

Define for any $\mu \in \mathbb{R}$, $q \geq 0$

$$\alpha_N(\mu; J) = \frac{1}{2|\Lambda_N|} \log \langle e^{\mu|\Lambda_N|(q_{12}-q)} \rangle_{\kappa, \beta\sqrt{q}, N}^{\otimes 2}$$

and $\alpha_N(\mu) = \mathbb{E} \alpha_N(\mu; J)$. We denote by $\langle \cdot \rangle_{\kappa, \beta\sqrt{q}, N}^{\otimes 2}$ the Gibbs measure for two replicas of the RFIM with parameters κ and $\gamma = \beta\sqrt{q}$ and volume Λ_N . Later we will also use the notation $\langle \cdot \rangle_{\kappa, \beta\sqrt{q}, N}^{(\mu)}$ for

$$\langle A \rangle_{\kappa, \beta\sqrt{q}, N}^{(\mu)} = \frac{\langle A e^{\mu|\Lambda_N|(q_{12}-q)} \rangle_{\kappa, \beta\sqrt{q}, N}^{\otimes 2}}{\langle e^{\mu|\Lambda_N|(q_{12}-q)} \rangle_{\kappa, \beta\sqrt{q}, N}^{\otimes 2}}.$$

Let $\bar{q}_N = \bar{q}_N(\beta, \kappa, h)$ be the value which minimizes

$$p_N^{\text{RFIM}}(\kappa, h, \beta\sqrt{q}) + \frac{\beta^2}{4}(1-q)^2$$

with respect to q , cf (5). Clearly, \bar{q}_N satisfies the “self-consistent equation”

$$q = \mathbb{E} \langle q_{12} \rangle_{\kappa, \beta\sqrt{q}, N}^{\otimes 2} = \frac{\sum_{i \in \Lambda_N} \mathbb{E} \langle \sigma_i \rangle_{\kappa, \beta\sqrt{q}, N}^2}{|\Lambda_N|}. \quad (18)$$

An analysis analogous to the one of Section 3.2 shows that the solution of (18) is unique for κ small enough, for N sufficiently large, and that $\bar{q}_N \rightarrow \bar{q}$ for $N \rightarrow \infty$. A Taylor expansion around $\mu = 0$ gives immediately

$$\alpha_N(\mu; J) = \alpha_N(0; J) + \mu \alpha'_N(0; J) + \int_0^\mu dy \int_0^y du \alpha''_N(u; J),$$

where

$$\alpha_N(0; J) = 0 \quad (19)$$

$$\alpha'_N(0; J) = \frac{1}{2} \left(\langle q_{12} \rangle_{\kappa, \beta \sqrt{q_N}, N}^{\otimes 2} - \mathbb{E} \langle q_{12} \rangle_{\kappa, \beta \sqrt{q_N}, N}^{\otimes 2} \right) \quad (20)$$

$$\alpha''_N(u; J) = \frac{|\Lambda_N|}{2} \left(\langle q_{12}^2 \rangle_{\kappa, \beta \sqrt{q_N}, N}^{(u)} - (\langle q_{12} \rangle_{\kappa, \beta \sqrt{q_N}, N}^{(u)})^2 \right) \quad (21)$$

$$= \frac{1}{2|\Lambda_N|} \sum_{i, j \in \Lambda_N} \langle (\sigma_i^1 \sigma_i^2 - \langle \sigma_i^1 \sigma_i^2 \rangle) (\sigma_j^1 \sigma_j^2 - \langle \sigma_j^1 \sigma_j^2 \rangle) \rangle_{\kappa, \beta \sqrt{q_N}, N}^{(u)} \quad (22)$$

We can view $\langle \cdot \rangle_{\kappa, \beta \sqrt{q_N}, N}^{(u)}$ as the Gibbs measure of a system with $2|\Lambda_N|$ spins, with exponentially decaying pair interactions. Since κ is small, taking μ itself sufficiently small, we keep this system inside the Dobrushin uniqueness region. Using exponential decay of correlations as in Section 3.1, we obtain that $\alpha''_N(u; J)$ is bounded above by a constant, uniformly in N, J and in $u \in [0, \mu]$, so that

$$\alpha_N(\mu; J) \leq \mu \alpha'_N(0; J) + C\mu^2 \quad (23)$$

for $\kappa < \kappa_0(h)$. Note that this bound holds for any μ , since it does for small μ and the function α can grow at most linearly at infinity.

With this in hand, we go back to proving (17). After a Gaussian transformation

$$\begin{aligned} \frac{Z^{(2)}(0, \lambda_0)}{Z^{(2)}(0, 0)} &= \int dz \sqrt{\frac{|\Lambda_N|}{2\pi}} e^{-|\Lambda_N| \frac{z^2}{2}} \left\langle e^{\beta z \sqrt{\lambda_0} |\Lambda_N| (q_{12} - \bar{q}_N)} \right\rangle_{\kappa, \beta \sqrt{q_N}, N}^{\otimes 2} \\ &= \int dz \sqrt{\frac{|\Lambda_N|}{2\pi}} e^{|\Lambda_N| (-z^2/2 + 2\alpha_N(\beta z \sqrt{\lambda_0}; J))}. \end{aligned}$$

Equations (23) and (20) imply that, for $\beta < \beta_0(h) = 1/\sqrt{4C\lambda_0}$

$$\begin{aligned} \frac{1}{2|\Lambda_N|} \mathbb{E} \log \frac{Z^{(2)}(0, \lambda_0)}{Z^{(2)}(0, 0)} &\leq \frac{1}{2|\Lambda_N|} \mathbb{E} \log \int dz \sqrt{\frac{|\Lambda_N|}{2\pi}} e^{|\Lambda_N| (-z^2/2 + 2z\beta\sqrt{\lambda_0}\alpha'_N(0; J))} \\ &\leq C' \mathbb{E} (\alpha'_N(0; J))^2 \\ &= \frac{C'}{4|\Lambda_N|^2} \sum_{i, j \in \Lambda_N} \mathbb{E} ((\langle \sigma_i \rangle^2 - \mathbb{E} \langle \sigma_i \rangle^2) (\langle \sigma_j \rangle^2 - \mathbb{E} \langle \sigma_j \rangle^2)), \quad (24) \end{aligned}$$

where for simplicity we have written $\langle \cdot \rangle$ for $\langle \cdot \rangle_{\kappa, \beta \sqrt{q_N}, N}^{\otimes 2}$. We now show that the last expression is of order $1/|\Lambda_N|$. Indeed, take two distinct sites i, j and consider the d -dimensional ball B_{ij} of radius $|i - j|/2$ centered at i . If

$\langle \sigma_i \rangle$ were depending only on $(J_k, k \in B_{ij})$ and $\langle \sigma_j \rangle$ on $(J_k, k \in B_{ij}^c)$ only, the corresponding term in (24) would be zero by independence. But in the Dobrushin region, we can approximate $\langle \sigma_i \rangle$ by the expectation of σ_i for the finite volume RFIM on B_{ij} with an error which is exponentially small in the radius $|i-j|/2$, uniformly in the J_k 's. Doing similarly with $\langle \sigma_j \rangle$, we conclude that

$$\mathbb{E}((\langle \sigma_i \rangle^2 - \mathbb{E} \langle \sigma_i \rangle^2)(\langle \sigma_j \rangle^2 - \mathbb{E} \langle \sigma_j \rangle^2)) \leq C' \rho^{-C''|i-j|}$$

for suitable constants $C', C'' > 0$. This, together with (24), immediately implies that

$$\frac{1}{2|\Lambda_N|} \mathbb{E} \log \frac{Z^{(2)}(0, \lambda_0)}{Z^{(2)}(0, 0)} \leq \frac{c}{|\Lambda_N|}.$$

At this point, recalling Eq. (16), one finds

$$p_N(\kappa, \beta, h) \geq p_N^{RFIM}(\kappa, h, \beta\sqrt{\bar{q}_N}) + \frac{\beta^2}{4}(1 - \bar{q}_N)^2 - (e^{1/(\lambda_0-1)} - 1) \frac{c}{|\Lambda_N|},$$

which, together with (13), proves the convergence in average of the pressure to the expression (5). Moreover, from Eqs. (15) and (12) one deduces that

$$\limsup_{N \rightarrow \infty} \sup_{0 \leq t \leq 1} |\Lambda_N| \mathbb{E} \langle (q_{12} - \bar{q}_N)^2 \rangle_t^{\otimes 2} < \infty, \quad (25)$$

which will be needed in next section, where we deal with pressure fluctuations.

3.5 Proof of Theorem 3

We follow the strategy which was introduced in [11], adapted to the present case where short-range interactions are also present. Let

$$\hat{f}_N(t) = \sqrt{|\Lambda_N|} \left(\frac{\log Z(t)}{|\Lambda_N|} - p_N(t) \right),$$

where $Z(t), p_N(t)$ were defined in Eqs. (11)-(12) and it is understood that q is taken to be $\bar{q}_N = \bar{q}_N(\beta, \kappa, h)$ as in Eq. (18), to be distinguished from $\bar{q} = \lim \bar{q}_N$.

We will prove that

$$\lim_{N \rightarrow \infty} \mathbb{E} e^{iu\hat{f}_N(t)} = \exp \left(-\frac{u^2}{2} \left(\Gamma - \frac{\beta^2}{2} t \bar{q}^2 \right) \right) \quad (26)$$

for any $u \in \mathbb{R}$, $0 \leq t \leq 1$, from which Eq. (6) follows for $t = 1$.

By means of integrations by parts one finds

$$\begin{aligned} \partial_t \mathbb{E} e^{iu\hat{f}_N(t)} &= iu \mathbb{E} e^{iu\hat{f}_N(t)} \frac{d}{dt} \hat{f}_N(t) \\ &= \frac{\beta^2}{4} u^2 \bar{q}_N^2 \mathbb{E} e^{iu\hat{f}_N(t)} - \frac{\beta^2}{4} u^2 \mathbb{E} e^{iu\hat{f}_N(t)} \langle (q_{12} - \bar{q}_N)^2 \rangle_t^{\otimes 2} \\ &\quad - i \frac{\beta^2}{4} u \sqrt{|\Lambda_N|} \mathbb{E} e^{iu\hat{f}_N(t)} \left(\langle (q_{12} - \bar{q}_N)^2 \rangle_t^{\otimes 2} - \mathbb{E} \langle (q_{12} - \bar{q}_N)^2 \rangle_t^{\otimes 2} \right) \end{aligned}$$

and, using Eq. (25),

$$\partial_t \mathbb{E} e^{iu\hat{f}_N(t)} = \frac{\beta^2 u^2 \bar{q}^2}{4} \mathbb{E} e^{iu\hat{f}_N(t)} + o(1).$$

Integrating in t , Eq. (26) is then proven provided that we show that

$$\lim_{N \rightarrow \infty} \mathbb{E} e^{iu\hat{f}_N(0)} = \exp\left(-\frac{u^2}{2}\Gamma\right), \quad (27)$$

i.e., a central limit theorem for pressure fluctuations of the RFIM at high temperature.

To this purpose, we employ the central limit theorem for martingales [13], which we recall for convenience. Let (Ω, \mathcal{A}, P) be a probability space, $\mathcal{F}^{(n)} = \{\mathcal{F}_k^n\}_{0 \leq k \leq n}$ a filtration of \mathcal{A} , for $n \in \mathbb{N}$, such that $\mathcal{F}_0^n = \{\emptyset, \Omega\}$, and $\xi^{(n)} = \{\xi_{n,k}\}_{1 \leq k \leq n}$ a sequence of random variables adapted to $\mathcal{F}^{(n)}$. We denote by $E^{n,k}$ (respectively $P^{n,k}$) the expectation (respectively the probability) conditioned to \mathcal{F}_k^n and by $V^{n,k}$ the conditional variance

$$V^{n,k}(X) = E^{n,k}(X^2) - (E^{n,k}X)^2,$$

of a random variable X . We say that the triangular array $\{\xi_{n,k}\}_{n>0, 1 \leq k \leq n}$ is asymptotically negligible if for any $\varepsilon > 0$

$$\sum_{k=1}^n P^{n,k-1}(|\xi_{n,k}| \geq \varepsilon) \xrightarrow{P} 0 \quad (28)$$

when $n \rightarrow \infty$. Then, the following holds:

Theorem 4. *Let $\{\xi_{n,k}\}_{n>0, 1 \leq k \leq n}$ be an asymptotically negligible triangular array of square integrable random variables, and assume that for some $\Gamma > 0$,*

$$\sum_{k=1}^n E^{n,k-1}(\xi_{n,k}) \xrightarrow{P} 0 \quad (29)$$

and

$$\sum_{k=1}^n V^{n,k-1}(\xi_{n,k}) \xrightarrow{P} \Gamma, \quad (30)$$

for $n \rightarrow \infty$. Then,

$$\sum_{k=1}^n \xi_{n,k} \xrightarrow{\text{law}} \mathcal{N}(0, \Gamma).$$

To simplify notations in our case, let $|\Lambda_N| = n$, $h + \beta\sqrt{q_N}J_i = h_i$ and

$$p_{n,\underline{h}} = p_N^{RFIM}(\kappa, h, \beta\sqrt{q_N}; J).$$

Introducing the usual lexicographic ordering of the sites in Λ_N , we define \mathcal{F}_k^n , for $k \in \Lambda_N$, as the σ -algebra generated by the random fields J_i for $i \in \Lambda_N, i \preceq k$, and

$$\xi_{n,k} = \sqrt{n}(E^{n,k}p_{n,\underline{h}} - E^{n,k-1}p_{n,\underline{h}})$$

so that

$$\sqrt{|\Lambda_N|}(p_N^{RFIM}(\kappa, h, \beta\sqrt{q_N}; J) - \mathbb{E}p_N^{RFIM}(\kappa, h, \beta\sqrt{q_N}; J)) = \sum_{k \in \Lambda_N} \xi_{n,k}. \quad (31)$$

Of course, $E^{n,k}(\cdot) = \mathbb{E}_{J_{\ell, \ell \in \Lambda_N, \ell \succ k}}(\cdot)$, and by convention $\mathcal{F}_k^n = \{\emptyset, \Omega\}$ if k precedes the first site in Λ_N . One can rewrite

$$\xi_{n,k} = -\frac{1}{\sqrt{n}} \mathbb{E}_{J'_k, J_{\ell, \ell \succ k}} \log \left\langle e^{(h'_k - h_k)\sigma_k} \right\rangle_{n,\underline{h}}$$

where J'_k is an independent copy of J_k – independent of $(J_l, l \in \mathbb{Z}^d)$ –, and $h'_k = h + \beta\sqrt{q_N}J'_k$, so that

$$|\xi_{n,k}| \leq n^{-1/2} \mathbb{E}_{J'_k} |h'_k - h_k| \leq Cn^{-1/2}(|h_k| + C)$$

for some constant C and

$$E^{n,k-1} |\xi_{n,k}|^3 \leq C'n^{-3/2}.$$

This implies asymptotic negligibility (28), since

$$\sum_{k \in \Lambda_N} P^{n,k-1} (|\xi_{n,k}| \geq \varepsilon) \leq \frac{1}{\varepsilon^3} \sum_{k \in \Lambda_N} E^{n,k-1} (|\xi_{n,k}|^3) \leq \frac{C'}{\varepsilon^3 \sqrt{n}}.$$

In order to apply Theorem 4, we have to check conditions (29) and (30). The first one is evident, since

$$E^{n,k-1}\xi_{n,k} = 0$$

identically. As for the second, notice that

$$V^{n,k-1}(\xi_{n,k}) = \frac{1}{n} \mathbb{E}_{J_k} \left(\mathbb{E}_{J'_k, J_\ell, \ell \succ k} \log \left\langle e^{(h'_k - h_k)\sigma_k} \right\rangle_{n, \underline{h}} \right)^2. \quad (32)$$

Let k correspond to a site “in the bulk” of Λ_N , i.e., assume that the distance between k and the boundary of Λ_N is larger than, say, $n^{1/(2d)}$. In this case, we will write $k \in B_N$. We want to replace $\langle \cdot \rangle_{n, \underline{h}}$ in (32) with the unique infinite-volume Gibbs measure $\langle \cdot \rangle_{\infty, \underline{h}}$. To this purpose, note preliminarily that

$$\left| \log \frac{\langle e^{(h'_k - h_k)\sigma_k} \rangle_{n, \underline{h}}}{\langle e^{(h'_k - h_k)\sigma_k} \rangle_{\infty, \underline{h}}} \right| \leq 2|h'_k - h_k|.$$

Moreover, thanks to Dobrushin’s theorem,

$$\frac{\langle e^{(h'_k - h_k)\sigma_k} \rangle_{n, \underline{h}}}{\langle e^{(h'_k - h_k)\sigma_k} \rangle_{\infty, \underline{h}}} = 1 + \frac{e^{(h'_k - h_k)} - e^{-(h'_k - h_k)}}{\langle e^{(h'_k - h_k)\sigma_k} \rangle_{\infty, \underline{h}}} \delta_{n, \underline{h}} \quad (33)$$

with some $\delta_{n, \underline{h}}$ such that

$$\lim_{n \rightarrow \infty} \sup_{k \in B_N} \sup_{\underline{h}} |\delta_{n, \underline{h}}| \equiv \lim_{n \rightarrow \infty} \epsilon_n = 0.$$

Denoting by $A_{n,k}$ the event

$$A_{n,k} = \left\{ |h'_k - h_k| \leq \frac{1}{2} \log \left(\frac{1}{2\epsilon_n} \right) \right\},$$

and using the fact that $|\log(1+x)| \leq D|x|$ for $|x| \leq 1/2$ for some finite constant D , one can write

$$\left| \log \frac{\langle e^{(h'_k - h_k)\sigma_k} \rangle_{n, \underline{h}}}{\langle e^{(h'_k - h_k)\sigma_k} \rangle_{\infty, \underline{h}}} \right| \leq D\epsilon_n e^{2|h'_k - h_k|} + 2|h'_k - h_k| 1_{A_{n,k}^c}.$$

Therefore one has

$$\begin{aligned} V^{n,k-1}(\xi_{n,k}) &= \frac{1}{n} \mathbb{E}_{J_k} \left(\mathbb{E}_{J'_k, J_\ell, \ell \succ k} \log \left\langle e^{(h'_k - h_k)\sigma_k} \right\rangle_{\infty, \underline{h}} \right)^2 + \frac{1}{n} o(1) \\ &= \frac{1}{n} (\phi(\theta_{-k} \underline{h}) + o(1)), \end{aligned}$$

where $o(1) \rightarrow 0$ for $n \rightarrow \infty$ uniformly in $k \in B_N$, θ_k is the shift of vector k and

$$\phi(\underline{h}) = \mathbb{E}_{J_0} \left(\mathbb{E}_{J'_0, J_\ell, \ell > 0} \log \left\langle e^{(h'_0 - h_0)\sigma_0} \right\rangle_{\infty, \underline{h}} \right)^2,$$

the subscript $_0$ referring of course to the origin of the lattice \mathbb{Z}^d . Note that there is a residual n -dependence in ϕ , since the fields h_i are defined through \bar{q}_N , but this dependence is easily seen to be harmless thanks to the exponential decay of correlations inside Dobrushin's uniqueness region, and to the fact that $\bar{q}_N \rightarrow \bar{q}$. Finally, defining $\tilde{h}_i = h + \beta\sqrt{\bar{q}}J_i$, the ergodic theorem implies that, for almost every J ,

$$\lim_{n \rightarrow \infty} \sum_{k \in \Lambda_N} V^{n, k-1}(\xi_{n, k}) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k \in \Lambda_N} \phi(\theta_{-k}\underline{h}) = \mathbb{E} \phi(\tilde{\underline{h}}) \equiv \Gamma(\kappa, \beta, h),$$

where we used the fact that the contribution of the spins $k \notin B_N$ vanishes for $|\Lambda_N| \rightarrow \infty$. At this point, all the conditions necessary to apply Theorem 4 are fulfilled and, recalling (31), one has (27), which concludes the proof of Theorem 3.

Acknowledgments

We would like to thank Giambattista Giacomin for many interesting discussions. F.G. and F.L.T. are grateful to the *Laboratoire de Probabilités et Modèles Aléatoires P6 & 7* for kind hospitality. F.L.T. was partially supported by Swiss Science Foundation Contract No. 20-100536/1.

References

- [1] M. Aizenman, J. L. Lebowitz, D. Ruelle: Some rigorous results on the Sherrington-Kirkpatrick spin glass model. *Comm. Math. Phys.* **112** (1987), 3–20.
- [2] M. Aizenman, J. Wehr: Rounding effects of quenched randomness on first-order phase transitions. *Comm. Math. Phys.* **130** (1990), 489–528.
- [3] F. Comets, G. Giacomin, J. Lebowitz: The Sherrington-Kirkpatrick model with short range ferromagnetic interactions. *C. R. Acad. Sci. Paris Sér. I Math.* **328** (1999), no. 1, 57–62.
- [4] F. Comets, J. Neveu: The Sherrington-Kirkpatrick model and stochastic calculus: the high temperature case. *Comm. Math. Phys.* **166** (1995), 459–464.

- [5] H. Föllmer: Random fields and diffusion processes. In *École d'été de Probabilités de Saint-Flour XV–XVII, 1985–87*, 101–203, Lecture Notes in Math. 1362, Springer, 1988.
- [6] Georgii, Hans-Otto: Gibbs measures and phase transitions. de Gruyter, Berlin, 1988.
- [7] F. Guerra: *Sum rules for the free energy in the mean field spin glass model*, in *Mathematical Physics in Mathematics and Physics: Quantum and Operator Algebraic Aspects*, Fields Institute Communications **30**, AMS, 2001.
- [8] F. Guerra: Replica Broken Bounds in the Mean Field Spin Glass Model. *Commun. Math. Phys.* **233** (2003), 1–12.
- [9] F. Guerra, F. L. Toninelli: Quadratic replica coupling in the Sherrington-Kirkpatrick mean field spin glass model. *J. Math. Phys.* **43** (2002), 3704–3716.
- [10] F. Guerra, F. L. Toninelli: The thermodynamic limit in mean field spin glass models. *Comm. Math. Phys.* **230** (2002), 71–79.
- [11] F. Guerra, F. L. Toninelli: Central limit theorem for fluctuations in the high temperature region of the Sherrington-Kirkpatrick spin glass model. *J. Math. Phys.* **43** (2002), 6224–6237.
- [12] M. Ledoux: The Concentration of Measure Phenomenon. *Mathematical Surveys and Monographs* **89**, AMS, 2002.
- [13] A. N. Shiryaev: Probability. Springer, 1996.
- [14] Talagrand, M: The Parisi Formula. *Ann. Math*, to appear.
- [15] Talagrand, Michel: On the high temperature phase of the Sherrington-Kirkpatrick model. *Ann. Probab.* **30** (2002), 364–381.
- [16] M. Talagrand: Spin Glasses: A Challenge for Mathematicians. Cavity and mean field models. Springer, 2003.
- [17] Vuillermot, P.: Thermodynamics of quenched random spin systems, and application to the problem of phase transitions in magnetic (spin) glasses. *J. Phys.* **A10** (1977) 1319–1333.

Francis Comets: Université Paris 7 – Denis Diderot, Mathématiques, case 7012, 2 place Jussieu, 75251 Paris Cedex 05, France. Email: comets@math.jussieu.fr
<http://www.proba.jussieu.fr/pageperso/comets/comets.html>

Francesco Guerra: Dipartimento di Fisica, Università di Roma “La Sapienza” and INFN, Sezione di Roma 1, P.le Aldo Moro 2, 00185 Roma, Italy. Email:

francesco.guerra@roma1.infn.it

Fabio Lucio Toninelli: Laboratoire de Physique, UMR-CNRS 5672, ENS Lyon, 46
Allée d'Italie, 69364 Lyon Cedex 07, France. Email: ftonine@ens-lyon.fr