



Near-IR spectroscopy of asteroids 21 Lutetia, 89 Julia, 140 Siwa, 2181 Fogelin, and 5480 (1989YK8), potential targets for the Rosetta mission; remote observations campaign on IRTF

Mirel Birlan, Schelte J. Bus, Irina Belskaya, Sonia Fornasier, Maria Antonietta Barucci, Richard P. Binzel, Pierre Vernazza, Marcello Fulchignoni

► **To cite this version:**

Mirel Birlan, Schelte J. Bus, Irina Belskaya, Sonia Fornasier, Maria Antonietta Barucci, et al.. Near-IR spectroscopy of asteroids 21 Lutetia, 89 Julia, 140 Siwa, 2181 Fogelin, and 5480 (1989YK8), potential targets for the Rosetta mission; remote observations campaign on IRTF. Sent: October 2003, Accepted: December, 2003. 2003. <hal-00000987>

HAL Id: hal-00000987

<https://hal.archives-ouvertes.fr/hal-00000987>

Submitted on 29 Dec 2003

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Near-IR spectroscopy of asteroids 21 Lutetia, 89 Julia, 140 Siwa, 2181 Fogelin, and 5480 (1989YK8), potential targets for the Rosetta mission; remote observations campaign on IRTF

Mirel Birlan^{a,b,*} Maria Antonietta Barucci^a Pierre Vernazza^a
 Marcello Fulchignoni^a Richard P. Binzel^c Schelte J. Bus^d
 Irina Belskaya^{e,a} Sonia Fornasier^f

^a*Observatoire de Paris-Meudon, LESIA, 5 Place Jules Janssen, 92195 Meudon Cedex, CNRS, France*

^b*Astronomical Institute of the Romanian Academy, Str Cutitul de Argint n 5, Bucharest 28 Romania*

^c*Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA, 02139 USA*

^d*Institute for Astronomy, 640 North A'ohoku Place, Hilo, HI 96720, USA*

^e*Astronomical Observatory of Kharkov University, Sumskaia str. 35, Kharkov 310022, Ukraine*

^f*Astronomical Department of Padova, Vicolo dell'Osservatorio 2, 35122 Padova, Italy*

Abstract

In the frame of the international campaign to observe potential target asteroids for the Rosetta mission, remote observations have been carried out between Observatoire de Paris, in Meudon-France, and the NASA Infrared Telescope Facility on Mauna Kea. The SpeX instrument was used in the 0.8-2.5 microns spectral region, for two observing runs in March and June 2003.

This paper presents near-IR spectra of the asteroids 21 Lutetia, 89 Julia, 140 Siwa, 2181 Fogelin, and 5480 (1989YK8).

Near-IR spectra of the asteroids 21 Lutetia and 140 Siwa are flat and featureless. The spectrum of 89 Julia reveals absorption bands around 1 and 2 microns, which may indicate the presence of olivine and olivine-pyroxene mixtures and confirm the S-type designation.

The small main-belt asteroids 2181 Fogelin and 5480 (1989YK8) are investigated spectroscopically for the first time. Near-IR spectra of these asteroids also

show an absorption feature around 1 micron, which could be an indicator of igneous/metamorphic surface of the objects; new observations in visible as well as thermal albedo data are necessary to draw a reliable conclusion on the surface mineralogy of both asteroids.

Key words: Remote observation, spectroscopy, Asteroid, Rosetta space mission
PACS: 95.45.+i, 95.85.Jq, 96.30.Ys

1 Introduction

The observational program presented in this paper is intimately linked to the scientific programme of the space mission Rosetta. The Rosetta mission was approved in 1993 as a Cornerstone Mission within the ESA's Horizons 2000 science programme. The original launch date for the mission was January 2003, and the scientific targets were assigned to be the nucleus of comet 46P/Wirtanen and two main-belt asteroids: 140 Siwa and 4979 Otawara. The main scientific goals of the Rosetta mission are to investigate 'in-situ' primitive objects of the Solar System and to find answers concerning the chemical composition of the primitive planetary nebula, thought to be 'frozen' in the comet nucleus and the asteroid mineralogical matrix.

Since 1993, several international campaigns of observations have been started in order to obtain a large amount of data for the targets of the Rosetta mission (Fornasier et al. (2003), LeBras et al. (2001), Barucci et al. (1998), etc). The groundbased knowledge of these objects is essential to optimise the science return as well as the mission trajectory.

In January 2003 the European Space Agency decided to postpone the launch of the Rosetta, with a new launch date set to February 2004. This decision implied a new mission baseline: rendezvous with the comet P/Churyumov-Grasimenko and one or two asteroid fly-bys to be defined after the spacecraft interplanetary orbit insertion manoeuvre, when the total amount of available δv will be known. Several fly-bys scenarios have been studied and several possible asteroid targets have been found. These potential targets are poorly known and for this reason systematic observations are needed in order to significantly improve on our knowledge of the physics and the mineralogy of these objects.

In this paper we present the spectroscopic results obtained for five asteroids (21 Lutetia, 89 Julia, 140 Siwa, 2181 Fogelin, and 5480 (1989YK8)), potential targets of the Rosetta mission. The 0.8-2.5 microns wavelength region was

* Corresponding author: Mirel.Birlan@obspm.fr

investigated using the SpeX instrument on IRTF-Hawaii, in 'remote observing' mode.

2 The Observing Protocol

The remote observations were conducted from Meudon-France, more than 12,000 km away from Hawaii, using several informational structures and networks. For the observers in Meudon, the observing hours occurred during relatively normal working daylight hours. Observation sessions began at 5 a.m. local time and ended at 5 p.m. local time. This type of observations between Meudon and IRTF/Hawaii was started in 2002 (Birlan & Binzel (2002)); since then, more than twenty nights of observations (in eight runs) were conducted from Meudon. The observations were realized through an ordinary network link, without the service quality warranty. Thus, the passband for our link was variable, a function of the traffic between Hawaii and Meudon.

During the remote observing run, the Observatoire de Paris team had the control of both the instrument/guider system and the spectrograph set-up and spectra acquisition. A permanent and constant audio/video link with the telescope operator is essential in order to administrate possible service interruptions. Two PC's running Linux were devoted to the telescope/spectrograph control. The X environment for the telescope and instrument control was exported from Mauna Kea to Meudon via two secure links (ssh tunnels). A third PC was used to keep the IP audio-video link open (webcam/Netmeeting at Meudon and Polycom ViewStation video-conference system on Mauna Kea). All software was re-initialized at the beginning of each night.

The communication lag was relatively small (approximately 0.5 seconds), and the image refresh time was 2 seconds on the average. Real-time image display was performed mainly for verification purposed and preliminary analysis. At the completion of each run, all files were transferred to Meudon

The SpeX instrument was utilized in low-resolution mode for this campaign. The observations were made in the 0.8-2.5 microns spectral interval. We used a 0.8 arcsec wide slit, with a 15 arcsec length and oriented North-South, which allowed us simultaneous measurements of the object and sky. The object position on the slit was alternated between two locations referred to as the A and B positions. The seeing varied between 0.7-1.8 arcsec during the observing runs, and the humidity was in the 25% - 55% range. The automatic guiding mode of the telescope was used for spectra acquisition.

The asteroids were observed in during two observing runs: 28-30 March 2003 and 5 July 2003. The observed objects are summarized in Table 1. We also

Table 1

Potential asteroid targets for the Rosetta mission. Observation date, semimajor axis, eccentricity, inclination, phase angle and the apparent magnitude are presented for each asteroid

| Object | Date | a (a.u) | e | i(deg) | Φ (deg) | V(mag) |
|--------------|---------------|---------|--------|---------|--------------|--------|
| 21 Lutetia | March 30 2003 | 2.4347 | 0.1636 | 3.0645 | 14.6 | 11.32 |
| 89 Julia | March 30 2003 | 2.5519 | 0.1825 | 16.1437 | 13.2 | 11.39 |
| 140 Siwa | March 30 2003 | 2.7365 | 0.2153 | 3.1882 | 21.2 | 13.72 |
| 2181 Fogelin | July 5 2003 | 2.5918 | 0.1177 | 13.0205 | 11.2 | 16.59 |
| 5480 1989YK8 | March 30 2003 | 3.1366 | 0.087 | 6.6717 | 12.7 | 16.25 |

observed the standard stars SA 98-978, SA 102-1081, SA 105-56, SA 107-684, SA 113-276, SA 115-271, HD 28099, HD 88618, and 16 CyB. Flat-fields and Argon lamp arc images were taken each night and used for data reduction.

Our strategy was to observe all asteroids as close to the zenith as possible. Thus, we managed to observe with an airmass less than 1.6 for all targets except 2181 Fogelin, which we could only observe with an airmass in the 1.7-1.78 range. Science exposures were alternated with standard stars spectra exposures, the latter taken to cover the 1-1.8 airmass range.

In order to obtain a S/N in the 80-150 range, we needed 20 to 40 minutes of exposure time, depending on the asteroid magnitude, and counting both the effective exposure and CCD camera readout time. This exposure time is unacceptable for a single near-infrared spectrum due to the large variations in the atmospheric conditions (a single near-infrared spectral exposure is usually no longer than 120 seconds). In order to obtain the required S/N, we obtained a number of 6 to 10 A and B exposure pairs (cycles) for both the asteroids -science exposures- and the standard stars. This constitutes *one series* of observations for each object. Details the science exposures are given in Table 2.

3 Data Reduction and Results

The major points of our reduction procedure are classic for the near-infrared spectroscopy. We started by combining and normalizing flat-fields for each observing night. The resulting flats were later used for the reduction of spectra of both the asteroids and the standard stars. In order to minimize the atmospheric and telescope influence and to eliminate the influence of electronic bias level and the dark current, we subtracted the B position spectra from the A position spectra for each pair of exposures (cycle), in the assumption of quasi-

Table 2

Exposure data for each asteroid. The columns show the mean UT value for each series, the individual time for each spectrum (Itime), the number of cycles and the mean airmass of each series.

| Object | UT (h m s) | Itime(s) | Cycles | Airmass |
|--------------|------------|----------|--------|---------|
| 21 Lutetia | 13 35 46 | 15 | 8 | 1.223 |
| 21 Lutetia | 11 28 17 | 20 | 4 | 1.329 |
| 21 Lutetia | 15 01 03 | 15 | 9 | 1.414 |
| 21 Lutetia | 10 20 21 | 20 | 3 | 1.635 |
| 89 Julia | 10 44 44 | 40 | 6 | 1.561 |
| 140 Siwa | 07 06 07 | 120 | 8 | 1.000 |
| 140 Siwa | 07 46 10 | 120 | 6 | 1.016 |
| 2181 Fogelin | 12 10 35 | 2x60 | 4 | 1.769 |
| 2181 Fogelin | 12 31 40 | 2x60 | 4 | 1.744 |
| 2181 Fogelin | 12 55 17 | 2x60 | 4 | 1.740 |
| 5480 1989YK8 | 12 41 01 | 120 | 6 | 1.151 |
| 5480 1989YK8 | 14 36 01 | 120 | 6 | 1.230 |

homogeneous sky background during A plus B exposure pair. The result of the subtraction was flat fielded. For each object, we median combined the result of all cycles in each observing series. This technique produces one positive and one negative spectrum on the same image. Next step was the construction of an accurate spatial profile for the extraction of the spectra. The final step was the wavelength calibration.

The Spextool package (the description of the procedures are presented in Cushing et al. (in press)) allowed us to perform following steps for both the asteroids and the standard stars: global flat-field and arc construction, possible non-linearity correction, addition of spectra of the same object, spatial profile determination of the spectrum in the image, aperture location, extraction of the spectrum, wavelength calibration and cleaning the spectra. The results are saved in both FITS and ASCII formats, as used by several image processing packages (like IDL, MIDAS, and IRAF), respectively dedicated plot/graphics software (Easyplot, Origin, Dataplot, etc).

The next step in the process of data reductions was the calculation of the extinction coefficients. The solar analogs spectra were used to find the correspondent extinction coefficient for each wavelength. A "superstar" was created by summing appropriately weighted contributions of the standard stars. The resulting "superstar", corrected for atmospheric extinction, was used to obtain the spectral reflectance of asteroids at different airmasses. Careful choice of

the "superstar", combined with several tests of shift between the spectrum of the asteroid and that of the superstar are very important in order to minimize the influence of the terrestrial atmosphere in the 1.4 and 1.9 microns spectral regions. In order to make more readable our spectra, we choose to eliminate these spectral regions. Moreover, the elimination of these spectral regions does not affect the conclusions of this article.

3.1 21 Lutetia

With an estimated diameter of 95.5 ± 4.1 km - for an IRAS albedo of 0.221 ± 0.020 (Tedesco and Veeder (1992)) - , 21 Lutetia belongs to the large asteroid class (diameter ≥ 100 km). Its synodic period has been computed from several lightcurve analysis and the value is 8.17 ± 0.01 hours (Zappala et al. (1984)). Color analysis of the ECAS data (Zellner et al. (1985)) designates the asteroid 21 Lutetia to be part of the X complex. Also, global ECAS and IRAS thermal albedo data analysis assign Lutetia to the M taxonomic type (Barucci et al. (1987), Tholen, (1989)). The M type asteroids are considered to be part of the core of differentiated asteroids and the parent bodies of metallic meteorites. On the basis of SMASS II spectroscopic data, Bus & Binzel(2002) proposed the new taxonomic class *Xk* for the asteroid 21 Lutetia.

21 Lutetia was observed on UT 2003 March 29 for the entire night. Four series of exposures resulted in four IR spectra with S/N in the 90-140 range. In order to detect possible surface spectral features variations, our observations covered 65% of the synodic period. These spectra are presented in Figure 1. However, a check of the physical ephemeris of 21 Lutetia revealed a close pole-on geometry of the asteroid, so our spectra are most probably dominated by the contribution of the same surface features on 21 Lutetia. There are no significant spectral features in the spectral region 0.8-2.5 micron, and the slightly increasing slope varies around 0.3%.

The lack of features in the spectrum of 21 Lutetia makes its taxonomic and compositional interpretation difficult. While the high IRAS albedo value of 0.221 leads to the previous M-type classification, we note there is an ambiguity in determining the albedo of Lutetia. An alternate low albedo value of 0.09 has been reported by Zellner et al. (1977) from polarimetric measurements and groundbased radiometric measurements.

In Figure 2 we compare our measurements of Lutetia with others published in the literature. We found a good match with both SMASS IR data (spectral interval 0.8-1.6 micron) and the 52 Color Asteroid Survey (Bell et al. (1988)) spectrophotometric data. We also compare in Figure 2 all of these Lutetia results with reflectance spectra of meteorites from Gaffey (1976). Our goal




Fig. 1. The spectra of the asteroid 21 Lutetia. The IRTF spectra are offset for clarity and presented in chronological order. The beginning of each series of spectra(UT) was marked on the graph.

is to find which meteorites may be most analogous with Lutetia in terms of near-infrared spectral properties. We note the spectrum of Lutetia is most qualitatively similar to the spectrum of the Vigarano meteorite, a CV3 carbonaceous chondrite, while being quite a poor match to the class IIIA iron meteorite Chulafinee. The purpose of our comparison is to note the difficulty of making a non-ambiguous interpretation of the composition of 21 Lutetia.

In Figure 3 we plot the negative polarization depth versus the inversion angle (Belskaya & Lagerkvist (1996)) for both 21 Lutetia and a sample asteroids and meteorites. The asteroid 21 Lutetia has the largest inversion angle ever observed for the asteroids. The same peculiarly large inversion angle were found in the laboratory for the carbonaceous chondrites of the CV3 and CO3 samples. These types of chondrites are characterized by a low carbon content and thus relatively a larger albedo when compared to other types (Zellner et al. (1977)).




Fig. 2. The average spectrum of the asteroid 21 Lutetia. The SMASS II and SMASS IR data are plotted as solid lines. Our data is in agreement with the SMASSIR data over the 0.75-1.6 microns spectral interval. The comparison with Chulafinnee (iron meteorite) and Vigarano (carbonaceous chondrite) reveal the good match with the IR spectrum associated to carbon rich surfaces.

3.2 89 Julia

The asteroid 89 Julia has an estimated diameter of 151.4 ± 3.1 km, for an IRAS albedo of 0.176 ± 0.007 . Photometry of 89 Julia yields a synodic period of 11.387 ± 0.002 hours (Schober and Lustig (1975)). Multivariate statistics classified the asteroid as part of the S cluster (Barucci et al. (1987), Tholen, (1989)). The mineralogical classification of the S-type asteroids (Gaffey et al. (1993)) based on the ECAS data and 52-Color spectrophotometric Survey (Bell et al. (1988)) data assigns 89 Julia (together with the asteroid 9 Metis) as ‘ungrouped’ S-asteroids. The main reason of this ‘unclassification’ is the long wavelength position of the 1-micron feature, which could be the presence of the abundant calcic clinopyroxene component (Gaffey et al. (1993)). Based on the SMASS II data, Bus and Binzel(2002) proposed a K cluster membership (derived from the S cluster) for 89 Julia.

The asteroid spectrum was obtained for an effective integration time of 10 minutes, with a S/N ratio of 90 in both A and B beams. This spectrum




Fig. 3. Diagram of the negative polarization depth and the inversion angle of asteroids and meteorites from Belskaya & Lagerkvist (1996).

presents a significant positive slope in the region 1.1-1.5 micron, and a plateau in the 1.7-1.9 microns and 2.2-2.5 microns spectral regions. In Figure 4 the IR spectrum of Julia was overlapped with the SMASS II spectrum in the 0.78-0.92 microns spectral region. We confirm a broad absorption band at 1 micron, typical of the silicate rich minerals. We also confirm a global trend of the spectral reflectance, which increases in the near-infrared. The influence of atmospheric water absorption on the spectra is visible around 1.4 and 1.9 microns. The geometry of observations was fairly unfavourable for this asteroid (airmass = 1.56).

In order to calculate the center of 1 micron absorption band, the continuum was defined between the local maximum of spectral reflectance at 0.76 microns and the small shoulder of the spectrum at 1.27 microns. The spectrum was then continuum subtracted and the result was fitted with three-degree polynomial function. We find the center of the feature localized at 1.01 ± 0.06 micron, which is a slightly shorter wavelength than the one found by Gaffey et al (1993).




Fig. 4. The spectrum of the asteroid 89 Julia. The IRTF spectrum was overlapped with the SMASS II visible spectrum in the region 0.78-0.92 micron.

3.3 140 Siwa

140 Siwa is one of the initial targets of the Rosetta mission (departure in January 2003). One of the proposed scenarios (February 2003) suggests keeping this asteroid as a candidate for fly-by. This was our motivation to observe it.

An IRAS albedo of 0.068 ± 0.004 for the asteroid 140 Siwa results in a of 109.8 ± 3.0 km diameter. Lightcurve analysis for two runs in 2000 (LeBras et al. (2001)) reveals a slow-rotator asteroid; the composite lightcurve presents an amplitude of 0.1 magnitudes and its synodic period was estimated at 18.495 ± 0.005 hours.

Two series of near-infrared spectra were obtained in March 30, 2003, with the ratio S/N of 50 and 70 (obtained in both A and B beams) for the spectrum obtained at airmass 1.001 and 1.018 respectively (Figure 5). The IR spectrum of 140 Siwa does not contain deep absorption features corresponding to mafic minerals. The spectrum slope is slightly positive, with a value 1% (0.8% and 0.7% respectively), which confirms the slope trend of its spectrum (LeBras et al. (2001)). We confirm a typical neutral spectrum of consistent with the a C-type asteroid; the IRAS albedo for 140 Siwa also fits very well the average value of C taxonomic class. The near-IR slope of our spectra is slightly different




Fig. 5. The spectra of the asteroid 140 Siwa. The spectra were offset for clarity and the UT represents the beginning of each series of exposures.

of those of a typical C-type asteroid but this alone cannot indicate a different mineralogy of the surface.

Figure 6 shows the computed a global IRTF spectrum for the asteroid 140 Siwa overlapped with SMASS II data (the overlap is only for the 0.7-0.85 microns region). This composite spectrum was compared to meteorite reflectance spectra from the Gaffey database. There is no meteorite spectrum that fits our spectrum very well. However, the carbonaceous chondrite meteorites spectra are similar to our spectrum, and the asteroid spectral reflectance spans a value in the range of CV3 meteorite class. The nearest spectra are those of Grosnaja and Mokoia meteorites.

3.4 2181 Fogelin

No physical data are available in the literature for the asteroid 2181 Fogelin. Thus, we can only roughly estimate its diameter. Following the asteroid mass distribution proposed by Kresak (1977) and the H magnitude of 2181 Fogelin, the diameter could be in the 12-18 km range. A second estimate can be made using an empirical relationship between the absolute magnitude, diameter and




Fig. 6. Cumulated spectral reflectance of the asteroid 140 Siwa obtained during the IRTF run. The spectrum was overlapped with the SMASS II data in the 0.70-0.85 microns spectral region. The spectrum is similar to CV3 meteorites Grosnaja and Mokoia. The surface of 140 Siwa shows similar properties with carbonaceous chondrite meteorites, with a reflectance trend between the curves for the two meteorites.

geometric albedo (Fowler and Chillemi (1992)):

$$\log D = 3.1236 - 0.5 \cdot \log p_v - 0.2 \cdot H \quad (1)$$

Using an IRAF albedo of 0.05-0.25 we find a 12-22 km diameter for the asteroid.

The observations of 2181 Fogelin were carried out in July, 5, 2003, with the ratio S/N of 80 in both A and B beams. The geometry of observation (airmass = 1.7) was unfavourable. However, the final spectrum is the average of 32 individual spectra, which can alleviate the high airmass problem. The IR spectrum presented in Figure 7 shows the large broad band absorption around 1 micron, typical of mafic minerals. The slightly positive slope in the 1.2-1.5 microns wavelength region doesn't allow us to classify this object as a S-type asteroid. Further observations in the visible region as well as thermal albedo data are necessary to make a taxonomic class assignment for this asteroid.




Fig. 7. The spectrum of the asteroid 2181 Fogelin, normalized to 1.275 microns, the maximum of J filter.

3.5 5480 1989YK8

There are no previous photometric or spectroscopic data available for the asteroid 5480 1989YK8. The same methods used for 2181 Fogelin have been used to compute its theoretical diameter. The diameter spans the 13-20 km range (following Kresak (1977)) and 13-31 km following the diameter empirical formula (1), used with IRAF albedos in the 0.05-0.25 range.

The asteroid was observed on March, 30, 2003, and the final spectrum S/N ratio is around 100 for the A and B positions in the slit. The analysis of the 0.8-2.3 microns region reveals an absorption band around 1 micron (Figure 8). The global tendency in near-infrared could justify a S-type asteroid classification; however, a visible spectrum and thermal albedo data are necessary for a taxonomical assignment.

4 Conclusions

Five asteroids potential targets for the Rosetta mission have been observed in the 0.8-2.5 microns spectral range: 21 Lutetia, 89 Julia, 140 Siwa, 2181 Fogelin




Fig. 8. The spectrum of the asteroid 5480 1989YK8, normalized to 1.275 microns, the maximum of J filter.

and 5480 (1989YK8). We confirm the flat near-IR spectrum of 21 Lutetia. The near-infrared spectrum of 89 Julia contains features confirming the initial idea that it belongs to the ‘heated’ bodies, possible to the S or K taxonomic class. A value 1.01 ± 0.06 microns was found for the center of the Band I. The asteroid 140 Siwa spectrum presents a slightly positive slope, and there are no major mineralogical signatures; we confirm the assignment of the asteroid as an ‘unheated’ asteroid.

The small asteroids 2181 Fogelin and 5480 (1989YK8) were observed spectroscopically for the first time. Our data shows the presence of a large (and weak) absorption band around 1 micron for both asteroids; new observations in visible and infrared are necessary to draw a reliable conclusion concerning their surface mineralogy.

The remote observations between IRTF and Observatoire de Paris-Meudon proved to be a robust and handy observing technique. It offered full access to the command line of the spectrograph and to several telescope controls (focus, tracking,...). We consider our observing program entirely accomplished without any discernable difference in the typical amount of spectra obtained in remote observations mode versus the local observing mode.

5 Acknowledgements

The authors are grateful to Paul Hardersen for useful comments that improved our article. We are indebted to Tony Denault who answered many questions prior to our runs about the Polycom system and exporting of X-windows for SpeX and guiding system, to Miranda Hawarden-Ogata for the administration system support, and to Paul Sears and Bill Golish, the telescope operators during the remote observing runs.

References

- Barucci, M.A., Doressoundiram, A., Fulchignoni, M., Florczak, M., Lazzarin, M., Angeli, C. - Compositional type characterization of Rosetta asteroid candidates *Planetary and Space Science*, v. 46, p. 75-82, 1998.
- Barucci, M.A., Capria, M.T., Coradini, A., Fulchignoni, M. - Classification of asteroids using G-mode analysis, *Icarus*, vol. 72, p. 304-324, 1987.
- Bell, J.F., Hawke, B.R., Owensby P.D., Gaffey, M.J. - The 52-Color Asteroid Survey; Final results and interpretation, *Lunar Planet. Sci. XIX*, 1988.
- Belskaya, I., Lagerkvist, C.-I. - Physical properties of M class asteroids, *Planet. Sp. Sci.* vol. 44, No. 8, 1996.
- Birlan, M., Binzel, R.P. - Paris Observatory Remote Observing January-May 2002: Sharing the Experience to Educational Astronomy, *Global Hands-On Universe Conference Proceedings*, Paris, July, 2002.
- Bus, S.J., Binzel, R.P. - Phase II of the Small Main-Belt Spectroscopic Survey: A Feature-Based Taxonomy, *Icarus*, v.158, issue 1, p 146-177, 2002.
- Cushing, M.C., Vacca, W.D., Rayner, J.T. - Spextool: a spectral extraction package for SpeX, a 0.8-5.5 micron cross-dispersed spectrograph, *PASP* (in press).
- Fornasier, S., Barucci, M. A., Binzel, R. P., Birlan, M., Fulchignoni, M., Barbieri, C., Bus, S. J., Harris, A. W., Rivkin, A. S., Lazzarin, M., Dotto, E.; Michalowski, T., Doressoundiram, A., Bertini, I., Peixinho, N. - A portrait of 4979 Otawara, target of the Rosetta space mission, *A&A* v.398, p.327-333, 2003.
- Fowler, J.W. & Chillemi, J.R. - IRAS asteroid data processing, in *The IRAS Minor Planet Survey*, Tech. Rep.PL-TR-92-2049, Phillips Laboratory, Hanscom AF Base, MA, 1992, Editors: Tedesco E.F., Veeder, G.J., Fowler, J.W., and Chillemi, J.R.
- Gaffey, M.J. - Spectral reflectance characteristics of meteorite classes, *J. Geophys. Res.* vol 81, 1976.
- Gaffey, M.J., Burbine, T.H., Piatek, J.L., Reed, K.L., Chaky, D.A., Bell, J.F., Brown, R.H. - Mineralogical variations within the S-type asteroid class, *Icarus*, vol. 106, 1993.

- Kresak, L. - Mass content and mass distribution of the asteroid system, *Bull. Astron. Insti. Czech*, n. 28, p 65-82, 1977.
- Le Bras, A., Dotto, E., Fulchignoni, M., Doressoundiram, A., Barucci, M. A., Le Mouelic, S., Forni, O., Quirico, E - The 2000 Rosetta asteroid targets observational campaign: 140 Siwa and 4979 Otawara, *A&A*, v.379, p.660-663, 2001.
- Schober, H.J., Lustig, G. - A photometric investigation of the asteroid 89 Julia, *Icarus*, vol. 25, p. 339-343, 1975.
- Tedesco, E.F., Veeder, G.J. - IMPS albedos and diameter catalog(FP102), in *The IRAS Minor Planet Survey*, Tech. Rep.PL-TR-92-2049, Phillips Laboratory, Hanscom AF Base, MA, 1992, Editors: Tedesco E.F., Veeder, G.J., Fowler, J.W., and Chillemi, J.R.
- Tholen, D. - Asteroid taxonomic classifications, in *Asteroids II*, p. 1139-1150, University of Arizona Press, 1989.
- Zappala, V., DiMartino, M., Knezevic, Z., Djurasevic, G. - New evidence for the effect of phase angle on asteroid lightcurve shape - 21 Lutetia, *A&A*, v.130, no. 1, p. 208-210, 1984.
- Zellner, B., Tholen, D.J., Tedesco, E. F. - The eight-color asteroid survey - Results for 589 minor planets, *Icarus*, vol. 61, p. 355-416, 1985.
- Zellner, B., Leake, M., LeBerte, T., Duseaux, M., Dollfus, A. - The asteroid albedo scale. II. Laboratory polarimetry of meteorites, *Proc. Lunar Sci. Conf.: 8th*, p. 1091-1110, Pergamon Press, Oxford, 1977.