

Title: Solution enthalpies of hydroxylic compounds

Author(s): Reis, Marina¹; Moreira, Luis^{1,2}; Nunes, Nélson³; Leitão, Ruben³; Martins, Filomena¹

Source: Journal of Thermal Analysis and Calorimetry

Volume: 108 **Issue:** 2 **Pages:** 761-767 **DOI:** 10.1007/s10973-011-1975-x **Published:** May 2012

Document Type: Article; Proceedings Paper

Language: English

Conference: 3rd Joint Czech-Hungarian-Polish-Slovak Thermoanalytical Conference

Location: Stara Lesna, Slovakia

Date: Jun 27-29, 2011

Sponsor(s): Netzsch; TA Instruments; Mettler-Toledo

Abstract: Solution enthalpies of adamantan-1-ol, 2-methylbutan-2-ol, and 3-methylbutan-1-ol have been measured at 298.15 K, in a set of 16 protogenic and non-protogenic solvents. The identification and quantification of solvent effects on the solution processes under study were performed using quantitative-structure property relationships. The results are discussed in terms of solute-solvent-solvent interactions and also in terms of the influence of compound's size and position of its hydroxyl group.

Author Keywords: QSPR; Solution Enthalpy; Solvent Effects; Adamantan-1-ol; 2-Methylbutan-2-ol; 3-Methylbutan-1-ol

KeyWords Plus: Solvatochromic Comparison Method; Solvent; Solvation; Scale; 1-Bromodamantane; Derivatives; Solvolysis; Models

Reprint Address: Martins, F (reprint author), Univ Lisbon, Fac Ciências, Dept Quim & Bioquim, CQB, Ed C8, P-1749016 Lisbon, Portugal.

Addresses:

1. Univ Lisbon, Fac Ciências, Dept Quim & Bioquim, CQB, P-1749016 Lisbon, Portugal
2. CQB, Inst Super Educ & Ciencias, P-1750 Lisbon, Portugal
3. CQB, Inst Politecn Lisboa, Inst Super Engrn Lisboa, Area Dept Engrn Quim, P-1950062 Lisbon, Portugal

E-mail Address: filomena.martins@fc.ul.pt

Publisher: Springer

Publisher Address: Van Godewijkstraat 30, 3311 GZ Dordrecht, Netherlands

ISSN: 1388-6150

Citation: Reis M, Moreira L, Nunes N, Leitão R, Martins F. Solution enthalpies of hydroxylic compounds. Journal of Thermal Analysis and Calorimetry. 2012; 2 (108).