

Title: Oxo-rhenium Complexes Bearing the Water-Soluble Tris(pyrazol-1-yl)methanesulfonate, 1,3,5-Triaza-7-phosphaadamantane, or Related Ligands, as Catalysts for Baeyer-Villiger Oxidation of Ketones

Author(s): Martins, Luisa M. D. R. S. ^[1,2]; Alegria, Elisabete C. B. A. ^[1,2]; Smolenski, Piotr ^[3]; Kuznetsov, Maxim L. ^[2]; Pombeiro, Armando J. L. ^[2]

Source: Inorganic Chemistry **Volume:** 52 **Issue:** 8 **Pages:** 4534-4546

DOI: 10.1021/ic400024r **Published:** Apr 15 2013

Document Type: Article

Language: English

Abstract: New rhenium(VII or III) complexes $[\text{ReO}_3(\text{PTA})(2)][\text{ReO}_4]$ (1) ($\text{PTA} = 1,3,5\text{-traza-7-phosphaadamantane}$), $[\text{ReO}_3(\text{mPTA})][\text{ReO}_4]$ (2) ($\text{mPTA} = \text{N-methyl-1,3,5-traza-7-phosphaadamantane cation}$), $[\text{ReO}_3(\text{HMT})(2)][\text{ReO}_4]$ (3) ($\text{HMT} = \text{hexamethylenetetramine}$), $[\text{ReO}_3(\text{eta}(2)\text{-Tpm})(\text{PTA})][\text{ReO}_4]$ (4) ($\text{Tpm} = \text{hydrotris(pyrazol-1-yl)methane}$, $\text{HC(pz)}(3)$, $\text{pz} = \text{pyrazolyl}$), $[\text{ReO}_3(\text{Hpz})(\text{HMT})][\text{ReO}_4]$ (5) ($\text{Hpz} = \text{pyrazole}$), $[\text{ReO}(\text{Tpms})(\text{HMT})]$ (6) ($\text{Tpms} = \text{tris(pyrazol-1-yl)methanesulfonate}$, $\text{O}_3\text{SC(pz)}(3)(-)$) and $[\text{ReCl}_2\{\text{N}_2\text{C(O)Ph}\}(\text{PTA})(3)]$ (7) have been prepared from the Re(VII) oxide Re_2O_7 (1-6) or, in the case of 7, by ligand exchange from the benzoyldiazenido complex $[\text{ReCl}_2(\text{N}_2\text{C(O)Ph})(\text{Hpz})(\text{PPh}_3)(2)]$, and characterized by IR and NMR spectroscopies, elemental analysis and electrochemical properties. Theoretical calculations at the density functional theory (DFT) level of theory indicated that the coordination of PTA to both Re(III) and Re(VII) centers by the P atom is preferable compared to the coordination by the N atom. This is interpreted in terms of the Re-PTA bond energy and hard-soft acid-base theory. The oxo-rhenium complexes 1-6 act as selective catalysts for the Baeyer-Villiger oxidation of cyclic and linear ketones (e.g., 2-methylcyclohexanone, 2-methylcyclopentanone, cyclohexanone, cyclopentanone, cyclobutanone, and 3,3-dimethyl-2-butanone or pinacolone) to the corresponding lactones or esters, in the presence of aqueous H_2O_2 . The effects of a variety of factors are studied toward the optimization of the process.

KeyWords Plus: Ray Crystal-Structures; Transition-Metal-Complexes; Hydrogen-Peroxide; Rhenium Complexes; Organometallic Compounds; Coordination Chemistry; Molecular-Oxygen; Mild Conditions; Ionic Liquids; Structural-Characterization

Reprint Address: Martins, LMDRS (reprint author) - ISEL, Chem Engeneering Dept Area, P-1959007 Lisbon, Portugal.

Addresses:

[1] ISEL, Chem Engeneering Dept Area, P-1959007 Lisbon, Portugal

[2] Univ Tecn Lisboa, Inst Super Tecn, Ctr Quim Estrutural, P-1049001 Lisbon, Portugal

[3] Univ Wroclaw, Fac Chem, PL-50383 Wroclaw, Poland

E-mail Addresses: lmartins@deq.isel.pt; pombeiro@ist.utl.pt

Funding:

Funding Agency	Grant Number
Fundacao para a Ciencia e a Tecnologia (FCT), Portugal	
FCT	
IST	
	PTDC/QUI-QUI/102150/2008
	PTDC/EQU-EQU/122025/2010
	PTDC/QUI-QUI/119561/2010
	PEst-OE/QUI/UI0100/2011

Publisher: Amer Chemical SOC

Publisher Address: 1155 16TH ST, NW, Washington, DC 20036 USA

ISSN: 0020-1669

Citation: MARTINS, Luisa M. D. R. S.; ALEGRIA, Elisabete C. B. A.; SMOLENSKI, Piotr; KUZNETSOV, Maxim L.; POMBEIRO, Armando J. L. - Oxorhenium Complexes Bearing the Water-Soluble Tris(pyrazol-1-yl)methanesulfonate, 1,3,5-Triaza-7-phosphaadamantane, or Related Ligands, as Catalysts for Baeyer-Villiger Oxidation of Ketones. Inorganic Chemistry. ISSN 0020-1669. Vol. 52, nr. 8 (2013), p. 4534-4546.