

# Ecotoxicological aspects related to the presence of pharmaceuticals in the aquatic environment

Lúcia H.M.L.M. Santos, A.N. Araújo, Adriano Fachini, A. Pena,  
C. Delerue-Matos, M.C.B.S.M. Montenegro

## ABSTRACT

Pharmaceuticals are biologically active and persistent substances which have been recognized as a continuing threat to environmental stability. Chronic ecotoxicity data as well as information on the current distribution levels in different environmental compartments continue to be sparse and are focused on those therapeutic classes that are more frequently prescribed and consumed. Nevertheless, they indicate the negative impact that these chemical contaminants may have on living organisms, ecosystems and ultimately, public health. This article reviews the different contamination sources as well as fate and both acute and chronic effects on non-target organisms. An extensive review of existing data in the form of tables, encompassing many therapeutic classes is presented.

*Keywords:* Pharmaceuticals, Sources, Environmental fate, Ecotoxicological effects




Fig. 1. Percentage of published studies on different therapeutic classes, expressed in relative percentage, described on 183 articles published between 1996 and 2009.

## 1. Introduction

The presence of medicines in the environment has become a recent research topic. Initially, the problem was highlighted in the US back in the 1970s [1,2] and almost a decade later in England (UK) [3–5]. Yet, it was only in the mid 90s with advances in analytical techniques that important knowledge on environmental contamination by those compounds grew. Powerful hyphenated chromatographic-detection techniques enabling detection limits within the  $\text{ng L}^{-1}$  to  $\mu\text{g L}^{-1}$  range allowed researchers to quantify a large number of medicines components (i.e. drugs and excipients) in the environment, thus compelling the scientific community to consider this contamination type as a potential issue meriting concern [6–8]. In fact, tons of them are produced annually worldwide to be consumed by humans or animals [9,10]. They are conceived primarily to have particular physiological modes of action and frequently to resist to inactivation before exerting their intended therapeutic effect. However, these same properties are paradoxically responsible either for bioaccumulation and toxic effects in aquatic and terrestrial ecosystems [10,11]. In a different way from some conventional pollutants (such as pesticides, detergents, fuels, among others), medicines are continuously delivered at low levels which might give rise to toxicity even without high persistence rates [11–13]. Wide dissemination at low concentrations mainly in the aquatic environment is evident today. Such concentrations have been detected in aquatic compartments such as influents [14–16] and effluents [17–19] from sewage treatment plants (STPs), surface waters (rivers, lakes, streams, estuaries, among others) [20–24], seawater [25], groundwater [26–28] and drinking water [29–32]. The scientific community is in broad agreement with the possibility that adverse effects may arise from the presence of pharmaceuticals not only for human health but also for aquatic organisms. Several, almost negligible effects have been shown to occur from continuous exposure during the life cycle of aquatic vertebrates and invertebrates to sub-therapeutic drug concentrations [33,34]. These effects slowly accumulate to manifest themselves into a final irreversible condition which is frequently only noticed several generations' later, affecting sustainability of aquatic organisms' populations [35].

This review presents an updated survey of the acquired knowledge regarding the sources, spreading conditions, occurrence and induced toxic effects on non-target organisms by drugs in the environment. Fig. 1 illustrates the clear predominance of studies on non-steroidal anti-inflammatory drugs (NSAIDs), antibiotics and




Fig. 2. Schematic representation of pharmaceutical biotransformation to increase their polarity (adapted from Reference [35]).

blood lipid lowering agents from the literature, drawn from human prescription and consumption. Most of the reported data concerns the occurrence of drugs of each therapeutic class in the aquatic environment and is included in the form of tables to facilitate easy comparison between regional sample sources and ecotoxicological data. Current EU and US legislation compels new medicines to undergo an environmental impact assessment and consequently, new evaluation methods for acute as well as chronic effects are being implemented. However, a significant lack of knowledge persists particularly concerning toxicological data from synergistic pharmaceutical interactions.

## 2. Sources of environmental contamination

The most obvious pathway for environmental contamination of medicines is via the unaltered excretion in urine and faeces although other anthropogenic mechanisms should be assumed, namely:

- Metabolism post-consumption; since many drugs are metabolised as the organism attempts to convert hydrophobic compounds into more easily excreted polar residues. Their bio-conversion into one or more metabolites can occur throughout Phase I<sup>1</sup> and Phase II<sup>2</sup> reactions as shown in Fig. 2 [36].
- Diagnostic compounds; such as X-ray contrast media are directly discharged in their native forms.
- Household Disposal; either topic formulations or unused medicines (out-of-date or unwanted) are discarded through the sink/toilet or via waste collection [9,37,38], before being taken to

<sup>1</sup> Phase I reactions include oxidation, reduction and hydrolysis to modify the original molecule structure by introducing functional groups more receptive to phase II reactions.

<sup>2</sup> Phase II reactions (or conjugation reactions) consist of the addition of endogenous groups (like glucuronic acid, sulphate, glutathione, etc.) to receptive functional groups present in the original molecule or in its metabolite derived from phase I.




Fig. 3. Representative sources and fate of pharmaceuticals in the environment (adapted from Reference [6]).

landfill sites where they appear as terrestrial ecosystem contaminants. Alternatively, they may possibly leak into surrounding water compartments [39,40].

d) Impacts due to anthropogenic activities; as, for instance, Sewage Treatment Plant (STP) sludge, which can carry non-suspected drugs and is frequently used as a fertilizer on agricultural land [41,42]; veterinary medicines, which are also excreted in urine and faeces by animals before being spread onto land via manure application as fertilisers. Apart from the potential for direct soil contamination, there is also the risk of run-off with heavy rain, thus potentially contaminating both the surrounding surface and groundwater [42–44]. Other example of an anthropogenic activity is aquaculture, whose pharmaceuticals employed, as well as their metabolites and degradation products, are directly discharged into surface waters [45,46]. Another important source of environmental contamination by pharmaceuticals is the effluents of pharmaceutical production facilities [47–49].

At a higher level, existing geographical information on environmental contamination sources is sparse and limited. Countries and regions worldwide differ concerning the prevalence of diseases, waste treatment processes, cultural habits or economic constraints related to the pharmaceutical market [8]. Nevertheless, it seems that urban regions are major sources of contamination due to the proximity of hospitals and STP facilities. Additionally, the contribution of rural regions where agriculture, animal husbandry and aquaculture represent important ways of life should be considered as important.

### 3. Environmental fate


The fate and behaviour of medicines in the environment still requires further elucidation. As previously stated, drugs (used in human and/or in veterinary medicine) and their metabolites are spread into the environment in different ways, namely through STP effluents, heavy rain on agricultural land provokes (surface) water run-off, and occasionally, through untreated sewage (domestic wastes and flooding, among others) (Fig. 3). Some of them do reach surface waters (rivers, lakes and estuaries, among others) and eventually groundwaters [11,35,39] after resisting the intended biological degradation. However, in surface waters they may be degraded through different processes such as photolysis whose

efficiency depends on factors such as intensity of solar irradiation, latitude, season of the year and presence of photosensitizers (e.g. nitrates, humic acids) [50,51].

In the case of drugs that have low volatility and high polarity distribution is mainly made by aqueous transport or even via food chain dispersion [35,52]. Usually, wastewaters are conducted to STPs, which play a key role in the entrance of pharmaceuticals in the environment. However, in some regions or even countries these kinds of facilities may not exist and the environmental problem is still worse. The evaluation of removal efficiency in STPs (by comparing influent and effluent contents) has been studied in detail, showing removal rates that can differ by up to 99% [22,53–55]. Depending both on the particular technology resorted to and the active substance properties they may undergo: (i) degradation (mineralization) to low molecular weight compounds (e.g. CO<sub>2</sub> and water); (ii) entrapment by suspended solids; (iii) discharge of the parent compound through chemical cleavage of the respective conjugate forms and (iv) conversion to a more hydrophilic, persistent form which will short-circuit the treatment process [39,41,56,57]. Thus, in hospitals use of specific antibiotics, antineoplastic or diagnostic agents subsequently requires a sewage treatment process more embracing and directed to these kind of drugs, which are only used in hospitals [35,58], and that must be different to the more specific procedure adopted at STPs receiving industrial discharges from drug manufactures [47–49,59]. In both, the form and extension of the final contamination risk will also depend on geographical location of the STP facility. Low adsorption coefficients that make active substances remain in the aqueous phase, favour their mobility through the STP and into nearby surface waters [53]. Adsorption to suspended solids depending on both hydrophobic and electrostatic interactions established between each will follow the same destiny [11,41]. On the other hand, hydrophobic metabolites will be held on STP sludge, provoking terrestrial contamination, thus affecting microorganisms and invertebrates. Aerobic/anaerobic biodegradation occurring either during sewage sludge digestion or during activated sludge treatment seems to be the most efficient process to eliminate chemical contaminants from the aquatic environment. Usually, the best biodegradation results are obtained when activated sludge treatment is conducted through an increase in hydraulic retention time and the use of mature sludge [10]. However, one should be aware of the fact that if a particular pharmaceutical is not detected in a STP effluent, this does not imply that it has been fully removed. On some occasions, it may have been degraded and give rise to unsuspecting metabolites that will subsequently contaminate surface waters [35,39,60]. Notwithstanding that some drugs and their metabolites show a stable nature, nowadays is still difficult to establish a complete contamination pattern in final receiving surface waters, due to the water dilution, the treatment and discharging processes [54].

### 4. Ecotoxicology

Continuous consumption of drugs even at sub-therapeutic concentrations represents a potential threat to public health although one should bear in mind that it is still impossible to evaluate the effects of exposure on human health [35,60,61]. In turn, many non-target organisms (which possess human- and animal-like metabolic pathways, similar receptors or biomolecules) are therefore inadvertently exposed to active substances released into the environment [10,35]. A comprehensive manner to evaluate the toxicity effects on non-target organisms must include the development of specific tests embracing either acute effects (where mortality rates are often registered) or chronic effects (by means of exposure to different concentrations of a chemical compound over a prolonged period of time). In the latter, effects are measured




**Fig. 4.** (a) Acute vs. chronic ecotoxicological studies. (b) Principal endpoints used in ecotoxicological studies, expressed in relative percentage (data collected from 94 articles published between 1996 and 2009).

through specific parameters such as growth index or reproduction rates [52]. Unfortunately, studies on acute effects in organisms belonging to different trophic levels (i.e. algae, zooplankton and other invertebrates and fish) predominate relatively to chronic ones (Fig. 4). Acute toxicity data is only valuable when accidental discharge of the drugs occurs, since the environmental concentrations usually reported for these compounds are low, typically in a factor of one thousand. Bioaccumulation and chronic toxicity tests are scarce [10,35] probably due to the complex experimental work involved. However, recent development of sensitive methods for identification and quantification of drugs enabled to devise their distribution patterns in several environmental samples, thus highlighting the more relevant therapeutic classes in terms of environmental contamination (Fig. 5). These data is useful to set out the most appropriate active substances to be used in ecotoxicity tests. According to data present in literature, scientific community has mainly concerned their attention on therapeutic classes such as, non-steroidal anti-inflammatory drugs, blood lipid lowering agents, antibiotics and sex hormones. By those reasons, this review will focus in the drugs belonging to those therapeutic classes.

Within this context, some of the acute and chronic toxicity effects caused by drugs belonging to different therapeutic classes and mixtures of them in non-targets organisms deserve further analysis and are discussed in the following section. For a critical analysis of the ecotoxicological data present in the literature relatively to different drugs, we decide to group them according to their main pharmacological activity. Therefore, toxicity data will be related to the environmental concentrations found by several authors, to establish the severity of the situation.

#### 4.1. Non-steroidal anti-inflammatory drugs

Non-steroidal anti-inflammatory drugs are weak acids acting by reversible or irreversible inhibition of one or both isoforms of the cyclooxygenase enzymes, COX-1 and COX-2, involved in the synthesis of different prostaglandins from arachidonic acid [62]. A cyclooxygenase enzyme similar to human COX-2 has been found in fish thereby making them a potential target for aquatic contamination [63]. Prostaglandins also play an important role in the synthesis of bird eggshells and from inhibiting its synthesis, shell thinning has been observed [64]. Among the NSAID, diclofenac showed the most acute toxic nature with effects being observed at concentrations below  $100 \text{ mg L}^{-1}$  [65]. Chronic toxicity trials performed on rainbow trout (*Oncorhynchus mykiss*) evidenced cytological changes in the liver, kidneys and gills after 28 days of exposure to just  $1 \mu\text{g L}^{-1}$  of diclofenac. For a concentration of  $5 \mu\text{g L}^{-1}$  renal lesions were evident as well as drug bioaccumulation in the liver, kidneys, gills and muscle [66,67]. Brown trout (*Salmo trutta f. fario*) showed similar cytological damage and a reduction of haematocrit values after 21 days of exposure to  $0.5 \text{ mg L}^{-1}$  of this active substance [68]. Schmitt-Jansen et al. [69] evaluated both diclofenac phytotoxicity and its photochemical products on the unicellular chlorophyte *Scenedesmus vacuolatus*. Inhibition of algal reproduction by the parent compound only occurred at a concentration of  $23 \text{ mg L}^{-1}$ , hence indicating no specific toxicity. However, the threat significantly increased when metabolites were produced from 53 h of exposure to daylight. Diclofenac also inhibited the growth of marine phytoplankton *Dunaliella tertiolecta* for concentrations of  $25 \text{ mg L}^{-1}$  and above [70]. For this organism, 96 h  $\text{EC}_{50}$  of



**Fig. 5.** Therapeutic classes detected in the environment, expressed in relative percentage. Data collected from 134 articles published between 1997 and 2009.

**Table 1**  
Examples of concentrations (ng L<sup>-1</sup>) of non-steroidal anti-inflammatory drugs measured in different aquatic environments.

Compound	CAS number	Sample	Country	Analytical procedure	LOD (ng L <sup>-1</sup> )	Concentration reported (ng L <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
Acetylsalicylic acid	50-78-2	Somes river water	Romania	SPE-GC-MS	30 (LOQ)	<30–37.2 (±4.6)	[20]	Algae	<i>D. subspicatus</i>	EC <sub>50</sub> (growth inhibition)	106.7 mg L <sup>-1</sup>	[95]
Acetylsalicylic acid		STP influent	Japan	SPE-GC-MS	10 (LOQ)	470–19,400	[86]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48h) (immobilization)	88.1 mg L <sup>-1</sup>	[95]
Salicylic acid	69-72-7	STP effluent	Canada	SPE-GC-MS/MS	0.1	38.0–111	[17]	Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (30 min)	90 mg L <sup>-1</sup>	[83]
		STP effluent				554.3–2178.2						
		River water				130.4–371.5						
		Lake water				286.7						
Salicylic acid		STP influent	Canada	SPE-GC-MS	10	2820–12,700	[18]	Algae	<i>Scenedesmus subspicatus</i>	EC <sub>50</sub> (72h)	>100 mg L <sup>-1</sup>	[83]
		STP effluent				10–320		Crustacean	<i>D. magna</i>	EC <sub>50</sub> (24h) (immobilization)	118 mg L <sup>-1</sup>	[83]
								Ciliates	<i>Tetrahymena pyriformis</i>	EC <sub>50</sub> (48 h) (growth inhibition)	>100 mg L <sup>-1</sup>	[83]
								Fish	<i>B. rerio</i> (zebra fish)	LC <sub>50</sub> (48 h)	37 mg L <sup>-1</sup>	[83]
Diclofenac	15307-79-6	STP influent	Spain	SPE-GC-MS	100	200–3600	[14]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48h) (immobilization)	68 mg L <sup>-1</sup>	[65]
		STP effluent				140–2200						
Diclofenac		STP influent	Switzerland	SPE-GC-MS	6	1300–2900	[15]	Algae	<i>D. subspicatus</i>	EC <sub>50</sub> (growth inhibition)	72 mg L <sup>-1</sup>	[65]
		STP effluent				1300–2400						
Diclofenac		STP effluent	Canada	SPE-GC-MS/MS	1.0	32–448	[17]	Duckweed	<i>L. minor</i>	EC <sub>50</sub> (7 d) (growth inhibition)	7.5 mg L <sup>-1</sup>	[65]
Diclofenac		STP influent	Canada	SPE-GC-MS	10	50–2450	[18]	Fish	<i>Oncorhynchus mykiss</i>	LOEC (28 days) (histopathological alterations)	5 μg L <sup>-1</sup>	[66]
		STP effluent				70–250						
Diclofenac		STP influent	Greece	SPE-GC-MS	1	12–560	[19]	Fish	<i>Oncorhynchus mykiss</i>	LOEC (28 days) (cytological alterations)	1 μg L <sup>-1</sup>	[67]
		STP effluent				10–365						
Diclofenac		STP influent	Sweden	SPE-GC-MS	— <sup>f</sup>	160	[21]	Fish	<i>Salmo trout f. fario</i>	NOEC (21 days) (histopathological alterations)	0.5 μg L <sup>-1</sup>	[68]
		STP effluent				120						
		Höje river water				10–120						
Diclofenac		Parafba do Sul river water	Brazil	SPE-GC-MS	10	20–60	[22]	Algae	<i>Dunaliella tertiolecta</i>	EC <sub>50</sub> (96 h) (growth inhibition)	185,690 μg L <sup>-1</sup>	[87]
		Drinking water										
		water										
Diclofenac		Groundwater	Germany	SPE-GC-MS	29	<10–50	[26]	Algae	<i>D. subspicatus</i>	EC <sub>50</sub> (growth inhibition)	71.9 mg L <sup>-1</sup>	[95]
						590		Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48h) (immobilization)	68.0 mg L <sup>-1</sup>	[95]
								Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (30 min)	11,454 μg L <sup>-1</sup>	[96]
Diclofenac		Drinking water	USA	SPE-LC-MS/MS	0.25	<0.25	[32]	Algae	<i>P. subcapitata</i>	NOEC (96h) (growth inhibition)	10,000 μg L <sup>-1</sup>	[96]
Diclofenac		Hospital effluent	Taiwan	SPE-HPLC-MS/MS	10	328	[47]			LOEC (96h) (growth inhibition)	20,000 μg L <sup>-1</sup>	[96]

Table 1 (Continued)

Compound	CAS number	Sample	Country	Analytical procedure	LOD( $\text{ngL}^{-1}$ )	Concentration reported ( $\text{ngL}^{-1}$ )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
		Pharmaceutical production facility effluent				53						
Diclofenac		STP influent	United Kingdom	SPE-HPLC-MS/MS	20	901–1036	[53]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48h) (immobilization)	22,430 $\mu\text{gL}^{-1}$	[96]
Diclofenac		STP effluent				261–598						
Diclofenac		STP influent	Spain	SPE-HPLC-MS/MS	7	21–148	[71]		<i>C. dubia</i>	EC <sub>50</sub> (48h) (immobilization)	22,704 $\mu\text{gL}^{-1}$	[96]
		STP effluent	Belgium			32–1420						
		River water	Germany			26–72						
		Drinking water	Slovenia			<7						
Diclofenac		Elber river water	Germany	SPE-GC-MS	0.08 (LOQ)	42–67	[72]			NOEC (7d) (reproduction)	1000 $\mu\text{gL}^{-1}$	[96]
Diclofenac		Alster lake water										
Diclofenac		Hospital effluent	Spain	SPE-HPLC-MS/MS	30	60–1900	[73]			LOEC (7d) (reproduction)	2000 $\mu\text{gL}^{-1}$	[96]
Diclofenac		STP influent	Taiwan	SPE-HPLC-MS/MS	— <sup>a</sup>	3–347	[87]	Fish	<i>D. rerio</i>	NOEC (10 d) (survival)	4000 $\mu\text{gL}^{-1}$	[96]
Diclofenac		STP effluent				4–101						
Diclofenac		Pearl Rivers water	China	GC-NCI-MS	1.1	ND–147 ( $\pm 5$ )	[88]			LOEC (10 d) (survival)	8000 $\mu\text{gL}^{-1}$	[96]
Diclofenac		STP influent	Luxembourg	SPE-LC-MS/MS	0.3	2–43	[89]	Fish	<i>O. mykiss</i>	LOEC (21 d) (liver cytopathology)	1 $\mu\text{gL}^{-1}$	[97]
		STP effluent				0.3–78						
		Alzette river water				0.3–55						
		Mess river water				0.3–19						
Diclofenac		STP effluent	South Korea	SPE-LC-MS/MS	1.0	8.8–127	[90]			LOEC (21 d) (kidney cytopathology)	1 $\mu\text{gL}^{-1}$	[97]
Diclofenac		Surface water				1.1–6.8						
Diclofenac		STP effluent	Spain	SPE-LC-QqLIT-MS	4 (LOQ)	890–1440	[91]			LOEC (21 d) (gills cytopathology)	1 $\mu\text{gL}^{-1}$	[97]
Diclofenac		STP effluent	United Kingdom	SPE-HPLC-MS/MS	20	350–460	[94]					
Fenoprofen	53746-45-5	Surface water				<20–91						
		STP influent	Japan	SPE-GC-MS	1 (LOQ)	9.68–80.6	[86]					
		STP effluent				1.59–9.22						
Ibuprofen	15687-27-1	STP influent	Spain	SPE-GC-MS	23	34,000–168,000	[14]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48h) (immobilization)	108 $\text{mgL}^{-1}$	[65]
Ibuprofen		STP effluent				240–28,000						
Ibuprofen		STP influent	Switzerland	SPE-GC-MS	8	1750–4500	[15]	Algae	<i>D. subspicatus</i>	EC <sub>50</sub> (growth inhibition)	315 $\text{mgL}^{-1}$	[65]
Ibuprofen		STP effluent				100–1200						
Ibuprofen		STP effluent	Canada	SPE-GC-MS	0.8	2235.2–6718.3	[17]	Duckweed	<i>L. minor</i>	EC <sub>50</sub> (7 d) (growth inhibition)	22 $\text{mgL}^{-1}$	[65]
Ibuprofen		STP influent	Canada	SPE-GC-MS	10	4100–10,210	[18]	Crustacean	<i>Daphnia magna</i>	EC <sub>50</sub> (48h) (immobilisation)	10–100 $\text{mgL}^{-1}$	[75]
Ibuprofen		STP effluent				110–2170						
Ibuprofen		Somes river water	Romania	SPE-GC-MS	30 (LOQ)	<30–115.2	[20]			EC <sub>50</sub> (14 d) (reproduction)	13.4 $\text{mgL}^{-1}$	[75]

Ibuprofen	STP influent	Sweden	SPE-GC-MS	— $\ddagger$	( $\pm 20.7$ ) 3590	[21]			NOEC (14 d) (survival)	20 mg L <sup>-1</sup>	[75]
	STP effluent Höje river water				150 10–220				LOEC (14 d) (survival)	80 mg L <sup>-1</sup>	
Ibuprofen	Paraíba do Sul river water Drinking water	Brazil	SPE-GC-MS	10	<10	[22]			LOEC (14 d) (population growth)	20 mg L <sup>-1</sup>	[75]
Ibuprofen	Po river water	Italy	SPE- HPLC-MS/MS	4.2 (LOQ)	<10	[24]	Crustacean	<i>Gammarus pulex</i>	LOEC (behaviour)	10 ng L <sup>-1</sup>	[76]
	Lambro river water				ND–9.76 78.50						
Ibuprofen	Groundwater	USA	SPE-LC-MS	18	3110	[28]	Duckweed	<i>Lemna minor</i>	EC <sub>50</sub> (7 d) (growth inhibition)	4.01 mg L <sup>-1</sup>	[77]
Ibuprofen	Hospital effluent	Taiwan	SPE- HPLC-MS/MS	25	119	[47]	Crustacean	<i>T. platyurus</i>	LC <sub>50</sub> (24 h) (mortality)	19.59 mg L <sup>-1</sup>	[78]
	Pharmaceutical production facility effluent				45,875						
Ibuprofen	STP influent	United Kingdom	SPE- HPLC-MS/MS	20	7741–33,764	[53]	Fish	<i>O. latipes</i>	LC <sub>50</sub> (96 h) (mortality)	>100 mg L <sup>-1</sup>	[79]
	STP effluent Tyne river water				1979–4239 144–2370						
Ibuprofen	STP influent	Spain	SPE- HPLC-MS/MS	12	37–860	[71]	Mollusc	<i>P. carinatus</i>	LC <sub>50</sub> (72 h) (survival)	17.1 mg L <sup>-1</sup>	[79]
	STP effluent River water	Belgium			18–1860						
	Drinking water	Germany Slovenia				60–152 <12					
Ibuprofen	Elber river water	Germany	SPE-GC-MS	0.05 (LOQ)	8.7–32	[72]			NOEC (21 d) (survival)	5.36 mg L <sup>-1</sup>	[79]
	Alster lake water				4.9						
Ibuprofen	Hospital effluent	Spain	SPE- HPLC-MS/MS	31	1500–151,000	[73]			NOEC (21 d) (growth)	1.02 mg L <sup>-1</sup>	[79]
Ibuprofen	STP effluent	USA	SPE-GC-MS	10	18 ( $\pm 14\%$ )	[81]			LOEC (21 d) (growth)	2.43 mg L <sup>-1</sup>	[79]
Ibuprofen	STP influent	Japan	SPE-GC-MS	1 (LOQ)	407–1130	[86]			NOEC (21 d) (reproduction)	2.43 mg L <sup>-1</sup>	[79]
	STP effluent STP influent				Taiwan		SPE-HPLC- MS/MS	— $\ddagger$	1.41–177 711–17,933	[87]	Algae
Ibuprofen	STP effluent STP influent	Luxembourg	SPE-LC-MS/MS	0.3	313–3777 82–3080	[89]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48h) (immobilization)	101.2 mg L <sup>-1</sup>	[95]
	STP effluent Alzette river water				3–359 10–295						
	Mess river water				9–2383						
Ibuprofen	STP effluent	South Korea	SPE-LC-MS/MS	1.0	10–137	[90]	Cnidarian	<i>Hydra attenuata</i>	LC <sub>50</sub> (96 h) (morphology)	22.36 mg L <sup>-1</sup>	[98]
	Surface water				11–38						

Table 1 (Continued)

Compound	CAS number	Sample	Country	Analytical procedure	LOD (ngL <sup>-1</sup> )	Concentration reported (ngL <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
		Drinking water				<1.0						
Ibuprofen		STP effluent	Spain	SPE-LC-QqLIT-MS	13 (LOQ)	100–340	[91]			EC <sub>50</sub> (96h) (morphology)	1.65 mgL <sup>-1</sup>	[98]
Ibuprofen		Mankyung river water	South Korea	SPE-LC-MS/MS	5	<5–414 (±13)	[92]			LOEC (96h) (morphology)	1 mgL <sup>-1</sup>	[98]
Ibuprofen		STP effluent	United Kingdom	SPE-HPLC-MS/MS	20	1700–3800	[94]			NOEC (96h) (morphology)	0.1 mgL <sup>-1</sup>	[98]
		Surface water				<20				EC <sub>50</sub> (96 h) (feeding)	3.85 mgL <sup>-1</sup>	[98]
Carboxy-ibuprofen*	— <sup>†</sup>	STP influent STP effluent Höje river water	Sweden	SPE-GC-MS	— <sup>†</sup>	10,750 430 230–680	[21]					
Carboxy-ibuprofen*		Elber river water Alster lake water	Germany	SPE-GC-MS	0.21	11–32 9.5	[72]					
Hydroxy-ibuprofen*	— <sup>†</sup>	STP influent STP effluent Höje river water	Sweden	SPE-GC-MS	— <sup>†</sup>	990 50 20–60	[21]					
Hydroxy-ibuprofen*		Elber river water Alster lake water	Germany	SPE-GC-MS	0.38	32–101 18	[72]					
Indomethacin	53-86-1	STP influent STP effluent	Canada	SPE-GC-MS	10	30–430 40–490	[18]					
Indomethacin		STP effluent	Spain	SPE-LC-QqLIT-MS	8 (LOQ)	160–390	[91]	Crustacean	<i>T. platyurus</i>	LC <sub>50</sub> (24 h) (mortality)	16.14 mgL <sup>-1</sup>	[78]
Indomethacin		Mankyung river water	South Korea	SPE-LC-MS/MS	1	<1–33.5 (±8)	[92]	Fish	<i>O. latipes</i>	LC <sub>50</sub> (96 h) (mortality)	81.92 mgL <sup>-1</sup>	[78]
Ketoprofen	22071-15-4	STP effluent	Canada	SPE-GC-MS/MS	1.0	8–351	[17]					
Ketoprofen		STP influent STP effluent	Canada	SPE-GC-MS	10	60–150 40–90	[18]					
Ketoprofen		STP influent STP effluent Höje river water	Sweden	SPE-GC-MS	— <sup>†</sup>	940 330 10–70	[21]					
Ketoprofen		Hospital effluent Pharmaceutical production facility effluent	Taiwan	SPE-HPLC-MS/MS	10	9.6 ND	[47]					
Ketoprofen		STP influent	Spain	SPE-HPLC-MS/MS	26	131	[71]					
		STP effluent	Belgium			<26						
		River water	Germany			<26						
		Drinking water	Slovenia			<26						
Ketoprofen		STP effluent	USA	SPE-GC-MS	9	23 (±6.8%)	[81]					
Ketoprofen		STP influent STP effluent	Japan	SPE-GC-MS	0.3 (LOQ)	108–369 68.1–219	[86]					



Ketorolac	74103-06-3	Hospital effluent	Spain	SPE-HPLC-MS/MS	26	200–59,500	[73]					
Mefenamic acid	61-68-7	STP influent	United Kingdom	SPE-HPLC-MS/MS	50	136–363	[53]	Crustacean	<i>T. platyurus</i>	LC <sub>50</sub> (24 h) (mortality)	3.95 mgL <sup>-1</sup>	[78]
Mefenamic acid		STPeffluent	Japan	SPE-GC-MS	1 (LOQ)	290–396	[86]	Fish	<i>O. latipes</i>	LC <sub>50</sub> (96 h) (mortality)	8.04 mgL <sup>-1</sup>	[78]
Mefenamic acid		Pearl Rivers water	China	GC-NCI-MS	2.2	ND–22.4 (±3.1)	[88]					
Mefenamic acid		STPeffluent	Spain	SPE-LC-QqLIT-MS	3 (LOQ)	40–60	[91]					
Mefenamic acid		Mankyung river water	South Korea	SPE-LC-MS/MS	10	<10–326 (±21)	[92]					
Mefenamic acid		STPeffluent	United Kingdom	SPE-HPLC-MS/MS	50	720–1100	[94]					
Naproxen	22204-53-1	Surface water	Canada	SPE-GC-MS/MS	0.5	<50–65	[17]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48h) (immobilization)	174 mgL <sup>-1</sup>	[65]
Naproxen		STP influent	Canada	SPE-GC-MS	10	1730–6030	[18]	Algae	<i>D. subspicatus</i>	EC <sub>50</sub> (growth inhibition)	>320 mgL <sup>-1</sup>	[65]
Naproxen		STPeffluent	Sweden	SPE-GC-MS	— <sup>f</sup>	360–2540	[21]	Duckweed	<i>L. minor</i>	EC <sub>50</sub> (7 d) (growth inhibition)	24.2 mgL <sup>-1</sup>	[65]
Naproxen		Höje river water				250						
Naproxen		Paraíba do Sul river water	Brazil	SPE-GC-MS	10	<10–50	[22]	Rotifers	<i>B. calyciflorus</i>	LC <sub>50</sub> (24 h)	62.48 mgL <sup>-1</sup>	[80]
Naproxen		Drinking water				90–250						
Naproxen		Drinking water	USA	SPE-LC-MS/MS	0.5	<10–30	[32]	Rotifers	<i>T. platyurus</i>	LC <sub>50</sub> (24 h)	84.09 mgL <sup>-1</sup>	[80]
Naproxen		Hospital effluent	Taiwan	SPE-HPLC-MS/MS	10	698	[47]	Crustaceans	<i>C. dubia</i>	EC <sub>50</sub> (24h) (immobilization)	66.37 mgL <sup>-1</sup>	[80]
Naproxen		Pharmaceutical production facility effluent				ND						
Naproxen		STP influent	Spain	SPE-HPLC-MS/MS	26	109–455	[71]	Algae	<i>P. subcapitata</i>	EC <sub>50</sub> (72 h) (growth inhibition)	31.82 mgL <sup>-1</sup>	[80]
Naproxen		STPeffluent	Belgium			625						
Naproxen		River water	Germany			70						
Naproxen		Drinking water	Slovenia			<26						
Naproxen		STPeffluent	USA	SPE-GC-MS	9	31 (±5.5%)	[81]	Rotifers	<i>B. calyciflorus</i>	EC <sub>50</sub> (48 h) (growth inhibition)	0.56 mgL <sup>-1</sup>	[80]
Naproxen		STP influent	Japan	SPE-GC-MS	0.3 (LOQ)	38.0–230	[86]	Crustaceans	<i>C. dubia</i>	EC <sub>50</sub> (7 d) (population growth inhibition)	0.33 mgL <sup>-1</sup>	[80]
Naproxen		STPeffluent	China	GC-NCI-MS	1.3	12.0–139	[88]	Algae	<i>D. subspicatus</i>	EC <sub>50</sub> (growth inhibition)	625.5 mgL <sup>-1</sup>	[95]
Naproxen		Pearl Rivers water				ND–118 (±10.1)						
Naproxen		STPeffluent	South Korea	SPE-LC-MS/MS	1.0	20–483	[90]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48h) (immobilization)	166.3 mgL <sup>-1</sup>	[95]
Naproxen		Surface water				1.8–18		Cnidarian	<i>Hydra attenuata</i>	LC <sub>50</sub> (96 h) (morphology)	22.36 mgL <sup>-1</sup>	[98]
Naproxen										EC <sub>50</sub> (96h) (morphology)	2.62 mgL <sup>-1</sup>	[98]

Table 1 (Continued)

Compound	CAS number	Sample	Country	Analytical procedure	LOD (ngL <sup>-1</sup> )	Concentration reported (ngL <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
										LOEC (96 h) (morphology)	5 mgL <sup>-1</sup>	[98]
										NOEC (96 h) (morphology)	1 mgL <sup>-1</sup>	[98]
Paracetamol	103-90-2	STP influent	Spain	SPE-GC-MS	32	29,000–246,000	[14]	Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (96 h) (feeding)	2.68 mgL <sup>-1</sup>	[98]
		STP effluent				<32–4300				EC <sub>50</sub> (15 min)	567.5 mgL <sup>-1</sup>	[82]
Paracetamol		Groundwater	USA	SPE-LC-MS	9	380	[28]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48 h) (immobilization)	30.1 mgL <sup>-1</sup>	[82]
Paracetamol		Hospital effluent	Taiwan	SPE-HPLC-MS/MS	2	62,250	[47]		<i>D. magna</i>	EC <sub>50</sub> (96 h) (immobilization)	26.6 mgL <sup>-1</sup>	[82]
		Pharmaceutical production facility effluent				124						
Paracetamol		STP influent	United Kingdom	SPE-HPLC-MS/MS	20	5529–69,570	[53]	Fish	<i>O. latipes</i>	LC <sub>50</sub> (48 h)	>160 mgL <sup>-1</sup>	[82]
Paracetamol		STP effluent				<20			<i>O. latipes</i>	LC <sub>50</sub> (96 h)	>160 mgL <sup>-1</sup>	[82]
Paracetamol		Hospital effluent	Spain	SPE-HPLC-MS/MS	47	500–29,000	[73]					
Paracetamol		Danube river water	Serbia	SPE-LC-MS/MS	0.50	78,170	[84]	Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (30 min)	650 mgL <sup>-1</sup>	[83]
		Sava river water				610						
		Tamis' river water				310						
Paracetamol		STP effluent	South Korea	SPE-LC-MS/MS	1.0	1.8–19	[90]	Algae	<i>Scenedesmus subspicatus</i>	EC <sub>50</sub> (72 h)	134 mgL <sup>-1</sup>	[83]
Paracetamol		Surface water				4.1–73						
Paracetamol		STP influent	Korea	SPE-LC-MS	5	13,046–56,944	[93]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (immobilization)	50 mgL <sup>-1</sup>	[83]
		STP effluent				<5–9						
		Han river water				<5–127						
Paracetamol		STP effluent	United Kingdom	SPE-HPLC-MS/MS	50	<50	[94]	Ciliates	<i>Tetrahymena pyriformis</i>	EC <sub>50</sub> (48 h) (growth inhibition)	112 mgL <sup>-1</sup>	[83]
		Surface water				<50		Fish	<i>B. rerio</i> (zebra fish)	LC <sub>50</sub> (48 h)	378 mgL <sup>-1</sup>	[83]

\*—Metabolite; †—Data not available; ND—Not detected; SPE—Solid Phase Extraction; GC-MS—Gas Chromatography with Mass Spectrometry Detection; GC-MS/MS—Gas Chromatography with Tandem Mass Spectrometry Detection; GC-NCI-MS—Gas Chromatography-Negative Chemical Ionization-Mass Spectrometry; HPLC-MS/MS—High Performance Liquid Chromatography with Tandem Mass Spectrometry Detection; LC-QqLIT-MS—Liquid chromatography-quadrupole-linear ion trap-mass spectrometry detection.

185.69 mg L<sup>-1</sup> was found [70]. Diclofenac was detected in STP effluents at maximum concentrations of 2.4 [15] and 1.42 µg L<sup>-1</sup> [71] in Switzerland and Belgium respectively (Table 1) which highlighted that the effects cited are of sufficient magnitude to suspect chronic toxicity in aquatic organisms. Diclofenac has also been found in rivers [21,22,72], groundwater [26], hospital effluents [47,73] and drinking water [22,32,71] but at concentrations in the order of ng L<sup>-1</sup>.

Ibuprofen is another NSAID with documented chronic toxicity. Female Japanese medaka (the Japanese killifish, *Oryzias latipes*) exposed to different concentrations of the drug over six weeks, showed a sharp rise in liver weight together with enhanced egg production, yet with a reduction in the number of weekly spawning events [74]. Authors associated these phenomena with changes in the spawning process and vitellogenin production, a glycoprotein precursor in yolk formation. With the water flea *Daphnia magna* population growth rate was significantly reduced for concentrations ranging from 0 to 80 mg L<sup>-1</sup> [75]. Reproduction was affected at all concentrations and completely inhibited at the highest pharmaceutical levels. An activity decrease of the freshwater amphipod *Gammarus pulex* was noticed when in contact with ibuprofen concentrations of 1 and 10 ng L<sup>-1</sup>, the latter value corresponding to the LOEC<sup>3</sup> obtained for behaviour change [76]. Regarding aquatic photosynthetic organisms, specific effects have been noticed. A 5-day exposure to concentrations in the 1–1000 µg L<sup>-1</sup> range stimulated the growth of the cyanobacterium *Synechocystis* sp. while inhibiting that of the duckweed plant *Lemna minor* after 7 days [77]. Ibuprofen has been detected in STP effluents at concentrations that can reach 28 µg L<sup>-1</sup> [14] (Spain) (Table 1). Two metabolites of ibuprofen (carboxyl-ibuprofen and hydroxyl-ibuprofen) were also found in surface waters and in a Swedish STP (influent and effluent) [21,72]. Due to demonstrable chronic toxicity, this may represent a real threat to non-target organisms, even at those lower concentrations. Ibuprofen was also found in rivers [20–22,24,72] and drinking water [22] which may broaden the scope of the problem to public health. However, effects in humans caused by chronic exposure to this active substance still remain unknown.

The ecotoxicity of naproxen and its photoderivative products have also been envisaged. Acute toxicity tests performed on the rotifer *Brachionus calyciflorus*, the water flea *Ceriodaphnia dubia* and the fairy shrimp *Thamnocephalus platyurus*, showed that naproxen had LC<sub>50</sub><sup>4</sup> and EC<sub>50</sub><sup>5</sup> values within the 1–100 mg L<sup>-1</sup> range, with the photolysis products being significantly more toxic [80]. Highly chronic toxic properties were equally noticed with algae being the less sensitive organisms. Yet again, degradation products were shown to be more toxic with EC<sub>50</sub> values of 26 and 62 µg L<sup>-1</sup> for *C. dubia*, relative to growth inhibition. Naproxen had been found in STP effluents in a concentration range between 31 ng L<sup>-1</sup> [81] and 7.96 µg L<sup>-1</sup> [17] and in surface waters [21,22,71], at concentration levels that can reach 250 ng L<sup>-1</sup> [21]. This active substance was also detected in drinking water [22,32,71].

The highly prescribed paracetamol (or acetaminophen) is a weak inhibitor of the cyclooxygenase enzyme, whose side effects are mainly associated with the formation of hepatotoxic metabolites, such as *N*-acetyl-*p*-benzoquinone imine (NAPQI) when the levels of liver glutathione are low [36]. Tests were carried out on algae, water fleas, fish embryos, luminescent bacteria and ciliates. The most sensitive species was shown to be *D. magna* for which EC<sub>50</sub> values of 30.1 [82] or 50 mg L<sup>-1</sup> [83] were reported. Some authors reported the presence of paracetamol in STP effluents at concentrations below to 20 ng L<sup>-1</sup> [53] to 4.3 µg L<sup>-1</sup> [14], and in surface

waters, values can reach 78.17 µg L<sup>-1</sup> [84] (Table 1), which are values higher than the predicted no-effect concentration (PNEC) of 9.2 µg L<sup>-1</sup> [85]. Hence, paracetamol might represent a threat for non-target organisms.

#### 4.2. Blood lipid lowering agents

Modulating drugs for lipid metabolism are frequently prescribed in the developed world and aim to decrease the concentration of blood circulating cholesterol and triglycerides. Pharmaceuticals belonging to this therapeutic class can be divided into two main groups: statins and the group most frequently detected in the environment, fibrates [99]. Statins act by inhibiting the 3-hydroxymethylglutaryl coenzyme A reductase (HMG-CoA), an enzyme involved in feedback control of cholesterol synthesis. In response, the number of LDL lipoprotein receptors at hepatocyte surfaces increases, thus lowering the circulating LDL cholesterol [100]. Toxicity data of statins on different organisms is very limited and restricted to the active substances simvastatin and atorvastatin. After an exposure of 96 h to simvastatin, larval and adult grass shrimp (*Palaemonetes pugio*) showed a LC<sub>50</sub> of 1.18 mg L<sup>-1</sup> and upper 10 mg L<sup>-1</sup>, respectively [101], while the harpacticoid copepod *Nitocra spinipes* had a 96-h LC<sub>50</sub> of 0.81 mg L<sup>-1</sup> [102]. Dahl et al. (2006) [102] also reported a significant increase in development time and body length of the copepod for a range of concentrations between 0.16 and 1.6 µg L<sup>-1</sup>. On the other hand, simvastatin exhibited an EC<sub>50</sub> of 22.8 mg L<sup>-1</sup>, after 96 h, for the marine phytoplankton *D. tertiolecta* [70]. Relatively to atorvastatin, this active substance can affect the development of the duckweed *Lemna gibba*, showing a LOEC of 300 µg L<sup>-1</sup> for parameters such as wet mass, frond number, chlorophyll-*a* and carotenoids content, for a time of exposure of 7 days [103]. Apart from statins had also the ability to suppress synthesis of the juvenile hormone in insects [104]. Statins were found in untreated sewage samples (Table 2) at concentrations between 4 and 117 ng L<sup>-1</sup> and in treated sewage samples at 1–59 ng L<sup>-1</sup> [105,106]. Additionally, they were also detected in surface water [105] and drinking water [32] at concentrations that can reach 1 ng L<sup>-1</sup>. In turn, fibrates act by activating specific transcription factors belonging to the nuclear hormone receptor super family, known as peroxisome proliferator-activated receptors (PPARs) [107]. There are three types of PPARs related to different cellular events. PPAR- $\alpha$  and PPAR- $\beta$  play key roles in catabolism and storage of fatty acids while PPAR- $\gamma$  plays an important role in cellular differentiation [108]. Some authors have reported a proliferation of peroxisomes in rodent livers caused by fibrates [10]. Embryonic development of non-target organisms that share these receptors can be stopped by simply inhibiting cellular differentiation. Fibrates present in the micromolar concentration range are sufficient to cause it in zebrafish (*Danio rerio*) [109,110] and amphibians [111]. Raldúa et al. [110] demonstrated that, when exposed to 0.5–1 mg L<sup>-1</sup> of clofibrate, zebrafish larvae had a significantly shorter body length and their morphologic characteristics were also altered. Clofibrate-exposed zebrafish larvae had also lethargic behaviour. It was evidenced that gemfibrozil and bezafibrate significantly affect feeding, attachment and hydrant growth of the cnidarian *Hydra attenuata* [98]. According to Quinn et al. [98], gemfibrozil could be classified as toxic (EC<sub>50</sub> between 1 and 10 mg L<sup>-1</sup>) and bezafibrate as harmful for non-target organisms (EC<sub>50</sub> between 10 and 100 mg L<sup>-1</sup>). Toxic properties of gemfibrozil were also respectively investigated on the inhibition of the bacterium *Vibrio fischeri* luminescence, growth inhibition of the alga *Chlorella vulgaris* and on the immobilization of the *D. magna*. In this study both the bacteria and the water flea were shown to be sensitive to gemfibrozil with the latter being the most sensitive, having an EC<sub>50</sub> of 30 mg L<sup>-1</sup> after 72 h [112]. Proliferative inhibition

<sup>3</sup> LOEC—Lowest Observed Effect Concentration.

<sup>4</sup> LC<sub>50</sub>—Half Maximal Lethal Concentration.

<sup>5</sup> EC<sub>50</sub>—Half Maximal Effective Concentration.

**Table 2**Examples of concentrations (ng L<sup>-1</sup>) of blood lipid lowering agents measured in different aquatic environments.

Compound	CAS number	Sample	Country	Analytical procedure	LOD(ngL <sup>-1</sup> )	Concentration reported(ngL <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
<i>Fibrates</i> Bezafibrate	41859-67-0	Paraíba do Sul river water Drinking water	Brazil	SPE-GC-MS	25	<25	[22]	Cnidarian	<i>Hydra attenuata</i>	LC <sub>50</sub> (96 h) (morphology)	70.71 mg L <sup>-1</sup>	[98]
Bezafibrate		Po river water Lambro river water	Italy	SPE-HPLC-MS/MS	0.3	<25 0.79–2.75 57.15	[24]			EC <sub>50</sub> (96h) (morphology)	25.85 mg L <sup>-1</sup>	[98]
Bezafibrate		STP effluent	Spain	SPE-LC-QqLIT-MS	3 (LOQ)	40–130	[91]			LOEC (96h) (morphology)	1 mg L <sup>-1</sup>	[98]
Bezafibrate		STP effluent	Italy	SPE-HPLC-MS/MS	0.1 (LOQ)	0.3–117	[118]	Rotifer	<i>B. calyciflorus</i>	NOEC (96 h) (morphology) EC <sub>50</sub> (96 h) (feeding) LC <sub>50</sub> (24 h) (mortality) EC <sub>50</sub> (48h) (population growth inhibition)	0.1 mg L <sup>-1</sup> 8.59 mg L <sup>-1</sup> 60.91 mg L <sup>-1</sup> 0.44 mg L <sup>-1</sup>	[98] [98] [113] [113]
								Crustacean	<i>T. platyurus</i> <i>D. magna</i> <i>C. dubia</i>	NOEC (48 h) LOEC (48 h) LC <sub>50</sub> (24 h) (mortality) EC <sub>50</sub> (24h) (immobilization) EC <sub>50</sub> (48h) (immobilization)	0.156 mg L <sup>-1</sup> 0.3125 mg L <sup>-1</sup> 39.69 mg L <sup>-1</sup> 100.08 mg L <sup>-1</sup> 75.79 mg L <sup>-1</sup>	[113] [113] [113] [113] [113]
Clofibrate	82115-62-6							Fish	<i>D. rerio</i>	EC <sub>50</sub> (7 d) (population growth inhibition) NOEC (7 d) LOEC (7 d) LC <sub>50</sub> (96 h) (mortality)	0.13 mg L <sup>-1</sup> 0.023 mg L <sup>-1</sup> 0.047 mg L <sup>-1</sup> 0.89 mg L <sup>-1</sup>	[113] [113] [113] [110]
Fenofibrate	49562-28-9							Rotifer	<i>B. calyciflorus</i>	LC <sub>50</sub> (24 h) (mortality) EC <sub>50</sub> (48h) (population growth inhibition)	64.97 mg L <sup>-1</sup> 1.44 mg L <sup>-1</sup>	[113] [113]
								Crustacean	<i>D. magna</i> <i>C. dubia</i>	NOEC (48 h) LOEC (48 h) EC <sub>50</sub> (24h) (immobilization) EC <sub>50</sub> (7 d) (population growth inhibition)	0.156 mg L <sup>-1</sup> 0.3125 mg L <sup>-1</sup> 50.12 mg L <sup>-1</sup> 0.76 mg L <sup>-1</sup>	[113] [113] [113] [113]
								Algae	<i>P. subcapitata</i>	NOEC (7 d) LOEC (7 d) EC <sub>50</sub> (72 h) (growth inhibition)	0.039 mg L <sup>-1</sup> 0.078 mg L <sup>-1</sup> 19.84 mg L <sup>-1</sup>	[113] [113] [113]
								Crustacean	<i>D. magna</i>	NOEC (72h) LOEC (72h) EC <sub>50</sub> (48h) (immobilization)	3.12 mg L <sup>-1</sup> 6.25 mg L <sup>-1</sup> 72 mg L <sup>-1</sup>	[113] [113] [65]
Clofibrac acid*	882-09-7	STP influent	Greece	SPE-GC-MS	1.8	ND	[19]	Crustacean	<i>D. magna</i>			

Clofibric acid*		STPeffluent Paraiba do Sul river water Drinking water	Brazil	SPE-GC-MS	10	5 <10–30	[22]	Algae	<i>D. subspicatus</i>	EC <sub>50</sub> (growth inhibition)	115 mg L <sup>-1</sup>	[65]
Clofibric acid*		Po river water Lambro river water	Italy	SPE- HPLC-MS/MS	0.3	<10–20 0.41–5.77 ND	[24]	Duckweed	<i>L. minor</i>	EC <sub>50</sub> (7 d) (growth inhibition)	12.5 mg L <sup>-1</sup>	[65]
Clofibric acid*		North Sea water	— <sup>†</sup>	SPE-GC-MS	0.008	ND–18.6	[25]	Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (30 min)	100 mg L <sup>-1</sup>	[83]
Clofibric acid*		STP influent	United Kingdom	SPE- HPLC-MS/MS	20	<20–651	[53]	Algae	<i>Scenedesmus subspicatus</i>	EC <sub>50</sub> (72h)	89 mg L <sup>-1</sup>	[83]
Clofibric acid*		STPeffluent STP influent	Spain	SPE- HPLC-MS/MS	17	<20–44 25–58	[71]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (immobilization)	106 mg L <sup>-1</sup>	[83]
Clofibric acid*		STPeffluent River water Drinking water Elbe river water Alster lake water	Belgium Germany Slovenia	SPE-GC-MS	0.26 (LOQ)	22–107 24–35 <17 3.2–7.6 2.4	[72]	Ciliates	<i>Tetrahymena pyriformis</i>	EC <sub>50</sub> (48 h) (growth inhibition)	175 mg L <sup>-1</sup>	[83]
Clofibric acid*		STP influent	Taiwan	SPE- HPLC- MS/MS	— <sup>†</sup>	36–2593	[87]	Fish	<i>D. rerio</i>	LC <sub>50</sub> (48 h)	86 mg L <sup>-1</sup>	[83]
Clofibric acid*		STPeffluent STPeffluent	Spain	SPE-LC-QqLIT- MS	4 (LOQ)	47–487 36–51	[91]	Algae	<i>Dunaliella tertiolecta</i>	EC <sub>50</sub> (96 h) (growth inhibition)	224,180 μg L <sup>-1</sup>	[87]
Clofibric acid*		STPeffluent	United Kingdom	SPE- HPLC-MS/MS	50	<50	[94]	Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (30 min)	91,827 μg L <sup>-1</sup>	[96]
Clofibric acid*		Surface water STPeffluent	Italy	SPE- HPLC-MS/MS	0.36 (LOQ)	<50 ND–82	[118]	Algae	<i>P. subcapitata</i>	NOEC (96h) (growth inhibition)	75,000 μg L <sup>-1</sup>	[96]
Clofibric acid*		Groundwater	Germany	SPE-GC-MS	2 (LOQ)	2–40	[119]	Crustacean	<i>D. magna</i> <i>C. dubia</i>	LOEC (96h) (growth inhibition) EC <sub>50</sub> (48h) (immobilization) EC <sub>50</sub> (48h) (immobilization)	150,000 μg L <sup>-1</sup> >200,000 μg L <sup>-1</sup> >200,000 μg L <sup>-1</sup>	[96] [96] [96]
										NOEC (7d) (reproduction) LOEC (7d) (reproduction)	640 μg L <sup>-1</sup> 2560 μg L <sup>-1</sup>	[96] [96]
								Fish	<i>D. rerio</i>	NOEC (10 d) (survival)	70,000 μg L <sup>-1</sup>	[96]
								Fish	<i>O. mykiss</i>	LOEC (10 d) (survival) LOEC (21 d) (liver cytopathology)	140,000 μg L <sup>-1</sup> >100 μg L <sup>-1</sup>	[96] [97]
										LOEC (21 d) (kidney cytopathology)	>100 μg L <sup>-1</sup>	[97]
										LOEC (21 d) (gills cytopathology)	5 μg L <sup>-1</sup>	[97]
Gemfibrozil	25812-30-0	STPeffluent  River water Lake water	Canada	SPE-GC-MS/MS	0.3	80.1–478.2 ND–18.4 ND	[17]	Cnidarian	<i>Hydra attenuata</i>	LC <sub>50</sub> (96 h) (morphology)	22.36 mg L <sup>-1</sup>	[98]

Table 2 (Continued)

Compound	CAS number	Sample	Country	Analytical procedure	LOD (ngL <sup>-1</sup> )	Concentration reported (ngL <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
Gemfibrozil		STP influent	Canada	SPE-GC-MS	10	120–36,530	[18]			EC <sub>50</sub> (96h) (morphology)	1.18 mgL <sup>-1</sup>	[98]
Gemfibrozil		STP effluent STP influent	Sweden	SPE-GC-MS	— <sup>a</sup>	80–2090 710	[21]			LOEC (96h) (morphology)	1 mgL <sup>-1</sup>	[98]
Gemfibrozil		STP effluent Höje river water				180 1–170						
Gemfibrozil		Drinking water	USA	SPE-LC-MS/MS	0.25	0.43	[32]			NOEC (96h) (morphology)	0.1 mgL <sup>-1</sup>	[98]
Gemfibrozil		Hospital effluent Pharmaceutical production facility effluent	Taiwan	SPE-HPLC-MS/MS	1.0	760 1795	[47]			EC <sub>50</sub> (96 h) (feeding)	1.76 mgL <sup>-1</sup>	[98]
Gemfibrozil		Pearl rivers water	China	SPE-GC-NCI-MS	1.8	ND–22.4 (±3.1)	[88]	Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (24h) (bioluminescence)	64.6 mgL <sup>-1</sup>	[112]
Gemfibrozil		STP effluent	South Korea	SPE-LC-MS/MS	1.0	3.9–17	[90]			EC <sub>50</sub> (48h) (bioluminescence)	45.1 mgL <sup>-1</sup>	[112]
Gemfibrozil		Surface water STP effluent	Spain	SPE-LC-QqLIT-MS	4 (LOQ)	1.8–9.1 470–3550	[91]	Algae	<i>Chlorella vulgaris</i>	EC <sub>50</sub> (24 h) (growth)	195 mgL <sup>-1</sup>	[112]
								Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48 h) (growth) EC <sub>50</sub> (72 h) (growth) EC <sub>50</sub> (24h) (immobilization)	161 mgL <sup>-1</sup> 150 mgL <sup>-1</sup> 57.1 mgL <sup>-1</sup>	[112] [112] [112]
										EC <sub>50</sub> (48h) (immobilization) EC <sub>50</sub> (72h) (immobilization)	42.6 mgL <sup>-1</sup> 30.0 mgL <sup>-1</sup>	[112] [112]
								Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (30 min) (bioluminescence)	85.74 mgL <sup>-1</sup>	[113]
								Rotifer	<i>B. calyciflorus</i>	LC <sub>50</sub> (24 h) (mortality) EC <sub>50</sub> (48h) (population growth inhibition)	77.30 mgL <sup>-1</sup> 0.44 mgL <sup>-1</sup>	[113] [113]
										NOEC (48h) LOEC (48h)	0.156 mgL <sup>-1</sup> 0.312 mgL <sup>-1</sup>	[113] [113]
								Crustacean	<i>T. platyurus</i>	LC <sub>50</sub> (24 h) (mortality)	161.05 mgL <sup>-1</sup>	[113]
									<i>D. magna</i>	EC <sub>50</sub> (24h) (immobilization)	74.30 mgL <sup>-1</sup>	[113]
									<i>C. dubia</i>	EC <sub>50</sub> (7 d) (population growth inhibition)	0.53 mgL <sup>-1</sup>	[113]
										NOEC (7d) LOEC (7d)	0.078 mgL <sup>-1</sup> 0.156 mgL <sup>-1</sup>	[113] [113]
								Algae	<i>P. subcapitata</i>	EC <sub>50</sub> (72 h) (growth inhibition) NOEC (72h) LOEC (72h)	15.19 mgL <sup>-1</sup> 3.125 mgL <sup>-1</sup> 6.25 mgL <sup>-1</sup>	[113] [113] [113]
<i>Statins</i> Atorvastatin	134523-03-8	Drinking water	USA	SPE-LC-MS/MS	0.25	<0.25	[32]	Duckweed	<i>L. gibba</i>	LOEC (7 d) (growth parameters)	300 µg L <sup>-1</sup>	[103]

Atorvastatin		STP influent STP effluent Otonabee river water	Canada	SPE-LC-MS/MS	0.1	76 ( $\pm 3$ ) 37 ( $\pm 2$ )	[105]						
Atorvastatin		STP effluent	Canada	SPE-LC-MS/MS	0.001	1 ( $\pm 0$ ) 22.4 ( $\pm 1.4$ )	[106]						
<i>o</i> -hydroxy atorvastatin*	214217-86-6	Drinking water	USA	SPE-LC-MS/MS	0.50	<0.50	[32]						
<i>p</i> -hydroxy atorvastatin*	214217-88-6	Drinking water	USA	SPE-LC-MS/MS	0.50	<0.50	[32]						
Lovastatin	81739-26-6	STP influent STP effluent Otonabee river water	Canada	SPE-LC-MS/MS	0.1	49 ( $\pm 2$ ) 14 ( $\pm 1$ )	[105]						
Pravastatin	81131-70-6	STP influent STP effluent Otonabee river water	Canada	SPE-LC-MS/MS	1.0	ND 117 ( $\pm 6$ ) 59 ( $\pm 2$ )	[105]						
Simvastatin	79902-63-9	STP influent	Canada	SPE-LC-MS/MS	0.1	ND 4 ( $\pm 0$ )	[105]	Algae	<i>Dunaliella tertiolecta</i>	EC <sub>50</sub> (96 h) (growth inhibition)	22,800 $\mu\text{g L}^{-1}$	[70]	
		STP effluent Otonabee river water				1 ( $\pm 0$ )							
						ND		Grass shrimp	<i>Palaemonetes pugio</i>	LC <sub>50</sub> (96 h) (larvae survival) NOEC (larvae survival) LOEC (larvae survival) LC <sub>50</sub> (96 h) (adult survival)	1.18 $\text{mg L}^{-1}$ 0.625 $\text{mg L}^{-1}$ 1.25 $\text{mg L}^{-1}$ >10 $\text{mg L}^{-1}$	[101] [101] [101] [101]	
								Copepod	<i>Nitocra spinipes</i>	NOEC (adult survival) LC <sub>50</sub> (96 h) (growth rate) LOEC (growth rate)	5.00 $\text{mg L}^{-1}$ 10.0 $\text{mg L}^{-1}$ 810 $\mu\text{g L}^{-1}$ 0.16 $\mu\text{g L}^{-1}$	[101] [101] [102] [102]	

\*—Metabolite; †—Data not available; ND—Not detected; SPE—Solid Phase Extraction; GC-MS—Gas Chromatography with Mass Spectrometry Detection; GC-MS/MS—Gas Chromatography with Tandem Mass Spectrometry Detection; GC-NCI-MS—Gas Chromatography-Negative Chemical Ionization-Mass Spectrometry; HPLC-MS/MS—High Performance Liquid Chromatography with Tandem Mass Spectrometry Detection; LC-MS/MS—Liquid Chromatography with Tandem Mass Spectrometry Detection.

of *C. vulgaris* was only observed for concentrations up to 150 mg L<sup>-1</sup> [112]. Isidori et al. [113] studied the acute and chronic toxicities caused by bezafibrate, fenofibrate and gemfibrozil and their photoproducts on non-target organisms, considering that they did not significantly affect the exposed organisms (LC<sub>50</sub> values ranged from 39.69 to 161.05 mg L<sup>-1</sup>). When goldfish (*Carassius auratus*) were exposed to 1.5 µg L<sup>-1</sup> of gemfibrozil for 14 days, a decrease of more than 50% in plasma testosterone levels was noticed [114], thereby proving that this pharmaceutical may also act as an endocrine disruptor. As the main active metabolite of several fibrate compounds, clofibric acid is frequently used to assess toxicity due to its high degree of persistence in the environment. In acute toxicity tests on bacteria, ciliates, daphnids and fish embryos, Ferrari et al. [96] noticed low toxicity when at concentrations up to 14 mg L<sup>-1</sup>. These results are in agreement with the tests performed on three estuarine species: algae *D. tertiolecta*, crustacean *P. pugio* and fish *Fundulus heteroclitus* [115]. For concentrations ≤1000 µg L<sup>-1</sup>, clofibric acid did not significantly affect cell density and growth rate of the first, neither did it affect the survival of the remainder. This is in agreement with the 96-h EC<sub>50</sub> of 224.18 mg L<sup>-1</sup> found for *D. tertiolecta* [87]. On the contrary, exposure to concentrations above 10 µg L<sup>-1</sup> and up to 100 µg L<sup>-1</sup> increased the proportion of male offspring produced by *D. magna* [116]. Rotifers have also shown to be sensitive and a NOEC<sup>6</sup> value of 250 µg L<sup>-1</sup> was deduced [96]. Fathead minnow fish (*Pimephales promelas*) showed alterations in reproductive function expressed by a reduction in sperm motility and plasma androgen concentration [117] while cytological changes in gills were noticed in rainbow trout exposed to 5 µg L<sup>-1</sup> of this metabolite [97]. Fibrates (as bezafibrate and gemfibrozil) have been detected in several environmental samples (Table 2). Bezafibrate was found in STP effluents [91,118] and in the Paraíba do Sul river (Brazil) [22] as was gemfibrozil [17,18,21] and further identified in surface waters [17,21,88]. Due to its greater persistence, clofibric acid has been found in STP influents [19,71] and effluents [19,53,71], surface water [22,24,71], drinking water [71,119] and North Sea water [25]. All of these pharmaceuticals were shown to be present at concentration levels in the order of ng L<sup>-1</sup> or low µg L<sup>-1</sup>, which indicates that their exposure may represent a threat for non-target organisms.

#### 4.3. Antibiotics

Antibiotics come within a therapeutic class where human health preservation and environmental disturbance are closely related. The major concern is associated with the development of resistance mechanisms by bacteria which can subsequently compromise public health by means of treatment effectiveness [52,108]. According to Jones et al. [120], antibiotics could be classified as extremely

toxic to microorganisms (EC<sub>50</sub> below 0.1 mg L<sup>-1</sup>) and very toxic to algae (EC<sub>50</sub> between 0.1 and 1 mg L<sup>-1</sup>). Chronic toxicity tests performed on algae have shown high sensitivity to antibacterial agents as deduced from growth inhibition measurements [121,122]. Vertebrates (such as fish) put directly in contact with low levels of antimicrobials apparently did not yield observable effects [123,124]. Accordingly, a LC<sub>50</sub> value above 100 mg L<sup>-1</sup> for Japanese medaka concerning sulfonamides was reported [81]. However, one must bear in mind that algae constitute the basis of the food chain and a decrease in their population will directly affect the entire aquatic ecosystem equilibrium [123,125]. Exposure of *D. magna* to erythromycin, lincomycin, sulfamethoxazole or trimethoprim

in a concentration ranging from 1 to 100 µg L<sup>-1</sup> did not affect the degree of survival, nor morphology in adults or neonates, nor fecun-

dity or sex ratio [116]. Similar results were obtained after chronic exposure to 10 µg L<sup>-1</sup> of sulfamethoxazole [116]. Amoxicillin concentrations ranging from 50 ng L<sup>-1</sup> to 50 mg L<sup>-1</sup> were tested on four different algae without observable effects, unless for the blue-green algae *Synechococcus leopoldensis* for which a NOEC of 0.78 µg L<sup>-1</sup> was achieved [126]. Isidori et al. [124] tested erythromycin, oxytetracycline, sulfamethoxazole, ofloxacin, lincomycin and clarithromycin on aquatic organisms belonging to different trophic levels (bacteria, algae, rotifers, crustaceans and fish). Results showed that acute toxicity level was in the order of mg L<sup>-1</sup> while chronic toxicity appeared at concentrations in the order of µg L<sup>-1</sup>, mainly for algae. The antibiotics tested were shown to be less active against rotifers, crustaceans and fish where no effect was noticed even for concentrations up to 1000 mg L<sup>-1</sup>. After a 48 h exposure period of the microalga *Scenedesmus obliquus* to a concentration range of norfloxacin between 0 and 60 mg L<sup>-1</sup> was noticed a growth inhibition (EC<sub>50</sub> = 38.49 mg L<sup>-1</sup>) and a reduction in chlorophyll-*a* concentration [127].

Most antibiotics used in veterinary medicine are aimed at preventing and treating diseases in livestock production or aquaculture. Even considering their use at sub-therapeutically concentrations, many studies suggest the development of bacterial resistance and further potential appearance of cross-resistance between different classes of antibiotics shared with humans [43,58,120,128]. Antibiotics used in livestock production are excreted in the urine and faeces of animals and often appear in manure. From here they can cause some problems in terrestrial ecosystems such as adverse effects on nitrifying bacteria [11] or growth inhibition of crop plants and weeds by bioaccumulation [129]. The presence of antibiotics in STP influents may also impair treatment processes that use bacteria and cause toxic effects in the downstream aquatic and/or terrestrial ecosystems at different trophic levels [11]. Bacterial cultures from sewage bioreactors receiving waters from a STP were tested for resistance against six antibiotics, showing that all were resistant to at least two of the antibiotics, whilst bacteria isolated from receiving waters were only resistant to erythromycin and ampicillin [130]. Aquatic photosynthetic organisms can also be affected. A study performed both on the cyanobacterium *Synechocystis* sp. and the duckweed *L. minor* showed growth inhibition in the presence of 1–1000 µg L<sup>-1</sup> erythromycin while another antibiotic, tetracycline, inhibited growth of the former when at concentrations between 10 and 100 µg L<sup>-1</sup> while stimulating the latter [77]. Eguchi et al. [131] studied the influence of several antimicrobial agents used as veterinary drugs in Japan on the growth of the green algae *Selenastrum capricornutum* and *C. vulgaris* by considering the growth inhibitory activity. Erythromycin showed

the strongest activity against *S. capricornutum* with an EC<sub>50</sub> value of 37 µg L<sup>-1</sup> followed by dihydrostreptomycin (EC<sub>50</sub> = 110 µg L<sup>-1</sup>), oxytetracycline (EC<sub>50</sub> = 340 µg L<sup>-1</sup>) and tylosin (EC<sub>50</sub> = 410 µg L<sup>-1</sup>). Sulfonamides exhibited lower inhibitory activity with EC<sub>50</sub> values between 1.53 and 2.30 mg L<sup>-1</sup>. In contrast, ampicillin and cefalozin did not show any effect even at concentrations as high as 1000 mg L<sup>-1</sup>. The authors also showed the arousal of a synergistic inhibitory growth activity from the very common combination of sulfamethoxazole with trimethoprim in medicines, when compared to the respective individual activities. Yamashita et al. [132] evaluated the growth inhibition of the algae *P. subcapitata* by two antibiotics, levofloxacin and clarithromycin, showing that the last one had a more pronounced toxic effect with an EC<sub>50</sub> of 11 µg L<sup>-1</sup> and a LOEC and a NOEC of 6.3 and 3.1 µg L<sup>-1</sup>, respectively. Toxic effects of sulfachlorpyridazine and oxytetracycline were also tested on the aquatic plant *L. minor*, showing EC<sub>50</sub> values of 2.33 and 4.92 mg L<sup>-1</sup>, respectively [133]. Assays on *D. magna* showed that following 48 h of exposure, oxolinic acid and tiamulin were the most toxic compounds, with EC<sub>50</sub> values of

<sup>6</sup> NOEC—Non-Observed Effect Concentration.



**Table 3**

Examples of concentrations (ng L<sup>-1</sup>) of antibiotics measured in different aquatic environments.

Compound	CAS number	Sample	Country	Analytical procedure	LOD(ngL <sup>-1</sup> )	Concentration reported(ngL <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
<i>(Fluor)quinolones</i>												
Ciprofloxacin	85721-33-1	Surface water	USA	SPE-LC-MS	20	20	[23]					
Ciprofloxacin		Po river water	Italy	SPE-HPLC-MS/MS	0.3	ND-26.15	[24]					
		Lambro river water				14.36						
Ciprofloxacin		STP influent	USA	SPE-HPLC-MS/MS	20	ND-1000	[138]					
		STP effluent				ND						
		Hospital effluent				ND-2000						
		Rio Grande river water				ND						
Ciprofloxacin		STP influent	Portugal	SPE-LC-FD	25 (LOQ)	418.8-667.1	[139]					
		STP effluent				100.8-309.2						
		Hospital effluent				127.0-10,962.5						
Ciprofloxacin		STP influent	USA	SPE-LC-MS	50	150	[140]					
		STP effluent				60						
Ciprofloxacin		STP influent	Sweden	SPE-LC-MS	6 (LOQ)	90-300	[141]					
		STP effluent				<6-60						
Ciprofloxacin		Mondego river water	Portugal	SPE-LC-FD	25	79.6-119.2	[142]					
Enrofloxacin	93106-60-6	STP influent	Portugal	SPE-LC-FD	50 (LOQ)	121.8-447.1	[139]	Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (15 min) (luminescence)	326.8 mg L <sup>-1</sup>	[13]6
		STP effluent					53.7-211.5					
		Hospital effluent				<50						
Enrofloxacin		STP influent	USA	SPE-LC-MS	50	250	[140]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (24h) (immobilization)	131.7 mg L <sup>-1</sup>	[136]
		STP effluent				270						
Enrofloxacin		Mondego river water	Portugal	SPE-LC-FD	25	67.0-102.5	[142]			EC <sub>50</sub> (48h) (immobilization)	56.7 mg L <sup>-1</sup>	[136]
										EC <sub>50</sub> (21 d) (adult survival)	11.47 mg L <sup>-1</sup>	[136]
										LOEC (21 d) (reproduction)	15 mg L <sup>-1</sup>	[136]
										NOEC (21 d) (reproduction)	5 mg L <sup>-1</sup>	[136]
									<i>M. macrocopa</i>	EC <sub>50</sub> (24h) (immobilization)	285.7 mg L <sup>-1</sup>	[136]
Levofloxacin	100986-34-5	Mankyoung river water	South Korea	SPE-LC-MS/MS	5	ND-87.4 (±13)	[92]	Crustacean	<i>T. platyurus</i>	LC <sub>50</sub> (24 h) (mortality)	>100 mg L <sup>-1</sup>	[78]
								Fish	<i>O. latipes</i>	LC <sub>50</sub> (96 h) (mortality)	>100 mg L <sup>-1</sup>	[78]
								Algae	<i>P. subcapitata</i>	EC <sub>50</sub> (96 h) (growth inhibition)	1200 μg L <sup>-1</sup>	[132]
										LOEC (96h) (growth inhibition)	630 μg L <sup>-1</sup>	[132]
										NOEC (96h) (growth inhibition)	310 μg L <sup>-1</sup>	[132]
								Crustacean	<i>D. magna</i>	EC <sub>50</sub> (21 d) (reproduction)	340 μg L <sup>-1</sup>	[132]
										LOEC (21 d) (reproduction)	63 μg L <sup>-1</sup>	[132]

Table 3 (Continued)

Compound	CAS number	Sample	Country	Analytical procedure	LOD (ngL <sup>-1</sup> )	Concentration reported (ngL <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
Nalidixic acid	389-08-2	STP influent	Taiwan	SPE-HPLC-MS/MS	— <sup>f</sup>	26–372	[87]			NOEC (21 d) (reproduction)	31 μg L <sup>-1</sup>	[132]
Norfloxacin	70458-96-7	STP effluent Surface water	USA	SPE-LC-MS	20	40–200 120	[23]	Algae	<i>S. obliquus</i>	IC <sub>50</sub> (48 h) (growth inhibition)	38.49 mg L <sup>-1</sup>	[127]
Norfloxacin		STP influent	Portugal	SPE-LC-FD	25 (LOQ)	191.2–455.0	[139]	Algae	<i>S. capricornutum</i>	EC <sub>50</sub> (growth inhibition)	16.6 mg L <sup>-1</sup>	[131]
Norfloxacin		STP effluent Hospital effluent STP influent	Sweden	SPE-LC-MS	7 (LOQ)	29.6–35.0 <25–334.0 72–174	[141]			NOEC (growth inhibition)	4.01 mg L <sup>-1</sup>	[131]
Norfloxacin		STP effluent Mondego river water	Portugal	SPE-LC-FD	25	<6–37 ND	[142]		<i>C. vulgaris</i>	EC <sub>50</sub> (growth inhibition)	10.4 mg L <sup>-1</sup>	[131]
Norfloxacin		Surface seawater	China (Hong Kong)	SPE-HPLC-MS/MS	13	<13–8.00	[144]			NOEC (growth inhibition)	4.02 mg L <sup>-1</sup>	[131]
Norfloxacin		Victoria Harbour seawater Pearl River water	China	SPE-HPLC-MS	3.2 (LOQ seawater) 10 (LOQ riverwater)	9.4–12.3	[145]	Rotifer	<i>B. calyciflorus</i>	LC <sub>50</sub> (24 h) (mortality)	29.88 mg L <sup>-1</sup>	[124]
Ofloxacin	82419-36-1	STP influent	Taiwan	SPE-HPLC-MS/MS	— <sup>f</sup>	12–150 115–1274	[87]			EC <sub>50</sub> (48h) (population growth inhibition)	0.53 mg L <sup>-1</sup>	[124]
Ofloxacin		STP effluent STP influent	USA	SPE-HPLC-MS/MS	10	53–991 ND–1000	[138]	Crustacean	<i>T. platyurus</i>	LC <sub>50</sub> (24 h) (mortality)	33.98 mg L <sup>-1</sup>	[124]
Ofloxacin		STP effluent Hospital effluent Rio Grande river water STP influent	Portugal	SPE-LC-FD	250	110 ND–35,500 ND	[139]		<i>D. magna</i>	EC <sub>50</sub> (24h) (immobilization)	31.75 mg L <sup>-1</sup>	[124]
Ofloxacin		STP effluent Hospital effluent STP influent	Sweden	SPE-LC-MS	6 (LOQ)	ND ND–10,675.5 <6–287	[141]		<i>C. dubia</i>	EC <sub>50</sub> (24h) (immobilization)	17.41 mg L <sup>-1</sup>	[124]
Ofloxacin		STP effluent Victoria Harbour seawater Pearl River water	China	SPE-HPLC-MS	2.6 (LOQ seawater) 10 (LOQ riverwater)	<6–52 5.2–10	[145]			EC <sub>50</sub> (7 d) (population growth inhibition)	3.13 mg L <sup>-1</sup>	[124]
Oxolinic acid	14698-29-4					11–77		Algae	<i>P. subcapitata</i>	EC <sub>50</sub> (72 h) (growth inhibition)	1.44 mg L <sup>-1</sup>	[124]
								Algae	<i>M. aeruginosa</i>	EC <sub>50</sub> (72 h) (growth inhibition)	0.180 mg L <sup>-1</sup>	[122]

										<i>R. salina</i>	EC <sub>50</sub> (72 h) (growth inhibition)	10 mg L <sup>-1</sup>	[122]
										<i>S. capricornutum</i>	EC <sub>50</sub> (72 h) (growth inhibition)	16 mg L <sup>-1</sup>	[122]
									Crustacean	<i>D. magna</i>	EC <sub>50</sub> (24h) (immobilization)	5.9 mg L <sup>-1</sup>	[134]
											EC <sub>50</sub> (48h) (immobilization)	4.6 mg L <sup>-1</sup>	[134]
											NOEC (21 d) (reproduction)	0.38 mg L <sup>-1</sup>	[134]
Sarafloxacin	98105-99-8								Algae	<i>R. salina</i>	EC <sub>50</sub> (72 h) (growth inhibition)	24 mg L <sup>-1</sup>	[122]
										<i>S. capricornutum</i>	EC <sub>50</sub> (72 h) (growth inhibition)	16 mg L <sup>-1</sup>	[122]
~Lactams									Algae	<i>M. aeruginosa</i>	EC <sub>50</sub> (72h) (growth inhibition)	0.0037 mg L <sup>-1</sup>	[122]
Amoxicillin	81030-75-3									<i>S. capricornutum</i>	NOEC (72h) (growth inhibition)	250 mg L <sup>-1</sup>	[122]
									Algae	<i>S. leopoliensis</i>	EC <sub>50</sub> (growth inhibition)	2.22 μg L <sup>-1</sup>	[126]
											NOEC (growth inhibition) LOEC	0.78 μg L <sup>-1</sup>	[126]
											(growth inhibition)	1.56 μg L <sup>-1</sup>	[126]
									Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (15 min) (luminescence)	3597 mg L <sup>-1</sup>	[136]
Ampicillin	69-53-4	Hospital effluent	Taiwan	SPE-HPLC-MS/MS	10	21	[47]		Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (15 min) (luminescence)	2627 mg L <sup>-1</sup>	[136]
		Pharmaceutical production facility effluent				ND							
Penicillin G (Benzylpenicillin)	69-57-8	STP influent	China	SPE-LC-MS	930 (LOQ)	153,000 ± 4000	[48]	Algae	<i>M. aeruginosa</i>		EC <sub>50</sub> (growth rate)	0.006 mg L <sup>-1</sup>	[121]
		STP effluent				1680 ± 480			<i>S. capricornutum</i>		NOEC (growth rate)	100 mg L <sup>-1</sup>	[121]
Cephalosporins													
Cephalexin	66905-57-5	STP influent	Taiwan	SPE-HPLC-MS/MS	— <sup>†</sup>	1563–4367	[87]						
		STP effluent				10–994							
Cephalexin		Surface seawater	China	SPE-HPLC-MS/MS	13	<13–182	[144]						
			(Hong Kong)										
Lincosamide													
Lincomycin	154-21-2	Surface water	USA	SPE-LC-MS	50	60	[23]	Rotifer	<i>B. calyciflorus</i>		LC <sub>50</sub> (24 h) (mortality)	24.94 mg L <sup>-1</sup>	[124]
		Po river water	Italy	SPE-HPLC-MS/MS	0.3	3.13–248.90	[24]				EC <sub>50</sub> (48h) (population growth inhibition)	0.68 mg L <sup>-1</sup>	[124]
		Lambro river water				24.40							
Lincomycin		Groundwater	USA	SPE-LC-MS	50	320	[28]	Crustacean	<i>T. platyurus</i>		LC <sub>50</sub> (24 h) (mortality)	30.00 mg L <sup>-1</sup>	[124]
Lincomycin		Hospital effluent	USA	SPE-HPLC-MS/MS	10	ND–2000	[138]		<i>D. magna</i>		EC <sub>50</sub> (24h) (immobilization)	23.18 mg L <sup>-1</sup>	[124]
		Livestock effluent				ND–6600			<i>C. dubia</i>		EC <sub>50</sub> (24h) (immobilization)	13.98 mg L <sup>-1</sup>	[124]

Table 3 (Continued)

Compound	CAS number	Sample	Country	Analytical procedure	LOD (ngL <sup>-1</sup> )	Concentration reported (ngL <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.	
<i>Macrolides</i>	Clarithromycin	81103-11-9	Po river water Lambro river water	Italy	SPE-HPLC-MS/MS	0.3	[24]	Algae	<i>P. subcapitata</i>	EC <sub>50</sub> (7 d) (population growth inhibition)	7.20 mgL <sup>-1</sup>	[124]	
										EC <sub>50</sub> (72 h) (growth inhibition)	0.07 mgL <sup>-1</sup>	[124]	
Clarithromycin		STP influent	Taiwan	SPE-HPLC-MS/MS	— <sup>†</sup>	59–1433	[87]	Rotifer	<i>B. calyciflorus</i>	LC <sub>50</sub> (24 h) (mortality)	35.46 mgL <sup>-1</sup>	[124]	
Clarithromycin		STP effluent	South Korea	SPE-LC-MS/MS	1	12–232 ND–443 (±14)	[92]	Crustacean	<i>T. platyurus</i>	LC <sub>50</sub> (24 h) (mortality)	94.23 mgL <sup>-1</sup>	[78]	
		Fish						<i>O. latipes</i>	LC <sub>50</sub> (96 h) (mortality)	>100 mgL <sup>-1</sup>	[78]		
		Crustacean						<i>T. platyurus</i>	LC <sub>50</sub> (24 h) (mortality)	33.64 mgL <sup>-1</sup>	[124]		
								<i>D. magna</i>	EC <sub>50</sub> (24h) (immobilization)	25.72 mgL <sup>-1</sup>	[124]		
								<i>C. dubia</i>	EC <sub>50</sub> (24h) (immobilization)	18.66 mgL <sup>-1</sup>	[124]		
									EC <sub>50</sub> (7 d) (population growth inhibition)	8.16 mgL <sup>-1</sup>	[124]		
								Algae	<i>P. subcapitata</i>	EC <sub>50</sub> (72 h) (growth inhibition)	0.0020 mgL <sup>-1</sup>	[124]	
								Algae	<i>P. subcapitata</i>	EC <sub>50</sub> (96 h) (growth inhibition)	11 μgL <sup>-1</sup>	[132]	
									LOEC (96h) (growth inhibition)	6.3 μgL <sup>-1</sup>	[132]		
									NOEC (96h) (growth inhibition)	3.1 μgL <sup>-1</sup>	[132]		
								Crustacean	<i>D. magna</i>	EC <sub>50</sub> (21 d) (reproduction)	40 μgL <sup>-1</sup>	[132]	
									LOEC (21 d) (reproduction)	6.3 μgL <sup>-1</sup>	[132]		
									NOEC (21 d) (reproduction)	3.1 μgL <sup>-1</sup>	[132]		
Erithromycin	114-07-8	Po river water Lambro river water	Italy	SPE-HPLC-MS/MS	0.3	1.40–15.90	[24]	Duckweed	<i>Lemna minor</i>	EC <sub>50</sub> (7 d) (growth inhibition)	5.62 mgL <sup>-1</sup>	[77]	
Erithromycin		STP effluent	South Korea	SPE-LC-MS/MS	1.0	8.9–294	[90]	Crustacean	<i>T. platyurus</i>	LC <sub>50</sub> (24 h) (mortality)	>100 mgL <sup>-1</sup>	[78]	
Erithromycin		Surface water	South Korea	SPE-LC-MS/MS	1	1.8–4.8 ND–137 (±15)	[92]	Fish	<i>O. latipes</i>	LC <sub>50</sub> (96 h) (mortality)	>100 mgL <sup>-1</sup>	[78]	
		Mankyung river water							Rotifer	<i>B. calyciflorus</i>	LC <sub>50</sub> (24 h) (mortality)	27.53 mgL <sup>-1</sup>	[124]
											EC <sub>50</sub> (48h) (population growth inhibition)	0.94 mgL <sup>-1</sup>	[124]

									Crustacean	<i>T. platyurus</i>	LC <sub>50</sub> (24 h) (mortality)	17.68 mg L <sup>-1</sup>	[124]
										<i>D. magna</i>	EC <sub>50</sub> (24h) (immobilization)	22.45 mg L <sup>-1</sup>	[124]
										<i>C. dubia</i>	EC <sub>50</sub> (24h) (immobilization)	10.23 mg L <sup>-1</sup>	[124]
											EC <sub>50</sub> (7 d) (population growth inhibition)	0.22 mg L <sup>-1</sup>	[124]
									Algae	<i>P. subcapitata</i>	EC <sub>50</sub> (72 h) (growth inhibition)	0.020 mg L <sup>-1</sup>	[124]
									Algae	<i>S. capricornutum</i>	EC <sub>50</sub> (growth inhibition)	0.0366 mg L <sup>-1</sup>	[131]
											NOEC (growth inhibition)	0.0103 mg L <sup>-1</sup>	[131]
										<i>C. vulgaris</i>	EC <sub>50</sub> (growth inhibition)	33.8 mg L <sup>-1</sup>	[131]
											NOEC (growth inhibition)	12.5 mg L <sup>-1</sup>	[131]
Erithromycin-H <sub>2</sub> O*	114-07-8	Hospital effluent Pharmaceutical production facility effluent	Taiwan	SPE-HPLC-MS/MS	1.0	938	[47]						
		STP influent	Taiwan	SPE-HPLC-MS/MS	—†	226–1537	[87]						
Erithromycin-H <sub>2</sub> O*		STP effluent	China	SPE-HPLC-MS/MS	13	361–811	[144]						
		Surface seawater	(Hong Kong)			9.50–486							
Erithromycin-H <sub>2</sub> O*		Victoria Harbour seawater	China	SPE-HPLC-MS	2.0 (LOQ seawater)	3.3–3.4	[145]						
		Pearl River water			5 (LOQ river water)								
Roxithromycin	80214-83-1	Surface water	USA	SPE-LC-MS	30	30–460	[23]						
Roxithromycin		Victoria Harbour seawater	China	SPE-HPLC-MS	2.0 (LOQ seawater)	50	[145]						
		Pearl River water			5 (LOQ river water)	5.1–6.1							
Spiramycin	67262-35-5	Po river water	Italy	SPE-HPLC-MS/MS	0.3	16–66	[24]	Algae	<i>M. aeruginosa</i>	EC <sub>50</sub> (growth rate)	0.005 mg L <sup>-1</sup>	[121]	
		Lambro river water				ND–43.80			<i>S. capricornutum</i>	EC <sub>50</sub> (growth rate)	2.3 mg L <sup>-1</sup>	[121]	
						74.20			<i>M. aeruginosa</i>	EC <sub>50</sub> (growth rate)	0.034 mg L <sup>-1</sup>	[121]	
Tylosin	1401-69-0	Surface water	USA	SPE-LC-MS	50	40	[23]	Algae	<i>S. capricornutum</i>	EC <sub>50</sub> (growth rate)	1.38 mg L <sup>-1</sup>	[121]	
Tylosin		Po river water	Italy	SPE-HPLC-MS/MS	0.3	ND–0.30	[24]		<i>S. capricornutum</i>	EC <sub>50</sub> (growth rate)			
		Lambro river water				2.77		Algae	<i>S. capricornutum</i>	EC <sub>50</sub> (growth inhibition)	0.411 mg L <sup>-1</sup>	[131]	
										NOEC (growth inhibition)	0.206 mg L <sup>-1</sup>	[131]	
								Crustacean	<i>D. magna</i>	LOEC (24h) (immobilization)	700 mg L <sup>-1</sup>	[134]	

Table 3 (Continued)

Compound	CAS number	Sample	Country	Analytical procedure	LOD (ng L <sup>-1</sup> )	Concentration reported (ng L <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
<i>Sulfonamides</i> Sulfachloropyridazine	80-32-0	STP influent STP effluent	Korea	SPE-LC-MS	30	<30-476 <30-149	[93]	Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (48h) (immobilization) NOEC (21 d) (reproduction)	680 mg L <sup>-1</sup> 45 mg L <sup>-1</sup>	[134] [134]
								Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48h) (immobilization)	375.3 mg L <sup>-1</sup>	[82]
									<i>D. magna</i>	EC <sub>50</sub> (96h) (immobilization)	233.5 mg L <sup>-1</sup>	[82]
								Fish	<i>O. latipes</i>	LC <sub>50</sub> (48 h)	589.3 mg L <sup>-1</sup>	[82]
									<i>O. latipes</i>	LC <sub>50</sub> (96 h)	535.7 mg L <sup>-1</sup>	[82]
Aquatic plant	<i>Lemna minor</i>	EC <sub>50</sub> (48 h) (n° of green fronds)	2.33 mg L <sup>-1</sup>	[133]								
Sulfadiazine	68-35-9	Tevere river water	Italy	SPE-LC-MS	21 (LOQ)	236	[143]	Algae	<i>M. aeruginosa</i>	EC <sub>50</sub> (72 h) (growth inhibition)	0.135 mg L <sup>-1</sup>	[122]
Sulfadiazine		Victoria Harbour seawater Pearl River water	China	SPE-HPLC-MS	0.5 (LOQ seawater)  1 (LOQ river water)	ND  38-209	[145]	S.	<i>capricornutum</i>	EC <sub>50</sub> (72 h) (growth inhibition)	7.8 mg L <sup>-1</sup>	[122]
								Algae	<i>S. capricornutum</i>	EC <sub>50</sub> (growth inhibition)	2.19 mg L <sup>-1</sup>	[131]
										NOEC (growth inhibition)	<1.00 mg L <sup>-1</sup>	[131]
								Crustacean	<i>D. magna</i>	LOEC (24h) (immobilization)	150 mg L <sup>-1</sup>	[132]
										EC <sub>50</sub> (48h) (immobilization)	221 mg L <sup>-1</sup>	[132]
EC <sub>50</sub> (21 d) (reproduction)	13.7 mg L <sup>-1</sup>	[132]										
Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48h) (immobilization)	212 mg L <sup>-1</sup>	[135]								
Sulfadimethoxine	122-11-2	Surface water	USA	SPE-LC-MS	50	60	[23]	Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (15 min)	>500 mg L <sup>-1</sup>	[82]
Sulfadimethoxine		Groundwater	USA	SPE-LC-MS/MS	30	46-68	[27]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48h) (immobilization)	248.0 mg L <sup>-1</sup>	[82]
Sulfadimethoxine		Hospital effluent Pharmaceutical production facility effluent	Taiwan	SPE-HPLC-MS/MS	1.0	ND 0.8	[47]		<i>D. magna</i>	EC <sub>50</sub> (96h) (immobilization)	204.5 mg L <sup>-1</sup>	[82]
Sulfadimethoxine		STP influent STP effluent Alzette river water Mess river water	Luxembourg	SPE-LC-MS/MS	0.3	0.3-26 0.3-9 0.3-3  <0.3	[89]	Fish	<i>O. latipes</i>	LC <sub>50</sub> (48 h)	>100 mg L <sup>-1</sup>	[82]
Sulfadimethoxine		STP influent STP effluent Han river water	Korea	SPE-LC-MS	10	<10-213 <10-70 <10-13	[93]		<i>O. latipes</i>	LC <sub>50</sub> (96 h)	>100 mg L <sup>-1</sup>	[82]
Sulfadimethoxine		Tevere river water	Italy	SPE-LC-MS	8	28	[143]	Algae	<i>S. capricornutum</i>	EC <sub>50</sub> (growth inhibition)	2.30 mg L <sup>-1</sup>	[131]

		Trigno river water				74								
		Drinking water				11								
												NOEC (growth inhibition)	0.529 mg L <sup>-1</sup>	[131]
										<i>C. vulgaris</i>		EC <sub>50</sub> (growth inhibition)	11.2 mg L <sup>-1</sup>	[131]
												NOEC (growth inhibition)	<20.3 mg L <sup>-1</sup>	[131]
								Crustacean		<i>D. magna</i>		EC <sub>50</sub> (48h) (immobilization)	270 mg L <sup>-1</sup>	[135]
								Crustacean		<i>D. magna</i>		EC <sub>50</sub> (24h) (immobilization)	639.8 mg L <sup>-1</sup>	[136]
										<i>M. macrocopa</i>		EC <sub>50</sub> (24h) (immobilization)	296.6 mg L <sup>-1</sup>	[136]
												EC <sub>50</sub> (48h) (immobilization)	183.9 mg L <sup>-1</sup>	[136]
Sulfamethazine	57-68-1	Groundwater	USA	SPE-LC-MS/MS	20	76-215	[27]	Bacteria		<i>V. fischeri</i>		EC <sub>50</sub> (15 min)	344.7 mg L <sup>-1</sup>	[82]
Sulfamethazine		Groundwater	USA	SPE-LC-MS	50	360	[28]	Crustacean		<i>D. magna</i>		EC <sub>50</sub> (48h) (immobilization)	174.4 mg L <sup>-1</sup>	[82]
Sulfamethazine		Hospital effluent	Taiwan	SPE-HPLC-MS/MS	0.5	ND	[47]			<i>D. magna</i>		EC <sub>50</sub> (96h) (immobilization)	158.8 mg L <sup>-1</sup>	[82]
		Pharmaceutical production facility effluent				178								
Sulfamethazine		STP influent	Luxembourg	SPE-LC-MS/MS	0.3	0.3-2	[89]	Fish		<i>O. latipes</i>		LC <sub>50</sub> (48 h)	>100 mg L <sup>-1</sup>	[82]
		STP effluent				<0.3								
		Alzette river water				<0.3								
		Mess river water				<0.3								
Sulfamethazine		STP influent	USA	SPE-LC-MS	50	160	[140]			<i>O. latipes</i>		LC <sub>50</sub> (96 h)	>100 mg L <sup>-1</sup>	[82]
		STP effluent				ND		Crustacean		<i>D. magna</i>		EC <sub>50</sub> (48h) (immobilization)	202 mg L <sup>-1</sup>	[135]
												EC <sub>50</sub> (21 d) (reproduction)	4.25 mg L <sup>-1</sup>	[135]
												LOEC (21 d) (reproduction)	3.125 mg L <sup>-1</sup>	[135]
												NOEC (21 d) (reproduction)	1.563 mg L <sup>-1</sup>	[135]
								Crustacean		<i>D. magna</i>		EC <sub>50</sub> (24h) (immobilization)	506.3 mg L <sup>-1</sup>	[136]
												EC <sub>50</sub> (48h) (immobilization)	215.9 mg L <sup>-1</sup>	[136]
										<i>M. macrocopa</i>		EC <sub>50</sub> (24h) (immobilization)	310.9 mg L <sup>-1</sup>	[136]
												EC <sub>50</sub> (48h) (immobilization)	110.7 mg L <sup>-1</sup>	[136]
Sulfamethoxazole	723-46-6	Surface water	USA	SPE-LC-MS	50	150	[23]	Bacteria		<i>V. fischeri</i>		EC <sub>50</sub> (15 min)	78.1 mg L <sup>-1</sup>	[82]
Sulfamethoxazole		Groundwater	USA	SPE-LC-MS	23	1110	[28]	Crustacean		<i>D. magna</i>		EC <sub>50</sub> (48h) (immobilization)	189.2 mg L <sup>-1</sup>	[82]
Sulfamethoxazole		Drinking water	USA	SPE-LC-MS/MS	0.25	0.32	[32]			<i>D. magna</i>		EC <sub>50</sub> (96h) (immobilization)	177.3 mg L <sup>-1</sup>	[82]
Sulfamethoxazole		Hospital effluent	Taiwan	SPE-HPLC-MS/MS	1.0	1335	[47]	Fish		<i>O. latipes</i>		LC <sub>50</sub> (48 h)	>750 mg L <sup>-1</sup>	[82]

Table 3 (Continued)

Compound	CAS number	Sample	Country	Analytical procedure	LOD (ngL <sup>-1</sup> )	Concentration reported (ngL <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
		Pharmaceutical production facility effluent				34						
Sulfamethoxazole		STP influent	Taiwan	SPE-HPLC-MS/MS	— <sup>†</sup>	179–1760	[87]		<i>O. latipes</i>	LC <sub>50</sub> (96 h)	562.5 mgL <sup>-1</sup>	[82]
Sulfamethoxazole		STP effluent	Luxembourg	SPE-LC-MS/MS	0.3	47–964	[89]	Cnidarian	<i>Hydra attenuata</i>	LC <sub>50</sub> (96 h) (morphology)	>100 mgL <sup>-1</sup>	[98]
		STP effluent				4–39						
		Alzette river water				1–22						
		Mess river water				0.3–5						
Sulfamethoxazole		STP effluent	South Korea	SPE-LC-MS/MS	1.0	3.8–407	[90]			LOEC (96 h) (morphology)	10 mgL <sup>-1</sup>	[98]
Sulfamethoxazole		Surface water	Korea	SPE-LC-MS	5	1.7–36	[93]			NOEC (96 h) (morphology)	5 mgL <sup>-1</sup>	[98]
		STP influent				156–984						
Sulfamethoxazole		STP effluent	USA	SPE-HPLC-MS/MS	12	25–492	[138]	Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (30 min) (luminescence)	23.3 mgL <sup>-1</sup>	[124]
		Han river water				<5–82						
		STP influent				ND–1000						
		Hospital effluent				310						
		Rio Grande river water				ND–2100						
Sulfamethoxazole		STP influent	USA	SPE-LC-MS	50	ND–300	[140]	Rotifer	<i>B. calyciflorus</i>	LC <sub>50</sub> (24 h) (mortality)	26.27 mgL <sup>-1</sup>	[124]
Sulfamethoxazole		STP effluent	Sweden	SPE-LC-MS	80 (LOQ)	300	[141]			EC <sub>50</sub> (48 h) (population growth inhibition)	9.63 mgL <sup>-1</sup>	[124]
		STP influent				200						
		STP influent				<80–304						
Sulfamethoxazole		Tevere river water	Italy	SPE-LC-MS	9	402	[143]	Crustacean	<i>T. platyurus</i>	LC <sub>50</sub> (24 h) (mortality)	35.36 mgL <sup>-1</sup>	[124]
		Drinking water				13–80						
Sulfamethoxazole		Victoria Harbour seawater	China	SPE-HPLC-MS	0.8 (LOQ seawater)	ND	[145]		<i>D. magna</i>	EC <sub>50</sub> (24 h) (immobilization)	25.20 mgL <sup>-1</sup>	[124]
		Pearl River water				1 (LOQ river water)						
						37–134						
									<i>C. dubia</i>	EC <sub>50</sub> (24 h) (immobilization)	15.51 mgL <sup>-1</sup>	[124]
										EC <sub>50</sub> (7 d) (population growth inhibition)	0.21 mgL <sup>-1</sup>	[124]
								Algae	<i>P. subcapitata</i>	EC <sub>50</sub> (72 h) (growth inhibition)	0.52 mgL <sup>-1</sup>	[124]
								Algae	<i>S. capricornutum</i>	EC <sub>50</sub> (growth inhibition)	1.53 mgL <sup>-1</sup>	[131]



									Crustacean	<i>D. magna</i>	NOEC (growth inhibition)	0.614 mg L <sup>-1</sup>	[131]
											EC <sub>50</sub> (48h) (immobilization)	123.1 mg L <sup>-1</sup>	[136]
										<i>M. macrocopa</i>	EC <sub>50</sub> (24h) (immobilization)	84.9 mg L <sup>-1</sup>	[136]
											EC <sub>50</sub> (48h) (immobilization)	70.4 mg L <sup>-1</sup>	[136]
Sulfapyridine	7238-91-7	Tevere river water Trigno river water	Italy	SPE-LC-MS	12	<12–121 66	[143]	Cnidarian	<i>Hydra attenuata</i>		LC <sub>50</sub> (96 h) (morphology)	>100 mg L <sup>-1</sup>	[98]
											EC <sub>50</sub> (96h) (morphology)	21.61 mg L <sup>-1</sup>	[98]
											LOEC (96h) (morphology)	5 mg L <sup>-1</sup>	[98]
											NOEC (96h) (morphology)	1 mg L <sup>-1</sup>	[98]
Sulfathiazole	72-14-0	STP influent STP effluent Alzette river water Mess river water	Luxembourg	SPE-LC-MS/MS	0.3	0.3–2 <0.3 <0.3 0.3–2	[89]	Bacteria	<i>V. fischeri</i>		EC <sub>50</sub> (15 min)	>1000 mg L <sup>-1</sup>	[82]
Sulfathiazole		STP influent STP effluent	Korea	SPE-LC-MS	30	<30–531 <30	[93]	Crustacean	<i>D. magna</i>		LOEC (21 d) (reproduction)	35 mg L <sup>-1</sup>	[136]
											NOEC (21 d) (reproduction)	11 mg L <sup>-1</sup>	[136]
								Crustacean	<i>D. magna</i>		EC <sub>50</sub> (48h) (immobilization)	149.3 mg L <sup>-1</sup>	[82]
									<i>D. magna</i>		EC <sub>50</sub> (96h) (immobilization)	85.4 mg L <sup>-1</sup>	[82]
								Fish	<i>O. latipes</i>		LC <sub>50</sub> (48 h)	>500 mg L <sup>-1</sup>	[82]
									<i>O. latipes</i>		LC <sub>50</sub> (96 h)	>500 mg L <sup>-1</sup>	[82]
								Crustacean	<i>D. magna</i>		EC <sub>50</sub> (24h) (immobilization)	616.7 mg L <sup>-1</sup>	[136]
									<i>M. macrocopa</i>		EC <sub>50</sub> (24h) (immobilization)	430.1 mg L <sup>-1</sup>	[136]
											EC <sub>50</sub> (48h) (immobilization)	391.1 mg L <sup>-1</sup>	[136]
<i>Tetracyclines</i>													
Chlortetracycline	57-62-5	Surface water	USA	SPE-LC-MS	100	420	[23]	Algae	<i>M. aeruginosa</i>		EC <sub>50</sub> (growth rate)	0.05 mg L <sup>-1</sup>	[121]
Chlortetracycline		Hospital effluent Pharmaceutical production facility effluent	Taiwan	SPE- HPLC-MS/MS	5.0	ND 5.7	[47]		<i>S. capricornutum</i>		EC <sub>50</sub> (growth rate)	3.1 mg L <sup>-1</sup>	[121]
								Bacteria	<i>V. fischeri</i>		EC <sub>50</sub> (15 min) (luminescence)	13.0 mg L <sup>-1</sup>	[136]
								Crustacean	<i>D. magna</i>		EC <sub>50</sub> (24h) (immobilization)	380.1 mg L <sup>-1</sup>	[136]
											EC <sub>50</sub> (48h) (immobilization)	225 mg L <sup>-1</sup>	[136]
									<i>M. macrocopa</i>		EC <sub>50</sub> (24h) (immobilization)	515 mg L <sup>-1</sup>	[136]
											EC <sub>50</sub> (48h) (immobilization)	272 mg L <sup>-1</sup>	[136]

Table 3 (Continued)

Compound	CAS number	Sample	Country	Analytical procedure	LOD( $\text{ngL}^{-1}$ )	Concentration reported ( $\text{ngL}^{-1}$ )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
								Fish	<i>O. latipes</i>	LC <sub>50</sub> (24 h) (mortality)	88.4 $\text{mgL}^{-1}$	[136]
										LC <sub>50</sub> (48 h) (mortality)	78.9 $\text{mgL}^{-1}$	[136]
Oxytetracycline	79-57-2	Surface water	USA	SPE-LC-MS	100	340	[23]	Cnidarian	<i>Hydra attenuata</i>	LC <sub>50</sub> (96 h) (morphology)	>100 $\text{mgL}^{-1}$	[98]
Oxytetracycline		Po river water	Italy	SPE-HPLC-MS/MS	0.3	ND-19.2	[24]			EC <sub>50</sub> (96h) (morphology)	40.13 $\text{mgL}^{-1}$	[98]
		Lambro river water				14.35						
Oxytetracycline		Hospital effluent	Taiwan	SPE-HPLC-MS/MS	2.0	2.9	[47]			LOEC (96h) (morphology)	100 $\text{mgL}^{-1}$	[98]
		Pharmaceutical production facility effluent				23						
Oxytetracycline		STP influent	Luxembourg	SPE-LC-MS/MS	0.3	0.3-7	[89]			NOEC (96h) (morphology)	50 $\text{mgL}^{-1}$	[98]
		STP effluent				0.3-5						
		Alzette river water				0.3-2						
		Mess river water				0.3-7						
								Algae	<i>M. aeruginosa</i>	EC <sub>50</sub> (72 h) (growth inhibition)	0.207 $\text{mgL}^{-1}$	[122]
									<i>R. salina</i>	EC <sub>50</sub> (72 h) (growth inhibition)	1.6 $\text{mgL}^{-1}$	[122]
									<i>S. capricornutum</i>	EC <sub>50</sub> (72 h) (growth inhibition)	4.5 $\text{mgL}^{-1}$	[122]
								Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (30 min) (luminescence)	64.50 $\text{mgL}^{-1}$	[124]
								Rotifer	<i>B. calyciflorus</i>	LC <sub>50</sub> (24 h) (mortality)	34.21 $\text{mgL}^{-1}$	[124]
										EC <sub>50</sub> (48h) (population growth inhibition)	1.87 $\text{mgL}^{-1}$	[124]
								Crustacean	<i>T. platyurus</i>	LC <sub>50</sub> (24 h) (mortality)	25.00 $\text{mgL}^{-1}$	[124]
									<i>D. magna</i>	EC <sub>50</sub> (24h) (immobilization)	22.64 $\text{mgL}^{-1}$	[124]
									<i>C. dubia</i>	EC <sub>50</sub> (24h) (immobilization)	18.65 $\text{mgL}^{-1}$	[124]
										EC <sub>50</sub> (7 d) (population growth inhibition)	0.18 $\text{mgL}^{-1}$	[124]
								Algae	<i>P. subcapitata</i>	EC <sub>50</sub> (72 h) (growth inhibition)	0.17 $\text{mgL}^{-1}$	[124]
								Algae	<i>S. capricornutum</i>	EC <sub>50</sub> (growth inhibition)	0.342 $\text{mgL}^{-1}$	[131]
										NOEC (growth inhibition)	0.183 $\text{mgL}^{-1}$	[131]
									<i>C. vulgaris</i>	EC <sub>50</sub> (growth inhibition)	7.05 $\text{mgL}^{-1}$	[131]
										NOEC (growth inhibition)	<3.58 $\text{mgL}^{-1}$	[131]

									Aquatic plant	<i>Lemna minor</i>	EC <sub>50</sub> (48 h) (n° of green fronds)	4.92 mg L <sup>-1</sup>	[133]
									Crustacean	<i>D. magna</i>	LOEC (48 h) (immobilization)	100 mg L <sup>-1</sup>	[134]
											EC <sub>50</sub> (21 d) (reproduction)	46.2 mg L <sup>-1</sup>	[134]
									Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (15 min) (luminescence)	87.0 mg L <sup>-1</sup>	[136]
									Crustacean	<i>D. magna</i>	EC <sub>50</sub> (24 h) (immobilization)	831.6 mg L <sup>-1</sup>	[136]
											EC <sub>50</sub> (48 h) (immobilization)	621.2 mg L <sup>-1</sup>	[136]
										<i>M. macrocopa</i>	EC <sub>50</sub> (24 h) (immobilization)	137.1 mg L <sup>-1</sup>	[136]
											EC <sub>50</sub> (48 h) (immobilization)	126.7 mg L <sup>-1</sup>	[136]
									Fish	<i>O. latipes</i>	LC <sub>50</sub> (24 h) (mortality)	215.4 mg L <sup>-1</sup>	[136]
											LC <sub>50</sub> (48 h) (mortality)	110.1 mg L <sup>-1</sup>	[136]
Tetracycline	60-54-8	Surface water	USA	SPE-LC-MS	100	110	[23]	Duckweed	<i>Lemna minor</i>	EC <sub>50</sub> (7 d) (growth inhibition)	1.06 mg L <sup>-1</sup>	[77]	
Tetracycline		Hospital effluent	Taiwan	SPE-HPLC-MS/MS	2.0	89	[47]	Algae	<i>M. aeruginosa</i>	EC <sub>50</sub> (growth rate)	0.09 mg L <sup>-1</sup>	[121]	
		Pharmaceutical production facility effluent				25							
Tetracycline		STP influent	Taiwan	SPE-HPLC-MS/MS	— <i>f</i>	46–234	[87]		<i>S. capricornutum</i>	EC <sub>50</sub> (growth rate)	2.2 mg L <sup>-1</sup>	[121]	
		STP effluent				16–38							
Tetracycline		STP influent	Luxembourg	SPE-LC-MS/MS	0.3	0.3–85	[89]	Crustacean	<i>D. magna</i>	NOEC (48 h) (immobilization)	340 mg L <sup>-1</sup>	[134]	
		STP effluent				0.3–24							
		Alzette river water				0.3–8							
		Mess river water				0.3–7							
Tetracycline		STP influent	USA	SPE-LC-MS	50	520	[140]			EC <sub>50</sub> (21 d) (reproduction)	44.8 mg L <sup>-1</sup>	[134]	
		STP effluent				170							
Tetracycline		Surface seawater	China (Hong Kong)	SPE-HPLC-MS/MS	13	<13–122	[144]						
<i>Others</i>													
Chloramphenicol	85666-84-8	Victoria Harbour seawater	China	SPE-HPLC-MS	4.1 (LOQ seawater)	ND	[145]						
		Pearl River water			5 (LOQ river water)								
						41–127							
Metronidazole	99616-64-5	STP influent	Taiwan	SPE-HPLC-MS/MS	— <i>f</i>	1–294	[87]	Crustacean	<i>D. magna</i>	LOEC (48 h) (immobilization)	1000 mg L <sup>-1</sup>	[134]	
		STP effluent				10–126				NOEC (21 d) (reproduction)	250 mg L <sup>-1</sup>	[134]	
Trimethoprim	738-70-5	Surface water	USA	SPE-LC-MS	30	150	[23]	Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (15 min)	176.7 mg L <sup>-1</sup>	[82]	
Trimethoprim		Drinking water	USA	SPE-LC-MS/MS	0.25	<0.25	[32]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48 h) (immobilization)	167.4 mg L <sup>-1</sup>	[82]	

Table 3 (Continued)

Compound	CAS number	Sample	Country	Analytical procedure	LOD (ngL <sup>-1</sup> )	Concentration reported (ngL <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.		
Trimethoprim		Danube river water	Serbia	SPE-LC-MS/MS	0.34	25	[84]		<i>D. magna</i>	EC <sub>50</sub> (96h) (immobilization)	120.7 mgL <sup>-1</sup>	[82]		
		Tamis' river water												
		Lake Oc'aga water Groundwater												
Trimethoprim		STP influent	Taiwan	SPE-HPLC-MS/MS	— <i>†</i>	259–949	[87]	Fish	<i>O. latipes</i>	LC <sub>50</sub> (48 h)	>100 mgL <sup>-1</sup>	[82]		
Trimethoprim		STP effluent	South Korea	SPE-LC-MS/MS	1.0	203–415	[90]		<i>O. latipes</i>	LC <sub>50</sub> (96 h)	>100 mgL <sup>-1</sup>	[82]		
Trimethoprim	STP effluent	Surface water				3.2–5.3								
Trimethoprim		STP influent	Korea	SPE-LC-MS	10	<10–496	[93]	Cnidarian	<i>Hydra attenuata</i>	LC <sub>50</sub> (96 h) (morphology)	>100 mgL <sup>-1</sup>	[98]		
Trimethoprim		STP effluent	USA	SPE-HPLC-MS/MS	10	<10–174	[138]			LOEC (96h) (morphology)	>100 mgL <sup>-1</sup>	[98]		
		Han river water				<10–26								
		STP influent				ND–1400								
Trimethoprim		STP effluent	USA	SPE-LC-MS	50	180	[140]			NOEC (96h) (morphology)	>100 mgL <sup>-1</sup>	[98]		
		Hospital effluent				ND–5000								
Trimethoprim		Rio Grande river water				ND								
Trimethoprim		STP influent	USA	SPE-LC-MS	50	330	[140]							
Trimethoprim		Surface seawater	China (Hong Kong)	SPE-HPLC-MS/MS	13	<13–21.8	[144]	Algae	<i>M. aeruginosa</i>	EC <sub>50</sub> (72 h) (growth inhibition)	112 mgL <sup>-1</sup>	[122]		
										<i>R. salina</i>	EC <sub>50</sub> (72 h) (growth inhibition)	16 mgL <sup>-1</sup>	[122]	
											<i>S. capricornutum</i>	EC <sub>50</sub> (72 h) (growth inhibition)	130 mgL <sup>-1</sup>	[122]
											<i>S. capricornutum</i>	EC <sub>50</sub> (growth inhibition)	80.3 mgL <sup>-1</sup>	[131]
									Algae	<i>S. capricornutum</i>	NOEC (growth inhibition)	25.5 mgL <sup>-1</sup>	[131]	
										Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48h) (immobilization)	149 mgL <sup>-1</sup>	[135]
									Crustacean		<i>D. magna</i>	EC <sub>50</sub> (24h) (immobilization)	155.6 mgL <sup>-1</sup>	[135]
										Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48h) (immobilization)	92.0 mgL <sup>-1</sup>	[135]
									<i>M. macrocopa</i>			EC <sub>50</sub> (24h) (immobilization)	144.8 mgL <sup>-1</sup>	[135]
									<i>D. magna</i>		EC <sub>50</sub> (48h) (immobilization)	54.8 mgL <sup>-1</sup>	[135]	
											LOEC (21 d) (reproduction)	20 mgL <sup>-1</sup>	[136]	
									Crustacean	<i>D. magna</i>	NOEC (21 d) (reproduction)	6 mgL <sup>-1</sup>	[136]	

ND—Not detected; *†*—Data not available; SPE—Solid Phase Extraction; GC-MS—Gas Chromatography with Mass Spectrometry Detection; HPLC-MS/MS—High Performance Liquid Chromatography with Tandem Mass Spectrometry Detection; LC-FD—Liquid Chromatography with Fluorescence Detection; LC-MS—Liquid Chromatography with Mass Spectrometry Detection; LC-MS/MS—Liquid Chromatography with Tandem Mass Spectrometry Detection.

4.6 and 40 mg L<sup>-1</sup> respectively [134], while sulfamethazine had an EC<sub>50</sub> of 202 mg L<sup>-1</sup> [135]. Reproduction was also impaired by oxytetracycline, sulfadiazine, tetracycline and tiamulin at concentrations between 5 and 50 mg L<sup>-1</sup>. Oxolinic acid, streptomycin and tylosin were revealed to be lethal after long-term exposure [134]. Chronic toxicity effects were also observed on the reproduction of the crustacean *D. magna*, when were exposed to levofloxacin and clarithromycin, with EC<sub>50</sub> values of 340 and 40 µg L<sup>-1</sup>, respectively [132]. Eleven commonly used antibiotics were evaluated in organisms belonging to different trophic levels (*V. fischeri*, *D. magna*, *Moina macrocopa*, and *O. latipes*). Neomycin showed significant effects on *D. magna* (EC<sub>50</sub> = 42.1 mg L<sup>-1</sup>), *M. macrocopa* (EC<sub>50</sub> = 34.1 mg L<sup>-1</sup>) and *O. latipes* (LC<sub>50</sub> = 80.8 mg L<sup>-1</sup>) while beta-lactam antibiotics (ampicillin and amoxycillin) were the less toxic to all tested organisms [136]. Neomycin showed chronic toxicity by affecting the reproduction and adult survival of *D. magna* and *M. macrocopa* even at low mg L<sup>-1</sup> levels of exposure (EC<sub>50</sub>s of 0.09 and 0.74 mg L<sup>-1</sup>, respectively). Other pharmaceuticals such as sulfathiazole, trimethoprim and enrofloxacin also showed similar effects on those two cladocerans in a dose-dependent manner. Luminescence inhibition on *V. fischeri* occurred after irradiation of tetracycline, proving that photolytic products become more toxic than the parent compound [137]. Antibiotics belonging to different classes have been found in different aquatic environments (Table 3). Lincomycin was detected in hospital and livestock effluents at concentrations of 2 and 6.6 µg L<sup>-1</sup>, respectively [138]. Fluorquinolone antibiotics as ciprofloxacin were found in hospital effluents [138,139] at values between 2 and 11 µg L<sup>-1</sup>, in STP influents (90–1000 ng L<sup>-1</sup>) and effluents (<6–310 ng L<sup>-1</sup>) [138–141] as well as in surface waters, i.e. the Lambro river (Italy) (14.36 ng L<sup>-1</sup>) [24] and Mondego river (Portugal) (79.6–119.2 ng L<sup>-1</sup>) [142]. Enrofloxacin, a fluorquinolone used by the veterinary medicine, was detected in STP influents (121.8–447.1 ng L<sup>-1</sup>) and effluents (53.7–270 ng L<sup>-1</sup>) in Portugal [139] and the US [140] as well as in surface waters from the Mondego river (Portugal) (67.0–102.5 ng L<sup>-1</sup>) [142]. Sulfonamides have been found in several aquatic systems as STP influents and effluents [138,140,141], surface waters [23,143], groundwaters [27,28] and drinking waters [143] in concentrations ranging from ng L<sup>-1</sup> to a few µg L<sup>-1</sup>. Regarding the tetracyclines, oxytetracycline was detected in the Po and Lambro rivers (Italy) at concentrations up to 248.90 and 24.40 ng L<sup>-1</sup> respectively [24], in combination with tetracycline [140] in American STP influent (47 µg L<sup>-1</sup>) and effluent (4.2 µg L<sup>-1</sup>) [140] and in surface waters (340 ng L<sup>-1</sup>) [23]. In addition to aquatic systems, antibiotics belonging to the fluorquinolones class have also been found in sediments at concentrations that can reach 4.8 mg kg<sup>-1</sup> [141]. This finding may represent a potential risk warning of persistence in the environment.

#### 4.4. Sex hormones

Sex hormones are extremely active biological compounds producing intense therapeutic effects even at very low doses. Today, they are commonly prescribed as oral contraceptives thus indirectly contributing to the increase in environmental concentrations [52,108]. Estrogens are the sex hormones most commonly found in the environment. These can exist as either natural or synthetic substances, mimicking the effects of endogenous estrogens as endocrine-disrupting compounds (EDCs) [146] through binding to specific receptors common to non-target organisms (invertebrates, fish, reptiles, birds and mammals) [108]. In fish, estrogens are involved in several physiological functions including: (i) vitellogenin synthesis; (ii) vitelline envelope (eggshell) protein production; (iii) gonadal differentiation; (iv) development of secondary sexual characteristics; (v) gonadotropin secretion; (vi) synthesis of estrogen receptors; (vii) pheromonal communi-

cation; (viii) bone formation; and (ix) calcium homeostasis [146]. The enhanced production of the vitellogenin found in the blood of male and juvenile fish provides a useful biomarker of aquatic contamination by compounds with estrogenic activity [52,146]. Wild fish (roach; *Rutilus rutilus*) exposed to such compounds in UK rivers receiving STP effluents suffered adverse reproductive effects. Male fish were shown to be intersex, i.e. they had simultaneous male and female gonadal characteristics besides a high plasma vitellogenin concentration [147]. Ethinylestradiol (EE<sub>2</sub>) is a synthetic estrogen found in oral contraceptive pills with marked estrogenic effects in fish. The life-cycle exposure of fathead minnows to EE<sub>2</sub> concentrations below 1 ng L<sup>-1</sup> caused a significant reduction in fertilization success, an increased egg production and decreased expression of secondary male sex characteristics [148]. Similar findings were obtained by Pawlowski et al. [149] in trials extended over a reduced period of three weeks. Concentrations below 1 ng L<sup>-1</sup> gave rise to an increased female population and for EE<sub>2</sub> concentrations above 3.5 ng L<sup>-1</sup>, fish became completely feminized [148]. Concentrations above 1 ng L<sup>-1</sup> of EE<sub>2</sub> also induced higher vitellogenin plasma levels in both males and females [149,150]. Nash et al. [151] registered similar findings for zebrafish males by simply performing the assay with 0.5 ng L<sup>-1</sup> of EE<sub>2</sub>. Life-long exposure of zebrafish to 5 ng L<sup>-1</sup> of EE<sub>2</sub> has led to reproductive failure due to the absence of secondary male sex characteristics and normal testes [151]. Exposure of juveniles to estrogen has caused skewed sex ratios in favour of females for concentrations of 1 ng L<sup>-1</sup> [150]. Sex reversal was complete at levels of 2 ng L<sup>-1</sup> [150]. Xu et al. [152] also exposed zebrafish to EE<sub>2</sub> during their period of sex differentiation, showing that, after 90 days post-hatch, there was already an increase in mortality rate and sex ratio for fish exposed to concentrations of 2 ng L<sup>-1</sup>. When the concentration was increased to 10 ng L<sup>-1</sup> was observed a significantly decrease in the weight and length body. On the other hand, 180 days post-hatch were found abnormal testicular morphologies in male fish, namely malformations of the sperm duct, an altered proportion of germ cell types, and a reduced number of spermatozoa, for those levels of EE<sub>2</sub> [152]. Exposure of male roach to EE<sub>2</sub> concentrations up to 4 ng L<sup>-1</sup> in early life disrupted normal sexual development causing a feminized response, characterized by the presence of an ovarian cavity and induced plasma vitellogenin production [153]. Kidd et al. [34] conducted a 7-year, whole-lake experiment, proving that chronic exposure of fathead minnow to concentrations of EE<sub>2</sub> in the order of 5–6 ng L<sup>-1</sup>, led to feminization of males fish, through production of vitellogenin and disruption in gonadal development, causing intersex, and altered oogenesis in females. Those reproductive alterations led to a collapse of the fathead minnow population due to the loss of the young generations, expressed in a loss of smaller sizes classes of fish, what contribute, in a last case, to leave this species from the lake near of extinction [34]. The natural estrogen 17β-estradiol (E<sub>2</sub>) can also negatively affect fish at low concentrations. Japanese medaka exposed to 33.5 ng L<sup>-1</sup> of this estrogen in early life enhanced their body length and body weight. Additionally, the males also exhibited testis-ova after 14 days of exposure [154]. When the E<sub>2</sub> concentration was increased to 140.6 ng L<sup>-1</sup>, testis-ova were observed in males (after 12 days exposure) and complete gonadal transformation to an ovary occurred after 20 days [154]. The exposure of adult fish to concentrations from 29.3 to 463 ng L<sup>-1</sup> over 21 days gave rise to testis-ova development and induced vitellogenin production in males to all tested concentrations [155]. At the higher level, a decrease in the number of eggs produced and fertility [155] was also observed. Amphibians and reptiles exposed to environmental estrogens showed sex reversal as well as significant changes in secondary sex characteristics [156,157]. Concerning invertebrates such as the amphipod *Hyalella azteca* it was observed that at sub-lethal concentrations of EE<sub>2</sub> (0.1–10 µg L<sup>-1</sup>) sexual development of males was affected

Table 4

Examples of concentrations ( $\text{ng L}^{-1}$ ) of sex hormones measured in different aquatic environments.

Compound	CAS number	Sample	Country	Analytical procedure	LOD ( $\text{ng L}^{-1}$ )	Concentration reported ( $\text{ng L}^{-1}$ )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
Diethylstilbestrol	8053-00-7	River water	China	SPME-GC-MS	2	20 ( $\pm$ 0)	[162]					
17 $\alpha$ -Estradiol	57-91-0	Surface water	USA	LLE-GC-MS	5	30	[23]					
17 $\alpha$ -Estradiol		Groundwater	France	SPE-LC-MS/MS	0.03	0.8–3.5	[164]					
17 $\beta$ -Estradiol	50-28-2	Surface water	USA	LLE-GC-MS	5	9	[23]	Fish	<i>O. latipes</i>	NOEC (21 d) (testis-ova induction)	<29.3 $\text{ng L}^{-1}$	[15]5
17 $\beta$ -Estradiol		Drinking water	USA	SPE-LC-MS/MS	0.50	<0.50	[32]			LOEC (21 d) (testis-ova induction)	<26.3 $\text{ng L}^{-1}$	[155]
17 $\beta$ -Estradiol		Hospital effluent	Taiwan	SPE-HPLC-MS/MS	25	25	[47]			NOEC (21 d) (VTG induction)	29.3 $\text{ng L}^{-1}$	[155]
		Pharmaceutical production facility effluent				ND						
17 $\beta$ -Estradiol		STP influent	Japan	SPE-GC-MS	0.1 (LOQ)	13.3–25.8	[86]					
		STP effluent				0.49–12.4						
17 $\beta$ -Estradiol		Pearl Rivers water	China	SPE-GC-MS	0.3	ND–7.5 ( $\pm$ 0.4)	[88]					
17 $\beta$ -Estradiol		STP influent	Luxembourg	SPE-LC-MS/MS	1.0	1.0–102	[89]					
		STP effluent				1.0–85						
		Alzette river water				1.0–35						
		Mess river water				1.0–6						
17 $\beta$ -Estradiol		STP effluent	South Korea	SPE-LC-MS/MS	1.0	<1.0	[90]					
		Surface water				ND						
17 $\beta$ -Estradiol		STP effluent	Japan	SPE-LC-MS/MS	0.3	0.3–2.5	[160]					
		Tamagawa river water				0.6–1.0						
		Kasumigaura lake water										
17 $\beta$ -Estradiol		STP influent	Germany	SPE-LC-MS/MS	2.0 (LOQ STP influent)	<0.3	[161]					
		STP effluent			0.4 (LOQ STP effluent)	11.8 ( $\pm$ 5.1)						
		Berlin surface water			0.2 (LOQ surface water)	0.8 ( $\pm$ 0.3)						
						<0.2						
17 $\beta$ -Estradiol		River water	China	SPME-GC-MS	9	100 ( $\pm$ 20)	[162]					
17 $\beta$ -Estradiol		STP influent	Italy	SPE-LC-MS/MS	1.9 (STP influent)	10–31	[163]					
		STP effluent			0.8 (STP effluent)	3–8						
		Tibre river water			0.2 (Tibre river water)	2–6						
17 $\beta$ -Estradiol		Groundwater	France	SPE-LC-MS/MS	0.01	0.3–1.3	[164]					
Estriol	50-27-1	Surface water	USA	LLE-GC-MS	5	19	[23]					
Estriol		STP influent	Japan	SPE-GC-MS	0.2 (LOQ)	83.0–255	[86]					
		STP effluent				0.31–0.84						
Estriol		STP effluent	South Korea	SPE-LC-MS/MS	5.0	8.9–25	[90]					
		Surface water				ND						
Estriol		STP influent	Italy	SPE-LC-MS/MS	7.0 (STP influent)	23–48	[163]					
		STP effluent			0.5 (STP effluent)	ND–1						
		Tibre river water			0.3 (Tibre river water)	2–5						
Estrone	53-16-7	Surface water	USA	LLE-GC-MS	5	27	[23]					
Estrone		Drinking water	USA	SPE-LC-MS/MS	0.20	<0.20	[32]					

Estrone		Hospital effluent Pharmaceutical production facility effluent STP influent	Taiwan	SPE-HPLC- MS/MS	25	126 ND	[47]					
Estrone		STP effluent	Japan	SPE-GC-MS	0.6 (LOQ)	28.7–197	[86]					
Estrone		Pearl Rivers water	China	SPE-GC-MS	0.2	2.80–110 ND–75.0 ( $\pm 5.3$ )	[88]					
Estrone		STP influent STP effluent Alzette river water Mess river water	Luxembourg	SPE-LC-MS/MS	0.3	0.3–9 0.3–14 0.3–6 0.3–27	[89]					
Estrone		STP effluent Surface water	South Korea	SPE-LC-MS/MS	1.0	2.2–36 1.7–5.0	[90]					
Estrone		STP effluent Tamagawa river water Kasumigaura lake water	Japan	SPE-LC-MS/MS	0.1	2.5–34 3.4–6.6 0.2–0.8	[160]					
Estrone		STP influent STP effluent Berlin surface water	Germany	SPE-LC-MS/MS	1.0 (LOQ STP influent) 0.2 (LOQ STP effluent) 0.1 (LOQ surface water)	188 ( $\pm 92$ ) 12.6 ( $\pm 7.0$ ) 0.16 ( $\pm 0.05$ )	[161]					
Estrone		River water	China	SPME-GC-MS	18	180 ( $\pm 20$ )	[162]					
Estrone		STP influent STP effluent Tibre river water	Italy	SPE-LC-MS/MS	1.2 (STP influent) 0.8 (STP effluent) 0.1 (Tibre river water)	15–60 5–30 5–12	[163]					
Estrone		Surface water Groundwater	France	SPE-LC-MS/MS	0.02	0.3 0.8–3.5	[164]					
17 $\alpha$ -Ethinylestradiol	57-63-6	Surface water	USA	LLE-GC-MS	5	73	[23]	Fish	<i>P. promelas</i>	LOEC (21 d) (plasma VTG induction)	1 ngL <sup>-1</sup>	[149]
17 $\alpha$ -Ethinylestradiol		Drinking water	USA	SPE-LC-MS/MS	1.0	<1.0	[32]			LOEC (21 d) (ultrastructure testes)	1 ngL <sup>-1</sup>	[149]
17 $\alpha$ -Ethinylestradiol		Hospital effluent Pharmaceutical production facility effluent	Taiwan	SPE- HPLC-MS/MS	25	32 ND	[47]			LOEC (21 d) (ultrastructure liver)	1 ngL <sup>-1</sup>	[149]
17 $\alpha$ -Ethinylestradiol		STP influent STP effluent Alzette river water Mess river water	Luxembourg	SPE-LC-MS/MS	2.0	2.0–24 <2.0 <2.0 <2.0	[89]			LOEC (21 d) (fertilization rate)	10 ngL <sup>-1</sup>	[149]
17 $\alpha$ -Ethinylestradiol		STP effluent	South Korea	SPE-LC-MS/MS	1.0	1.3	[90]	Fish	<i>D. rerio</i>	LOEC (38 dph) (plasma VTG induction)	2 ngL <sup>-1</sup>	[150]

Table 4 (Continued)

Compound	CAS number	Sample	Country	Analytical procedure	LOD (ng L <sup>-1</sup> )	Concentration reported (ng L <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
17 $\alpha$ -Ethinylestradiol		Surface water	Germany	SPE-LC-MS/MS	2.0 (LOQ STP influent)	ND	[161]					
		STP influent				8.8 ( $\pm$ 8.0)						
		STP effluent Berlin surface water				1.7 ( $\pm$ 1.3)						
17 $\alpha$ -Ethinylestradiol		STP influent	Italy	SPE-LC-MS/MS	1.6 (STP influent)	ND	[163]					
		STP effluent				ND						
		Tibre river				ND-1						
17 $\alpha$ -Ethinylestradiol Mestranol	72-33-3	Groundwater	France USA	SPE-LC-MS/MS LLE-GC-MS	0.20 5	0.5-3.0	[164] [23]					
		Surface water				74						

ND—Not detected; SPE—Solid Phase Extraction; SPME—Solid Phase Microextraction; LLE—Liquid-Liquid Extraction; GC-MS—Gas Chromatography with Mass Spectrometry Detection; LC-MS/MS—Liquid Chromatography with Tandem Mass Spectrometry Detection; dph—days post-hatch.

[158]. On the other hand, the estrogens E<sub>2</sub> and EE<sub>2</sub> did not show significant effects on reproduction or survival of *C. dubia* even at concentrations of 1 and 0.5 mg L<sup>-1</sup>, respectively [159]. According to many authors, the concentrations of estrogens detected in the environment may not pose a threat to humans. However regarding these compounds, there is the possibility of bioaccumulation within aquatic organisms, thereby reaching humans through the food chain or directly through drinking water [146]. Estrogens have been found in water samples (Table 4) at low ng L<sup>-1</sup> concentrations but they represent a greater risk for non-target organisms as already proved. For example, 17 $\beta$ -estradiol was detected in rivers [23,160–163] at levels ranging from 0.6 to 100 ng L<sup>-1</sup> and in STP effluents at concentrations between 0.3 [160] and 85 ng L<sup>-1</sup> [89]. Ethinylestradiol was also found in surface waters in the US (73 ng L<sup>-1</sup>) [23] and Italy (the Tibre river) at 1 ng L<sup>-1</sup> [163].

#### 4.5. Antiepileptics

Antiepileptic drugs act in the central nervous system (CNS) by reducing the overall neuronal activity. This can be achieved either by blocking voltage-dependent sodium channels (e.g. carbamazepine) or by enhancement of the inhibitory effects of the  $\gamma$ -aminobutyric acid (GABA) neurotransmitter (e.g. benzodiazepines) [99]. Carbamazepine is carcinogenic to rats but does not have mutagenic properties in mammals [165]. Moreover, this drug is lethal to zebrafish at the 43  $\mu$ g L<sup>-1</sup> level and produces sub-lethal changes in *Daphnia* sp. at 92  $\mu$ g L<sup>-1</sup> [165]. Regarding aquatic organisms, it can be deduced that carbamazepine does have harmful proclivity since most of the acute toxicity data were harvested from trial concentrations between 10 and 100 mg L<sup>-1</sup> [98]. In fact, *D. magna* growth was shown to be sensitive to this compound, being inhibited for concentrations of carbamazepine above 12.7 mg L<sup>-1</sup> and with acute toxicity being evident at 17.2 mg L<sup>-1</sup> [165]. The EC<sub>50</sub> value (considering the motility as indicator) was approximately 13.8 mg L<sup>-1</sup> after 48 h of exposure [96]. Female *D. pulex* exposed to 1  $\mu$ g L<sup>-1</sup> of carbamazepine showed a tendency to mature and reproduce earlier (with more offspring), suggesting that this pharmaceutical may slightly induce stimulatory effects [166]. For *C. dubia*, chronic toxicity studies revealed a NOEC of 25  $\mu$ g L<sup>-1</sup> [96] while the activity of *G. pulex* was slightly reduced by exposure to a concentration range from 1 to 10 ng L<sup>-1</sup> [76]. Continuous exposure of *H. attenuata* to carbamazepine caused a significant reduction in feeding, with an EC<sub>50</sub> of 3.76 mg L<sup>-1</sup> [98]. Japanese medaka showed a LC<sub>50</sub> of 35.4 mg L<sup>-1</sup> [82] and ultrastructural changes in the liver, kidney and gill tissues of carps were induced by this pharmaceutical [97]. The changes observed in the kidney were shown to occur as a cellular response to impaired kidney function. In gills, the effects were more pronounced for concentrations above 20  $\mu$ g L<sup>-1</sup>. Another important issue concerning carbamazepine is that it can adsorb to sediments, in this way threatening aquatic organisms which feed on organic matter. Oetken et al. [167] showed that exposure of the invertebrate *Chironomus riparius* to this pharmaceutical through sediments caused a blockade of pupation and decreased emergence with EC<sub>50</sub> values of 160 and 280  $\mu$ g kg<sup>-1</sup> of dry weight, respectively. Carbamazepine is ubiquitously present in the environment, having an extremely low removal rate in STPs (7%) [54] and consequently being detected in rivers [16,20,21,54,92] at concentrations up to 595 ng L<sup>-1</sup> [92] (Table 5). In addition to surface waters, carbamazepine has also been found in groundwater [26,119] at concentrations that can reach 900 ng L<sup>-1</sup>. A monitoring programme performed on the river Rhine (Germany) over a decade, showed the regular detection of carbamazepine, with an annual average concentration of 100 ng L<sup>-1</sup> [168]. These results support the idea that the presence of carbamazepine in the environment may represent a real threat.



Table 5

Examples of concentrations (ng L<sup>-1</sup>) of antiepileptic drugs measured in different aquatic environments.

Compound	CAS number	Sample	Country	Analytical procedure	LOD (ng L <sup>-1</sup> )	Concentration reported (ng L <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
Carbamazepine	298-46-4	STP influent	Spain	SPE-GC-MS	30	120–310	[14]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48 h) (immobilization)	>100 mg L <sup>-1</sup>	[65]
Carbamazepine		STP effluent	Finland	SPE-HPLC-MS/MS	1.4	110–230	[16]	Algae	<i>D. subspicatus</i>	EC <sub>50</sub> (growth inhibition)	74 mg L <sup>-1</sup>	[65]
Carbamazepine		STP influent				290–400						
		STP effluent	Romania	SPE-GC-MS	30	380–470	[20]	Duckweed	<i>L. minor</i>	EC <sub>50</sub> (7 d) (growth inhibition)	25.5 mg L <sup>-1</sup>	[65]
Carbamazepine		Vantaa river water				<1.4–66						
		Luhtajoki river water				23						
Carbamazepine		Somes river water				<30–75.1 (±6.1)	[20]					
Carbamazepine		STP influent	Sweden	SPE-LC-MS/MS	— <sup>†</sup>	1680	[21]	Crustacean	<i>Gammarus pulex</i>	LOEC (behaviour)	10 ng L <sup>-1</sup>	[76]
		STP effluent				1180						
		Höje river water				<1–500						
Carbamazepine		Groundwater	Germany	SPE-GC-MS	32	900	[26]	Crustacean	<i>T. platyurus</i>	LC <sub>50</sub> (24 h) (mortality)	>100 mg L <sup>-1</sup>	[78]
Carbamazepine		Drinking water	USA	SPE-LC-MS/MS	0.5	6.8	[32]	Fish	<i>O. latipes</i>	LC <sub>50</sub> (96 h) (mortality)	45.87 mg L <sup>-1</sup>	[78]
Carbamazepine		STP effluent	Germany	SPE-LC-MS/MS	50 (STP effluent)	2100	[54]	Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (15 min)	52.2 mg L <sup>-1</sup>	[82]
		Surface water			30 (surface water)	250						
Carbamazepine		Hospital effluent	Spain	SPE-HPLC-MS/MS	7	30–70	[73]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48 h) (immobilization)	>100 mg L <sup>-1</sup>	[82]
Carbamazepine		Danube river water	Serbia	SPE-LC-MS/MS	0.27	8–130	[84]		<i>D. magna</i>	EC <sub>50</sub> (96 h) (immobilization)	76.3 mg L <sup>-1</sup>	[82]
		Sava river water				29–50						
		Tamis <sup>†</sup> river water				30						
		Lake Oc <sup>†</sup> aga water				30						
Carbamazepine		Groundwater				6–23						
		STP influent	Japan	SPE-GC-MS	6	14.9–270	[86]	Fish	<i>O. latipes</i>	LC <sub>50</sub> (48 h)	35.4 mg L <sup>-1</sup>	[82]
		STP effluent				10.8–163						
Carbamazepine		STP influent	Taiwan	SPE-HPLC-MS/MS	— <sup>†</sup>	82–357	[87]		<i>O. latipes</i>	LC <sub>50</sub> (96 h)	35.4 mg L <sup>-1</sup>	[82]
Carbamazepine		STP effluent	South Korea	SPE-LC-MS/MS	1.0	93–214	[90]	Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (30 min)	>81,000 μg L <sup>-1</sup>	[96]
		STP effluent				73–729						
		Surface water				4.5–61						
		Drinking water				<1.0						
Carbamazepine		Mankyung river water	South Korea	SPE-LC-MS/MS	1	ND–595 (±14)	[92]	Algae	<i>P. subcapitata</i>	NOEC (96 h) (growth inhibition)	>100,000 μg L <sup>-1</sup>	[96]
Carbamazepine		STP influent	Korea	SPE-LC-MS	5	<5–451	[93]			LOEC (96 h) (growth inhibition)	>100,000 μg L <sup>-1</sup>	[96]
		STP effluent				<5–195						
		Han river water				<5–36						
Carbamazepine		STP effluent	Italy	SPE-HPLC-MS/MS	1.3	ND–1318	[118]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48 h) (immobilization)	>13,800 μg L <sup>-1</sup>	[96]
Carbamazepine		Groundwater	Germany	SPE-GC-MS	2 (LOQ)	45	[119]		<i>C. dubia</i>	EC <sub>50</sub> (48 h) (immobilization)	77,700 μg L <sup>-1</sup>	[96]
Carbamazepine		STP influent	France	SPE-LC-MS	2.4	193–420	[169]			NOEC (7 d) (reproduction)	25 μg L <sup>-1</sup>	[96]
		STP effluent				86–258						

Table 5 (Continued)

Compound	CAS number	Sample	Country	Analytical procedure	LOD (ng L <sup>-1</sup> )	Concentration reported (ng L <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
Carbamazepine-10,11-epoxide*	-†	STP influent STP effluent	Spain	SPE-GC-MS	70	300–500 <70–300	[14]	Fish	<i>D. rerio</i>	LOEC (7 d) (reproduction) NOEC (10 d) (survival)	100 µg L <sup>-1</sup> 25,000 µg L <sup>-1</sup>	[96]
Carbamazepine-10,11-epoxide*		STP influent STP effluent	France	SPE-LC-MS	5.2	ND–27 <5.2–29	[169]	Fish	<i>O. mykiss</i>	LOEC (10 d) (survival) LOEC (21 d) (liver cytopathology) LOEC (21 d) (kidney cytopathology)	50,000 µg L <sup>-1</sup> >100 µg L <sup>-1</sup> 1 µg L <sup>-1</sup>	[96] [97] [97]
			Cnidarian						<i>Hydra attenuata</i>	LC <sub>50</sub> (96 h) (morphology) EC <sub>50</sub> (96 h) (morphology) LOEC (96 h) (morphology) NOEC (96 h) (morphology) FC <sub>50</sub> (96 h) (fecundity)	29.4 mg L <sup>-1</sup> 15.52 mg L <sup>-1</sup> 5 mg L <sup>-1</sup> 1 mg L <sup>-1</sup> 3.76 mg L <sup>-1</sup>	[98] [98] [98] [98] [98]

\* – Metabolite; † – Data not available; ND – Not detected; SPE – Solid Phase Extraction; GC-MS – Gas Chromatography with Mass Spectrometry Detection; HPLC-MS/MS – High Performance Liquid Chromatography with Tandem Mass Spectrometry Detection; LC-MS – Liquid Chromatography with Mass Spectrometry Detection; IC-MS/MS – Liquid Chromatography with Tandem Mass Spectrometry Detection.

#### 4.6. Beta-blockers

Beta-blockers act by competitive inhibition of beta-adrenergic receptors, a class of receptors critical for normal functioning in the sympathetic branch of the vertebrate autonomic nervous system in vertebrates. Within the most commonly used  $\beta$ -blockers propranolol is a non-specific antagonist, blocking both  $\beta_1$  and  $\beta_2$ -receptors while metoprolol and atenolol present  $\beta_1$ -receptor specificity [99]. Fish, like other vertebrates, possess  $\beta$ -receptors in the heart, liver and reproductive system [170,171] so that prolonged exposure to drugs belonging to this therapeutic class may cause deleterious effects. From a two weeks study, it was observed that exposure to 500 µg L<sup>-1</sup> of propranolol reduced growth rates of Japanese medaka [172]. Plasma steroid levels were altered in both male and female fish even at concentrations as low as 1 µg L<sup>-1</sup> propranolol. Exposure to concentrations of 0.5 and 1 µg L<sup>-1</sup> resulted in a decreased egg production. On the other hand, acute exposure of rainbow trout to 70.9 µg L<sup>-1</sup> of propranolol showed no significant reduction in its heart rate [173]. However, for concentrations of metoprolol of 1 µg L<sup>-1</sup>, ultrastructural changes in the liver and kidney were observed as well in gills if the concentration rose above 20 µg L<sup>-1</sup> [97]. Fathead minnows exposed to atenolol during embryo-larval development showed NOEC and LOEC values for growth rate of 3.2 and 10 mg L<sup>-1</sup>, respectively [174]. Furthermore, a reproduction study performed in adults over a 21-day exposure period demonstrated that the male fish condition index was the most sensitive endpoint with NOEC and LOEC values of 1.0 and 3.2 mg L<sup>-1</sup>, respectively [174]. These data suggest that atenolol has a low chronic toxicity to fish when compared to propranolol.

As invertebrates do not possess  $\beta$ -receptors a different potential impact on these organisms would be expected. Accordingly, the acute toxicity of propranolol, metoprolol and nadolol was assessed on the invertebrates *H. azteca*, *D. magna*, *D. lumholtzi* and *C. dubia*. Following a 48-h exposure to propranolol, LC<sub>50</sub> values of 29.8, 1.6 and 0.8 mg L<sup>-1</sup> were obtained for *H. azteca*, *D. magna* and *C. dubia* respectively [172] while acute exposure to nadolol did not affect the survival of the invertebrates [172]. Regarding metoprolol, *D. magna* and *C. dubia* exhibited LC<sub>50</sub> values of 63.9 and 8.8 mg L<sup>-1</sup>, respectively [172]. However, Cleuvers [175] obtained a higher EC<sub>50</sub> value (438 mg L<sup>-1</sup>) in an acute toxicity test performed on *D. magna*. Reproduction in invertebrates decreased following propranolol exposure with NOEC values of 1 and 125 µg L<sup>-1</sup> for *H. azteca* and *C. dubia* respectively [172]. Propranolol inhibited the growth of the green algae *Desmodesmus subspicatus*, showing an EC<sub>50</sub> of 7.7 mg L<sup>-1</sup> [175] while atenolol almost failed to register a toxic effect (EC<sub>50</sub> of 620 mg L<sup>-1</sup>). Chronic exposure of *D. magna* to propranolol (9 days) resulted in a significant reduction in heart rate, fecundity and biomass with LOECs values of 55, 110 and 440 µg L<sup>-1</sup> respectively [176] while chronic exposure to metoprolol showed LOECs of 12.5 mg L<sup>-1</sup> (body mass) and 6.15 mg L<sup>-1</sup> (reproduction). At the highest concentrations (25 and 50 mg L<sup>-1</sup>) reproduction ceased and at the highest levels, all organisms died before the end of the test. A reduced heart rate for *D. magna* was evident for a 3.2 mg L<sup>-1</sup> level of metoprolol. Chronic toxicity tests performed in algae also evidenced their sensitivity to  $\beta$ -blockers with NOEC values below 1 mg L<sup>-1</sup> [52].

Collectively, this data might indicate a possible environmental risk since propranolol has been detected in STP effluents [21,53,94] at concentrations from 30 to 373 ng L<sup>-1</sup> and in surface waters [21,53,92,94] at levels of ng L<sup>-1</sup> (Table 6). This pharmaceutical has also been found in hospital effluent (Spain) at concentrations that can reach 6.5 µg L<sup>-1</sup> [73]. Other  $\beta$ -blockers such as atenolol, metoprolol and solatol have also been detected in environmental samples [16,21,24,73,118] including groundwater [26] at concentrations up to 122 µg L<sup>-1</sup>.

**Table 6**

Examples of concentrations (ng L<sup>-1</sup>) of  $\beta$ -blockers agents measured in different aquatic environments.

Compound	CAS number	Sample	Country	Analytical procedure	LOD (ng L <sup>-1</sup> )	Concentration reported (ng L <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
Acebutolol	37517-30-9	STP influent	Finland	SPE-HPLC-MS/MS	0.8	390–510	[16]					
		STP effluent				80–230						
		Vantaa river water				<0.8–8						
Atenolol	29122-68-7	Luhtajoki river water	Finland	SPE-HPLC-MS/MS	11.8	8	[16]	Crustacean	<i>T. platyurus</i>	LC <sub>50</sub> (24 h) (mortality)	>100 mg L <sup>-1</sup>	[78]
		STP influent				510–800						
		STP effluent				40–440						
Atenolol		Vantaa river water	Sweden	SPE-LC-MS/MS	— <i>f</i>	<11.8–25	[21]	Fish	<i>O. latipes</i>	LC <sub>50</sub> (96 h) (mortality)	>100 mg L <sup>-1</sup>	[78]
		Luhtajoki river water				30						
		STP influent				160						
Atenolol		Höje river water	Italy	SPE-HPLC-MS/MS	0.3 (LOQ)	10–60	[24]	Algae	<i>D. subspicatus</i>	EC <sub>50</sub> (growth inhibition)	620 mg L <sup>-1</sup>	[17]5
		Po river water				3.44–39.43						
		Lambro river water				241						
Atenolol		Drinking water	USA	SPE-LC-MS/MS	0.25	0.47	[32]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48h) (immobilization)	313 mg L <sup>-1</sup>	[175]
Atenolol		Hospital effluent	Spain	SPE-HPLC-MS/MS	28	100–122,000	[73]	Fish	<i>P. promelas</i>	NOEC (28 d) (growth)	3.2 mg L <sup>-1</sup>	[174]
Atenolol		Mankyung river water	South Korea	SPE-LC-MS/MS	30	ND–690 (±26)	[92]			LOEC (28 d) (growth)	10 mg L <sup>-1</sup>	[174]
Atenolol		STP influent	Taiwan	SPE-HPLC-MS/MS	— <i>f</i>	738–2883	[87]			NOEC (21 d) (condition index)	1.0 mg L <sup>-1</sup>	[174]
Atenolol		STP effluent	Italy	SPE-HPLC-MS/MS	1.07 (LOQ)	210–681	[118]			LOEC (21 d) (condition index)	3.2 mg L <sup>-1</sup>	[174]
		STP effluent				27–1168				NOEC (21 d) (reproduction)	10 mg L <sup>-1</sup>	[174]
										LOEC (21 d) (reproduction)	>10 mg L <sup>-1</sup>	[174]
Metoprolol	83-43-2	STP influent	Finland	SPE-HPLC-MS/MS	3.8	980–1350	[16]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48h) (immobilization)	>100 mg L <sup>-1</sup>	[65]
		STP effluent				910–1070						
		Vantaa river water				<3.8–116						
Metoprolol		Luhtajoki river water	Sweden	SPE-LC-MS/MS	— <i>f</i>	38	[21]	Algae	<i>D. subspicatus</i>	EC <sub>50</sub> (growth inhibition)	7.3 mg L <sup>-1</sup>	[65]
		STP influent				160						
		STP effluent				190						
Metoprolol		Höje river water	Taiwan	SPE-HPLC-MS/MS	— <i>f</i>	30–70	[87]	Duckweed	<i>L. minor</i>	EC <sub>50</sub> (7 d) (growth inhibition)	>320 mg L <sup>-1</sup>	[65]
		STP influent				14–597						

Table 6 (Continued)

Compound	CAS number	Sample	Country	Analytical procedure	LOD (ngL <sup>-1</sup> )	Concentration reported (ngL <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
		STP effluent				12–199		Fish	<i>O. mykiss</i>	LOEC (21 d) (liver cytopathology)	1 μg L <sup>-1</sup>	[97]
										LOEC (21 d) (gills cytopathology)	20 μg L <sup>-1</sup>	[97]
								Crustacean	<i>H. azteca</i>	LC <sub>50</sub> (48 h) (mortality)	>100 mg L <sup>-1</sup>	[172]
									<i>C. dudia</i>	LC <sub>50</sub> (48 h) (mortality)	8.8 mg L <sup>-1</sup>	[172]
									<i>D. magna</i>	LC <sub>50</sub> (48 h) (mortality)	63.9 mg L <sup>-1</sup>	[172]
								Fish	<i>O. latipes</i>	LC <sub>50</sub> (48 h) (mortality)	>100 mg L <sup>-1</sup>	[172]
								Algae	<i>D. subspicatus</i>	EC <sub>50</sub> (48 h) (growth inhibition)	7.9 mg L <sup>-1</sup>	[177]
								Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48 h) (immobilization)	438 mg L <sup>-1</sup>	[175]
								Crustacean	<i>D. magna</i>	NOEC (9 d) (body mass)	6.15 mg L <sup>-1</sup>	[176]
										LOEC (9 d) (body mass)	12.5 mg L <sup>-1</sup>	[176]
										LOEC (9 d) (reproduction)	6.15 mg L <sup>-1</sup>	[176]
										LOEC (9 d) (heart rate)	3.2 mg L <sup>-1</sup>	[176]
Propranolol	525-66-6	STP influent	Sweden	SPE-LC-MS/MS	— <sup>†</sup>	50	[21]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48 h) (immobilization)	7.5 mg L <sup>-1</sup>	[65]
		STP effluent				30						
		Höje river water				<1–10						
Propranolol		Hospital effluent	Taiwan	SPE-HPLC-MS/MS	0.5	54	[47]	Algae	<i>D. subspicatus</i>	EC <sub>50</sub> (growth inhibition)	5.8 mg L <sup>-1</sup>	[65]
		Pharmaceutical production facility effluent				ND						
Propranolol		STP influent	United Kingdom	SPE-HPLC-MS/MS	10	60–119	[53]	Duckweed	<i>L. minor</i>	EC <sub>50</sub> (7 d) (growth inhibition)	114 mg L <sup>-1</sup>	[65]
		STP effluent				195–373						
		Tyne river water				35–107						
Propranolol		Hospital effluent	Spain	SPE-HPLC-MS/MS	8	200–6500	[73]	Crustacean	<i>T. platyurus</i>	LC <sub>50</sub> (24 h) (mortality)	10.31 mg L <sup>-1</sup>	[78]
Propranolol		Mankyung river water	South Korea	SPE-LC-MS/MS	10	ND–40.1 (±3)	[92]	Fish	<i>O. latipes</i>	LC <sub>50</sub> (96 h) (mortality)	11.40 mg L <sup>-1</sup>	[78]
Propranolol		STP effluent	United Kingdom	SPE-HPLC-MS/MS		130–180	[94]	Crustacean	<i>H. azteca</i>	LC <sub>50</sub> (48 h) (mortality)	29.8 mg L <sup>-1</sup>	[172]
		Surface water				<10–37				NOEC (27 d) (reproduction)	0.001 mg L <sup>-1</sup>	[172]

									LOEC (27 d) (reproduction)	0.1 mgL <sup>-1</sup>	[172]	
									<i>C. dudia</i>	LC <sub>50</sub> (48 h) (mortality)	0.8 mgL <sup>-1</sup>	[172]
										NOEC (7 d) (reproduction)	0.125 mgL <sup>-1</sup>	[172]
										LOEC (7 d) (reproduction)	0.25 mgL <sup>-1</sup>	[172]
									<i>D. magna</i>	LC <sub>50</sub> (48 h) (mortality)	1.6 mgL <sup>-1</sup>	[172]
								Fish	<i>O. latipes</i>	LC <sub>50</sub> (48 h) (mortality)	24.3 mgL <sup>-1</sup>	[172]
								Algae	<i>D. subspicatus</i>	EC <sub>50</sub> (48 h) (growth inhibition)	0.7 mgL <sup>-1</sup>	[175]
								Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48 h) (immobilization)	7.7 mgL <sup>-1</sup>	[175]
								Duckweed	<i>L. minor</i>	EC <sub>50</sub> (growth rate)	113 mgL <sup>-1</sup>	[175]
								Crustacean	<i>D. magna</i>	NOEC (9 d) (body mass)	0.22 mgL <sup>-1</sup>	[176]
										LOEC (9 d) (body mass)	0.44 mgL <sup>-1</sup>	[176]
										NOEC (9 d) (reproduction)	0.055 mgL <sup>-1</sup>	[176]
										LOEC (9 d) (reproduction)	0.11 mgL <sup>-1</sup>	[176]
										LOEC (9 d) (heart rate)	0.055 mgL <sup>-1</sup>	[176]
Sotalol	959-24-0	STP influent	Finland	SPE- HPLC-MS/MS	3.9	640–830	[16]					
		STP effluent				160–300						
		Vantaa river water				<3.9–52						
		Luhtajoki river water				37						
Sotalol		Groundwater	Germany	SPE- HPLC-MS/MS	8.0	560	[26]					

†—Data not available; ND—Not Detected; SPE—Solid Phase Extraction; HPLC-MS/MS—High Performance Liquid Chromatography with Tandem Mass Spectrometry Detection; LC-MS/MS—Liquid Chromatography with Tandem Mass Spectrometry Detection.

#### 4.7. Antidepressants

Serotonin (or 5-hydroxytryptamine) is an important neurotransmitter in hormonal and neuronal mechanisms. It participates in different regulatory and endocrine functions so that altered levels may cause changes in appetite, immune system, reproduction and other behavioural functions [10,35]. It is also important to lower vertebrates and invertebrates though being associated with different physiological mechanisms from those observed for mammals. In therapeutics, the selective serotonin reuptake inhibitors (SSRIs) fluoxetine, fluvoxamine, paroxetine and sertraline are the most widely used synthetic antidepressants. They act by inhibiting the reuptake of serotonin from the pre-synaptic nerve cleft. It is thus obvious that from the presence of SSRIs in the environment (even at low concentrations ( $\text{ng}$  or  $\mu\text{g L}^{-1}$ )), adverse effects on aquatic organisms could arise [177]. In fact, fluvoxamine at a concentration of  $0.32 \mu\text{g L}^{-1}$  or fluoxetine at higher concentrations were capable of inducing spawning and oocyte maturation of zebra mussels (*Dreissena polymorpha*) [178]. On the contrary, a NOEC value of  $0.47 \mu\text{g L}^{-1}$  was deduced for the ability of fluoxetine to reduce reproduction of the freshwater mudsnail *Potamopyrgus antipodarum* [179]. Japanese medaka were exposed to a range of fluoxetine from  $0.1$  to  $5 \mu\text{g L}^{-1}$  over four weeks, revealing that fecundity, egg fertilization and hatching success were unaffected. However, an increase in developmental abnormalities in fish embryos was observed and plasma estradiol concentrations were significantly raised in females [180]. Following an one-week exposure of western mosquitofish (*Gambusia affinis*) neonates to fluoxetine, a  $\text{LC}_{50}$  value of  $546 \mu\text{g L}^{-1}$  was obtained [181]. Although chronic exposure to concentrations from  $0.05$  to  $5 \mu\text{g L}^{-1}$  increased lethargy, it did not affect survival, growth or sex ratio [181]. In turn,

*G. affinis* exposed to  $71 \mu\text{g L}^{-1}$  of fluoxetine from juvenile through adult life stages showed a delay in the development of mature sexual morphology in both male and female fish [181].

Another SSRI, sertraline, exhibits highly toxic properties. Following a 96-h exposure of rainbow trout to sertraline, a  $\text{LC}_{50}$  of  $0.38 \text{ mg L}^{-1}$  was deduced [182]. The same authors also found that those surviving fish exposed to  $0.32 \text{ mg L}^{-1}$  of sertraline for 72 h, died following irreparable physiological damage after being removed to control water. Fish exposed to higher concentrations of this pharmaceutical showed a decreased respiration and a loss of movement coordination.

SSRIs were also tested on algae by evaluating the growth inhibition induced. Chronic toxicity tests proved that the organisms were sensitive with NOEC values below  $1 \text{ mg L}^{-1}$  [52]. *C. vulgaris* was shown to be the least sensitive species for all SSRIs tested [183]. On the contrary, *Pseudokirchneriella subcapitata* was the most sensitive species mainly regarding fluoxetine with a reported  $\text{EC}_{50}$  of  $24 \mu\text{g L}^{-1}$  after 48 h [177,184] or  $45 \mu\text{g L}^{-1}$  when the exposure time was increased to 96 h [183]. Cell deformities in these green algae were noticed with just  $13.6 \mu\text{g L}^{-1}$  of fluoxetine. Similar  $\text{EC}_{50}$  values were determined for acute toxic effects caused by sertraline on *P. subcapitata* and *Scenedesmus acutus* ( $12.1$  and  $99 \mu\text{g L}^{-1}$  respectively) [183]. By reducing the exposure time from 96 to 72 h, *P. subcapitata* showed an  $\text{EC}_{50}$  of  $0.14 \text{ mg L}^{-1}$  [182]. Fluvoxamine gave rise to the highest  $\text{EC}_{50}$  values for all algae species tested ( $3563$ – $10,208 \mu\text{g L}^{-1}$ ) [183]. An exposure of 96 h of the marine phytoplankton *D. tertiolecta* to fluoxetine showed an  $\text{EC}_{50}$  of  $169.81 \mu\text{g L}^{-1}$  [70], which is higher than growth rate  $\text{EC}_{50}$ s reported previously to algae species.

Tests performed on the invertebrates *C. dubia*, *D. magna* and on fathead minnow fish showed  $\text{LC}_{50}$  values of  $234$ ,  $820$  and  $705 \mu\text{g L}^{-1}$  respectively, after 48 h of exposure to fluoxetine [184]. On the other hand, for paroxetine, *D. magna* showed an  $\text{EC}_{50}$  of  $2.5 \text{ mg L}^{-1}$  [185]. Regarding the invertebrates, fluoxetine may cause a stimulation of reproduction as is the case of *C. dubia* when exposed to

$56 \mu\text{g L}^{-1}$  of this pharmaceutical [184]. This same effect was also found for *D. magna* after 30 days of exposure to a concentration of  $36 \mu\text{g L}^{-1}$  [116] which resulted in an increase in total number of offspring produced. Higher concentrations of fluoxetine were tested (e.g.  $223 \mu\text{g L}^{-1}$ ) and proven to exert the opposite effect [184] in a similar way to that observed for sertraline, exhibiting an  $\text{EC}_{50}$  of  $0.066 \text{ mg L}^{-1}$  and a LOEC of  $0.1 \text{ mg L}^{-1}$  [182]. A multi-generational study was performed by exposing *D. magna* and their newborns to fluoxetine [33]. The highest effects were found on the development of the embryos. The newborns length was affected (NOEC =  $8.9 \mu\text{g L}^{-1}$  and LOEC =  $31 \mu\text{g L}^{-1}$ ), what had consequences in their future reproduction, that was significantly reduced for a concentration of  $31 \mu\text{g L}^{-1}$  [33]. The exposure of the invertebrate

*P. antipodarum* to fluoxetine caused a decrease in reproduction, resulting in a NOEC of  $13 \mu\text{g L}^{-1}$  and a LOEC of  $69 \mu\text{g L}^{-1}$  [33]. In contrast, *H. azteca* reproduction was not affected by this SSRI, but a significant effect on growth was noticed, showing a NOEC and a LOEC of  $33$  and  $100 \mu\text{g L}^{-1}$ , respectively [33].

The behaviour of aquatic invertebrates was also shown to be affected by SSRIs as illustrated by the amphipod *G. pulex* in the presence of  $10$  and  $100 \text{ ng L}^{-1}$  of fluoxetine [76]. Fairy shrimps *T. platyurus* are more sensitive to sertraline compared to *D. magna*. For the former an  $\text{EC}_{50}$  of  $0.6 \text{ mg L}^{-1}$  after 24 h was obtained and with *D. magna* corresponding  $\text{EC}_{50}$  values were  $3.1$  and  $1.3 \text{ mg L}^{-1}$  after 24 and 48 h, respectively [182]. Nematoceran flies *Chironomus tentans* and hydras *H. azteca* were exposed to fluoxetine by sediments, showing growth inhibition with LOECs of  $1.3$  and

$5.6 \text{ mg kg}^{-1}$  respectively [184]. However, hydras reproduction was stimulated for all concentrations tested ( $1.4$ – $22.4 \text{ mg kg}^{-1}$ ) as well as blackworms *Lumbriculus variegatus* when exposed to  $0.94$  and  $2.34 \text{ mg kg}^{-1}$  of fluoxetine [179]. In *C. tentans*, this kind of exposure caused a reduction in emergence with a LOEC of  $1.12 \text{ mg kg}^{-1}$ . On the other hand, Péry et al. [33] did not observe toxic effects on *C. riparius* growth, emergence and reproduction, even when exposed to  $59.5 \text{ mg kg}^{-1}$  of fluoxetine.

SSRIs contaminate different aquatic environments at concentrations in the order of  $\text{ng L}^{-1}$  (Table 7). Fluoxetine is a typical example, being detected in STP influents at concentrations of  $0.4$ – $18.7 \text{ ng L}^{-1}$  and in effluents in the lower range of  $0.12$ – $8.4 \text{ ng L}^{-1}$  [186–188]. This pharmaceutical was also detected in surface waters [23,188], groundwaters [28] and drinking water [32]. Other SSRIs, such as fluvoxamine, sertraline and paroxetine have also been detected in STP influents and effluents [186–188] as well as seawater (Norway) [187]. Antidepressants were detected at low concentrations ( $\text{ng L}^{-1}$ ) which may not represent isolated threats to non-target organisms when considering the respective contribution. However, since they exert similar effects and are present in the environment as a mixture, it is possible that chronic exposure of aquatic organisms may induce toxicity.

#### 4.8. Antineoplastics

Antineoplastic drugs are designed to kill cells that are proliferating excessively such as those found in pathological cancer conditions. Therefore, a similar effect on any other growing eukaryotic organisms is expected [189]. Pharmaceuticals belonging to this therapeutic class possess genotoxic, mutagenic, carcinogenic, teratogenic and fetotoxic properties and can constitute (in their native form) from  $14$  to  $53\%$  of the administered drug excreted in urine [108]. Cyclophosphamide and ifosfamide ecotoxicity predicted by ECOSAR have yielded  $\text{EC}_{50}$  values of  $8.2$  and  $70 \text{ mg L}^{-1}$  for algae and fish respectively, whereas the freshwater flea *D. magna* registered a  $\text{LC}_{50}$  of  $1795 \text{ mg L}^{-1}$  [108]. Toxicity tests performed on the algae *P. subcapitata* and the invertebrate *D. magna* showed that cyclophosphamide slightly increased the growth of the former (NOEC above  $100 \text{ mg L}^{-1}$ ) and reduced offspring number in the lat-

Table 7

Examples of concentrations (ng L<sup>-1</sup>) of antidepressants measured in different aquatic environments.

Compound	CAS number	Sample	Country	Analytical procedure	LOD(ngL <sup>-1</sup> )	Concentration reported(ngL <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
Amitriptyline	— <sup>†</sup>	STP influent	Canada	SPE-LC-MS/MS	0.077	17.6 (±0.8)–20.8 (±1.2)	[188]					
		STP effluent				15.6 (±0.8)–21.0 (±1.5)						
		St. Lawrence River water				0.87 (±0.07)–3.7 (±0.2)						
Nortriptyline*	— <sup>†</sup>	STP influent	Canada	SPE-LC-MS/MS	0.057	3.1 (±0.1)–4.5 (±0.4)	[188]					
		STP effluent				1.5 (±0.1)–3.8 (±0.4)						
		St. Lawrence River water				0.41 (±0.02)–0.73 (±0.06)						
Citalopram	59729-33-8	STP influent	Norway	SPE-HPLC-MS	0.16	13.0–612	[186]					
STP effluent												
Citalopram		STP influent	Norway	HF-LPME-HPLC-MS	0.017	62.9 (±30.7)–303.6 (±4.3)	[187]					
		STP effluent				21.9 (±13.5)–238.4 (±23.6)						
Citalopram		STP influent	Canada	SPE-LC-MS/MS	0.077	52.2 (±3.7)–52.7 (±4.9)	[188]					
		STP effluent				46.8 (±1.2)–57.8 (±0.3)						
		St. Lawrence River water				3.4 (±0.2)–11.5 (±0.8)						
Fluoxetine	54910-89-3	Surface water	USA	SPE-LC-MS	18	12	[23]	Amphipod	<i>H. azteca</i>	LOEC (28 d) (growth)	100 μg L <sup>-1</sup>	[33]
Fluoxetine		Groundwater	USA	SPE-HPLC-MS	18	56	[28]			NOEC (28 d) (growth)	33 μg L <sup>-1</sup>	[33]
Fluoxetine		Drinking water	USA	SPE-LC-MS/MS	0.50	0.64	[32]	Crustacean	<i>D. magna</i>	NOEC (21 d) (newborns length)	8.9 μg L <sup>-1</sup>	[33]
Fluoxetine		STP effluent	South Korea	SPE-LC-MS/MS	1.0	1.7	[90]			LOEC (21 d) (newborns length)	31 μg L <sup>-1</sup>	[33]
		Surface water				ND						
Fluoxetine		STP influent	Norway	SPE-HPLC-MS	0.12	0.4–2.4	[186]	Freshwater snail	<i>P. antipodarum</i>	NOEC (reproduction)	13 μg L <sup>-1</sup>	[33]
		STP effluent				<0.12–1.3						
Fluoxetine		STP influent	Norway	HF-LPME-HPLC-MS	0.15	1.1 (±22.9)–18.7 (±0.9)	[187]			LOEC (reproduction)	69 μg L <sup>-1</sup>	[33]
		STP effluent				0.6 (±20.0)–8.4 (±22.9)						
Fluoxetine		STP influent	Canada	SPE-LC-MS/MS	0.05	3.1 (±0.3)–3.5 (±0.3)	[188]	Crustacean	<i>Gammarus pulex</i>	LOEC (behaviour)	100 ng L <sup>-1</sup>	[76]
		STP effluent				2.0 (±0.1)–3.7 (±0.1)						
		St. Lawrence River water				0.42 (±0.01)–1.3 (±0.1)		Algae	<i>Dunaliella tertiolecta</i>	EC <sub>50</sub> (96 h) (growth inhibition)	169.81 μg L <sup>-1</sup>	[70]
									<i>P. subcapitata</i>	EC <sub>50</sub> (120 h) (growth)	24 μg L <sup>-1</sup>	[177]
										LOEC (growth)	13.6 μg L <sup>-1</sup>	[177]
								Crustacean	<i>C. dubia</i>	LC <sub>50</sub> (48 h)	234 μg L <sup>-1</sup>	[177]
										NOEC	56 μg L <sup>-1</sup>	[177]
										LOEC	112 μg L <sup>-1</sup>	[177]

Table 7 (Continued)

Compound	CAS number	Sample	Country	Analytical procedure	LOD( $\text{ngL}^{-1}$ )	Concentration reported ( $\text{ngL}^{-1}$ )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.	
								Fish	<i>D. magna</i>	LC <sub>50</sub> (48 h)	820 $\mu\text{gL}^{-1}$	[177]	
									<i>P. promelas</i>	LC <sub>50</sub> (48 h)	705 $\mu\text{gL}^{-1}$	[177]	
									Midge	<i>C. tentans</i>	LC <sub>50</sub> (10 d)	15.2 $\text{mg kg}^{-1}$	[177]
										LOEC (10 d) (growth)	1.3 $\text{mg kg}^{-1}$	[177]	
								Amphipod	<i>H. azteca</i>	LOEC (growth)	5.4 $\text{mg kg}^{-1}$	[177]	
									Freshwater snail	<i>P. antipodarum</i>	EC <sub>10</sub> (56 d) (n° embryos whitout shell)	0.81 $\mu\text{gL}^{-1}$	[179]
										NOEC (56 d) (n° embryos whitout shell)	0.47 $\mu\text{gL}^{-1}$	[179]	
								Midge		<i>C. riparius</i>	LOEC (28 d) (emergence)	1.12 $\text{mg kg}^{-1}$	[179]
								Mosquitofish		<i>Gambusia affinis</i>	LC <sub>50</sub> (7 d) (lethality)	546 $\mu\text{gL}^{-1}$	[181]
											Algae	<i>P. subscapitata</i>	IC <sub>50</sub> (96 h) (growth inhibition)
								<i>S. acutus</i>	IC <sub>50</sub> (96 h) (growth inhibition)	91.23 $\mu\text{gL}^{-1}$		[183]	
								<i>S. quadricauda</i>	IC <sub>50</sub> (96 h) (growth inhibition)	212.98 $\mu\text{gL}^{-1}$		[183]	
								<i>C. vulgaris</i>	IC <sub>50</sub> (96 h) (growth inhibition)	4339.25 $\mu\text{gL}^{-1}$		[183]	
								Algae	<i>P. subscapitata</i>	EC <sub>50</sub> (120 h) (growth)		39 $\mu\text{gL}^{-1}$	[184]
									Crustacean	<i>C. dubia</i>	LC <sub>50</sub> (48 h) (survival)	234 $\mu\text{gL}^{-1}$	[184]
Fish	<i>D. magna</i>	LC <sub>50</sub> (48 h) (survival)	820 $\mu\text{gL}^{-1}$	[184]									
	Midge	<i>P. promelas</i>	LC <sub>50</sub> (48 h) (survival)	705 $\mu\text{gL}^{-1}$	[184]								
		<i>C. tentans</i>	LC <sub>50</sub> (10 d) (survival)	15.2 $\text{mg kg}^{-1}$	[184]								
	Amphipod		LOEC (10 d) (growth)	1.3 $\text{mg kg}^{-1}$	[184]								
		<i>H. azteca</i>	LOEC (10 d) (growth)	5.6 $\text{mg kg}^{-1}$	[184]								
Norfluoxetine*	83891-03-6	Drinking water	USA	SPE-LC-MS/MS	0.50	0.77	[32]						
Norfluoxetine*		STP influent	Norway	HF-LPME-HPLC-MS	0.16	0.7 ( $\pm 13.1$ )–9.3 ( $\pm 4.6$ )	[187]						
		STP effluent				0.54 (LOQ)		<0.54–2.4 ( $\pm 14.5$ )					
Norfluoxetine*		STP influent	Canada	SPE-LC-MS/MS	0.087	1.8 ( $\pm 0.3$ )–4.2 ( $\pm 0.5$ )	[188]						
		STP effluent				1.7 ( $\pm 0.1$ )–1.8 ( $\pm 0.3$ )							
		St. Lawrence River water				1.2 ( $\pm 0.1$ )–1.3 ( $\pm 0.1$ )							
Fluvoxamine	54739-18-3	STP influent	Norway	SPE-HPLC-MS	0.15	0.4–3.9	[186]	Algae	<i>P. subscapitata</i>	IC <sub>50</sub> (96 h) (growth inhibition)	4002.88 $\mu\text{gL}^{-1}$	[183]	
Fluvoxamine		STP effluent	Norway	HF-LPME-HPLC-MS	0.129	<0.15–0.8	[187]						
		STP influent				0.8 ( $\pm 38.2$ )–1.7 ( $\pm 18.6$ )			<i>S. acutus</i>	IC <sub>50</sub> (96 h) (growth inhibition)	3620.24 $\mu\text{gL}^{-1}$	[183]	
		STP effluent				0.379 (LOQ)		<0.379–0.8 ( $\pm 38.2$ )					
		Seawater				0.5 ( $\pm 0.5$ )–0.8 ( $\pm 0.3$ )							
									<i>S. quadricauda</i>	IC <sub>50</sub> (96 h) (growth inhibition)	3563.34 $\mu\text{gL}^{-1}$	[183]	
									<i>C. vulgaris</i>	IC <sub>50</sub> (96 h) (growth inhibition)	10,208.47 $\mu\text{gL}^{-1}$	[183]	



Paroxetine	61869-08-7	STP influent	Norway	SPE-HPLC-MS	0.12	0.6–12.3	[186]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48h) (immobilization)	2.5 mgL <sup>-1</sup>	[185]
Paroxetine		STP effluent				0.5–1.6						
Paroxetine		STP influent	Norway	HF-LPME- HPLC-MS	0.053	2.9 (±19.0)–12.9 (±29.4)	[187]					
		STP effluent				1.0 (±15.7)–11.7 (±36.8)						
		Seawater				0.6 (±0.4)–1.4 (±0.4)						
Paroxetine		STP influent	Canada	SPE-LC-MS/MS	0.096	4.6 (±0.3)–5.3 (±0.2)	[188]					
		STP effluent				4.3 (±0.2)–5.2 (±0.5)						
		St. Lawrence River water				1.3 (±0.1)–3.0 (±0.1)						
Sertraline	79617-96-2	STP influent	Norway	SPE-HPLC-MS	0.29	1.8–2.5	[186]	Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (30 min) (inhibition)	10.72 mgL <sup>-1</sup>	[182]
		STP effluent				0.9–2.0						
Sertraline		STP influent	Norway	HF-LPME- HPLC-MS	0.16	8.4 (±4.5)–19.8 (±10.8)	[187]			NOEC (30 min) (inhibition)	2.25 mgL <sup>-1</sup>	[182]
		STP effluent			0.52 (LOQ)	3.7 (±16.3)–14.6 (±4.2)						
		Seawater				<0.52						
Sertraline		STP influent	Canada	SPE-LC-MS/MS	0.048	6.0 (±0.4)–6.1 (±0.3)	[188]			LOEC (30 min) (inhibition)	4.5 mgL <sup>-1</sup>	[182]
		STP effluent				5.1 (±0.3)–5.8 (±0.8)						
		St. Lawrence River water				0.84 (±0.09)–2.4 (±0.1)						
								Algae	<i>P. subcapitata</i>	EC <sub>50</sub> (72h) (inhibition)	0.14 mgL <sup>-1</sup>	[182]
										NOEC (72h) (inhibition)	0.05 mgL <sup>-1</sup>	[182]
										LOEC (72h) (inhibition)	0.075 mgL <sup>-1</sup>	[182]
								Shrimp	<i>T. platyurus</i>	LC <sub>50</sub> (24 h)(lethality)	0.6 mgL <sup>-1</sup>	[182]
										NOEC (24h) (lethality)	0.4 mgL <sup>-1</sup>	[182]
								Crustacean	<i>D. magna</i>	LOEC (24h)(lethality)	0.6 mgL <sup>-1</sup>	[182]
										EC <sub>50</sub> (48h) (immobilization)	1.3 mgL <sup>-1</sup>	[182]
										NOEC (48h) (immobilization)	0.10 mgL <sup>-1</sup>	[182]
										LOEC (48h) (immobilization)	0.18 mgL <sup>-1</sup>	[182]
										EC <sub>50</sub> (21 d) (reproduction)	0.066 mgL <sup>-1</sup>	[182]
										NOEC (21 d) (reproduction)	0.032 mgL <sup>-1</sup>	[182]
										LOEC (21 d) (reproduction)	0.1 mgL <sup>-1</sup>	[182]
										LC <sub>50</sub> (21 d)(lethality)	0.12 mgL <sup>-1</sup>	[182]
										NOEC (21 d) (lethality)	0.032 mgL <sup>-1</sup>	[182]
								Fish	<i>O. mykiss</i>	LOEC (21 d)(lethality)	0.1 mgL <sup>-1</sup>	[182]
										LC <sub>50</sub> (96 h)(lethality)	0.38 mgL <sup>-1</sup>	[182]
										NOEC (96h) (lethality)	0.1 mgL <sup>-1</sup>	[182]

Table 7 (Continued)

Compound	CAS number	Sample	Country	Analytical procedure	LOD (ngL <sup>-1</sup> )	Concentration reported (ngL <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
										NOEC (96 h) (lethality)	0.32 mgL <sup>-1</sup>	[182]
								Algae	<i>P. subcapitata</i>	IC <sub>50</sub> (96 h) (growth inhibition)	12.10 μg L <sup>-1</sup>	[183]
									<i>S. acutus</i>	IC <sub>50</sub> (96 h) (growth inhibition)	98.92 μg L <sup>-1</sup>	[183]
									<i>S. quadricauda</i>	IC <sub>50</sub> (96 h) (growth inhibition)	317.02 μg L <sup>-1</sup>	[183]
									<i>C. vulgaris</i>	IC <sub>50</sub> (96 h) (growth inhibition)	763.66 μg L <sup>-1</sup>	[183]
Desmethylsertraline*	87857-41-8	STP influent	Canada	SPE-LC-MS/MS	0.072	4.2 (±0.6)–5.0 (±0.8)	[188]					
		STP effluent				3.6 (±0.3)–4.7 (±0.5)						
		St. Lawrence River water				2.3 (±0.1)–4.5 (±0.4)						
Venlafaxine	99300-78-4	STP influent	Canada	SPE-LC-MS/MS	0.10	195.7 (±25.3)–213.0 (±8.2)	[188]					
		STP effluent				175.9 (±12.7)–214.6 (±3.6)						
		St. Lawrence River water				12.9 (±0.1)–45.9 (±2.0)						
Desmethylvenlafaxine*	— <sup>†</sup>	STP influent	Canada	SPE-LC-MS/MS	0.097	274.3 (±26.5)–345.9 (±19.8)	[188]					
		STP effluent				222.5 (±16.8)–330.0 (±9.8)						
		St. Lawrence River water				21.0 (±0.5)–68.7 (±3.1)						

\*—Metabolite; ND—Not Detected; <sup>†</sup>—Data not available; SPE—Solid Phase Extraction; HF-LPME—Hollow Fibre Supported Liquid Phase Microextraction; HPLC-MS—High Performance Liquid Chromatography with Mass Spectrometry Detection; LC-MS—Liquid Chromatography with Mass Spectrometry Detection; LC-MS/MS—Liquid Chromatography with Tandem Mass Spectrometry Detection.

**Table 8**

Examples of concentrations (ng L<sup>-1</sup>) of antineoplastic drugs measured in different aquatic environments.

Compound	CAS number	Sample	Country	Analytical procedure	LOD (ng L <sup>-1</sup> )	Concentration reported (ng L <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
Cyclophosphamide	50-18-0	Somes river water	Romania	SPE-GC-MS	30 (LOQ)	<30–64.8 (±8.0)	[20]	Algae	<i>P. subcapitata</i>	EC <sub>50</sub> (72 h) (growth inhibition)	>100 mg L <sup>-1</sup>	[190]
Cyclophosphamide		STP effluent	Italy	SPE-HPLC-MS/MS	1.9 (LOQ)	<1.9–9.0	[118]			NOEC (72h) (growth inhibition)	>100 mg L <sup>-1</sup>	[190]
Cyclophosphamide		STP influent	—†	SPE-GC-MS	6	<6–143	[192]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (21 d) (reproduction)	>100 mg L <sup>-1</sup>	[190]
		STP effluent				6–15						
		Hospital effluent				19–4486						
Cyclophosphamide		STP influent	Switzerland	SPE-LC-MS/MS	0.3	2.0–6	[193]			LOEC (21 d) (reproduction)	100 mg L <sup>-1</sup>	[190]
		STP effluent				2.1–4				NOEC (21 d) (reproduction)	56 mg L <sup>-1</sup>	[190]
Ifosfamide	84711-20-6	STP influent	Switzerland	SPE-LC-MS/MS	0.3	<0.3–5	[193]					
		STP effluent				1.7–6						
Methotrexate	59-05-2	STP effluent	Italy	SPE-HPLC-MS/MS	0.83 (LOQ)	<0.83–12.6	[118]	Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (30 min)	1220 mg L <sup>-1</sup>	[83]
								Algae	<i>Scenedesmus subspicatus</i>	EC <sub>50</sub> (72h)	260 mg L <sup>-1</sup>	[83]
								Crustacean	<i>D. magna</i>	EC <sub>50</sub> (immobilization)	>1000 mg L <sup>-1</sup>	[83]
								Ciliates	<i>Tetrahymena pyriformis</i>	EC <sub>50</sub> (48 h) (growth inhibition)	45 mg L <sup>-1</sup>	[83]
								Fish	<i>D. rerio</i>	LC <sub>50</sub> (48 h)	85 mg L <sup>-1</sup>	[83]
Tamoxifen	74899-71-1	STP influent	United Kingdom	SPE-HPLC-MS/MS	10	143–215	[53]	Rotifer	<i>B. calyciflorus</i>	LC <sub>50</sub> (24 h) (mortality)	0.97 mg L <sup>-1</sup>	[191]
		STP effluent				146–369						
		Tyne river water				27–212						
Tamoxifen		STP effluent	United Kingdom	SPE-HPLC-MS/MS	10	<10	[94]			EC <sub>50</sub> (48h) (population growth inhibition)	0.25 mg L <sup>-1</sup>	[191]
		Surface water				<10						
Tamoxifen		STP influent	United Kingdom	SPE-LC-MS/MS	0.003	0.2–15	[194]	Crustacean	<i>T. platyurus</i>	LC <sub>50</sub> (24 h) (mortality)	0.40 mg L <sup>-1</sup>	[191]
		STP effluent				0.2–0.7						
									<i>D. magna</i>	EC <sub>50</sub> (24h) (immobilization)	1.53 mg L <sup>-1</sup>	[191]
									<i>C. dubia</i>	EC <sub>50</sub> (7 d) (population growth inhibition)	8.1 × 10 <sup>-4</sup> mg L <sup>-1</sup>	[191]

†—Data not available; SPE—Solid Phase Extraction; GC-MS—Gas Chromatography with Mass Spectrometry Detection; HPLC-MS/MS—High Performance Liquid Chromatography with Tandem Mass Spectrometry Detection; LC-MS/MS—Liquid Chromatography with Tandem Mass Spectrometry Detection.

ter at all tested concentrations of the drug (10–100 mg L<sup>-1</sup>), with a NOEC of 56 mg L<sup>-1</sup> [190]. Methotrexate revealed teratogenicity for fish embryos with an EC<sub>50</sub> of 85 mg L<sup>-1</sup> after 48 h of exposure [83] and acute effects in the ciliate *Tetrahymena pyriformis* with an EC<sub>50</sub> for 48 h of 45 mg L<sup>-1</sup> [83]. Acute and chronic toxicity of tamoxifen and its photoproducts was studied by DellaGreca et al. [191], showing that both the active pharmaceutical and its photoproducts affected the rotifer *B. calyciflorus* and crustacean *T. platyurus* with LC<sub>50</sub> values ranging from 0.95 to 1.31 mg L<sup>-1</sup> and 0.40 to 1.59 mg L<sup>-1</sup> respectively. In chronic toxicity tests, *C. dubia* proved the most sensitive organism. An EC<sub>50</sub> value of 0.81 μg L<sup>-1</sup> for tamoxifen and EC<sub>50</sub> values ranging from 0.41 to 2.8 μg L<sup>-1</sup> for its photoproducts, relative to population growth inhibition, were found after a 7-day trial [191].

The antineoplastic drug cyclophosphamide has been detected in hospital effluents at concentrations ranging from 19 ng L<sup>-1</sup> to 4.5 μg L<sup>-1</sup> [192], in STP influents [192,193] and effluents [118,192,193] and in surface waters [20] (Table 8). Other antineoplastic pharmaceuticals detected to date have been in the order of ng L<sup>-1</sup>. However, as chronic toxicity data is very sparse, further studies are required to elucidate the potential effect of life-cycle exposure to these compounds in aquatic organisms.

#### 4.9. X-ray contrast media

Contrast media are used as diagnostic tools for capturing detailed X-ray images of soft tissues. Iodinated X-ray contrast media are highly hydrophilic substances that are widely used and eliminated almost non-metabolised. STP removal processes are usually ineffective and for this reason they persist for a long time in the environment. As X-ray contrast media do not show biological activity, their presence might not represent a threat to public health [35,195,196]. Toxicity tests have shown that iopromide or its main metabolite do not have a toxic effect in luminescent bacteria, algae (*Scenedesmus subspicatus*), daphnids or fish (*D. rerio*, *Leuciscus idus*) even at concentrations as high as 1 g L<sup>-1</sup> [196,197]. Contamination by X-ray contrast media has been reported in different aquatic environments (Table 9). Media have been detected in STP influents and effluents [198–201], surface waters [199,201–203], groundwaters [26,199,200] and even drinking water [200,202,203] at concentrations that can reach few μg L<sup>-1</sup>. Although accepting that X-ray contrast media do not exhibit toxic effects at high concentration levels, additional studies should be undertaken with a view to evaluating chronic effects, due to continuous exposure of aquatic organisms to these pharmaceuticals.

#### 4.10. Mixture effects

Presently environmental risk assessment of pharmaceuticals is based on single compounds ecotoxicity studies. However, pharmaceuticals do not occur alone in the environment, but as a mixture of different active substances, their metabolites and transformation products [23,205,206]. Ecotoxicological data showed that mixtures might have different effects than single compounds [65,70,207], but in general knowledge about the toxicity of the mixture of active substances is still sparse. There are some examples of toxicity studies in literature showing that mixture of pharmaceuticals at environmentally relevant concentrations may exhibit additive effects [70]. In some cases, lower levels than expected may lead to toxic effects when in the presence of a mixture of active substances [70]. For instance, Cleuvers [65] showed that a mixture of diclofenac and ibuprofen had a stronger toxicity than predicted in

*D. magna*, and when the author added more two NSAIDs (naproxen and acetylsalicylic acid) to the last two, a considerable toxicity on *Daphnia* was also reported, even at concentrations at which

the single NSAIDs do not exhibit effects [95]. The exposure of the cnidarian *H. attenuata* to a mixture of eleven pharmaceuticals, belonging to different therapeutic classes, showed also sub-lethal effects for environmentally relevant concentrations (μg–ng L<sup>-1</sup>) [207]. Acute exposure of *D. magna* to a mixture of 36 μg L<sup>-1</sup> of fluoxetine and 100 μg L<sup>-1</sup> of clofibrac acid caused a significant mortality and malformation, while there are no apparent effects for the same concentrations of individual pharmaceuticals [116]. The mixture of trimethoprim with sulfamethoxazole and sulfadiazine increased significantly the growth inhibition of the algae *S. capricornutum* [131]. On the other hand, the exposure of *H. azteca* to a mixture of seven pharmaceuticals did not reveal significant effects on their survival, mating, body size or reproduction, but there was a slight increase in the number of males [208]. Identical results were observed for fish. Apparently, a life-cycle exposure of fathead minnows to a mixture of six pharmaceuticals, in the order of ng L<sup>-1</sup>, did not affect their survival, growth or egg production, however it increased the number of deformities in their offspring [209]. The examples here highlighted showed that the simultaneous presence of several pharmaceuticals in the environment might result in a greater toxicity to non-target organisms than the predicted one for individual active substances. However, more ecotoxicological studies should be done to evaluate the impact of different mixtures of pharmaceuticals in non-target organisms, once that most of the published studies are focused on mixture of NSAIDs, antibiotics and blood lipid lowering agents.

### 5. Pharmaceuticals and legislation: what does legislation say?

Every day an increasing number of pharmaceuticals reach the environment all over the world. However, there is a gap in legislation regarding environmental contamination by pharmaceuticals. This probably arises because available data is insufficient to quantify a precise contamination profile. Abundant conclusive studies concerning chronic toxicity are also lacking so that it becomes impossible to infer the risks of long-term exposure of pharmaceuticals and their metabolites on fauna and flora. In this section, EU and US laws concerning the necessity of environmental risk assessment studies to obtain a marketing authorisation for pharmaceuticals is approached.

The European Union Directive 92/18/EEC [210] introduced for the first time, the requirement for an environmental risk assessment, as a prerequisite to obtain marketing authorization for veterinary pharmaceuticals. For this purpose, the European Agency for the Evaluation of Medicinal Products (EMA) published a “Note for Guidance” [211] where guidelines to assess the environmental risk of veterinary medicinal products are established. The European Commission extended its concerns to pharmaceuticals for human use by publishing Directive 2001/83/EC which was subsequently amended by Directive 2004/27/EC [212]. These directives established that marketing authorization for new medical products for human use should be accompanied by an environmental risk assessment, whose guidelines were set out by EMA [213]. Nevertheless, the environmental impact does not provide sufficient grounds for a refusal. Environmental risk assessment of both veterinary and human pharmaceuticals is assessed in a step-wise approach, divided into two phases. In Phase I, environmental exposure to the pharmaceutical or its metabolites is estimated. Phase II comprises its fate and effects in the environment. For this reason, Phase II is sub-divided into two parts: Tier A, in which possible fate and effects of the pharmaceutical and/or its major metabolites are evaluated; and Tier B, which focuses on the effects on fauna and flora within environmental compartments that are likely to be affected [211,213]. However, medicinal products for human use

Table 9

Examples of concentrations (ng L<sup>-1</sup>) of X-ray contrast media measured in different aquatic environments.

Compound	CAS number	Sample	Country	Analytical procedure	LOD (ng L <sup>-1</sup> )	Concentration reported (ng L <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
Diatrizoate	131-49-7	STP effluent	Germany	SPE-LC-MS/MS	50 (LOQ STP effluent)	250	[199]					
		Surface water										
Diatrizoate		Groundwater	Germany	SPE-HPLC-MS	— <sup>f</sup>	30	[203]					
		Surface water										
Diatrizoate		Drinking water	Germany	SPE-LC-MS/MS	50 (LOQ STP effluent)	1140	[204]					
		STP effluent										
Iohexol	66108-95-0	Rhine river water	Australia	DI-LC-MS/MS	800	60	[201]					
		Drinking water										
Iohexol		STP influent	Germany	SPE-HPLC-MS/MS	40	40–86	[202]					
		STP effluent										
Iomeprol	78649-41-9	Danube river water	Germany	SPE-LC-MS/MS	50 (LOQ STP effluent)	370	[199]					
		STP effluent										
Iomeprol		STP influent	Australia	DI-LC-MS/MS	730	<730	[201]					
		STP effluent										
Iomeprol		Danube river water	Germany	SPE-HPLC-MS/MS	40	100–160	[202]					
		STP effluent										
Iopamidol	60166-93-0	Groundwater	Germany	SPE-HPLC-MS/MS	14	300	[26]					
		STP effluent										
Iopamidol		Surface water	Germany	SPE-LC-MS/MS	50 (LOQ STP effluent)	660	[199]					
		Groundwater										
Iopamidol		STP influent	Australia	DI-LC-MS/MS	220	160	[201]					
		STP effluent										
Iopamidol		Danube river water	Germany	SPE-HPLC-MS/MS	40	210	[202]					
		STP effluent										
Iopamidol		STP effluent	Germany	SPE-LC-MS/MS	50 (LOQ STP effluent)	590	[204]					
		Rhine river water										
Iopromide	73334-07-3	Drinking water	South Korea	SPE-LC-MS/MS	1.0	70	[90]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (48h) (immobilization)	>1 g L <sup>-1</sup>	[157]
		STP effluent										
Iopromide		Surface water	Spain	SPE-LC-MS/MS	6.7	20–361	[198]	Fish	<i>D. rerio</i>	NOEC (28 d) (hatchability, post-hatch survival, body length, weight)	>100 mg L <sup>-1</sup>	[157]
		Drinking water										
Iopromide		STP influent	Germany	SPE-LC-MS/MS	50 (LOQ STP effluent)	4400	[199]	Bacteria	<i>V. fischeri</i>	EC <sub>50</sub> (30 min) (luminescence)	>10.0 g L <sup>-1</sup>	[158]
		STP effluent										
Iopromide		STP effluent	USA	SPE-LC-MS/MS	0.577	11–910	[200]		<i>P. putida</i>	EC <sub>10</sub> (16 h) (growth inhibition)	>10.0 g L <sup>-1</sup>	[158]
		Groundwater										
Iopromide		STP influent				ND–17						
		Surface water										
Iopromide		Groundwater				4.6						
		STP influent										
Iopromide		Ohio river water				4.6						
		Drinking water										

Table 9 (Continued)

CAE compound	Sample	Country	Analytical procedure	LOD (ng L <sup>-1</sup> )	Concentration reported (ng L <sup>-1</sup> )	Ref.	Taxon	Species	Toxicological endpoint	Ecotoxicity data	Ref.
Iopromide	Groundwater	Australia	SPE-LC-MS/MS	0.577	168	[200]	Algae	<i>S. subspicatus</i>	EC <sub>50</sub> (72 h) (growth inhibition)	>10.0 g L <sup>-1</sup>	[158]
Iopromide	STP effluent	South Korea	SPE-LC-MS/MS	0.577	152–2670	[200]	Crustacean	<i>D. magna</i>	EC <sub>50</sub> (24 h) (immobilization)	>10.0 g L <sup>-1</sup>	[158]
Iopromide	STP influent	Australia	DL-LC-MS/MS	200	430–1350	[201]					
Iopromide	STP effluent			<200							
Iopromide	Danube river water	Germany	SPE- HPLC-MS/MS	40	76–100	[202]					
Iopromide	Surface water	Germany	HPPLC-MS	50	1600	[203]					
	Drinking water				<50						
Iopromide	STP effluent	Germany	SPE-LC-MS/MS	50 (LOQ STP effluent)	3070	[204]			EC <sub>50</sub> (22 d) (reproduction)	>1.0 g L <sup>-1</sup>	[158]
	Rhine river water			10 (LOQ surface and drinking water)	150						
	Drinking water				40		Fish	<i>D. rerio</i>	LC <sub>50</sub> (96 h) (mortality)	>10.0 g L <sup>-1</sup>	[158]
	Drinking water								LC <sub>50</sub> (48 h) (mortality)	>10.0 g L <sup>-1</sup>	[158]

† – Data not available; ND – Not detected; DL-LC-MS/MS – Direct Injection Liquid Chromatography-Tandem Mass Spectrometry; SPE – Solid Phase Extraction; HPLC-MS – High Performance Liquid Chromatography with Mass Spectrometry Detection; HPPLC-MS/MS – High Performance Liquid Chromatography with Tandem Mass Spectrometry Detection; LC-MS/MS – Liquid Chromatography with Tandem Mass Spectrometry Detection.

only require Phase II studies if the predicted environmental concentration in surface water is equal to or above 0.01  $\mu\text{g L}^{-1}$  [213].

In the US, issues concerning pharmaceuticals in the environment are regulated by the U.S. Food and Drug Administration (FDA). This institution requires environmental assessments to obtain marketing authorisations which are specified in the “Guidance for Industry—Environmental Assessment of Human Drug and Biologic Applications” [214]. However, an environmental assessment is required only if the estimated environmental concentration of the pharmaceutical at the point of the entry is above 1  $\mu\text{g L}^{-1}$  [214]. As EMEA, the FDA also requires environmental assessments for veterinary medicinal products, using a tiered approach. With a view to harmonising the guidelines that govern these environmental impact assessments, the EU, US and Japan elaborated two guidelines: “Environmental Impact Assessment (EIAs) for Veterinary Medicinal Products (VMPs)—Phase I” [215] and “Environmental Impact Assessment for Veterinary Medicinal Products—Phase II Guidance” [216] so that environmental fate and toxicity data obtained could be used to obtain marketing authorisation in all these regions.

## 6. Conclusions

Today, the presence of pharmaceuticals in the environment is being reported worldwide. Furthermore, new data on the sources, fate and effects of pharmaceuticals in the environment, seems to indicate the possibility of a negative impact on different ecosystems and imply a threat to public health. For this assumption, data from acute and chronic ecotoxicity tests on species belonging to different trophic levels such as bacteria, algae, crustaceans and fish among others, is relevant to illustrate the several adverse effects that environmental exposure to measured concentrations of these contaminants can have. On literature, the principal toxicological endpoints/studies that are described are growth, survival, reproduction and immobilization of species, comparatively to transgenerational and population level studies that are still sparse. This demonstrates the lack of data relatively to long-term exposure of non-target organisms and principally how a continuous exposure, during several generations, may affect a whole population. To our knowledge, just one work followed the impact of a pharmaceutical in a fish population throughout seven years, showing how ethinylestradiol negatively affect the fish population, leaving them near of the extinction. In the near future, the evaluation of chronic toxicity effects should be set out as a priority for the scientific community since simultaneous exposure to pharmaceuticals, metabolites and transformation products of several therapeutic classes are unknown and whose probable effects on subsequent generations should be assumed. Another example of missing data is what occurs with statins. Nowadays, they are the blood lipid lowering agents most used all over the world, although toxicity data relatively to them is almost non-existent and limited to the active substances simvastatin and atorvastatin. It is also important to assess the presence of pharmaceuticals and/or their metabolites and transformation products in several environmental compartments in different countries with a view to gaining reliable knowledge of the contamination levels. Only with further available information will be easier to improve existing legislation in order to protect humans, animals and ecosystems from the threat posed by the presence of pharmaceuticals in the environment.

## Acknowledgements

Lúcia H.M.L.M. Santos thanks to FCT the PhD grant (SFRH/BD/48168/2008).

## References

- [1] A.W. Garrison, J.D. Pope, F.R. Allen, GC/MS analysis of organic compounds in domestic wastewaters, in: C.H. Keith (Ed.), *Identification and Analysis of Organic Pollutants in Water*, Ann Arbor Science Publishers, Ann Arbor, MI, 1976, pp. 517–556.
- [2] C. Hignite, D.L. Azarnoff, Drugs and drugs metabolites as environmental contaminants: chlorophenoxyisobutyrate and salicylic acid in sewage water effluent, *Life Sci.* 20 (1977) 337–341.
- [3] M.L. Richardson, J.M. Bowron, The fate of pharmaceutical chemicals in the aquatic environment, *J. Pharm. Pharmacol.* 37 (1985) 1–12.
- [4] G.W. Aherne, J. English, V. Marks, The role of immunoassay in the analysis of microcontaminants in water samples, *Ecotoxicol. Environ. Saf.* 9 (1985) 79–83.
- [5] G.W. Aherne, A. Hardcastle, A.H. Nield, Cytotoxic drugs and the aquatic environment: estimation of bleomycin in river and water samples, *J. Pharm. Pharmacol.* 42 (1990) 741–742.
- [6] K. Kümmerer, Introduction: pharmaceuticals in the environment, in: K. Kümmerer (Ed.), *Pharmaceuticals in the Environment: Sources, Fate, Effects and Risks*, Springer, Berlin, 2001, pp. 1–8.
- [7] P. Pflüger, D.R. Dietrich, Effects on pharmaceuticals in the environment—an overview and principle considerations, in: K. Kümmerer (Ed.), *Pharmaceuticals in the Environment: Sources, Fate, Effects and Risks*, Springer, Berlin, 2001, pp. 11–17.
- [8] E. Zuccato, S. Castiglioni, R. Fanelli, G. Reitano, R. Bagnati, C. Chiabrando, F. Pomati, C. Rossetti, D. Calamari, Pharmaceuticals in the environment in Italy: causes, occurrence, effects and control, *Environ. Sci. Pollut. Res.* 13 (2006) 15–21.
- [9] S.T. Glassmeyer, E.H. Hinchey, S.E. Boehme, C.G. Daughton, I.S. Ruhoy, O. Conerly, R.L. Daniels, L. Lauer, M. McCarthy, T.G. Nettesheim, K. Sykes, V.G. Thompson, Disposal practices for unwanted residential medications in the United States, *Environ. Int.* 35 (2009) 566–572.
- [10] K. Fent, A.A. Weston, D. Caminada, *Ecotoxicology of human pharmaceuticals*, *Aquat. Toxicol.* 76 (2006) 122–159.
- [11] B. Halling-Sørensen, S. Nors Nielsen, P.F. Lanzky, F. Ingerslev, H.C. Holten Lützhøft, S.E. Jørgensen, Occurrence, fate and effects of pharmaceutical substances in the environment—a review, *Chemosphere* 36 (1998) 357–393.
- [12] J.L.C.M. Dorne, A.M.J. Ragas, G.K. Frampton, D.S. Spurgeon, D.F. Lewis, Trends in human risk assessment of pharmaceuticals, *Anal. Bioanal. Chem.* 387 (2007) 1167–1172.
- [13] S.T. Glassmeyer, D.W. Kolpin, E.T. Furlong, M.J. Focazio, Environmental presence and persistence of pharmaceuticals: an overview, in: D.S. Aga (Ed.), *Fate of Pharmaceuticals in the Environment and in Water Treatment Systems*, CRC Press, Taylor and Francis, 2008, pp. 3–52.
- [14] M.J. Gómez, M.J. Martínez Bueno, S. Lacorte, A.R. Fernández-Alba, A. Agüera, Pilot survey monitoring pharmaceuticals and related compounds in a sewage treatment plant located on the Mediterranean coast, *Chemosphere* 66 (2007) 993–1002.
- [15] A. Tauxe-Wuersch, L.F. De Alencastro, D. Grandjean, J. Tarradellas, Occurrence of several acidic drugs in sewage treatment plants in Switzerland and risk assessment, *Water Res.* 39 (2005) 1761–1772.
- [16] N.M. Vieno, T. Tuhkanen, L. Kronberg, Analysis of neutral and basic pharmaceuticals in sewage treatment plants and in recipient rivers using solid phase extraction and liquid chromatography-tandem mass spectrometry detection, *J. Chromatogr. A* 1134 (2006) 101–111.
- [17] S.S. Veremitch, C.J. Lowe, A. Mazumder, Determination of acidic drugs and caffeine in municipal wastewaters and receiving waters by gas chromatography-ion trap tandem mass spectrometry, *J. Chromatogr. A* 1116 (2006) 193–203.
- [18] H.-B. Lee, T.E. Peart, M.L. Svoboda, Determination of endocrine-disrupting phenols, acidic pharmaceuticals, and personal-care products in sewage by solid-phase extraction and gas chromatography-mass spectrometry, *J. Chromatogr. A* 1094 (2005) 122–129.
- [19] V. Koutsouba, Th. Heberer, B. Fuhrmann, K. Schmidt-Baumler, D. Tsipi, A. Hiskia, Determination of polar pharmaceuticals in sewage water of Greece by gas chromatography-mass spectrometry, *Chemosphere* 51 (2003) 69–75.
- [20] Z. Moldovan, Occurrences of pharmaceutical and personal care products as micropollutants in rivers from Romania, *Chemosphere* 64 (2006) 1808–1817.
- [21] D. Bendz, N.A. Paxéus, T.R. Ginn, F.J. Loge, Occurrence and fate of pharmaceutically active compounds in the environment, a case study: Høje River in Sweden, *J. Hazard. Mater.* 122 (2005) 195–204.
- [22] M. Stumpf, T.A. Ternes, R.-D. Wilken, S.V. Rodrigues, W. Baumann, Polar drug residue in sewage and natural waters in the state of Rio de Janeiro, Brazil, *Sci. Total Environ.* 225 (1999) 135–141.
- [23] D.W. Kolpin, E.T. Furlong, M.T. Meyer, E.M. Thurman, S.D. Zaugg, L.B. Barber, H.T. Buxton, Pharmaceuticals, hormones, and other organic wastewater contaminants in U.S. streams, 1999–2000: a national reconnaissance, *Environ. Sci. Technol.* 36 (2002) 1202–1211.
- [24] D. Calamari, E. Zuccato, S. Castiglioni, R. Bagnati, R. Fanelli, Strategic survey of therapeutic drugs in the Rivers Po and Lambro in Northern Italy, *Environ. Sci. Technol.* 37 (2003) 1241–1248.
- [25] S. Weigel, J. Kuhlmann, H. Hühnerfuss, Drugs and personal care products as ubiquitous pollutants: occurrence and distribution of clofibric acid, caffeine and DEET in the North Sea, *Sci. Total Environ.* 295 (2002) 131–141.
- [26] F. Sacher, F.T. Lange, H.-J. Brauch, I. Blankenhorn, Pharmaceuticals in groundwaters. Analytical methods and results of a monitoring program in Baden-Württemberg, Germany, *J. Chromatogr. A* 938 (2001) 199–210.
- [27] A.L. Batt, D.D. Snow, D.S. Aga, Occurrence of sulfonamide antimicrobials in private water wells in Washington County, Idaho, USA, *Chemosphere* 64 (2006) 1963–1971.
- [28] K.K. Barnes, D.W. Kolpin, E.T. Furlong, S.D. Zaugg, M.T. Meyer, L.B. Barber, A national reconnaissance of pharmaceuticals and other organic wastewater contaminants in the United States—I Groundwater, *Sci. Total Environ.* 402 (2008) 192–200.
- [29] C. Potera, Drugged drinking water, *Environ. Health Perspect.* 108 (2000) A446–A449.
- [30] R. Loos, J. Wollgast, T. Huber, G. Hanke, Polar herbicides, pharmaceutical products, perfluorooctanesulfonate (PFOS), perfluorooctanoate (PFOA), and nonylphenol and its carboxylates and ethoxylates in surface and tap waters around Lake Maggiore in Northern Italy, *Anal. Bioanal. Chem.* 387 (2007) 1469–1478.
- [31] M.J. Focazio, D.W. Kolpin, K.K. Barnes, E.T. Furlong, M.T. Meyer, S.D. Zaugg, L.B. Barber, M.E. Thurman, A national reconnaissance for pharmaceuticals and other organic wastewater contaminants in the United States—II) Untreated drinking water sources, *Sci. Total Environ.* 402 (2008) 201–216.
- [32] M.J. Benotti, R.A. Trenholm, B.J. Vanderford, J.C. Holady, B.D. Stanford, S.A. Snyder, Pharmaceuticals and endocrine disrupting compounds in U.S. drinking water, *Environ. Sci. Technol.* 43 (2009) 597–603.
- [33] A.R.R. Péry, M. Gust, B. Vollat, R. Mons, M. Ramil, G. Fink, T. Ternes, J. Garric, Fluoxetine effects assessment on the life cycle of aquatic invertebrates, *Chemosphere* 73 (2008) 300–304.
- [34] K.A. Kidd, P.J. Blanchfield, K.H. Mills, V.P. Palace, R.E. Evans, J.M. Lazorchak, R.W. Flick, Collapse of a fish population after exposure to a synthetic estrogen, *Proc. Natl. Acad. Sci. U.S.A.* 104 (2007) 8897–8901.
- [35] C.G. Daughton, T.A. Ternes, Pharmaceuticals and personal care products in the environment: agents of subtle changes? *Environ. Health Perspect.* 107 (1999) 907–938.
- [36] J. Timbrell, *Principles of Biochemical Toxicology*, third ed., Taylor & Francis, London, 2002.
- [37] R. Braund, B.M. Peake, L. Schieffelin, Disposal practices for unused medications in New Zealand, *Environ. Int.* 35 (2009) 952–955.
- [38] M. Persson, E. Sabelström, B. Gunnarsson, Handling of unused prescription drugs—knowledge, behaviour and attitude among Swedish people, *Environ. Int.* 35 (2009) 771–774.
- [39] T. Heberer, Occurrence, fate, and removal of pharmaceutical residues in the aquatic environment: a review of recent research data, *Toxicol. Lett.* 131 (2002) 5–17.
- [40] J.P. Bound, N. Voulvoulis, Household disposal of pharmaceuticals as a pathway for aquatic contamination in the United Kingdom, *Environ. Health Perspect.* 113 (2005) 1705–1711.
- [41] K. Xia, A. Bhandari, K. Das, G. Pillar, Occurrence and fate of pharmaceuticals and personal care products (PPCPs) in biosolids, *J. Environ. Qual.* 34 (2005) 91–104.
- [42] E. Topp, S.C. Monteiro, A. Beck, B.B. Coelho, A.B.A. Boxall, P.W. Duenk, S. Kleywegt, D.R. Lapen, M. Payne, L. Sabourin, H. Li, C.D. Metcalfe, Runoff of pharmaceuticals and personal care products following application of biosolids to an agricultural field, *Sci. Total Environ.* 396 (2008) 52–59.
- [43] N. Kemper, Veterinary antibiotics in the aquatic and terrestrial environment, *Ecol. Indic.* 8 (2008) 1–13.
- [44] P. Kay, P.A. Blackwell, A.B.A. Boxall, Transport of veterinary antibiotics in overland flow following the application of slurry to arable land, *Chemosphere* 59 (2005) 951–959.
- [45] T.X. Le, Y. Munkage, Residues of selected antibiotics in water and mud from shrimp ponds in mangrove areas in Viet Nam, *Marine Pollut. Bull.* 49 (2004) 922–929.
- [46] G.M. Lalumera, D. Calamari, P. Galli, S. Castiglioni, G. Crosa, R. Fanelli, Preliminary investigation on the environmental occurrence and effects of antibiotics used in aquaculture in Italy, *Chemosphere* 54 (2004) 661–668.
- [47] A.Y.-C. Lin, Y.-T. Tsai, Occurrence of pharmaceuticals in Taiwan's surface waters: impact of waste streams from hospitals and pharmaceutical production facilities, *Sci. Total Environ.* 407 (2009) 3793–3802.
- [48] D. Li, M. Yang, J. Hu, Y. Zhang, H. Chang, F. Jin, Determination of penicillin G and its degradation products in a penicillin production wastewater treatment plant and the receiving river, *Water Res.* 42 (2008) 307–317.
- [49] D.G.J. Larsson, C. Pedro, N. Paxéus, Effluent from drug manufactures contains extremely high levels of pharmaceuticals, *J. Hazard. Mater.* 148 (2007) 751–755.
- [50] A.L. Boreen, W.A. Arnold, K. McNeill, Photodegradation of pharmaceuticals in the aquatic environment: a review, *Aquat. Sci.* 65 (2003) 320–341.
- [51] P. Bartels, W. Tümping Jr., Solar radiation influence on the decomposition process of diclofenac in surface waters, *Sci. Total Environ.* 374 (2007) 143–155.
- [52] M. Crane, C. Watts, T. Boucard, Chronic aquatic environmental risks from exposure to human pharmaceuticals, *Sci. Total Environ.* 367 (2006) 23–41.
- [53] P.H. Roberts, K.V. Thomas, The occurrence of selected pharmaceuticals in wastewater effluent and surface waters of the lower Tyne catchment, *Sci. Total Environ.* 356 (2006) 143–153.
- [54] T.A. Ternes, Occurrence of drugs in German sewage treatment plants and rivers, *Water Res.* 32 (1998) 3245–3260.

- [55] N. Lindqvist, T. Tuhkanen, L. Kronberg, Occurrence of acidic pharmaceuticals in raw and treated sewages and in receiving waters, *Water Res.* 39 (2005) 2219–2228.
- [56] D.L. Sedlak, K.E. Pinkston, Factors affecting the concentrations of pharmaceuticals released to the aquatic environment, *Water Resour. Update* 120 (2001) 56–64.
- [57] M. Cirja, P. Ivashechkin, A. Schäffer, P.F.X. Corvini, Factors affecting the removal of organic micropollutants from wastewater in conventional treatment plants (CTP) and membrane bioreactors (MBR), *Rev. Environ. Sci. Biotechnol.* 7 (2008) 61–78.
- [58] K. Kümmerer, Drugs in the environment: emission of drugs, diagnostic aids and disinfectants into wastewater by hospitals in relation to other sources—a review, *Chemosphere* 45 (2001) 957–969.
- [59] A.Y.-C. Lin, T.-H. Yu, C.-F. Lin, Pharmaceutical contamination in residential, industrial, and agricultural waste streams: risk to aqueous environments in Taiwan, *Chemosphere* 74 (2008) 131–141.
- [60] C. Zwiener, T.J. Gremm, F.H. Frimmel, Pharmaceutical residues in the aquatic environment and their significance for drinking water production, in: K. Kümmerer (Ed.), *Pharmaceuticals in the Environment: Sources, Fate, Effects and Risks*, Springer, Berlin, 2001, pp. 81–89.
- [61] P.E. Stackelberg, E.T. Furlong, M.T. Meyer, S.D. Zaugg, A.K. Henderson, D.B. Reissman, Persistence of pharmaceutical compounds and other organic wastewater contaminants in a conventional drinking-water-treatment plant, *Sci. Total Environ.* 329 (2004) 99–113.
- [62] J.R. Vane, R.M. Botting, Mechanism of action of antiinflammatory drugs, *Int. J. Tissue React.* 20 (1998) 3–15.
- [63] J. Zou, N.F. Neumann, J.W. Holland, M. Belosevic, C. Cunningham, C.J. Secombes, A.F. Rowley, Fish macrophages express a cyclo-oxygenase-2 homologue after activation, *Biochem. J.* 340 (1999) 153–159.
- [64] C.E. Lundholm, DDE-induced eggshell thinning in birds: effects of p,p'-DDE on the calcium and prostaglandin metabolism of the eggshell gland, *Comp. Biol. Physiol.* 118C (1997) 113–128.
- [65] M. Cleuvers, Aquatic ecotoxicity of pharmaceuticals including the assessment of combination effects, *Toxicol. Lett.* 142 (2003) 185–194.
- [66] J. Schwaiger, H. Ferling, U. Mallow, H. Wintermayr, R.D. Negele, Toxic effects of the non-steroidal anti-inflammatory drug diclofenac. Part I: histopathological alterations and bioaccumulation in rainbow trout, *Aquat. Toxicol.* 68 (2004) 141–150.
- [67] R. Triebkorn, H. Casper, A. Heyd, R. Eikemper, H.-R. Köhler, J. Schwaiger, Toxic effects of the non-steroidal anti-inflammatory drug diclofenac. Part II. Cytological effects in liver, kidney, gills and intestine of rainbow trout (*Oncorhynchus mykiss*), *Aquat. Toxicol.* 68 (2004) 151–166.
- [68] B. Hoeger, B. Köllner, D.R. Dietrich, B. Hitzfeld, Water-borne diclofenac affects kidney and gill integrity and selected immune parameters in brown trout (*Salmo trutta f. fario*), *Aquat. Toxicol.* 75 (2005) 53–64.
- [69] M. Schmitt-Jansen, P. Bartels, N. Adler, R. Altenburger, Phytotoxicity assessment of diclofenac and its phototransformation products, *Anal. Bioanal. Chem.* 387 (2007) 1389–1396.
- [70] M.E. DeLorenzo, J. Fleming, Individual and mixture effects of selected pharmaceuticals and personal care products on the marine phytoplankton species *Dunaliella tertiolecta*, *Arch. Environ. Contam. Toxicol.* 54 (2008) 203–210.
- [71] M.D. Hernando, E. Heath, M. Petrovic, D. Barceló, Trace-level determination of pharmaceuticals residues by LC–MS/MS in natural and treated waters. A pilot-survey study, *Anal. Bioanal. Chem.* 385 (2006) 985–991.
- [72] S. Weigel, R. Kallenborn, H. Hühnerfuss, Simultaneous solid-phase extraction of acidic, neutral and basic pharmaceuticals from aqueous samples at ambient (neutral) pH and their determination by gas chromatography-mass spectrometry, *J. Chromatogr. A* 1023 (2004) 183–195.
- [73] M.J. Gómez, M. Petrovic, A.R. Fernández-Alba, D. Barceló, Determination of pharmaceuticals of various therapeutic classes by solid-phase extraction and liquid chromatography-tandem mass spectrometry analysis in hospital effluent wastewaters, *J. Chromatogr. A* 1114 (2006) 224–233.
- [74] J.L. Flippin, D. Huggett, C.M. Foran, Changes in the timing of reproduction following chronic exposure to ibuprofen in Japanese medaka, *Oryzias latipes*, *Aquat. Toxicol.* 81 (2007) 73–78.
- [75] L.-H. Heckmann, A. Callaghan, H.L. Hooper, R. Connon, T.H. Hutchinson, S.J. Maund, R.M. Sibly, Chronic toxicity of ibuprofen to *Daphnia magna*: effects on life history traits and population dynamics, *Toxicol. Lett.* 172 (2007) 137–145.
- [76] H.J. De Lange, W. Noordoven, A.J. Murk, M. Lüring, E.T.H.M. Peeters, Behavioural responses of *Gammarus pulex* (Crustacea, Amphipoda) to low concentrations of pharmaceuticals, *Aquat. Toxicol.* 78 (2006) 209–216.
- [77] F. Pomati, A.G. Netting, D. Calamari, B.A. Neilan, Effects of erythromycin, tetracycline and ibuprofen on the growth of *Synechocystis* sp. and *Lemma minor*, *Aquat. Toxicol.* 67 (2004) 387–396.
- [78] J.-W. Kim, H. Ishibashi, R. Yamauchi, N. Ichikawa, Y. Takao, M. Hirano, M. Koga, K. Arizono, Acute toxicity of pharmaceutical and personal care products on freshwater crustacean (*Thamnocephalus platyurus*) and fish (*Oryzias latipes*), *J. Toxicol. Sci.* 34 (2009) 227–232.
- [79] N. Pounds, S. Maclean, M. Webley, D. Pascoe, T. Hutchinson, Acute and chronic effects of ibuprofen in the mollusc *Planorbis carinatus* (Gastropoda: Planorbidae), *Ecotoxicol. Environ. Saf.* 70 (2008) 47–52.
- [80] M. Isidori, M. Lavorgna, A. Nardelli, A. Parrella, L. Previtera, M. Rubino, Ecotoxicity of naproxen and its phototransformation products, *Sci. Total Environ.* 348 (2005) 93–101.
- [81] P.M. Thomas, G.D. Foster, Determination of nonsteroidal anti-inflammatory drugs, caffeine, and triclosan in wastewater by gas chromatography-mass spectrometry, *J. Environ. Sci. Health. Part A Toxic/Hazard. Subst. Environ. Eng.* A39 (2004) 1969–1978.
- [82] Y. Kim, K. Choi, J. Jung, S. Park, P.-G. Kim, J. Park, Aquatic toxicity of acetaminophen, carbamazepine, cimetidine, diltiazem and six major sulfonamides, and their potential ecological risks in Korea, *Environ. Int.* 33 (2007) 275–370.
- [83] K.-P. Henschel, A. Wenzel, M. Diedrich, A. Fliedner, Environmental hazard assessment of pharmaceuticals, *Regul. Toxicol. Pharm.* 25 (1997) 220–225.
- [84] S. Grujić, T. Vasiljević, M. Lausević, Determination of multiple pharmaceutical classes in surface and ground waters by liquid chromatography-ion trap-tandem mass spectrometry, *J. Chromatogr. A* 1216 (2009) 4989–5000.
- [85] C. Carlsson, A.-K. Johansson, G. Alvan, K. Bergman, T. Kühler, Are pharmaceuticals potent environmental pollutants? Part I: environmental risk assessments of selected active pharmaceutical ingredients, *Sci. Total Environ.* 364 (2006) 67–87.
- [86] N. Nakada, T. Tanishima, H. Shinohara, K. Kiri, H. Takada, Pharmaceutical chemicals and endocrine disruptors in municipal wastewater in Tokyo and their removal during activated sludge treatment, *Water Res.* 40 (2006) 3297–3303.
- [87] A.Y.-C. Lin, T.-H. Yu, S.K. Lateef, Removal of pharmaceuticals in secondary wastewater treatment processes in Taiwan, *J. Hazard. Mater.* (2009), doi:10.1016/j.jhazmat.2009.01.108.
- [88] J.-L. Zhao, G.-G. Ying, L. Wang, J.-F. Yang, X.-B. Yang, L.-H. Yang, X. Li, Determination of phenolic endocrine disrupting chemicals and acidic pharmaceuticals in surface water of the Pearl Rivers in South China by gas chromatography-negative chemical ionization-mass spectrometry, *Sci. Total Environ.* 407 (2009) 962–974.
- [89] J.-Y. Pailler, K.L. Pfister, L. Hoffmann, C. Guignard, Solid phase extraction coupled to liquid chromatography-tandem mass spectrometry analysis of sulfonamides, tetracyclines, analgesics and hormones in surface water and wastewater in Luxembourg, *Sci. Total Environ.* 407 (2009) 4736–4743.
- [90] S.D. Kim, J. Cho, I.S. Kim, B.J. Vanderford, S.A. Snyder, *Water Res.* 41 (2007) 1013–1021.
- [91] M.J.M. Bueno, A. Agüera, M.D. Hernando, M.J. Gómez, A.R. Fernández-Alba, Evaluation of various liquid chromatography-quadrupole-linear ion trap-mass spectrometry operation modes applied to the analysis of organic pollutants in wastewaters, *J. Chromatogr. A* 1216 (2009) 5995–6002.
- [92] J.-W. Kim, H.-S. Jang, J.-G. Kim, H. Ishibashi, M. Hirano, K. Nasu, N. Ichikawa, Y. Takao, R. Shinohara, K. Arizono, Occurrence of pharmaceutical and personal care products (PPCPs) in surface water from Mankyung River, South Korea, *J. Health Sci.* 55 (2009) 249–258.
- [93] K. Choi, Y. Kim, J. Park, C.K. Park, M. Kim, H.S. Kim, P. Kim, Seasonal variations of several pharmaceutical residues in surface water and sewage treatment plants of Han River, Korea, *Sci. Total Environ.* 405 (2008) 120–128.
- [94] M.J. Hilton, K.V. Thomas, Determination of selected human pharmaceutical compounds in effluent and surface water samples by high-performance liquid chromatography-electrospray tandem mass spectrometry, *J. Chromatogr. A* 1015 (2003) 129–141.
- [95] M. Cleuvers, Mixture toxicity of anti-inflammatory drugs diclofenac, ibuprofen, naproxen, and acetylsalicylic acid, *Ecotoxicol. Environ. Saf.* 59 (2004) 309–315.
- [96] B. Ferrari, N. Paxéus, R.L. Giudice, A. Pollio, J. Garric, Ecotoxicological impact of pharmaceuticals found in treated wastewaters: study of carbamazepine, clofibrac acid, and diclofenac, *Ecotoxicol. Environ. Saf.* 55 (2003) 359–370.
- [97] R. Triebkorn, H. Casper, V. Scheil, J. Schwaiger, Ultrastructural effects of pharmaceuticals (carbamazepine, clofibrac acid, metoprolol, diclofenac) in rainbow trout (*Oncorhynchus mykiss*) and common carp (*Cyprinus carpio*), *Anal. Bioanal. Chem.* 387 (2007) 1405–1416.
- [98] B. Quinn, F. Gagné, C. Blaise, An investigation into the acute and chronic toxicity of eleven pharmaceuticals (and their solvents) found in wastewater effluent on the cnidarian, *Hydra attenuate*, *Sci. Total Environ.* 389 (2008) 306–314.
- [99] H.P. Rang, M.M. Dale, J.M. Ritter, *Pharmacology*, fourth ed., Churchill Livingstone, Edinburgh, 1999.
- [100] C.A. Aguilar-Salinas, H. Barrett, G. Schonfeld, Metabolic modes of action of the statins in the hyperlipoproteinemias, *Atherosclerosis* 141 (1998) 203–207.
- [101] P.B. Key, J. Hoguet, L.A. Reed, K.W. Chung, M.H. Fulton, Effects of the statin antihyperlipidemic agent simvastatin on grass shrimp, *Palaemonetes pugio*, *Environ. Toxicol.* 23 (2008) 153–160.
- [102] U. Dahl, E. Gorokhova, M. Breitholtz, Application of growth-related sublethal endpoints in ecotoxicological assessments using a harpacticoid copepod, *Aquat. Toxicol.* 77 (2006) 433–438.
- [103] R.A. Brain, D.J. Johnson, S.M. Richards, M.L. Hanson, H. Sanderson, M.W. Lam, C. Young, S.A. Mabury, P.K. Sibley, K.R. Solomon, Microcosm evaluation of the effects of an eight pharmaceutical mixture to the aquatic macrophytes *Lemma gibba* and *Myriophyllum sibiricum*, *Aquat. Toxicol.* 70 (2004) 23–40.
- [104] S. Debernard, F. Rossignol, F. Couillaud, The HMG-CoA reductase inhibitor fluvastatin inhibits insect juvenile hormone biosynthesis, *Gen. Comp. Endocrinol.* 95 (1994) 92–98.
- [105] X.-S. Miao, C.D. Metcalfe, Determination of cholesterol-lowering statin drugs in aqueous samples using liquid chromatography-electrospray ionization tandem mass spectrometry, *J. Chromatogr. A* 998 (2003) 133–141.



- [106] X.-S. Miao, C.D. Metcalfe, Determination of pharmaceuticals in aqueous samples using positive and negative voltage switching microbore liquid chromatography/electrospray ionization tandem mass spectrometry, *J. Mass Spectrom.* 38 (2003) 27–34.
- [107] B. Staels, J. Dallongeville, J. Auwerx, K. Schoonjans, E. Leitersdorf, J.-C. Fruchart, Mechanism of action of fibrates on lipid and lipoprotein metabolism, *Circulation* 98 (1998) 2088–2093.
- [108] H. Sanderson, R.A. Brain, D.J. Johnson, C.J. Wilson, K.R. Solomon, Toxicity classification and evaluation of four pharmaceutical classes: antibiotics, anti-neoplastics, cardiovascular, and sex hormones, *Toxicology* 203 (2004) 27–40.
- [109] A. Ibabe, A. Herrero, M.P. Cajaraville, Modulation of peroxisome proliferator-activated receptors (PPARs) by PPAR $\alpha$ - and PPAR $\gamma$ -specific ligands and by 17 $\beta$ -estradiol in isolated zebrafish hepatocytes, *Toxicol. In Vitro* 19 (2005) 725–735.
- [110] D. Raldúa, M. André, P.J. Babin, Clofibrate and gemfibrozil induce an embryonic malabsorption syndrome in zebrafish, *Toxicol. Appl. Pharmacol.* 228 (2008) 301–314.
- [111] S.A. Kliewer, S.S. Sundseth, S.A. Jones, P.J. Brown, G.B. Wisely, C.S. Koble, P. Devchand, W. Wahli, T.M. Willson, J.M. Lenhard, J.M. Lehmann, Fatty acids and eicosanoids regulate gene expression through direct interactions with peroxisome proliferation-activated receptors  $\alpha$  and  $\gamma$ , *Proc. Natl. Acad. Sci. U.S.A.* 94 (1997) 4318–4323.
- [112] J.L. Zurita, G. Repetto, A. Jos, M. Salguero, M. López-Artigues, A.M. Cameán, Toxicological effects of the lipid regulator gemfibrozil in four aquatic systems, *Aquat. Toxicol.* 81 (2007) 106–115.
- [113] M. Isidori, A. Nardelli, L. Pascarella, M. Rubino, A. Parrella, Toxic and genotoxic impact of fibrates and their photoproducts on non-target organism, *Environ. Int.* 33 (2007) 635–641.
- [114] C. Mimeault, A.J. Woodhouse, X.-S. Miao, C.D. Metcalfe, T.W. Moon, V.L. Trudeau, The human lipid regulator, gemfibrozil bioconcentrates and reduces testosterone in the goldfish, *Carassius auratus*, *Aquat. Toxicol.* 73 (2005) 44–54.
- [115] J.P. Emblidge, M.E. DeLorenzo, Preliminary risk assessment of the lipid-regulating pharmaceutical clofibrate, for three estuarine species, *Environ. Res.* 100 (2006) 216–226.
- [116] C.M. Flaherty, S.I. Dodson, Effects of pharmaceuticals on *Daphnia* survival, growth, and reproduction, *Chemosphere* 61 (2005) 200–207.
- [117] T.J. Runnalls, D.N. Hala, J.P. Sumpter, Preliminary studies into the effects of the human pharmaceutical Clofibrate acid on sperm parameters in adult Fathead minnow, *Aquat. Toxicol.* 84 (2007) 111–118.
- [118] S. Castiglioni, R. Bagnati, D. Calamari, R. Fanelli, E. Zuccato, A multiresidue analytical method using solid-phase extraction and high-pressure chromatography tandem mass spectrometry to measure pharmaceuticals of different therapeutic classes in urban wastewaters, *J. Chromatogr. A* 1092 (2005) 206–215.
- [119] K. Reddersen, T. Heberer, Multi-compound methods for the detection of pharmaceutical residues in various waters applying solid phase extraction (SPE) and gas chromatography with mass spectrometric (GC–MS) detection, *J. Sep. Sci.* 26 (2003) 1443–1450.
- [120] O.A.H. Jones, N. Voulvoulis, J.N. Lester, Aquatic environmental assessment of the top 25 English prescription pharmaceuticals, *Water Res.* 36 (2002) 5013–5022.
- [121] B. Halling-Sørensen, Algal toxicity of antibacterial agents used in intensive farming, *Chemosphere* 40 (2000) 731–739.
- [122] H.C. Holten Lützhøft, B. Halling-Sørensen, S.E. Jørgensen, Algal toxicity of antibacterial agents applied in Danish farming, *Arch. Environ. Contam. Toxicol.* 36 (1999) 1–6.
- [123] P.F. Lanzky, B. Halling-Sørensen, The toxic effect of the antibiotic metronidazole on aquatic organisms, *Chemosphere* 35 (1997) 2553–2561.
- [124] M. Isidori, M. Lavorgna, A. Nardelli, L. Pascarella, A. Parrella, Toxic and genotoxic evaluation of six antibiotics on non-target organisms, *Sci. Total Environ.* 346 (2005) 87–98.
- [125] T. Kumpel, R. Alexy, K. Kümmerer, What do we know about antibiotics in the environment? in: K. Kümmerer (Ed.), *Pharmaceuticals in the Environment: Sources, Fate, Effects and Risks*, Springer, Berlin, 2001, pp. 67–76.
- [126] R. Andreozzi, V. Caprio, C. Ciniglia, M. De Champdoré, R. Lo Giudice, R. Marotta, E. Zuccato, Antibiotics in the environment: occurrence in Italian STPs, fate, and preliminary assessment on algal toxicity of amoxicillin *Environ. Sci. Technol.* 38 (2004) 6832–6838.
- [127] X. Nie, J. Gu, J. Lu, W. Pan, Y. Yang, Effects of norfloxacin and butylated hydroxyanisole on the freshwater microalga *Scenedesmus obliquus*, *Ecotoxicology* 18 (2009) 677–684.
- [128] A.B. Boxall, D.W. Kolpin, B. Halling-Sørensen, J. Tolls, Are veterinary medicines causing environmental risks? *Environ. Sci. Technol.* 37 (2003) 286A–294A.
- [129] L. Migliore, C. Civitareale, S. Cozzolino, P. Casoria, G. Brambilla, L. Gaudio, Laboratory models to evaluate phytotoxicity of sulphadimethoxine on terrestrial plants, *Chemosphere* 37 (1998) 2957–2961.
- [130] S.D. Costanzo, J. Murby, J. Bates, Ecosystem response to antibiotics entering the aquatic environment, *Mar. Pollut. Bull.* 51 (2005) 218–223.
- [131] K. Eguchi, H. Nagase, M. Ozawa, Y.S. Endoh, K. Goto, K. Hirata, K. Miyamoto, H. Yoshimura, Evaluation of antimicrobial agents for veterinary use in the ecotoxicity test using microalgae, *Chemosphere* 57 (2004) 1733–1738.
- [132] N. Yamashita, M. Yasojima, K. Miyajima, Y. Suzuki, H. Tanaka, Effects of antibacterial agents, levofloxacin and clarithromycin, on aquatic organisms, *Water Sci. Technol.* 53 (2006) 65–72.
- [133] J. Pro, J.A. Ortiz, S. Boleas, C. Fernández, G. Carbonell, J.V. Tarazona, Effect assessment of antimicrobial pharmaceuticals on the aquatic plant *Lemna minor*, *Bull. Environ. Contam. Toxicol.* 70 (2003) 290–295.
- [134] L. Wollenberger, B. Halling-Sørensen, K.O. Kusk, Acute and chronic toxicity of veterinary antibiotics to *Daphnia magna*, *Chemosphere* 40 (2000) 723–730.
- [135] M. De Liguoro, B. Fioretto, C. Poltronieri, G. Gallina, The toxicity of sulfamethazine to *Daphnia magna* and its additivity to other veterinary sulfonamides and trimethoprim, *Chemosphere* 75 (2009) 1519–1524.
- [136] S. Park, K. Choi, Hazard assessment of commonly used agricultural antibiotics on aquatic ecosystems, *Ecotoxicology* 17 (2008) 526–538.
- [137] S. Jiao, S. Zheng, D. Yin, L. Wang, L. Chen, Aqueous photolysis of tetracycline and toxicity of photolytic products to luminescent bacteria, *Chemosphere* 73 (2008) 377–382.
- [138] K.D. Brown, J. Kulis, B. Thomson, T.H. Chapman, D.B. Mawhinney, Occurrence of antibiotics in hospital, residential, and dairy effluent, municipal wastewater, and the Rio Grande in New Mexico, *Sci. Total Environ.* 366 (2006) 772–783.
- [139] M. Seifrtová, A. Pena, C.M. Lino, P. Solich, Determination of fluorquinolone antibiotics in hospital and municipal wastewaters in Coimbra by liquid chromatography with a monolithic column and fluorescence detection, *Anal. Bioanal. Chem.* 391 (2008) 799–805.
- [140] K.G. Karthikeyan, M.T. Meyer, Occurrence of antibiotics in wastewater treatment facilities in Wisconsin, USA, *Sci. Total Environ.* 361 (2006) 196–207.
- [141] R.H. Lindberg, P. Wennberg, M.I. Johansson, M. Tysklind, B.A.V. Andersson, Screening of human antibiotic substances and determination of weekly mass flows in five sewage treatment plants in Sweden, *Environ. Sci. Technol.* 39 (2005) 3421–3429.
- [142] A. Pena, D. Chmielova, C.M. Lino, P. Solich, Determination of fluorquinolone antibiotics in surface waters from Mondego River by high performance liquid chromatography using a monolithic column, *J. Sep. Sci.* 30 (2007) 2924–2928.
- [143] D. Perret, A. Gentili, S. Marchese, A. Greco, R. Curini, Sulphonamide residues in Italian surface and drinking waters: a small scale reconnaissance, *Chromatographia* 63 (2006) 225–232.
- [144] A. Gulkowska, Y. He, M.K. So, L.W.Y. Yeung, H.W. Leung, J.P. Giesy, P.K.S. Lam, M. Martin, B.J. Richardson, The occurrence of selected antibiotics in Hong Kong coastal waters, *Mar. Pollut. Bull.* 54 (2007) 1287–1306.
- [145] W.-H. Xu, G. Zhang, S.-C. Zou, X.-D. Li, Y.-C. Liu, Determination of selected antibiotics in the Victoria Harbour and the Pearl River, South China using high-performance liquid chromatography-electrospray ionization tandem mass spectrometry, *Environ. Pollut.* 145 (2007) 672–679.
- [146] D.G.J. Larsson, M. Adolfsson-Erici, J. Parkkonen, M. Pettersson, A.H. Berg, P.-E. Olsson, L. Förlin, Ethinylestradiol—an undesired fish contraceptive? *Aquat. Toxicol.* 45 (1999) 91–97.
- [147] S. Jobling, M. Nolan, C.R. Tyler, G. Brighty, J.P. Sumpter, Widespread sexual disruption in wild fish, *Environ. Sci. Technol.* 32 (1998) 2498–2506.
- [148] J.L. Parrott, B.R. Blunt, Life-cycle exposure of fathead minnows (*Pimephales promelas*) to an ethinylestradiol concentration below 1 ng/L reduces egg fertilization success and desmasculinizes males, *Environ. Toxicol.* 20 (2005) 131–141.
- [149] S. Pawlowski, R. van Aerle, C.R. Tyler, T. Braunbeck, Effects of 17 $\alpha$ -ethinylestradiol in a fathead minnow (*Pimephales promelas*) gonadal recrudescence assay, *Ecotoxicol. Environ. Saf.* 57 (2004) 330–345.
- [150] S. Örn, H. Holbech, T.H. Madsen, L. Norrgren, G.I. Petersen, Gonad development and vitellogenin production in zebrafish (*Danio rerio*) exposed to ethinylestradiol and methyltestosterone, *Aquat. Toxicol.* 65 (2003) 397–411.
- [151] J.P. Nash, D.E. Kime, L.T.M. Van der Vem, P.W. Wester, F. Brion, G. Maack, P. Stahlschmidt-Allner, C.R. Tyler, Long-term exposure to environmental concentrations of the pharmaceutical ethinylestradiol causes reproductive failure in fish, *Environ. Health Perspect.* 112 (2004) 1725–1733.
- [152] H. Xu, J. Yang, Y. Wang, Q. Jiang, H. Chen, H. Song, Exposure to 17 $\alpha$ -ethinylestradiol impairs reproductive functions of both male and female zebrafish (*Danio rerio*), *Aquat. Toxicol.* 88 (2008) 1–8.
- [153] Y. Katsu, A. Lange, H. Urushitani, R. Ichikawa, G.C. Paull, L.L. Cahill, S. Jobling, C.R. Tyler, T. Iguchi, Functional associations between two estrogen receptors, environmental estrogens, and sexual disruption in the roach (*Rutilus rutilus*), *Environ. Sci. Technol.* 41 (2007) 3368–3374.
- [154] N. Hirai, A. Nanba, M. Koshio, T. Kondo, M. Morita, N. Tatarazako, Feminization of Japanese medaka (*Oryzias latipes*) exposed to 17 $\beta$ -estradiol: Formation of testis-ova and sex-transformation during early-ontogeny, *Aquat. Toxicol.* 77 (2006) 78–86.
- [155] I.J. Kang, H. Yokota, Y. Oshima, Y. Tsuruda, T. Yamaguchi, M. Maeda, N. Imada, H. Tadokoro, T. Honjo, Effect of 17 $\beta$ -estradiol on the reproduction of Japanese medaka (*Oryzias latipes*), *Chemosphere* 47 (2002) 71–80.
- [156] E.F. Orlando, L.J. Guillelte Jr., Sexual dimorphic responses in wildlife exposed to endocrine disrupting chemicals, *Environ. Res.* 104 (2007) 163–173.
- [157] I. Gyllenhammar, L. Holm, R. Eklund, C. Berg, Reproductive toxicity in *Xenopus tropicalis* after developmental exposure to environmental concentrations of ethinylestradiol, *Aquat. Toxicol.* 91 (2009) 171–178.
- [158] G.F. Vandenbergh, D. Adriaens, T. Verslycke, C.R. Janssen, Effects of 17 $\alpha$ -ethinylestradiol on sexual development of the amphipod *Hyalella azteca*, *Ecotoxicol. Environ. Saf.* 54 (2003) 216–222.
- [159] J.A. Jukosky, M.C. Watzin, J.C. Leiter, Elevated concentrations of ethinylestradiol, 17 $\beta$ -estradiol, and medroxyprogesterone have little effect on reproduction and survival of *Ceriodaphnia dubia*, *Bull. Environ. Contam. Toxicol.* 81 (2008) 230–235.

- [160] T. Isobe, H. Shiraishi, M. Yasuda, A. Shinoda, H. Suzuki, M. Morita, Determination of estrogens and their conjugates in water using solid-phase extraction followed by liquid chromatography-tandem mass spectrometry, *J. Chromatogr. A* 984 (2003) 195–202.
- [161] S. Zuehlke, U. Duennbier, T. Heberer, Determination of estrogenic steroids in surface water and wastewater by liquid chromatography-electrospray tandem mass spectrometry, *J. Sep. Sci.* 28 (2005) 52–58.
- [162] L. Yang, T. Luan, C. Lan, Solid-phase microextraction with on-fiber silylation for simultaneous determinations of endocrine disrupting chemicals and steroid hormones by gas chromatography-mass spectrometry, *J. Chromatogr. A* 1104 (2006) 23–32.
- [163] A. Laganà, A. Bacaloni, I. De Leva, A. Faberi, G. Fago, A. Marino, Analytical methodologies for determining the occurrence of endocrine disrupting chemicals in sewage treatment plants and natural waters, *Anal. Chim. Acta* 501 (2004) 79–88.
- [164] E. Vulliet, L. Wiest, R. Baudot, M.-F. Grenier-Loustalot, Multi-residue analysis of steroids at sub-ng/L levels in surface and ground-waters using liquid chromatography coupled to tandem mass spectrometry, *J. Chromatogr. A* 1210 (2008) 84–91.
- [165] P.D. Thacker, Pharmaceutical data elude researchers, *Environ. Sci. Technol.* 39 (2005) 193A–194A.
- [166] M. Lüring, E. Sargant, I. Roessink, Life-history consequences for *Daphnia pulex* exposed to pharmaceutical carbamazepine, *Environ. Toxicol.* 21 (2006) 172–180.
- [167] M. Oetken, G. Nentwig, D. Löffler, T. Ternes, J. Oehlmann, Effects of pharmaceuticals on aquatic invertebrates. Part I. The antiepileptic drug carbamazepine, *Arch. Environ. Contam. Toxicol.* 49 (2005) 353–361.
- [168] F. Sacher, M. Ehmann, S. Gabriel, C. Graf, H.-J. Brauch, Pharmaceutical residues in the river Rhine—results of a one-decade monitoring programme, *J. Environ. Monit.* 10 (2008) 664–670.
- [169] M. Leclercq, O. Mathieu, E. Gomez, C. Casellas, H. Fenet, D. Hillaire-Buys, Presence and fate of carbamazepine, oxcarbazepine, and seven of their metabolites at wastewater treatment plants, *Arch. Environ. Contam. Toxicol.* 56 (2009) 408–415.
- [170] J.G. Nickerson, S.G. Dugan, G. Drouin, T.W. Moon, A putative  $\alpha$ -adrenoceptor from the rainbow trout (*Oncorhynchus mykiss*). Molecular characterization and pharmacology, *Eur. J. Biochem.* 268 (2001) 6465–6472.
- [171] S. Haider, S.S.R. Baqri,  $\alpha$ -Adrenoceptor antagonists reinitiate meiotic maturation in *Clarias batrachus* oocytes, *Comp. Biochem. Physiol. A* 126 (2000) 517–525.
- [172] D.B. Huggett, B.W. Brooks, B. Peterson, C.M. Foran, D. Schlenk, Toxicity of select beta adrenergic receptor-blocking pharmaceuticals (B-Blockers) on aquatic organisms, *Arch. Environ. Contam. Toxicol.* 43 (2002) 229–235.
- [173] D.G.J. Larsson, S. Fredriksson, E. Sandblom, N. Paxeus, M. Axelsson, Is heart rate in fish a sensitive indicator to evaluate acute effects of  $\beta$ -blockers in surface water? *Environ. Toxicol. Pharmacol.* 22 (2006) 338–340.
- [174] M.J. Winter, A.D. Lillicrap, J.E. Caunter, C. Schaffner, A.C. Alder, M. Ramil, T.A. Ternes, E. Giltrow, J.P. Sumpter, T.H. Hutchinson, Defining the chronic impacts of atenolol on embryo-larval development and reproduction in the fathead minnow (*Pimephales promelas*), *Aquat. Toxicol.* 86 (2008) 361–369.
- [175] M. Cleuvers, Initial risk assessment for three  $\beta$ -blockers found in the aquatic environment, *Chemosphere* 59 (2005) 199–205.
- [176] E.M. Dzialowski, P.K. Turner, B.W. Brooks, Physiological and reproductive effects of beta adrenergic receptor antagonists in *Daphnia magna*, *Arch. Environ. Contam. Toxicol.* 50 (2006) 503–510.
- [177] B.W. Brooks, C.M. Foran, S.M. Richards, J. Weston, P.K. Turner, J.K. Stanley, K.R. Solomon, M. Slattery, T.W. LaPoint, Aquatic ecotoxicology of fluoxetine, *Toxicol. Lett.* 142 (2003) 169–183.
- [178] P.P. Fong, Zebra mussel spawning is induced in low concentrations of putative serotonin reuptake inhibitors, *Biol. Bull.* 194 (1998) 143–149.
- [179] G. Nentwig, Effects of pharmaceuticals on aquatic invertebrates. Part II: the antidepressant drug fluoxetine, *Arch. Environ. Contam. Toxicol.* 52 (2007) 163–170.
- [180] C.M. Foran, J. Weston, M. Slattery, B.W. Brooks, D.B. Huggett, Reproductive assessment of Japanese medaka (*Oryzias latipes*) following a four-week fluoxetine (SSRI) exposure, *Arch. Environ. Contam. Toxicol.* 46 (2004) 511–517.
- [181] T.B. Henry, M.C. Black, Acute and chronic toxicity of fluoxetine (selective serotonin reuptake inhibitor) in western mosquitofish, *Arch. Environ. Contam. Toxicol.* 54 (2008) 325–330.
- [182] E. Minagh, R. Herman, K. O'Rourke, F.M. Lyng, M. Davoren, Aquatic ecotoxicity of the selective serotonin reuptake inhibitor sertraline hydrochloride in a battery of freshwater test species, *Ecotoxicol. Environ. Saf.* 72 (2009) 434–440.
- [183] D.J. Johnson, H. Sanderson, R.A. Brain, C.J. Wilson, K.R. Solomon, Toxicity and hazard of selective serotonin reuptake inhibitor antidepressants fluoxetine, fluvoxamine, and sertraline to algae, *Ecotoxicol. Environ. Saf.* 67 (2007) 128–139.
- [184] B.W. Brooks, P.K. Turner, J.K. Stanley, J.J. Weston, E.A. Glidewell, C.M. Foran, M. Slattery, T.W. LaPoint, D.B. Huggett, Waterborne and sediment toxicity of fluoxetine to select organisms, *Chemosphere* 52 (2003) 135–142.
- [185] V.L. Cunningham, D.J.C. Constable, R.E. Hannah, Environmental risk assessment of paroxetine, *Environ. Sci. Technol.* 38 (2004) 3351–3359.
- [186] T. Vasskog, U. Berger, P.-J. Samuelsen, R. Kallenborn, E. Jensen, Selective serotonin reuptake inhibitors in sewage influents and effluents from Tromsø, Norway, *J. Chromatogr. A* 1115 (2006) 187–195.
- [187] T. Vasskog, T. Anderssen, S. Pedersen-Bjergaard, R. Kallenborn, E. Jensen, Occurrence of selective serotonin reuptake inhibitors in sewage and receiving waters at Spitsbergen and in Norway, *J. Chromatogr. A* 1185 (2008) 194–205.
- [188] A. Lajeunesse, C. Gagnon, S. Sauvé, Determination of basic antidepressants and their *N*-desmethyl metabolites in raw sewage and wastewater using solid-phase extraction and liquid chromatography-tandem mass, *Anal. Chem.* 80 (2008) 5325–5333.
- [189] A.C. Johnson, M.D. Jürgens, R.J. Williams, K. Kümmerer, A. Kortenkamp, J.P. Sumpter, Do cytotoxic chemotherapy drugs discharged into rivers pose a risk to the environment and human health? An overview and UK case study, *J. Hydrol.* 348 (2008) 167–175.
- [190] M. Grung, T. Källqvist, S. Sakshaug, S. Skurtveit, K.V. Thomas, Environmental assessment of Norwegian priority pharmaceuticals based on the EMEA guideline, *Ecotoxicol. Environ. Saf.* 71 (2008) 328–340.
- [191] M. DellaGreca, M.R. Iesce, M. Isidori, A. Nardelli, L. Previtera, M. Rubino, Phototransformation products of tamoxifen by sunlight in water. Toxicity of the drug and its derivatives on aquatic organisms, *Chemosphere* 67 (2007) 1933–1939.
- [192] T. Steger-Hartmann, K. Kümmerer, A. Hartmann, Biological degradation of cyclophosphamide and its occurrence in sewage water, *Ecotoxicol. Environ. Saf.* 36 (1997) 174–179.
- [193] I.J. Buerge, H.-R. Buser, T. Poiger, M.D. Müller, Occurrence and fate of the cytostatic drugs cyclophosphamide and ifosfamide in wastewater and surface waters, *Environ. Sci. Technol.* 40 (2006) 7242–7250.
- [194] J.L. Zhou, Z.L. Zhang, E. Banks, D. Grover, J.Q. Jiang, Pharmaceutical residues in wastewater treatment works effluents and their impact on receiving river water, *J. Hazard. Mater.* 166 (2009) 655–661.
- [195] S. Pérez, D. Barceló, Fate and occurrence of X-ray contrast media in the environment, *Anal. Bioanal. Chem.* 387 (2007) 1235–1246.
- [196] T. Steger-Hartmann, R. Länge, H. Schweinfurth, M. Tschampel, I. Rehmann, Investigations into the environmental fate and effects of iopromide (ultravist), a widely used iodinated X-ray contrast medium, *Water Res.* 36 (2002) 266–274.
- [197] T. Steger-Hartmann, R. Länge, H. Schweinfurth, Environmental risk assessment for the widely used iodinated X-ray contrast agent iopromide (Ultravist), *Ecotoxicol. Environ. Saf.* 42 (1999) 274–281.
- [198] M. Carballa, F. Omil, J.M. Lema, M. Llompert, C. Garcia-Jares, I. Rodríguez, M. Gómez, T. Ternes, Behavior of pharmaceuticals, cosmetics and hormones in a sewage treatment plant, *Water Res.* 38 (2004) 2918–2926.
- [199] T.A. Ternes, R. Hirsch, Occurrence and behavior of X-ray contrast media in sewage facilities and the aquatic environment, *Environ. Sci. Technol.* 34 (2000) 2741–2748.
- [200] R.A. Trenholm, B.J. Vanderford, J.C. Holady, D.J. Rexing, S.A. Snyder, Broad range analysis of endocrine disruptors and pharmaceuticals using gas chromatography and liquid chromatography tandem mass spectrometry, *Chemosphere* 65 (2006) 1990–1998.
- [201] F. Busetti, K.L. Linge, J.W. Blythe, A. Heitz, Rapid analysis of iodinated X-ray contrast media in secondary and tertiary treated wastewater by direct injection liquid-chromatography-tandem mass spectrometry, *J. Chromatogr. A* 1213 (2008) 200–208.
- [202] W. Seitz, W.H. Weber, J.-Q. Jiang, B.J. Lloyd, M. Maier, D. Maier, W. Schulz, Monitoring of iodinated X-ray contrast media in surface water, *Chemosphere* 64 (2006) 1318–1324.
- [203] A. Putschew, S. Wischnack, M. Jekel, Occurrence of triiodinated X-ray contrast agents in the aquatic environment, *Sci. Total Environ.* 255 (2000) 129–134.
- [204] R. Hirsch, T.A. Ternes, A. Lindart, K. Haberer, R.-D. Wilken, A sensitive method for the determination of iodine containing diagnostic agents in aqueous matrices using LC-electrospray-tandem-MS detection, *Fresenius J. Anal. Chem.* 366 (2000) 835–841.
- [205] M. Farré, S. Pérez, L. Kantiani, D. Barceló, Fate and toxicity of emerging pollutants, their metabolites and transformation products in the aquatic environment, *Trends Anal. Chem.* 27 (2008) 991–1007.
- [206] M.J. García-Galán, M.S. Díaz-Cruz, D. Barceló, Identification and determination of metabolites and degradation products of sulfonamide antibiotics, *Trends Anal. Chem.* 27 (2008) 1008–1022.
- [207] B. Quinn, F. Gagné, C. Blaise, Evaluation of the acute, chronic and teratogenic effects of a mixture of eleven pharmaceuticals on the cnidarian, *Hydra attenuata*, *Sci. Total Environ.* 407 (2009) 1072–1079.
- [208] U. Borgmann, D.T. Bennie, A.L. Ball, V. Palabrica, Effect of a mixture of seven pharmaceuticals on *Hyalella azteca* over multiple generations, *Chemosphere* 66 (2007) 1278–1283.
- [209] J.L. Parrott, D.T. Bennie, Life-cycle exposure of fathead minnows to a mixture of six common pharmaceuticals and tri-closan, *J. Toxicol. Environ. Health A* 72 (2009) 633–641.
- [210] Commission Directive 92/18/EEC, Modifying the Annex to Council Directive 81/852/EEC on the Approximation of the Laws of Member States Relating to Analytical, Pharmacotoxicological and Clinical Standards and Protocols in Respect of the Testing of Veterinary Medicinal Products, 1992.
- [211] EMEA, Note for Guidance: Environmental Risk Assessment for Veterinary Medicinal Products Other Than GMO-Containing and Immunological Products, The European Agency for the Evaluation of Medicinal Products: Veterinary Medicines Evaluation Unit, EMEA/CVMP/055/96-FINAL, 1998.
- [212] EudraLex – Volume 1 – Pharmaceutical Legislation Medicinal Products for Human Use, On line at: <http://ec.europa.eu/enterprise/pharmaceuticals/eudralex/vol1/en.htm> (accessed in 16 February 2009).

- [213] EMEA, Guideline on the Environmental Risk Assessment of Medicinal Products for Human Use, The European Agency for the Evaluation of Medicinal Products: Committee for Medicinal Products for Human Use, EMEA/CHMP/SWP/4447/00, 2006.
- [214] FDA, Guidance for Industry: Environmental Assessment of Human Drug and Biologics Applications, Food and Drug Administration (Center for Drug Evaluation and Research), CMC 6, Revision 1, 1998.
- [215] EMEA, Guideline on Environmental Impact Assessment (EIAs) for Veterinary Medicinal Products—Phase I, The European Agency for the Evaluation of Medicinal Products: Committee for Medicinal Products for Veterinary Use, CVMP/VICH/592/98-FINAL, 2000.
- [216] EMEA, Guideline on Environmental Impact Assessment for Veterinary Medicinal Products—Phase II, The European Agency for the Evaluation of Medicinal Products: Committee for Medicinal Products for Veterinary Use, CVMP/VICH/790/03-FINAL, 2005.