

Is there sufficient training of healthcare staff on noise reduction in Neonatal Intensive Care Units? A Pilot Study from NeoNoise Project

Carlos Carvalhais^{a,b*}, Joana Santos^{a,b}, Manuela Vieira da Silva^{a,b}, Ana Xavier^{a,b}

^a *Environmental Health Department, School of Allied Health Technologies of Polytechnic Institute of Porto (ESTSP.IPP), R. Valente Perfeito, no. 322, 4400-330 Vila Nova de Gaia, Portugal*

Phone: +351. 22 206 10 00; Fax: +351. 22 206 10 01; Email: caa@estsp.ipp.pt / jds@estsp.ipp.pt / m.silva@eu.ipp.pt / arfx@estsp.ipp.pt

^b *Research Centre on Health and Environment (CISA), School of Allied Health Technologies of Polytechnic Institute of Porto (ESTSP.IPP), R. Valente Perfeito, no. 322, 4400-330 Vila Nova de Gaia, Portugal*

Phone: +351. 22 206 10 00; Fax: +351. 22 206 10 01; Email: caa@estsp.ipp.pt / jds@estsp.ipp.pt / m.silva@eu.ipp.pt / arfx@estsp.ipp.pt

* *Contact person*

Running head title: Impact of a training program on noise reduction

ABSTRACT

Evidence indicates that exposure to high levels of noise adversely affects human health, and these effects are dependent upon various factors. In hospitals, there are many sources of noise, and high levels exert an impact on patients and staff, increasing both recovery time and stress, respectively. The goal of this pilot study was to develop, implement and evaluate the effectiveness of a training program (TP) on noise reduction in a Neonatal Intensive Care Units (NICU) by comparing the noise levels before and after the implementation of the program. A total of 79 health professionals participated in the study. The measurements of sound pressure levels took into account the layout of the unit and location of the main sources of noise. General results indicated that L_{Aeq} levels before implementation of the training program were often excessive, ranging from 48.7 ± 2.94 dBA to 71.7 ± 4.74 dBA, exceeding international guidelines. Similarly following implementation of the training program noise levels remained unchanged (54.5 ± 0.49 dBA to 63.9 ± 4.37 dBA), despite a decrease in some locations. There was no significant difference before and after the implementation of TP. However a significant difference was found for $L_{p, Cpeak}$, before and after training staff, suggesting greater care by healthcare professionals performing their tasks. Even recognizing that a TP is quite important to change behaviors, this needs to be considered in a broader context to effectively control noise in the NICU.

INTRODUCTION

Evidence indicates that human exposure to high levels of noise produces physiological and psychological disorders, and that these effects are dependent upon various factors (Guthrie et al. 2014). Noise in neonatal intensive care units (NICU) is recognized as an agent with negative implications on health and well-being of premature infants (Nicolau et al. 2005) and health professionals. Philbin and Gray (2002) documented that sound pressure levels in intensive care units ranged between 55 to 75 A-weighted decibels (dBA). These results are higher than the limit recommended by the World Health Organization (WHO), which recommends that the average background noise in hospitals should not exceed 35 dB L_{Aeq} for areas where patients are treated or observed. For wardrooms in hospitals the guideline values indoors are 30 dBA L_{Aeq} (A-weighted equivalent sound pressure level) with a corresponding L_{Amax} (maximum A-weighted sound pressure level) of 40 dBA (Berglund et al. 1999). These levels are influenced by the equipment (including alarms, monitors, ventilators, infusion pumps, nebulizers) and by health professionals/visitors behavior (with tasks and conversation) (Short et al. 2011), as shown in Table 1 (Pugh and Griffiths 2007).

In general high-intensity noise levels may induce physiological instabilities in newborns, such as apnea, bradycardia and abrupt fluctuations in heart rate, respiratory rate, blood pressure and oxygen saturation (Philbin and Klaas 2000; Wachman and Lahav 2011). Complex exposures to multiple chemical and physical agents, such noise, have the potential to produce several different sorts of interaction with regard to health outcomes (Fechter 2004; Guthrie et al. 2014). Actually, noise acting in synergy with ototoxic drugs may increase the risk of sensorineural hearing loss in premature infants (AAP 1997; Surenthiran et al. 2003). Other long-term negative effects include language difficulties and altered brain development (Brown 2009), abnormal auditory development as well as suggestion of a link between

excessive noise and attention deficit hyperactivity disorder (Bremmer et al.2003). Health professionals are concerned about this issue and identified noise as a barrier to work performance (Gurses and Carayon 2009; Sampaio Neto et al. 2010). In fact, noise may induce extra-auditory effects in professionals including burnout, stress and fatigue, which results in errors (Mahmood et al. 2011).

In Portugal there has been a considerable increase in preterm births, which in 2004 increased from 6.7% to 8.8% in 2009 (Machado et al. 2011). Thus, it is essential to promote a quiet environment to reduce the impact of noise levels on health and well-being of premature infants and health professionals. Environmental modifications might effectively decrease noise levels (Philbin and Gray 2002; Philbin and Klaas 2000; Ramesh et al. 2009); however, the process of caring for hospitalized patients, require frequent and ongoing interpersonal discussions. Minimizing patient exposure to interpersonal communications between healthcare staff requires a behavioral change. A well-structured training program seems to be a low-cost measure to begin noise reduction process in a hospital environment (Tsunemi et al. 2012).

The aims of this study were to (1) assess the levels of noise by measuring the equivalent sound pressure levels in zones located in NICU of a Portuguese hospital and (2) examine the influence of implementation of a training program (TP) for staff on noise reduction.

MATERIALS AND METHODS

Clinical Settings

All measurements were performed in a NICU of a hospital located in Porto, Portugal, between July 2011 (first phase – measurements taken prior to implementation of TP) and July 2012 (second phase – measurements obtained 6 months after the implementation of TP).

The clinical/technical area of the unit consists of two rooms (A – Intensive Care and B – Special Care), without total separation between infrastructures. Room A includes the integrated set of physical, technical and human expertise, where newborns in critical condition with failure of vital bodily functions are assisted by advanced life support for 24 hr/day. Room B also includes an integrated set of physical, technical and human expertise, intended to provide care for infants with failure of an organ or system, but not intended for neonates requiring mechanical ventilation. The existing physical infrastructure separating the compartments consists of plywood with glazed surface on top. The floor is concrete with vinyl covering, walls are made of painted plasterboard with three glass windows and the roof is concrete. NICU (rooms A and B) has capacity to provide care for approximately 19 patients with a total of 14 incubators and 5 nurseries. Both preparation of parenteral nutrition and medication are located in a common area of the two rooms, but the entrance is accessed through room A, and consequently underwent greatest amount of staff/traffic activity.

Noise Measurements

Noise level measurements were performed using a sound level meter class 1 (01 dB®, model Solo-Premium). In accordance with Robertson et al. (1998), a preliminary survey was performed in order to identify noise sources. In both rooms (A and B) measurements were made continuously over 24 hours in two areas: work stations and traffic zones (at least 1m away from the walls at a height between 1m and 1.65m). In room A noise was also determined inside an incubator. The measurements of peak sound pressure level ($L_{p, Cpeak}$) were made using the C filter and the A-weighted equivalent sound pressure level (L_{Aeq}) were obtained using the A filter, which is a frequency weighting filter that simulates human hearing. Slow response time averaging (1 sec) was also used because is the most appropriate response for the majority of the applications in hospitals and provide stable readings,

according to Philbin and Gray (2002). To ensure accurate measurement, recording was preceded by calibration of the sound level meter (Kent et al. 2002), with an acoustic calibrator class 1 (RION®, model NC-74). In analysis and interpretation of results reference values given by WHO were used (Berglund et al. 1999).

Training Program (TP)

The TP was performed through a lecture of approximately 60 min and conducted by the investigators. In order to ensure that all the staff of the NICU under study such as physicians, nursing staff and auxiliary staff attended the lecture (n=79), 14 training sessions were given. The lecture included the results of the sound pressure levels obtained in the first phase and comparing these to the recommended values suggested by WHO and other regulatory agencies. The negative impact of noise on health, both for neonates and professionals, was also discussed and some actions that needed to be implemented to ensure noise reduction were undertaken. Regarding these actions, the health professionals were led to discuss and reflect on current practice, framing the problem. The health professionals were encouraged to develop an action plan to address specific noise issues, in order to be involved in the process and obtain their commitment for future implementation of noise reduction protocols. Without their commitment, the transfer of knowledge may not be effective in changing practices and behavior.

Statistical Analysis

The processing and data analysis involved descriptive statistics, with analysis of L_{Aeq} and L_p, C_{peak} values. All tests considered a 95% confidence interval. The normality Shapiro-Wilk test and the Student's t test for paired samples were applied. The software IBM SPSS™ (Statistical Package for the Social Sciences) 20th version and MS Excel® 2013 were used for the analysis.

RESULTS

The training content and the staff perception regarding the main sources of noise in the NICU and suggestions to decrease noise in these environments are presented in Table 2. In general, the healthcare staff identified equipment, visitors, healthcare procedures, traffic inside the rooms and team conversation as the main sources of noise in the NICU.

The results obtained in the two rooms of NICU before and after implementation of TP are shown in Table 3. Before implementation of TP, L_{Aeq} (dBA) values ranged between 60.4 to 71.7 dBA in Room A and between 58.1 to 59.9 dBA in Room B. Inside the incubator, L_{Aeq} was 48.7 dBA. After implementation of TP, L_{Aeq} values in Room A areas ranged between 58.8 to 59.5 dBA and in Room B, ranged between 60.3 to 63.9 dBA. The L_{Aeq} values inside the incubator increased to 54.5 dBA. The results demonstrated no significant differences between L_{Aeq} values before and after TP implementation, although some relevant work practices concerning the impact of noise were adjusted.

The highest $L_{p, Cpeak}$ (dBC) value was found in the “Work Station” area of Room A before TP implementation (143.3 dBC). Data showed that significant differences were found between $L_{p, Cpeak}$ (dBC) values before and after TP. Table 4 shows the frequencies spectrum in octave bands among the areas under study. The 500 Hz was the frequency which had higher levels in the majority of the areas before and after the implementation of TP (Room A (“Traffic Zone” – 56.7 dBA and 56.5 dBA); Room B (“Work Station” – 56.7 dBA and 58.7 dBA; “Traffic Zone” – 59.1 dBA and 53.7 dBA)).

DISCUSSION

During TP sessions, noise was identified by professionals as a disturbing agent and exerted a negative impact (Table 2), is in agreement with data presented by Gurses and Carayon (2009) and by Santos et al. (2014) who demonstrated that health professionals

perceived “equipment’s” as one of the most annoying noise sources followed by "team conversation" and “visits”.

The “Work Station” of Room A had a decrease on L_{Aeq} and $L_{p, Cpeak}$ values, 71.7 to 58.8 dBA and 143.3 to 102.8 dBC, respectively (Table 3). It was apparent by professionals, that greater care was being undertaken in carrying out their tasks regarding noise production. However, in the “Traffic Zone” of Room B, the noise level increase almost 6 dB after the TP, probably attributed to the presence of visitors and other staff (from ancillary departments that did not participated in the TP) and might be the source of this rise. The L_{Aeq} values obtained in the “Work Station” and “Traffic Zone” before and after the implementation of TP exceed the recommended values given by WHO for day and night periods, indicating more attention needs to be taken. Regarding the values inside the incubator, despite the elevation of L_{Aeq} values after the TP (48.7 dBA to 54.4 dBA), the levels obtained exceeded the recommended threshold. In general, the results obtained (Table 3) may be attributed to the number of newborns that were in NICU before (14 without specific care needs) and after TP (10, which two were helped by an oximeter and ventilator, that may produce 60-78 and 60-80 dBA respectively (Pugh and Griffiths 2007)). Similar data were reported by Tsunemi et al. (2012).

Data analysis revealed that low frequencies tend to have more influence on noise produced in the NICU than higher frequencies (Table 4). These results are in agreement with Gray and Philbin (2000), who stated that noise in nurseries is dominated by low frequencies, with some exceptions due to loud mid-frequencies alarms. Kellam and Bhatia (2008) suggested that human speech contribute to the spike in sound energy at 500 Hz.

Despite adherence by healthcare staff to TP, there was no significant reduction of noise after TP implementation. These results suggest that it is necessary to consider several factors that

may ensure the effectiveness of TP. It seems that TP impact was lost over time, but diminution of L_p, C_{peak} values may indicate that health professionals undertake their activities more carefully. In fact, the effectiveness of training healthcare professionals presents some contradictory findings in literature (Oliveira et al. 2013). Several studies showed that the implementation of TP in this area contributed to reduction of sound average levels, although by itself does not yield the recommended threshold levels and its impact tended to decrease over time (Philbin and Klaas 2000). It is suggested that a TP incorporated into a more comprehensive quiet protocol, involving low cost environmental (or others) modifications, seems to ensure decrease of noise levels (Ramesh et al. 2009) affecting positively patient well-being and improvement of satisfaction levels (Connor and Ortiz 2009). Other studies also reported some resistance from health professionals to noise reduction programs, being a factor that needs to be considered in future programs (Taylor-Ford et al. 2008). Connor and Ortiz (2009), noted in their study that staff also believed that noise affected the physiologic, psychologic, and overall health of patients. In this study, the impact of staff education was measured by patient-satisfaction scores surveys. In surveys after staff education, improvement was expressed in fewer poor ratings and an increase in good to very good ratings (Connor and Ortiz 2009).

CONCLUSIONS

Noise is a common problem in NICU, and exerts significant adverse implications for health and well-being of patients and staff. Results showed that after 6 months of TP implementation, there was no significant noise reduction in the NICU and inside the incubator. However, L_p, C_{peak} data are an indication that in conjunction with other factors, a TP may be an excellent action to reduce noise levels. While recognizing the importance of TP in order to promote changes in the team's attitudes, it needs to be recognized that the

effects are not long lasting. The training sessions needs to be repeated more often, and physicians, nursing staff, supervisors, senior leadership, staff from related departments and family members/visitors need to attend. Noise in all the rooms of a NICU might be reduced considerably by incorporating affordable behavioral and environmental modifications, and by renovation and/or preventive maintenance of equipment. Training the staff in order to implement quiet work behaviors is essential and needs to be seen as a first step to implement a quiet time protocol in neonatal intensive units.

ACKNOWLEDGMENTS

This work is part of the “NeoNoise” project, which is being conducted and supported by the Research Centre on Health and Environment (CISA) of the School of Allied Health Technologies of Polytechnic Institute of Porto (ESTSP.IPP). The authors would like to thank the assistance given by the Occupational Health Service of the Hospital Centre.

REFERENCES

- AAP (American Academy of Pediatrics: Committee on Environmental Health). 1997. Noise: A hazard for the fetus and newborn. *Pediatrics* 100 : 724–727
- Berglund, B., L. Thomas, and H. S. Dietrich. 1999. *Guidelines for Community Noise. Guidelines for Community Noise.* Geneva, Switzerland.
<http://www.who.int/docstore/peh/noise/guidelines2.html>.
- Bremmer, P., J. F. Byers, and E. Kiehl. 2003. Noise and the premature infant: Physiological effects and practice implications. *J Obstet Gynecol Neonatal Nurs* 32: 447–454. .
- Brown, G. 2009. NICU noise and the preterm infant. *J Neonatal Nurs* 28 : 165–173.
- Connor, A., and E. Ortiz. 2009. Staff solutions for noise reduction in the workplace. *Permanente J* 13: 23–27.
- Fechter, LD. 2004. Promotion of noise-induced hearing loss by chemical contaminants. *J Toxicol Environ Health A* 67: 727–740.
- Gray, L., and M. K. Philbin. 2000. Measuring sound in hospital nurseries. *J Perinatol* 20: S100–S104.
- Gurses, A. P., and P. Carayon. 2009. Exploring performance obstacles of intensive care nurses. *Appl Ergonom* 40: 509–518.
- Guthrie, O.W., H. Xu, B.A. Wong, S.M., McIturf, J.E. Reboulet, P.A. Ortiz and D.R. Mattie. 2014. Exposure to low levels of jet-propulsion fuels impairs brainstem encoding of stimulus intensity. *J Toxicol Environ Health A* 77: 261-280.
- Kellam, B., and J. Bhatia. 2008. Sound spectral analysis in the intensive care nursery: Measuring high-frequency sound. *J Pediat Nurs* 23: 317–323.

- Kent, W. T., A. W. Tan, M. C. Clarke, and T. Bardell. 2002. Excessive noise levels in the neonatal ICU: Potential effects on auditory system development. *J Otolaryngol* 31: 355–360.
- Machado, M. C., M. I. Alves, and M. L. Couceiro. 2011. “Saúde infantil e juvenil em Portugal: Indicadores do plano Nacional de Saúde. *Acta Pediát Portuguesa* 42 : 195–204.
- Mahmood, A., H. Chaudhury, and M. Valente. 2011. Nurses’ perceptions of how physical environment affects medication errors in acute care settings. *Appl Nurs Res* 24: 229–237.
- Nicolau, A. S., D. Casal, P. M. Lopes, and P. Kronenberg. 2005. O Ruído Nas Unidades Neonatais de Lisboa E Vale Do Tejo. *Acta Pediát Portuguesa* 36: 15–21.
- Oliveira, D., M. Figueiredo, and V. Batista. 2013. Ubiquidade do ruído em neonatologia: Efeitos e efectividade de medidas de controlo. *Acta Pediát Portuguesa* 44 : 234–241.
- Philbin, M. K., and L. Gray. 2002. Changing levels of quiet in an intensive care nursery. *J Perinatol* : 22: 455–460. .
- Philbin, M. K., and P. Klaas. 2000. Evaluating studies of the behavioral effects of sound on newborns. *J Perinatol* 20: S61–S67.
- Pugh, R. J., and R. Griffiths. 2007. Noise in critical care. *Care Critically Ill* 23: 105–109.
- Ramesh, A., P. N. S. Rao, G. Sandeep, M. Nagapoornima, V. Srilakshmi, M. Dominic, and Swarnarekha. 2009. Efficacy of a low cost protocol in reducing noise levels in the neonatal intensive care unit. *Indian J Pediat* 76: 475–478
- Robertson, A., J. Kohn, P. Vos, and C. Cooper-Peel. 1998. Establishing a noise measurement protocol for neonatal intensive care units. *J Perinatol* 18: 126–130.

- Sampaio Neto, R. A., F. O. S. Mesquita, M. D. S. Paiva Junior, F. F. Ramos, F. M. D. Andrade, and M. A. V. Correia Junior. 2010. Ruídos Na Unidade de Terapia Intensiva: Quantificação E Percepção Dos Profissionais de Saúde. *Revista Brasil Terap Intensiv* 22 :369–374.
- Santos, J., C. Carvalhais, M. Nóia, J. S. Baptista, and M. V. da Silva. 2014. “NeoNoise Project: Preliminary results of sound pressure levels in a neonatal intensive care unit.” In *SHO2014: Proceedings Book of the International Symposium on Occupational Safety and Hygiene*, edited by P. Arezes, Baptista, J.S., Barroso, M.P., Carneiro, P., Cordeiro, P., Costa, N., Melo, R., Miguel, A.S. and G. Perestrelo, 189–93. Guimarães, Portugal: Portuguese Society of Occupational Safety & Hygiene.
- Short, A. E., K. T. Short, A. Holdgate, N. Ahern, and J. Morris. 2011. Noise levels in an Australian emergency department. *Aust Emerg Nurs J* 14: 26–31.
- Surenthiran, S. S., K. Wilbraham, J. May, T. Chant, A. J. B. Emmerson, and V. E. Newton. 2003. Noise levels within the ear and post-nasal space in neonates in intensive care. *Arch Dis Child. Fetal Neonatal Edition* 88: F315–F318. .
- Taylor-Ford, R., A. Catlin, M. LaPlante, and C. Weinke. 2008. Effect of a noise reduction program on a medical-surgical unit. *Clin Nurs Res* 17: 74–88.
- Tsunemi, M. H., T. Y. Kakehashi, and E. M. Pinheiro. 2012. Noise at the neonatal intensive care unit after the implementation of an educational program. *Texto Contexto - Enfermagem* 21 : 775–782.
- Wachman, E. M., and A. Lahav. 2011. The effects of noise on preterm infants in the NICU. *Arch Dis Child. Fetal Neonatal Edition* 96: F305–F309.

Table 1 – Equipment and behavioral causes of noise in intensive care units. Adapted from Pugh and Griffiths (2007)

Source of noise	
Items falling onto the floor	Up to 92 dB(A)
Equipment movement (e.g. bed)	90 dB(A)
Connection of gas supply	88 dB(A)
Door closure	85 dB(A)
Pager	84 dB(A)
Talking	75 - 85 dB(A)
Ventilator alarm	70 - 85 dB(A)
Nebulizer	80 dB(A)
Telephone	70 - 80 dB(A)
Television	79 dB(A)
Oximeter	60 - 80 dB(A)
Monitor Alarm	79 dB(A)
Ventilator	60 - 78 dB(A)
IV infusion alarm	65 - 77 dB(A)
Endotracheal aspiration unit	50 - 75 dB(A)

Table 2 – Training content and feedback from healthcare staff.

Training content	Healthcare staff feedback	
	<i>Main factors that contribute to the noise level on unit</i>	<i>Suggestions to help decrease the noise level</i>
1. General concepts of noise; 2. The impact of noise exposure in a hospital setting; 3. National legislation and international standards regarding noise exposure in hospitals; 4. Main health effects of noise exposure of neonates and health professionals; 5. Presentation and discussion of results for the study developed in the NICU; 6. Recommendations to reduce noise in neonatal units.	1. Traffic in the room; 2. Visitors behavior; 3. Equipment / alarms; 4. Team conversation; 5. Some healthcare procedures.	1. Limit number of visitors; 2. Keep voices down; 3. Improve the technology regarding the implementation of a centralized control of all the alarms of NICU in workstation area; 4. More quiet at change of shift; 5. Substitute metallic materials (drug transport cars for eg.) by others made of washable plastic; 6. Perform some tasks more carefully to avoid noise production.

Table 3 - Values of mean L_{Aeq} (dBA) and $L_{p, Cpeak}$ (dBC), before and after implementation of the training program.

Room	Area	L_{Aeq} (dBA) – Before Mean \pm SD (Min-Max)	$L_{p, Cpeak}$ (dBC) – Before	L_{Aeq} (dBA) - After Mean \pm SD (Min-Max)	$L_{p, Cpeak}$ (dBC) - After
A	Work Station	71.7 \pm 4.74 (47.8-114.6)	143.3	58.8 \pm 2.72 (47.6-76.4)	102.8
	Traffic Zone	60.4 \pm 5.32 (43.6-91.5)	115.8	59.5 \pm 1.95 (53.0-75.0)	101.5
	Inside Incubator	48.7 \pm 2.94 (42.2-68.1)	104.1	54.5 \pm 0.49 (53.3-65.7)	92.8
B	Work Station	59.9 \pm 6.01 (39.5-85.8)	106.3	60.3 \pm 3.09 (46.4-79.2)	99.7
	Traffic Zone	58.1 \pm 3.07 (43.8-82.0)	113.2	63.9 \pm 4.37 (47.0-82.2)	98.9

SD – Standard Deviation; $p = 0.917$ L_{Aeq} (dBA); $p = 0.043$ $L_{p, Cpeak}$ (dBC).

Table 4 – Spectral analysis of noise generated in rooms A and B, before and after the implementation of the TP.

	Room	Area	Frequencies (Hz)							
			63	125	250	500	1000	2000	4000	8000
dB										
Before TP	A	Work Station	78.3	75.3	71.6	68.8	65.9	63.1	59.8	56.2
		Traffic Zone	52.9	50.7	56.0	56.7	52.6	52.9	50.7	48.6
		Inside Incubator	61.8	61.8	58.2	47.2	49.9	46.2	39.1	31.3
	B	Work Station	54.5	53.1	55.1	56.7	54.6	53.2	52.6	46.0
		Traffic Zone	53.8	49.4	57.6	59.1	59.1	57.8	54.8	48.4
		Work Station	52.7	55.7	54.8	55.5	53.3	51.4	50.6	45.1
After TP	A	Traffic Zone	48.3	50.8	53.2	56.5	53.1	55.8	51.0	44.9
		Inside Incubator	55.1	53.9	48.0	46.8	44.5	37.7	36.9	27.2
		Work Station	53.9	49.6	54.2	58.7	54.6	52.4	48.3	40.8
	B	Traffic Zone	51.2	47.3	51.4	53.7	50.9	50.5	52.2	47.6