



THE AGA KHAN UNIVERSITY

eCommons@AKU

Department of Medicine

Department of Medicine

September 2009

Low prevalence of the intact cag pathogenicity island in clinical isolates of *Helicobacter pylori* in Karachi, Pakistan

Javed Yakoob

Aga Khan University, javed.yakoob@aku.edu

Wasim Jafri

Aga Khan University, wasim.jafri@aku.edu

Zaigham Abbas

Aga Khan University, zaigham.abbas@aku.edu

Shahab Abid

Aga Khan University, shahab.abid@aku.edu

R. Khan

Aga Khan University

See next page for additional authors

Follow this and additional works at: https://ecommons.aku.edu/pakistan_fhs_mc_med_med



Part of the [Digestive System Diseases Commons](#), and the [Gastroenterology Commons](#)

Recommended Citation

Yakoob, J., Jafri, W., Abbas, Z., Abid, S., Khan, R., Jafri, N., Ahmad, Z. (2009). Low prevalence of the intact cag pathogenicity island in clinical isolates of *Helicobacter pylori* in Karachi, Pakistan. *British Journal of Biomedical Science*, 66(3), 137-142.

Available at: https://ecommons.aku.edu/pakistan_fhs_mc_med_med/155

Authors

Javed Yakoob, Wasim Jafri, Zaigham Abbas, Shahab Abid, R. Khan, N. Jafri, and Z. Ahmad



Low prevalence of the intact *cag* pathogenicity island in clinical isolates of *Helicobacter pylori* in Karachi, Pakistan

J. Yakoob, W. Jafri, Z. Abbas, S. Abid, R. Khan, N. Jafri & Z. Ahmad

To cite this article: J. Yakoob, W. Jafri, Z. Abbas, S. Abid, R. Khan, N. Jafri & Z. Ahmad (2009) Low prevalence of the intact *cag* pathogenicity island in clinical isolates of *Helicobacter pylori* in Karachi, Pakistan, British Journal of Biomedical Science, 66:3, 137-142, DOI: 10.1080/09674845.2009.11730260

To link to this article: <https://doi.org/10.1080/09674845.2009.11730260>



Published online: 23 May 2016.



Submit your article to this journal [↗](#)



Article views: 3



Citing articles: 5 View citing articles [↗](#)

Low prevalence of the intact *cag* pathogenicity island in clinical isolates of *Helicobacter pylori* in Karachi, Pakistan

J. YAKOOB, W. JAFRI, Z. ABBAS, S. ABID, R. KHAN, N. JAFRI and Z. AHMAD*

Departments of Medicine and Pathology, The Aga Khan University, Karachi, Pakistan

Accepted: 29 May 2009

Introduction

Helicobacter pylori is a Gram-negative microaerophilic bacterium that infects human gastric mucosa and is recognised as a major cause of chronic active gastritis, peptic ulcer disease, gastric adenocarcinoma and gastric mucosa-associated lymphoid tissue lymphoma.^{1,2} It also has a relationships with gastro-oesophageal reflux disease and adenocarcinoma of the gastric cardia.³ It is a class I carcinogen that colonises the human stomach of more than 50% of the world's population. However, only a small subset of infected people experience *H. pylori*-associated illnesses.

Although the pathogenesis of *H. pylori* infection is not well understood, there are several virulence factors that may contribute to mucosal damage. The *cagA* gene is conventionally used as a marker of pathogenic strains. However, the literature suggests that the *cagA* gene alone is not a suitable marker of *H. pylori* virulence. A strong predictor of severe disease outcome is infection with a bacterial strain harbouring the *cag* pathogenicity island (*cag*-PAI), a 40 kb segment of DNA.⁴

H. pylori *cag*-PAI is horizontally acquired and follows free recombinations within the species. The chromosomal integrity of this island, or the lack thereof, is thought to play an important role in the progress of the gastroduodenal pathology caused by *H. pylori*. These genes encode a type IV secretion system that forms a syringe-like structure to translocate the immunodominant *cagA* protein into the gastric epithelial cells.⁵ *cag*-PAI has also been implicated in the induction of interleukin-8 (IL-8) in cultured gastric cells.⁶

At least 17 out of 27 genes were found to be essential for translocation of *cagA* into host cells (syringe-like function), and 14 were required to fully induce transcription of IL-8.⁶ The *cag*-PAI and vacuolating toxin A (*vacA*) gene are located far apart on the chromosome, but there is a statistical linkage between presence of the *cag*-PAI and the s1 genotype of

ABSTRACT

Clinical diseases that follow *Helicobacter pylori* infection are associated with expression of the *cagA* gene, a part of cytotoxin-associated gene pathogenicity island (*cag*-PAI). This study aims to determine whether or not the presence of *cagA* is associated with the presence of complete *cag*-PAI and to evaluate inflammatory changes associated with the five loci in the *cag*-PAI of *H. pylori* comprising *cagA*, *cagA* promoter region (*cagAP*), *cagE*, *cagT* and the left end of the *cagA* gene (LEC). *H. pylori* isolates were obtained from patients with dyspeptic symptoms. Clinical strains of *H. pylori* were screened by the polymerase chain reaction (PCR) for respective genes of the *cag*-PAI. Of 115 *H. pylori* isolates, 31 (28%) were positive for the five *cag*-PAI loci. *H. pylori* isolates with intact *cag*-PAI were associated with gastric carcinoma (GC; $n=9$ [60%]) and gastric ulcer (GU; $n=5$ [45%]) compared to non-ulcer dyspepsia (NUD; $n=14$ [18%]) ($P=0.001$ and $P=0.049$, respectively). In patients with intact *cag*-PAI, acute on chronic inflammation was present in 25 (81%) and was more common than chronic inflammation ($P=0.013$). The *cagE* and *cagAP* had deletions in 25 (37%) and 23 (35%) cases, respectively. The *cagAP* region was significantly associated with GC ($n=12$ [80%], $P<0.001$) and GU ($n=9$ [82%], $P=0.001$) compared to NUD ($n=24$ [30%]) and with significant acute on chronic inflammation ($n=40$ [80%], $P=0.007$). The distribution of *vacAs1a* with intact *cag*-PAI in GC was 9(60%) and in NUD was 10(13%) ($P<0.001$). The presence of the *cagA* gene does not signify presence of an intact *cag*-PAI. Most of the *H. pylori* isolates studied had partial *cag*-PAI with missing *cagE* and *cagA* promoter regions.

KEY WORDS: Duodenal ulcer.
Genes.
Genomic islands.
Helicobacter pylori.
Non-ulcer dyspepsia.

vacA.⁷ Strains with this genotype are called type I strains and are generally more virulent than type II strains that lack the *cag*-PAI and harbour the less virulent *vacA* s2 genotype.⁸

The aim of this study is to determine the association of *cagA* with intact *cag*-PAI to evaluate inflammatory changes associated with intact *cag*-PAI and determine any association between *cag*-PAI and the *vacA* allele s1. The completeness of *cag*-PAI is determined using specific flanking polymerase chain reaction (PCR) primers from five loci in the *H. pylori* *cag*-PAI comprising *cagA*, *cagA* promoter region (*cagAP*),

Correspondence to: Dr Javed Yakoob

Department of Medicine, The Aga Khan University
Stadium Road, PO Box 3500, Karachi 74800, Pakistan.
Email: yakooobjaved@hotmail.com

cagE, *cagT* and the left end of the *cag*-PAI (LEC) in clinical isolates of *H. pylori*. The absence of *cag*-PAI is detected by the PCR for the *cag* empty site flanking the left and right ends of the *cag*-PAI indicating that they truly lacked the *cagA* gene.⁹

Materials and methods

Patients and *H. pylori* strains

A total of 115 *H. pylori* isolates were obtained from patients undergoing gastroscopy and gastric biopsy for dyspeptic symptoms attending the endoscopy suite of the gastroenterology department at the Aga Khan University from July 2005 to December 2007 (Table 1). Mean age was 40±14 years (range: 18–78 years). Informed consent was obtained for oesophagogastroduodenoscopy (EGD) and biopsy from the antrum and body of the stomach. The study was approved by the institution's ethics review committee.

Sterilised biopsy forceps were used to obtain gastric biopsy specimens from an area of inflammation for rapid urease test, histopathology, *cag*-PAI-associated virulence genes, *cagA* empty site and *vacA* PCR.

After primary isolation and identification of the gastric organisms as *H. pylori*, the strains were frozen at -70°C until their use in the study. Subsequently, bacteria were cultured on Columbia blood agar (Oxoid) plates containing 7% sheep blood for 3–5 days at 37°C under microaerobic conditions using Campygen strips in anaerobic jars (Oxoid).

Several small, rounded colonies were picked from different parts of the culture plate and subcultured twice to obtain a pure culture. Gram's stain, urease and catalase tests were used to confirm identification and to exclude the possibility of contamination.

H. pylori cells were harvested from plates by suspension in 2-mL sterile phosphate-buffered saline (PBS) and were pelleted by centrifugation at 3000 *xg* for 20 min. The clinical symptoms at the time of presentation, diagnosis, drug treatment dosage and duration were noted, together with endoscopy findings.

Rapid urease test

Pronto Dry, a commercial rapid urease test (Medical Instrument Corp., France), results were read in 30 min after sampling, in accordance with the manufacturer's instructions. Colour change from yellow to pink was indicative of a positive reaction.¹⁰

Histopathology

Gastric antral biopsy specimens stained with haematoxylin and eosin (H&E) were examined by a pathologist unaware of the patient's clinical diagnosis. In doubtful cases, Giemsa staining was performed to ascertain the presence of *H. pylori*. Degree of gastritis on H&E stain was scored in accordance with the Sydney system¹¹ for the level of inflammation and neutrophil infiltration (0=none, 1=mild, 2=moderate, 3=severe).

Extraction of genomic DNA

DNA was extracted from confluent cultures with fewer than three *in vitro* passages.¹² The bacterial cells from a chocolate agar plate were washed (x2) with PBS and centrifuged at 3000 *xg* for 20 min. *H. pylori* DNA was extracted by a modified phenol/chloroform method.¹³ Briefly, a bacterial

Table 1. Demographic and clinical findings of patients (n=115).

Gender	Male: Female	77:38
Age	Range	18–78 years
	Mean age±SD	40±14 years
Symptoms	Abdominal pain	92 (82%)
	Vomiting	13 (10%)
	Weight loss	10 (8%)
Diagnosis	Non-ulcer dyspepsia	79 (69%)
	Gastric ulcer	11 (9%)
	Gastric carcinoma	15 (13%)
	Duodenal ulcer	10 (9%)
Histopathology	Acute inflammation	10 (9%)
	Chronic inflammation	19 (16%)
	Acute on chronic inflammation	86 (75%)

pellet was resuspended in Tris-HCl buffer containing ethylenediaminetetraacetate (TE, pH 8.0) and lysozyme (30 mg/mL) and was incubated at 37°C for 30 min. The suspension was treated with sodium dodecyl sulphate (SDS, 10%), proteinase K (2.5 mg/mL) and RNase A (40 mg/mL). DNA was extracted with phenol/chloroform/isoamyl alcohol, precipitated by sodium acetate and ice-cold absolute alcohol, and washed with ice-cold alcohol (70%). The DNA pellet was finally resuspended in TE buffer. DNA content and purity were determined by measuring absorbance at 260 nm and 280 nm using a spectrophotometer (Beckman DU-600, USA).

cag-PAI PCR analysis

The PCR analyses were carried out with oligonucleotide pairs (Table 2A), as described previously,¹⁴ to amplify five different loci spread over the *cag* I and *cag* II regions. *cagA*, the *cagA* promoter region and *cagE* represented *cagI*, while the left end of *cag* (LEC) and *cagT* were selected to represent *cagII*.

Pairs of oligonucleotide primers were used to detect the presence of the *cag*-PAI genes (*cagA*, the *cagAP* region, *cagE*, *cagT* and the LEC, containing both inside and outside genes of *cag*-PAI). These primer pairs were designed on the basis of published sequences.^{4,9,15} To test for the absence of the *cag*-PAI, PCR for the *cag* empty-site was used with primers that flanked the left and right ends of the *cag*-PAI.⁹

H. pylori strains ATCC 43526 and ATCC 43504, which have been determined to have the entire *cag*-PAI, were used as positive controls, and ATCC 51932 was used as a negative control as it lacks *cagA*. Amplification specificity for the different loci was assessed by testing *H. pylori* strains ATCC 43526, ATCC 43504 and ATCC 51932 with each experiment. Amplification by PCR was performed in 25- μ L volumes containing 10 mmol/L Tris-HCl (pH 8.3), 50 mmol/L KCl, 1.5–2.5 mmol/L MgCl₂, 200 μ mol/L deoxynucleoside triphosphates, 2 units *Thermus aquaticus* (*Taq*) DNA polymerase (Promega) and 25 μ mol both forward and reverse primers (synthesised using an MWG automatic synthesiser). The total volume was made up with double-distilled water (DDW).

Amplification was performed in a Perkin Elmer 9700 thermal cycler (Applied Biosystems). The cycles comprised

Table 2A. Sequences and locations of oligonucleotide primers.

Primer	Primer sequence	Amplicon (bp)	Location ^a
cagA-F1	5'-AACAGGACAAGTAGCTAGCC-3'	349	2700–2719'
cagA-R1	5'-TATTAATGCGTGTGTGGCTG-3'		3400–3381'
cagA-F2	5'-GATAACAGGCAAGCTTTTGA-3'	701	157–176'
cagA-R2	5'-CTGCAAAGATTGTTGGCAGA-3'		505–484'
cagE-F1	5'-GCGATTGTTATTGTGCTGTAG-3'	329	16,891–16,870'
cagE-R1	5'-GAAGTGGTTAAAAATCAATGCCCC-3'		16,563–16,587'
cagT-F1	5'-CCATGTTTATACGCCTGTGT-3'	301	442–461'
cagT-R1	5'-CATCACCACACCCTTTTGTAT-3'		723–742'
LEC-F1	5'-ACATTTTGGCTAAATAAACGCTG-3'	384	3920–3942 [‡]
LEC-R1	5'-TCTCCATGTTGCCATTATGCT-3'		4303–4283 [‡]
LEC-F2	5'-ATAGCGTTTTGTGCATAGAA-3'	877	3856–3875 [‡]
LEC-R2	5'-ATCTTTAGTCTCTTTAGCTT-3'		4732–4713 [‡]
cagAP-F1	5'-GTGGGTAAAAATGTGAATCG-3'	730	18,738–18,757'
cagAP-R1	5'-CTGCAAAGATTGTTGGCAGA-3'		505–484'
cagAP-F2	5'-CTACTTGTCCTCAACATTTT-3'	1181	18,495–18,514'
cagAP-R2	5'-CTGCAAAGATTGTTGGCAGA-3'		505–484'
cagA empty site [‡]		360	
2	5'-ACATTTTGGCTAAATAAACGCTG-3'		
25	5'-TCATGCGAGCGCGATGTG-3'		

^aGenBank Accession No: [‡]AF001357; [†]U60176; [‡]AC000108; [‡]regions flanking the left and right ends of the cag-PAI insert location for NCTC 12455.

an initial denaturation of target DNA at 94°C for 5 min and then denaturation at 94°C for 1 min, primer annealing at 55°C for 1 min and extension at 72°C for 1 min. The final cycle included an extension step for 7 min at 72°C to ensure full extension of the product. Samples were amplified through 35 consecutive cycles.

The amplified products were resolved on 2% agarose gel in 1x TAE buffer (40 mmol/L Tris-acetate, 1 mmol/L EDTA [pH 8.0]) containing ethidium bromide (0.5 µg/mL) and then visualised under ultraviolet (UV) light.

vacA PCR analysis

Amplification of *vacA* alleles by PCR was performed in a 25 µL volume containing 10 mmol/L Tris-HCl (pH 8.3), 50 mmol/L KCl, 1.5–2.5 mmol/L MgCl₂, 200 µmol/L deoxynucleoside

triphosphates, 2 units *Taq* DNA polymerase (Promega) and 25 µmol forward and reverse primers (synthesised by an MWG automatic synthesiser) (Table 2B).¹⁶

The amplification cycle for *vacA* alleles consisted of an initial denaturation of target DNA at 95°C for 5 min and then denaturation at 95°C for 1 min, primer annealing at 52°C for 1 min and an extension at 72°C for 1 min. The final cycle included an extension step for 7 min at 72°C to ensure full extension of the product. Samples were amplified through 35 consecutive cycles.

Positive and negative reagent controls were performed with each batch of amplifications. DNA from *H. pylori* strains ATCC 43504 (*vacAs1am1*, *cagA*-positive), ATCC 51932 (*vacAs2m2*, *cagA*-negative) and ATCC 43526 (*vacAs1bm1*, *cagA*-positive) was used to define the accuracy of the *vacA* PCR.

Table 2B. Oligonucleotide primers used in typing of *H. pylori vacA* alleles.

Region amplified	Primer	Primer sequence	Size and location of PCR product
m1	VA3-F	5'-GGTCAAATGCGGTCATGG-3'	290 bp (2741–3030)
	VA3-R	5'-CCATTGGTACTGTAGAAAC-3'	
m2	VA4-F	5'-GGAGCCCCAGGAAACATTG-3'	352 bp (976–1327)
	VA4-R	5'-CATACTAGCGCCTTGAC-3'	
S1a	SS1-F	5'-GTCAGCATCACACCGCAAC-3'	190 bp (866–1055)
	VA1-R	5'-CTGCTTGAATGCGCCAAAC-3'	
S1b	SS3-F	5'-AGCGCCATACCGCAAGAG-3'	187 bp (869–1055)
	VA1-R	5'-CTGCTTGAATGCGCCAAAC-3'	

a. Location in published *cagA* sequence (ref. 14).
b. Location in published *vacA* sequence (ref. 16).

Statistical assessment

The Statistical Package for Social Science SPSS (Release 11.5, standard version, SPSS; 1989–99) was used for data analysis. Descriptive analysis was performed for demographic and clinical features. Results were presented as mean±SD for quantitative variables and number (%) for qualitative variables. Differences in proportion were assessed using Pearson's χ^2 test, Fisher's exact or likelihood ratio test, where appropriate. $P < 0.05$ was considered significant.

Results

The clinical features and diagnosis of patients included in the study are given in Table 1. Partial deletions involved *cag* I and *cag* II regions of the *cag*-PAI equally. In isolates from NUD, deletions of *cag*-PAI genes were more frequent (Table 3A). These deletions predominantly involved *cagE* and *cagA* promoter region of the *cag*-PAI (Table 3B). The product corresponding to the *cagA* empty site was obtained from five patients with partially intact *cagA* PAI, indicating that they had mixed infections (i.e., a mixture of *H. pylori* strains with and without the *cagA* gene).

Presence of *cag*-PAI genes

The *cagA* gene was positive in 66 (57%), *cagA* LEC in 63 (55%), *cagAT* in 63 (55%), *cagAP* region in 50 (43%) and *cagAE*




Fig. 1. Distribution of *cag*-PAI gene amplified products in patients with non-ulcer dyspepsia. Lane M: molecular weight marker 50 bp (Promega); Lane A and B: *cagA* gene present; Lane C: *cagA* promoter absent; Lane D: LEC present; Lane E: *cagE* absent; Lane F: *cagT* present.




Fig. 2. Distribution of *cag*-PAI gene amplified products in patients with duodenal ulcer. Lane M: molecular weight marker 50 bp (Promega); Lane A and B: *cagA* gene present; Lane C: *cagA* promoter absent; Lane D: LEC present; Lane E: *cagE* present; Lane F: *cagT* present.




Fig. 3. Distribution of *cag*-PAI gene amplified products in patients with gastric carcinoma. Lane M: molecular weight marker 50 bp (Promega); Lane A: *cagA* gene; Lane B: *cagA* promoter; Lane C: LEC; Lane D: negative control; Lane E: *cagE*; Lane F: *cagAT*.

Table 3A. Correlation of *cag*PAI with diagnosis and histopathology.

	<i>cag</i> PAI		
	Intact <i>n</i> (%)	Partial <i>n</i> (%)	Deleted <i>n</i> (%)
	31(28)	42(36)	42(36)
<i>Diagnosis</i>			
Non-ulcer dyspepsia	14 (18)	28(35)	37(47)
Gastric ulcer	5(45)	6(55)	0
Gastric carcinoma	9 (60)	5(33)	1(7)
Duodenal ulcer	3 (30)	3(30)	4(40)
<i>Histopathology</i>			
<i>Inflammation</i>			
Acute inflammation	5(50)	3(30)	2(20)
Chronic inflammation	1(5)	12(63)	6(32)
Acute on chronic inflammation	25(29)	27(31)	34(39)
<i>Lymphoid aggregates</i>			
Positive	26(31)	26(34)	30(35)
Negative	5(17)	13(43)	12(40)
<i>Neutrophil infiltration</i>			
Mild	1(7)	7(50)	6(43)
Moderate	25(30)	27(33)	30(37)
Severe	5(26)	8(42)	6(32)

in 48 (42%) *H. pylori* isolates. In 31 (28%) isolates *cag*-PAI was intact, it had partial deletions with some genes missing in 42 (36%), and was completely deleted in 42 (36%). Deletions were present in *cagE* ($n=23$ [37%], $P < 0.001$) and in *cagAP* (23 [35%], $P < 0.001$) (Figs. 1 and 2).

Associations of *cag*-PAI genes with gastroduodenal disorders

Association of *H. pylori* isolates with intact *cag*-PAI was seen in GC ($n=9$ [60%]), GU ($n=5$ [45%]) and NUD ($n=14$ [18%]) (Table 3A, Fig. 3). The *cagA* gene was associated with GU ($n=9$ [82%]) and GC ($n=14$ [93%]), compared to NUD ($n=38$ [48%]) ($P=0.049$ and $P=0.001$, respectively). The *cagAP* region was significantly associated with GC ($n=14$ [93%], $P < 0.001$), GU ($n=9$ [82%], $P=0.001$) and DU ($n=7$ [70%], $P=0.016$) compared to NUD ($n=24$ [30%]). *cagA* LEC was associated with GU ($n=11$ [100%]) and GC ($n=13$ [87%]) compared to NUD ($n=40$ [51%]) ($P < 0.001$ and $P=0.010$, respectively). *cagA* LEC was also associated with GU ($n=11$ [100%]) compared to DU ($n=5$ [50%]) ($P=0.002$) and GC ($n=13$ [87%]) compared to DU ($n=5$ [50%]) ($P=0.045$). *cagE* was associated with GU ($n=7$ [64%], $P=0.043$) and GC ($n=13$ [87%], $P < 0.001$) compared to NUD ($n=25$ [32%]). *cagT* was associated with GU ($n=10$ [91%], $P=0.006$) and GC ($n=14$ [93%], $P=0.002$) compared to NUD ($n=40$ [51%]).

Associations of *cag*-PAI genes with histology

In patients infected with *H. pylori* with intact *cag*-PAI, acute on chronic inflammation was present in 25 (81%) ($P=0.013$, Table 3A). In these patients, acute on chronic inflammation was more common than chronic inflammation ($P=0.014$), and *cagAP* ($n=40$ [47%], $P=0.007$) was associated with acute on chronic inflammation. Intestinal metaplasia was present in four (57%) isolates with intact *cag*-PAI ($P=0.082$).

Table 3B. Distribution of *cag*-PAI genes in the clinical isolates.

		Intact n(%) 31(28)	Partial n(%) 42(36)	Deleted n(%) 42(36)
<i>cagA</i>	Positive	31(46)	35(52)	0(0)
	Negative	0(0)	7(14)	40(86)
<i>cagAP</i>	Positive	31(62)	19(38)	0(0)
	Negative	0(0)	23(35)	42(65)
<i>cagLEC</i>	Positive	31(48)	32(49)	0(0)
	Negative	0(0)	10(19)	42(81)
<i>cagE</i>	Positive	31(65)	17(35)	0(0)
	Negative	0(0)	23(37)	42(63)
<i>cagT</i>	Positive	31(49)	32(51)	0(0)
	Negative	0(0)	10(19)	42(81)
<i>cagA</i> empty site	Positive	0(0)	5(11)	41(89)
	Negative	31(45)	37(54)	1(1)

cagAP: *cagA* promoter region; *LEC*: left end of *cag*-PAI.

Associations of *cag*-PAI genes with *vacAs1a* and *m1*

The presence of *vacAs1a* was seen in 80(70%) isolates and that of *vacAm1* was seen in 64(56%). Distribution of *vacAs1a* with intact *cag*-PAI was GC ($n=11$ [73%]), NUD ($n=10$ [13%], $P<0.001$), GU ($n=4$ [36%], $P=0.05$) and DU ($n=3$ [30%] $P=0.03$). It was associated with acute inflammation in three (30%) cases and acute on chronic inflammation in 20 (23%) ($P=0.164$). Distribution of *vacAm1* with intact *cag*-PAI was GC ($n=8$ [53%]), DU ($n=3$ [30%]) and GU ($n=3$ [27%]) compared to NUD ($n=5$ [6%]) ($P<0.001$, $P=0.038$, $P=0.050$,

respectively). Acute on chronic inflammation was more frequent ($n=16$ [19%]) than chronic inflammation ($P=0.008$).

Discussion

The epidemiology of *H. pylori* infection remains important in public health because of its high prevalence and its association with gastric cancer and lymphoma. The seroprevalence of *H. pylori* in the adult population of Pakistan is reported as 58%.¹⁸ A recent study revealed early colonisation/infection of infants with *H. pylori* and a prevalence of 67% at nine months of age in a peri-urban community in Karachi, Pakistan.¹⁹

In the present study, *H. pylori* isolates from patients with NUD constituted the majority, which reflects the high numbers in this group attending the gastroenterology clinic and having endoscopy. *H. pylori* isolates with intact *cag*-PAI were associated with GC, GU and DU and on histopathology with acute and acute on chronic inflammation. *cagA* gene, *cagAP* region, *cagLEC*, *cagE* and *cagT* were all associated with GC, GU and DU compared to NUD.

H. pylori isolates from NUD lacked the *cagAP* region and *cagE* (Fig. 1) while they were intact in those from GC (Fig. 2). There was a significant difference in inflammatory infiltration of the gastric mucosa between patients with intact *cag*-PAI and those with partial *cag*-PAI type strains (Table 3A). These findings suggest that the strains with partial *cag*-PAI have less ability to cause inflammation than those with intact *cag*-PAI.

The implications of the present study are that *cag*-PAI is partially intact in the majority of our clinical isolates. This is consistent with a previous study in which *cag*-PAI appeared

Table 4. Relationship between intact *cag*-PAI and *vacA* subtypes in *H. pylori* strains.

	<i>cag</i> -PAI s1a			<i>cag</i> -PAI m1		
	Positive n(%)	Negative n(%)	<i>P</i> value	Positive n(%)	Negative n(%)	<i>P</i> value
<i>Diagnosis</i>						
Non-ulcer dyspepsia	10(13)	69(87)	0.002	5(6)	74(94)	<0.001
Gastric ulcer	3(27)	8(73)		3(27)	8(73)	
Duodenal ulcer	2(20)	8(80)		3(30)	7(70)	
Gastric carcinoma	9(60)	6(40)		8(53)	7(47)	
<i>Histopathology</i>						
<i>Inflammation</i>						
Acute	3(30)	7(70)	0.164	3(30)	7(70)	0.016
Chronic	1(5)	18(95)		0(0)	19(100)	
Acute on chronic	20(23)	66(77)		16(19)	70(81)	
<i>Lymphoid aggregates</i>						
Positive	20(23)	65(77)	0.237	3(10)	27(90)	0.243
Negative	4(13)	26(87)		16(19)	69(81)	
<i>Neutrophil infiltration</i>						
Mild	1(7)	13(93)	0.302	0(0)	14(100)	0.011
Moderate	18(22)	64(78)		18(22)	64(78)	
Severe	5(26)	14(74)		1(5)	18(95)	

cag-PAI: *cag* pathogenicity island.

to be partial in the majority of clinical isolates around the world.¹⁹ Only 12% of the Indian isolates carried an intact cag-PAI. Deletion frequencies of *cagA*, *cagE*, and *cagT* genes were higher in benign cases than in isolates from severe ulcers and gastric cancer.¹⁹

In contrast, another study from Calcutta, India, showed that 80–90% of Calcutta strains carried cag-PAI and the potentially toxigenic *vacAs1* alleles of *vacA*, independently of disease status.²⁰ In the present study, the clinical outcome of *H. pylori* infection was predicted by analysing genes of the cag-PAI, with the *cagAP* region and *cagE* more commonly associated with GC and GU compared to NUD, which is consistent with a previous study.²¹

Jenks *et al.* described the presence of *cagE* and correlated it with that of *cagA*.²² However, in the present study it was associated with the *cagA* promoter region. Also, the majority of partial deletions of cag-PAI were observed in *cagE*, followed by the *cagAP* region.

The *cagA* gene promoter region appeared to be a better marker of *H. pylori* virulence; however, it is known for the diversity of its sequences in isolates. The *cagE* gene, which is consistently retained near the *cagAP* region, appeared to be a better substitute for the *cagAP* region. Also, *cagE* gene PCR can be used to screen the cag-PAI structure, as suggested previously.²²

An association was noted between *vacAs1a* and *vacAm1*, with intact cag-PAI. In the present study, *vacAs1* with intact cag-PAI was associated with GC, GU and DU. This is in keeping with a previous study that showed *vacAs1* occurs significantly more often in isolates from patients with peptic ulceration or gastric adenocarcinoma, while *vacAs2* invariably occurs in patients with gastritis alone.²³

A shortcoming of the present study is the small number of patients in the different groups. The study, however, is ongoing but the results presented here indicate that the presence of the *cagA* gene does not signify the presence of an intact cag-PAI, as most of the *H. pylori* isolates had partial cag-PAI with missing *cagE* and *cagAP* regions. □

*This work was supported by an Aga Khan University Research Grant SM 100 327*11 431 to JY. The authors are grateful to staff at the Juma Research Building, The Aga Khan University, for their assistance during this work.*

References

- Blaser MJ. Ecology of *Helicobacter pylori* in the human stomach. *J Clin Invest* 1997; **100**: 759–62.
- Blaser MJ, Perez-Perez GI, Kleanthous H *et al.* Infection with *Helicobacter pylori* strains possessing *cagA* is associated with an increased risk of developing adenocarcinoma of the stomach. *Cancer Res* 1995; **55**: 2111–5.
- Cover TL, Blaser MJ. *Helicobacter pylori* factors associated with disease. *Gastroenterology* 1999; **117**: 257–61.
- Censini S, Lange C, Xiang Z *et al.* Cag pathogenicity island of *Helicobacter pylori* encodes type-specific and disease-associated virulence factors. *Proc Natl Acad Sci USA* 1996; **93**: 14648–53.
- Karnholz A, Hoefler C, Odenbreit S, Fischer W, Hofreuter D, Haas R. Functional and topological characterization of novel components of the comB DNA transformation competence system in *Helicobacter pylori*. *J Bacteriol* 2006; **188**: 882–93.
- Fischer W, Jürgen PL, Buhrdorf R, Gebert B, Odenbreit S, Haas R. Systematic mutagenesis of the *Helicobacter pylori* cag pathogenicity island: essential genes for cagA translocation in host cells and induction of interleukin-8. *Mol Microbiol* 2001; **42**: 1337–48.
- van Doorn LJ, Figueiredo C, Sanna R, Blaser MJ, Quint WG. Distinct variants of *Helicobacter pylori* cagA are associated with *vacA* subtypes. *J Clin Microbiol* 1999; **37**: 2306–11.
- Xiang Z, Censini S, Bayeli PF *et al.* Analysis of expression of CagA and VacA virulence factors in 43 strains of *Helicobacter pylori* reveal that clinical isolates can be divided into two major types and that cagA is not necessary for expression of the vacuolating cytotoxin. *Infect Immun* 1995; **63**: 94–8.
- Akopyants NS, Clifton SW, Kersulyte D *et al.* Analyses of the cag pathogenicity island of *Helicobacter pylori*. *Mol Microbiol* 1998; **28**: 37–53.
- Said RM, Cheah PL, Chin SC, Goh KL. Evaluation of a new biopsy urease test: Pronto Dry for the diagnosis of *Helicobacter pylori* infection. *Eur J Gastroenterol Hepatol* 2004; **16**: 195–9.
- Price AB. The Sydney System: histological division. *J Gastroenterol Hepatol* 1991; **6**: 209–22.
- Matteo MJ, Granados G, Pérez CV, Olmos M, Sanchez C, Catalano M. *Helicobacter pylori* cag pathogenicity island genotype diversity within the gastric niche of a single host. *J Med Microbiol* 2007; **56**: 664–9.
- Yakoob J, Hu GL, Fan XG *et al.* Diversity of *Helicobacter pylori* in Chinese persons with *H. pylori* infection. *APMIS* 2000; **108**: 482–6.
- Ikenoue T, Maeda S, Gura KO *et al.* Determination of *Helicobacter pylori* virulence by simple gene analysis of the cag pathogenicity island. *Clin Diagn Lab Immunol* 2001; **8**: 181–6.
- Tomb JE, White O, Kerlavage AR *et al.* The complete genome sequence of the gastric pathogen *Helicobacter pylori*. *Nature* 1997; **338**: 539–47.
- Covacci A, Rappuoli R. PCR amplification of gene sequences from *Helicobacter pylori* strains. In: *Helicobacter pylori: techniques for clinical diagnosis and basic research*. Philadelphia: WB Saunders, 1996: 94–109.
- Abbas Z, Jafri W, Khan AH, Shah MA. Prevalence of *Helicobacter pylori* antibodies in endoscopy personnel and non-medical volunteers of Karachi. *J Pak Med Assoc* 1998; **48**: 201–3.
- Nizami SQ, Bhutta ZA, Weaver L, Preston T. *Helicobacter pylori* colonization in infants in a peri-urban community in Karachi, Pakistan. *J Pediatr Gastroenterol Nutr* 2005; **4**: 191–4.
- Kausar F, Khan AA, Hussain MA *et al.* The cag pathogenicity island of *Helicobacter pylori* is disrupted in the majority of patient isolates from different human populations. *J Clin Microbiol* 2004; **42**: 5302–8.
- Mukhopadhyay AK, Kersulyte D, Jeong JY *et al.* Distinctiveness of genotypes of *Helicobacter pylori* in Calcutta, India. *J Bacteriol* 2000; **182**: 3219–27.
- Blaser MJ. Role of *vacA* and the *cagA* locus of *Helicobacter pylori* in human disease. *Aliment Pharmacol Ther* 1996; **10** (Suppl 1): 73–8.
- Jenks PJ, Megraud F, Labigne A. Clinical outcome after infection with *Helicobacter pylori* does not appear to be reliably predicted by the presence of any of the genes of the cag pathogenicity island. *Gut* 1998; **43**: 752–8.
- Kidd M, Lastovica A, Atherton J, Louw JA. Heterogeneity in the *Helicobacter pylori* genes *vacA* and *cagA*: associated with gastroduodenal disease in South Africa? *Gut* 1999; **45**: 499–503.