



THE AGA KHAN UNIVERSITY

eCommons@AKU

Department of Pathology and Laboratory Medicine

Medical College, Pakistan

July 1998

Well differentiated hepatocellular carcinoma metastasizing to jaw and oral cavity within six months of primary diagnosis.

Nazma Afroze
Aga Khan University

Irshad N. Soomro
Aga Khan University

Shahid Pervez
Aga Khan University, shahid.pervez@aku.edu

Naila Kayani
Aga Khan University

Sheema H. Hasan
Aga Khan University

Follow this and additional works at: http://ecommons.aku.edu/pakistan_fhs_mc_pathol_microbiol

 Part of the [Oncology Commons](#), [Oral Biology and Oral Pathology Commons](#), [Orthopedics Commons](#), and the [Pathology Commons](#)

Recommended Citation

Afroze, N., Soomro, I. N., Pervez, S., Kayani, N., Hasan, S. H. (1998). Well differentiated hepatocellular carcinoma metastasizing to jaw and oral cavity within six months of primary diagnosis. *Journal of Pakistan Medical Association*, 48(7), 219-220.
Available at: http://ecommons.aku.edu/pakistan_fhs_mc_pathol_microbiol/302

Well Differentiated Hepatocellular Carcinoma Metastasizing to Jaw and Oral Cavity within Six Months of Primary Diagnosis

Pages with reference to book, From 219 To 220

Nazma Afroze, Irshad N. Soomro, Shahid Pervez, Naila Kayani, Sheema H. Hasan (Department of Pathology, The Aga Khan University Hospital, Karachi.)

Introduction

Metastasis of hepatocellular carcinoma to the soft tissue and bone of the jaw and to oral cavity is a rare phenomenon. One such case is described in this report. Fine needle aspiration cytology is a useful procedure for detection of hepatocellular carcinoma at this site. However, at times cytologic interpretation of swelling in this region may give rise to diagnostic dilemma and clinical history may be essential in establishing the diagnosis.

Case Report

A sixty-seven year old male, was referred by an E.N.T. surgeon in May, 1997 for FNA of left parotid region swelling. Clinical history obtained revealed that the lump appeared three months back and was hard and non tender on palpation.




Figure 1. Photomicrograph of smear from swelling of parotid region showing metastatic hepatocellular carcinoma (Mag. X 100)

The clinical impression was parotid adenoma with a suspicion of malignancy. The smear showed

clusters of large cells with abundant eosinophilic granular cytoplasm and central round vesicular nuclei. Considering the swelling arising from parotid gland itself, a possibility of oncocytoma was raised. Surgical removal and histological evaluation was recommended.




Figure 2. Photomicrograph of histology of swelling around parotid region showing metastatic well differentiated hepatocellular carcinoma (Mag X 200).

The patient underwent surgical removal of the jaw swelling in June, 1997. Frozen section showed a large cell carcinoma with features suggesting metastatic hepatocellular carcinoma. The clinicians were informed and past medical history revealed that in October, 1996 the patient presented with liver mass and got an FNAB done on that mass by another surgeon. The clinical impression at that time was cavernous haemangioma. Cytology slides and aspirated tissue fragments were sent to our department. A differential diagnosis of well differentiated hepatocellular carcinoma and hepatic adenoma was given favouring the former. After that, the patient was lost to follow-up. Four months later he developed a parotid swelling and reported to E.N.T. Department in May, 1997. After frozen section diagnosis of the parotid mass, a radical parotidectomy was performed with removal of left superficial and deep parotid gland and ramus of left mandible. The permanent sections revealed the typical histology of a well differentiated hepatocellular carcinoma which had metastasized into the soft tissue around the parotid gland. The adjacent bony tissue of ramus of mandible and surrounding skeletal muscle was invaded by the tumour. In December, 1997 the patient again developed a mass in oral cavity with a single cervical lymph node enlargement. The oral mass proved to be a metastatic hepatocellular carcinoma on histology while the lymph node was benign.

Discussion

Metastatic tumours involving head and neck region are unusual. Hepatocellular carcinoma metastasis to jaw and soft tissue of the oral cavity is a rare occurrence¹. A review of literature revealed occasional case reports. Hepatocellular carcinoma metastasizes by direct extension, lymphatic route or hematogenously. Distant metastasis occur in 50% of cases. Bony metastasis are relatively infrequent and are involved in 11% of cases with the spine being the most important site of localization². One study³ reported seven cases of metastatic tumours in head and neck region over a period of eighteen years. Out of these three had arisen from primary hepatocellular carcinoma with metastasis to paranasal sinuses. In another study⁴ fine needle aspiration findings of fifteen hepatocellular carcinomas were described, four in musculoskeletal, four in adrenal, four in regional lymph node, two in pancreas and one in pelvic region. Eleven of these cases had a biopsy proven hepatocellular carcinoma. In the remaining patients, the FNA diagnosis of hepatocellular carcinoma in metastatic sites was the initial diagnosis. In our case the metastatic tumour had involved the left jaw including the mandibular ramus, skeletal muscle and soft tissue around parotid gland. Involvement of the major salivary gland by malignant neoplasm whose origin is outside the salivary gland occurs by direct extension, hematogenous route or lymphatic route. The intimate relationship of the paraparotid and intraparotid lymph nodes with the parotid parenchyma frequently results in lymphatic migration of extrasalivary neoplasm to parotid region. Direct extension of non salivary gland tumours into the parotid regions occurs primarily from squamous cell carcinoma and basal cell carcinoma of overlying skin. Eighty percent of metastasis to this site have been from primary tumour elsewhere in the head-neck region and 20% have been from the infraclavicular site⁵. Hematogenous pathway is the route of metastasis from infraclavicular areas. In descending order of frequency the most common infraclavicular sites of primary tumour that metastasized to major salivary glands are lung, kidney and breast. Other infrequent sites include colon, rectum, prostate, ovary, pancreas, stomach, uterus and urinary bladder⁵. For most patients the primary tumour is known at the time of diagnosis of their salivary gland metastasis. In this case the FNA diagnosis of parotid swelling was suggestive of oncocytoma. Oncocytoma is a neoplasm made up of large polygonal cells with granular eosinophilic cytoplasm and comparatively small round nuclei. These cells can have cytologic resemblance to well differentiated hepatocytes. The FNA cytology slides were reviewed. On re-examination the cell clusters showed striking similarity to well differentiated hepatocytes but the characteristic FNA features of hepatocellular carcinoma like presence of naked nuclei, endothelial cells derived from capillary vessels surrounding clusters of hepatocytes and intracytoplasmic bile pigment were not seen in this case. Initial diagnostic dilemma was due to lack of communication to the cytopathologist about the previous diagnosis of malignancy. The most important and helpful factor in resolving this dilemma was previous history. The importance of clinical history and radiographic and biochemical findings in the diagnosis of metastatic hepatocellular carcinoma cannot be overstated⁴. Patient may occasionally develop more than one primary malignant neoplasm, however in this case histologic similarity between the parotid tumour and a previously diagnosed neoplasm was a strong evidence in favour of metastasis. Familiarity with the FNA cytology of hepatocellular carcinoma and proper medical history should allow its diagnosis in rare metastatic sites. Degree of differentiation in a hepatocellular carcinoma varies and may predict more or less aggressive behaviour. However, metastasis to head and neck area of this very well differentiated hepatocellular carcinoma within six months of initial diagnosis is another question mark about the reliability of tumour grading in predicting its histological behaviour.

References

1. Marker P, Clausen PP Metastasis to the mouth and jaws from hepatocellular carcinoma *Acascrepfl Int. J. Oral. Maxillofac, Surg*, 1991;20:371-4
2. Welhk K, Uayduk K, Pitzer P Fine needle puncture cytology in metastatic hepatocellular carcinoma.

Z. Gastroenterol., 1989;27:225-7.

3. Mochimatsu J, Tsukuda M, Furukawa S et al. Tumours metastasizing to the head and neck: A report of seven cases. *J. Laryngol. Otol.*, 1993;107:1171-3.

4. Dusenbery D, Carr BI. Fine needle aspiration diagnosis of hepatocellular carcinoma in metastatic sites. *Acta Cytol.*, 1996;40:443-9

5. Gary L, Ellis DDS, Paul L, Auselair. Turnouts of salivary gland. 1996. Third series, Fascicle-17 Washington DC. Armed Forces Institute of Pathology. 1996;pp:403-410.