



THE AGA KHAN UNIVERSITY

eCommons@AKU

Section of Urology

Department of Surgery

September 1991

Sertoli cell only syndrome (SECOS): lessons from case studies

Jamsheer J Talati

Aga Khan University, jamsheer.talati@aku.edu

Hizbullah Sheikh

Aga Khan University

Follow this and additional works at: http://ecommons.aku.edu/pakistan_fhs_mc_surg_urol



Part of the [Urology Commons](#)

Recommended Citation

Talati, J., Sheikh, H. (1991). Sertoli cell only syndrome (SECOS): lessons from case studies. *Journal of Pakistan Medical Association*, 41(9), 219-223.

Available at: http://ecommons.aku.edu/pakistan_fhs_mc_surg_urol/74

SERTOLI CELL ONLY SYNDROME (SECOS): LESSONS FROM CASE STUDIES

Pages with reference to book, From 219 To 223

Jamsheer Talati (Departments of Surgery (Urology), The Aga Khan University, Faculty of Health Sciences, Karachi.)
Hizbullah Sheikh (Departments of Pathology, The Aga Khan University, Faculty of Health Sciences, Karachi.)

ABSTRACT

Between June 85 and December 87, 69 testicular biopsies were submitted for histopathological examination during investigation of infertility; ten (14%) patients had a Sertoli cell only syndrome. The history, clinical features, and hormonal profiles were analyzed in an attempt to categorize these patients on aetiological basis. Two followed treatment of malignancy - one by radiation for testicular cancer and one by cyclophosphamide for a lymphoma. One had unilateral cryptorchidism. Mumps was etiological factor in one patient. FSH levels determined in 6 patients were elevated in all suggesting a possible dependence of (sick) Sertoli cells on spermatogenic cells for production of inhibin. Alternative explanations include changes in sertoli cell enzymes or FSH receptors. Testosterone levels are in the low normal range suggesting that Leydig cells may also be affected by the etiological factor producing the syndrome. Two patients who had earlier received a higher Johansen score were found to have a sertoli cell only syndrome on expert review of testicular biopsies. It is suggested that the condition is more common than hitherto reported and is often confused with maturation arrest. Testicular histopathology should be done by specialists in testicular pathology (JPMA 41: 219, 1991).

INTRODUCTION

The Sertoli cell only syndrome (SECOS) is a condition in which there is an absence of germ cells and the seminiferous tubules contain only Sertoli cells. Reproduction is therefore impossible and the patient is sterile. Thirty percent of our referred infertile population has azoospermia as compared to 5% in the west¹. Accurate determination of the cause of azoospermia is necessary as treatment is helpful in the obstructive azoospermia but impossible in the Sertoli cell only syndrome.

A retrospective review of all testicular biopsies was undertaken by a clinician and histopathologists with a view to categorizing testicular pathology accurately. The records of the subset of 10 patients with SECOS were reviewed retrospectively.

MATERIALS AND METHODS

All (69) testicular biopsies done during investigation of infertility on patients attending the Aga Khan University Hospital infertility clinic between June 85 and December 87 were reviewed by two histopathologists and a urologist with special interest in andrology. Ten patients were detected as having SECOS. (Figure).




Figure . Histopathological appearance of Sertoli cell only syndrome.

The History, Physical Findings, Hormonal status and semen analysis of these 10 patients were reviewed in order to detect etiological factors and clinicopathological features of this syndrome. Testicular measurements were done by transparent ruler and hormone analysis by radioimmunoassay.

Results

TABLE I. Duration of infertile marriage before discovery of sertoli cell only syndrome.

Duration of marriage		Husband's age	Wife's age
1	3	26	27
2	3	33	29
3	4	30	22
4	3	27	22
5	7	36	28
6	2	31	21
7	NR	27	25
8	4	28	20
9	6	28	NR
10	5	35	25
Mean	4.1	30.1	24.3
Range	2-7	26-36	20-29

Table 1 gives the patient's and spouse age and duration of infertility.

TABLE II. Sertoli Cell only Syndrome: etiological factors.

	Mumps	Irradiation	Chemotherapy	Smoking	Insecticide	Cryptorchidism	Estrogen	Occupation
1	0	0	0	0	0	0	0	BSNS
2	0	0	0	0	0	0	0	SERV
3	0	+	0	+	0	+	0	SERV
4	0	0	0	+	0	0	0	BSNS
5	+	0	0	0	0	0	0	NAVY
6	0	0	0	0	0	0	0	SERV
7	+	0	0	0	0	0	0	BSNS
8	0	0	0	0	0	0	0	ENG
9	0	+	+	0	0	0	0	BSNS
10	0	0	0	+	+	0	0	Farmer

TABLE III. Sertoli Cell only Syndrome: Past history of illness, medications & operations.

1	Typhoid 5 years previously, Jaundice 10 years age, Tagamate & antibodies
2	Hyperlipidemia, hypertension (age 33 years) (Inderal)
3	Mumps at age 12
4	Fracture tibia, Peptic ulcer, (Tagamate for 60 days) Appendicitis
5	Neonatal pneumonia, typhoid
6	Tonsillectomy
7	Gonococcal urethritis, Mumps
8	0
9	0
10	0

Tables II and III mention associated conditions which may contribute as possible etiological factors. All testes (Table IV)

TABLE IV. Sertoli Cell only Syndrome Testicular size & Testosterone levels.

	Test. Size		Testosterone ng/dl	Mumps	Patients age history
	R	L			
1	3.5	(Orchidectomy)	600	0	26
2	3.5	3.5	526	0	33
3	'V. small'	4.0	457	0	30
4	3.5	3.5	456	0	27
5	"Small"	"Small"	ND	+	36
6	3.5	4.0	452	0	31
7	5	5	ND	+	27
8	4	3	ND	0	28
9	4	4	390	0	28
10	"Small"	"Small"	ND	0	35

except six were 3.5 cm or larger in their long axis as measured by transparent plastic ruler. Five of the 12 testes were 4 cm or larger. Of the two patients who had mumps, one had normal sized testes indicating that mumps had probably not affected the testes. All the patients were young and hence any reduction in testicular size was not related to age.

TABLE V. Sertoli Cell only Syndrome: Hormonal profile.

Hormonal Normal Values	FSH upto 20m IU/ml	LH upto 25m IU/ml	PRL upto 20ng/ml	Testosterone 300-1200 ng/dL	Remarks
1	21.7	ND	ND	600	
2	24.7	9.7	7.8	526	
3	42.8	29.7	8.8	457	Smoker
4	ND	26.7	ND	456	Smoker, Irradiation, Cryptorchidism
5	41.4	17.8	ND	ND	Mumps
6	36.4	19.4	15.4	52.7	
7	ND	ND	ND	ND	Mumps
8	ND	ND	ND	ND	
9	57.3	13.0	33.4	390	Irradiation, Cyclophosphamide for lymphoma
10	ND	ND	ND	ND	Smoker, Exposure to insecticides

ND - Not determined

Hormonal Status

FSH was determined in 6 patients (Table V) and ranged from 21.7 - 57.32 mIU/ml (Maximum normal 20 mIU/ml. LH levels (determined in 6 patients) were normal in four and marginally raised in two patients. Testosterone levels (estimated in 6) were normal in all patients but remained in the lower 1/2 of normal range. Prolactin level was elevated in one of the 4 patients in whom it was estimated.

DISCUSSION

Spermatogenic tubules of all vertebrates have 2 types of cells - the spermatogenic cell and the supportive Sertoli cell. For various reasons spermatogenic cells may in the embryo fail to develop or migrate, or, later under varying Chemotherapy Smoking Insecticide Cryptorchidism Estrogen Occupation circumstances be destroyed resulting in the isolation of Sertoli cells to producing a picture of Sertoli Cell only Syndrome (SECOS).

SECOS was found in 29% of testicular biopsies done for Suspected obstructive azoospermia². It is difficult to find figures about incidence of SECOS in Pakistani literature. In our series 14% had SECOS. Usually testicular biopsy is not done in patients with obvious non-obstructive azoospermia in whom a diagnosis of testicular failure is made on the basis of hormonal evaluation. Sertoli Cell Only Syndrome is seen in patients with intraabdominal cryptorchidism (detected in late life), with viral infections, in patients who have received radiotherapy, chemotherapy or oestrogens and in an idiopathic variety³.

Idiopathic (congenital) forms of SECOS may result from failure of migration of the primordial germ cells into gonadal anlage or to early destruction of the germinal epithelium by noxious substances. The etiology however remains obscure in most patients. One of our patients developed SECOS following radiotherapy to the abdomen for a seminoma; another after chemotherapy with cyclophosphamide. Two patients had a history of mumps but testicular atrophy was present in only one patient. Pakistan's population has been unable to gain access to modern medical care until recently. We have noted a large number of men with disparity in testicular size, one testes much smaller than normal. One possibility is that they have had unreported intermittent testicular torsion in childhood. SECOS could also perhaps be explained on the basis of bilateral recurrent torsion in which death of the sensitive germ cells has occurred but the more resistant Leydig cells and Sertoli cells have just managed to survive. Additionally roadside quacks prescribe medications which often contain heavy metals which may produce germ cell damage. Three of our patients were heavy smokers, but the effect of smoking on the testes is variable⁴⁻⁷ and it is not known to produce testicular atrophy. One of our patients had an intra-abdominal testis and seminoma, the opposite testis being normal. None of our patients had estrogens. One patient (a farmer) had repeatedly sprayed insecticides on his farms.

Testicular size

Despite the absence of germ cells, the testes have retained a good size in many patients. The clinical presentation is variable, at times suggestive of obstructive azoospermia with well developed testes measuring around 4.5 to 5 cm in the longitudinal axis, but many manifest as intermediate sized testes (3.5 cms) and a few have small testes. Clinical features do not help detect this syndrome which is diagnosed by biopsy alone.

Histological patterns

Histological patterns were initially misdiagnosed as maturation arrest in two patients. Diagnosis of SECOS must be water tight as patients need to be told that they are incapable of reproduction. Whilst the psychological effect of informing the patient that he cannot produce Off spring is devastating it is inappropriate to shy away from the issue and treat such patients in vain hope that they will improve on therapy.

Histopathological appearances vary according to the cause. In the idiopathic form also called the del Costello Syndrome⁸ there is great uniformity of appearance of pyramidal cells with a resulting adenoma like pattern. The cells contain a small number of inclusion bodies and elongated nuclei. FSH levels are consistently raised in this form³.

In the form seen after chemotherapy or radiotherapy, the nuclei are of normal configuration, the cytoplasm contains phagocytosed material and large amounts of glycogen. In cryptorchids the nuclei of the sertoli cells are rounded and contain nucleoli. In some patients the cytoplasm is filled with characteristic pleomorphic inclusions.

Hormonal levels

As Sertoli cells produce inhibin, one would expect normal inhibin production (in this syndrome) and hence normal FSH levels. This is not so. Our findings of elevated FSH levels are in keeping with others^{3,9,10} as also the normal LH levels and normal to low Testosterone levels¹¹. This had been noted even in the idiopathic variety which has been presumed to result from failure of migration of germ cells from the yolk sac.

The high levels of FSH could be explained by the following hypotheses:

1. germ cell cooperation is necessary for inhibin production.
2. Sertoli cell damage co-exists with germ cell damage.
3. Leydig cells are unable to provide enough testosterone for optimal sertoli function.

1. Germ cell cooperation

Steinberger and Steinberger¹² have shown that Sertoli Cells in culture can secrete a factor which selectively inhibits FSH. Galdieri et al¹³ have shown that A.B.P. secretion by the Sertoli cell is influenced by contact with germ cells, and Ritzen et al¹⁴ do not rule out the possibility that germ cell cooperation may enhance inhibin production, or that cells other than Sertoli are capable of producing inhibin. It is possible that potent inhibin is produced by cells other than sertoli cells.

2. Sertoli Cell Damage

This may manifest as a decreased production or altered direction of secretion of inhibin; or may result in loss of FSH receptors.

a. Decreased production of Inhibin:

Whilst Sertoli Cells may escape total annihilation, there is reason to suspect subtle damage at enzyme levels in these cells. It is known that post-irradiation damage to the sertoli cell may reduce locally produced GnRH like factor³. It is possible that though the cells look morphologically normal, the enzymes for production of inhibin are destroyed in much the same way as has been demonstrated for GnRH. De Kretser et al¹⁰ have shown decreased inhibin levels in patients with gonadal damage, inhibin levels are low even though FSH rises and they propose that in many men with testicular damage the inverse relationship between inhibin and FSH is not evident.

b. Luminal vs. interstitial secretion of Inhibin:

It has recently been shown that the Sertoli Cell products can be secreted bidirectionally both into the seminiferous tubular fluid (STF) and into testicular interstitial fluid (TIF) and hence into blood¹⁵.

Following hypophysectomy cryptorchidism, exposure to ethane dimethane sulphonate¹⁶ or busulfation¹⁷, the major part of androgen binding protein is secreted into testicular interstitial fluid and this has been related to absence of the pachytene spermatocytes and elongated spermatids from the tubule¹⁵. Selective enzyme damage may alter capacity of the Sertoli cell to secrete substances preferentially either into the lumen or blood stream. It has been shown that failure of glycosylation results in loss of the ability to secrete some of Sertoli cell proteins into the blood stream¹⁸. Inhibin secretion has also been shown to be bi-directional¹⁹. In the Sertoli Cell Only Syndrome, it is possible that intraluminal secretion of inhibin into STF occurs instead of passage of inhibin into the blood stream via

TIF from the basal surface of the cell. This may be because of some yet undiscovered enzyme defect. Lack of secretion into the blood stream may allow FSH levels to rise unchecked.

c. Decreased FSH receptors:

FSH and testosterone are the two main stimuli for the Sertoli cell function¹⁴. Namiki et al²⁰ have shown that FSH binding sites were not affected by testicular organ culture at 37°C (as compared to 33°C) and they felt that short term temperature changes may not affect FSH binding sites. Presumably one would not expect a reduction in FSH receptors. Hagenas et al²¹ have however shown a dramatic decrease in testicular receptors for FSH in cryptorchid rats. As Sertoli cell function is dependent on FSH, understandably the inhibin levels would fall in the absence of these receptors.

d. Age related changes in Sertoli cell:

A drop in the inhibin level would be expected with age. Our oldest patient was 36 years old and hence this is unlikely to be a cause.

3. The Leydig Cell

Whilst many authors have noted normal LH and T levels in patients with SECOS²², we feel that the Leydig cells may also be affected to some extent in SECOS as suggested by the fact that testosterone levels are in the lower half of normal range, in fact below 600 ng/dl. Leiper et al²³ have shown decreased Leydig cell function after testicular irradiation for acute lymphoblastic leukemia. There was little increase of plasma testosterone on HCG stimulation in 10 patients, 7 of whom were pubertal. This reduction remained evident 4 years after testicular irradiation and the testicular volume remained lower than normal. Chemotherapy, however, was not as toxic. Forty six boys previously treated with chemotherapy had normal testosterone response to HCG stimulation² suggesting normal Leydig cell function.

It is possible that disordered Leydig cell function is responsible for the low normal testosterone levels in our patients. Sertoli cells do have a biological threshold for testosterone. In combined cell culture of pen-tubular and Sertoli cells 10 nmol/L concentrations of testosterone or dihydrotestosterone are the minimal levels needed for Sertoli cell function²³. It is possible that in vivo, larger peaks are needed for adequate function. The role of testosterone in controlling the Sertoli cell is still ill understood²⁴ and could involve the peritubular myoid cells²³ which may also be susceptible to radiation damage.

REFERENCES

1. Jouannet, P., Ducot, B., Fencux, D. and Spira, A. Male factors and likelihood of pregnancy in infertile couples. I. Study of sperm characteristics. *Int. J. Androl.*, 1988; 11:379.
2. Akerman, L. and Rosai, J. The testis, in surgical pathology. 5th ed. St. Louis, Mosby, 1974.
3. Schulze, C. Sertoli cells and Leydig-cells in man, in advances in anatomy, embryology and cell biology. No. 88 Berlin, Springer Verlag, 1984.
4. Klaiber, E.L., Broverman, D.M., Pokoly, T.B., Albert, A.I., Howard, P.J.Jr. and Sherer, J.F. Interrelationships of cigarette smoking, testicular varicoceles, and seminal fluid indexes. *Fertil. Steril.*, 1987; 47:481.
5. Dikshit, R.K., Buch, J.G. and Mansuri, S.M. Effect of tobacco consumption on semen quality of a population of hypofertile males. *Fertil. Steril.*, 1987; 48:334.
6. Purl, V.N. Smoking and seminal fluid index (letter). *Fertil. Steril.*, 1987; 48:702.
7. Lahdetie, J. and Husgafvel-Pursiainen, K. Cigarette smoking a possible genotoxic hazard to male germ cells? *Int. J. Androl.*, 1987; 10:431.
8. del Castillo, E.B., Trabucco, A. and de la Blaz, F.A. Syndrome produced by absence of germinal epithelium without impairment of Sertoli and Leydig-cells. *Clin. Endocrinol. Metabol.*, 1947; 7:493.
9. de Kretser, D.M., Kerr, J.B. and Paulsen, C.A. Evaluation of the ultrastructural changes in the human

- Sertoli cells in testicular disorders and the relationship of the changes to the levels of serum FSH. *Int. J. Androl.*, 1981; 4:129.
10. de Kretser, D.M., Robertson, D.M., Burger, H.G., Risbridger, G.P., Hodgson, Y., McLochlan, R.M. and Healy, D. Changes in inhibin levels in reproductive states. 3. *Endocrinol.* 1988; 117 supplement Abstract, 7th Joint meeting of British Endocrine Societies 18-21 April.
 11. Ishida, H., Isurugi, K., Aso, Y., Takayasu, H. and Tamaoki, B. Endocrine studies in Sertoli-cell-only syndrome. *Urol.*, 1976; 116:56.
 12. Steinberger, A. and Steinberger, E. Secretion of FSH-inhibiting factor by cultured Sertoli cell. *Endocrinology*, 1976; 99:918.
 13. Galdieri, M., Monaco, L and Stefanini, M. Secretion of androgen binding protein by Sertoli cell is influenced by contact with germ cells. *Androl.*, 1984; 5:409.
 14. Ritzen, E.M., Hanson, V. and French, F.S. The Sertoli cell, in the testis. Edited by H. Burger and D.M. de Kretser. New York, Raven press 1981, p. 180.
 15. Sharpe, R.M. Bidirectional secretion by the Sertoli cell. *Int. J. Androl.*, 1988; 11:87.
 16. Sharpe, R.M. and Bartlett, M. Change in secretion of ABP into testicular interstitial fluid with age and in situations of impaired spermatogenesis. *Androl.*, 1987; 10:701.
 17. Morris, I.D., Barden, C.W., Musto, N.A., Thau, R.B. and Gunsalus, G.L. Evidence suggesting that sperm cells influence the bidirectional secretion of androgen binding protein by the seminiferous epithelium demonstrated by selective impairment of spermatogenesis with busuiphan. *Int. J. Androl.*, 1987; 10:691.
 18. Cheng, C.Y., Gunsulas, G.L., Morris, I.D., Turner, T.T and Barden, C.W. The heterogeneity of rat androgen binding protein in the vascular compartment differs from that in the testicular tubular lumen. Further evidence for bidirectional secretion of androgen binding protein. *J. Androl.*, 1986; 7:175.
 19. Baker, H.W., Burger, H., de Kretser, D.M., Eddie, L.W., Higginson, R.E., Hudson, B. and Lee, V.W. Studies on purification of inhibin from bovine testicular secretions using an *in vitro* bioassay. *Int. J. Androl. Suppl.*, 1978; 2:115.
 20. Namiki, M., Nakamura, N.I., Okuyama, A., Sonoda, T., Nishimune, Y., Tskatsuka, D. and Matsumoto, K. Influence of temperature on function of Sertoli and Leydig-cell of human testes. *Fertil. Steril.*, 1987; 47:475.
 21. Hagenas, L. and Ritzen, E.M. Impaired Sertoli cells function in experimental cryptorchidism in the rat. *Mol. Cell Endocrinol.*, 1976; 4:25.
 22. Hagenas, U., Ritzen, E.M., Svensson, J., Haneson, V. and Purvis, K. Temperature dependence of Sertoli cell function. *Int. J. Androl. (Suppl.)*, 1978; 2:449.
 23. Leiper, A.D., Grant, D.B. and Cessels, J.M. Gonadal function after testicular radiation for acute lymphoblastic leukemia. *Arch. Dis. Child.*, 1986; 61:53.
 24. Wheeler, I.E. and Rudy, F.R. The testes and testicular structures and male external genitalia, in principles and practice of surgical pathology. Edited by Steven Silverberg. New York, Wiley, 1983, vol 2, p. 1159.
 25. Shalet, S.M., Hann, M., Tendon, M., Mona Jones, P.H. and Beardwell, C. G. Testicular function after combination chemotherapy in childhood for acute lymphoblastic leukemia. *Arch. Dis. Child.*, 1981; 56:275.
 26. Groosegoed, J.A. Towards a better understanding of Sertoli cell function (editorial). *Int. J. Androl.*, 1978; 10:727.
 27. de Kretser, D.M., Berger, H.G., Fortune, D., Hudson, B., Long, A.R., Paulsen, C.A. and Taft, H.P. Hormonal, histological and chromosomal studies in adult males with testicular disorders. *Clin. Endocrinol. Metab.*, 1972; 35:392.