



THE AGA KHAN UNIVERSITY

eCommons@AKU

Paediatrics and Child Health, East Africa

Medical College, East Africa

January 2014

# Pediatric cerebral palsy in Africa: A systematic review

Kirsten A. Donald  
*University of Cape Town*

Pauline Samia  
*Aga Khan University, pauline.samia@aku.edu*

Angelina Kakooza-Mwesige  
*Makerere University College of Health Sciences*

David Bearden  
*Children's Hospital of Philadelphia*

Follow this and additional works at: [http://ecommons.aku.edu/eastafrica\\_fhs\\_mc\\_paediatr\\_child\\_health](http://ecommons.aku.edu/eastafrica_fhs_mc_paediatr_child_health)

 Part of the [Pediatrics Commons](#)

## Recommended Citation

Donald, K. A., Samia, P., Kakooza-Mwesige, A., Bearden, D. (2014). Pediatric cerebral palsy in Africa: A systematic review. *Seminars in Pediatric Neurology*, 21(1), 30-35.

**Available at:** [http://ecommons.aku.edu/eastafrica\\_fhs\\_mc\\_paediatr\\_child\\_health/65](http://ecommons.aku.edu/eastafrica_fhs_mc_paediatr_child_health/65)

# Pediatric Cerebral Palsy in Africa: A Systematic Review

Kirsten A. Donald, MBChB, MPhil (Paed Neuro),\* Pauline Samia, MBChB, MPhil (Paed Neuro),†  
Angelina Kakooza-Mwesige, MBChB, MMed,‡,§ and David Bearden, MD||

Cerebral palsy is a common neurologic problem in children and is reported as occurring in approximately 2-2.5 of 1000 live births globally. As is the case with many pediatric neurologic conditions, very little has been reported on this condition in the African context. Resource-limited settings such as those found across the continent are likely to result in a different spectrum of etiologies, prevalence, severity as well as management approaches. This review aims to establish what has been reported on this condition from the African continent so as to better define key clinical and research questions.

Semin Pediatr Neurol 21:30-35 © 2014 Elsevier Inc. All rights reserved.

## Introduction

The term cerebral palsy (CP) describes a group of permanent disorders of the development of movement and posture causing activity limitation, which are attributed to nonprogressive disturbances that occurred in the developing fetal or infant brain. It is an umbrella term encompassing a range of different etiologies and phenotypes as well as associated with a variety of comorbid conditions (including epilepsy, intellectual disability, sensory disorders, and behavioral problems).<sup>1</sup> CP is a common neurologic problem in children and is reported as occurring in approximately 2-2.5 of 1000 live births globally.<sup>2</sup> As is the case with many pediatric neurologic conditions, very little has been reported on this condition in the African context. This review aims to establish what has been reported on this condition from the African continent so as to better define key clinical and research questions.

## Methods

We performed a systematic review of available literature to identify data on the prevalence, etiology, comorbidities, outcomes, screening tools, and treatment of CP in Africa. Abstracts from any study from or involving an African country were reviewed for possible inclusion. Studies reporting more general neurodevelopmental disabilities were included only if they specifically mentioned CP as an outcome. Studies that focused on a subgroup of children with CP (eg, premature infants or children with genetic disorders) were excluded.

## Search Strategy

Online databases were searched using specific search strategies to identify relevant articles published in English. Ovid Medline, PubMed, and the Cochrane database were searched using combinations of the terms “cerebral palsy,” “Africa,” “developing countries,” “developmental disabilities,” “neurological impairment,” and “childhood disability.” Web sites from African and international disability organizations as well as the World Health Organization were scanned for further references missed through the aforementioned search. The abstracts of any articles identified in this search strategy were screened for relevance, and electronic or hard copies of the full text were obtained for all relevant articles. References from each relevant article were scanned for further relevant articles and a snowballing search was performed. Commercial search engines Google and Bing were then used to check for any missing publications. Additional local journals not on the PubMed database (such as the East African Medical Journal) were searched for relevant articles. For the final analysis, only original research publications published after 1980 were

\*Division of Developmental Paediatrics, School of Child and Adolescent Health, University of Cape Town, Cape Town, South Africa.

†Department of Paediatrics and Child Health, Aga Khan University, Nairobi, Kenya.

‡Department of Paediatrics & Child Health, Makerere University College of Health Sciences, Kampala, Uganda.

§Neuropediatric Research Unit, Department of Women's & Children's Health, Astrid Lindgren Children's Hospital, Karolinska Institutet, Stockholm, Sweden.

||Division of Child Neurology, Children's Hospital of Philadelphia, Philadelphia, PA.

Address reprint requests to K.A. Donald, MBChB, MPhil (Paed Neuro), Red Cross War Memorial Children's Hospital, 5th Floor ICH Building, Klipfontein Rd, Rondebosch, Cape Town 7700, South Africa. E-mail: Kirsty.Donald@uct.ac.za

included. When multiple publications existed on the same study cohort, only the most recent or most relevant publication was included, and studies were included only if they had some special relevance to the African setting.

## Results

We identified 16 publications on prevalence or etiology or both of cerebral palsy in Africa (4 from Nigeria,<sup>3-6</sup> 4 from South Africa,<sup>7-10</sup> 2 from Ethiopia,<sup>11,12</sup> Ghana,<sup>13,14</sup> and Tanzania,<sup>15,16</sup> and 1 each from Egypt<sup>17</sup> and Kenya<sup>18</sup>). A summary of key prevalence and etiology studies is presented in Table 1.

There were 6 studies that examined screening instruments and strategies in an African setting for developmental disabilities including CP.<sup>19-24</sup> In addition, 2 articles provided an overview of screening strategies in low-resource settings for neurodevelopmental disabilities including CP.<sup>25,26</sup>

A total of 4 studies assessed the effect of caring for children with CP on the caregiver,<sup>27-30</sup> and 10 studies assessed specific aspects of treatment of children with CP<sup>31-36</sup> and comorbidities.<sup>37-40</sup> There were no studies that addressed broad longitudinal outcomes in children with CP in Africa.

## Prevalence

Methodologies for determining prevalence varied widely between studies, varying from door-to-door surveys to hospital-based record reviews. Almost all studies were cross-sectional. A common technique was a 2-phase design using the World Health Organization Ten Questions screen administered by a community health worker to identify possible cases followed by confirmation by a clinician.<sup>26</sup> Prevalence varied widely from country to country from approximately 2-10 per 1000<sup>9,25,41</sup> children in community-based samples, although owing to heterogeneity between studies, it is difficult to

attribute this to true differences in prevalence or to differences in methodology (Table 1).

## Etiology or Risk Factors

The most common reported etiologies identified in African cohorts were birth asphyxia, kernicterus, and neonatal infections, with prematurity or low birth weight identified as a major etiology in only 2 studies (Table). This is in contrast with most studies in the United States and Europe in which prematurity or low birth weight is one of the major risk factors identified in almost all studies.<sup>42</sup>

## Identification and Classification

Several studies on prevalence attempted to assess the sensitivity and specificity of their screening methodology and found that the Ten Questions Screen followed by clinician confirmation had good sensitivity and reasonable specificity. A study conducted in Uganda and Sri Lanka attempted to develop and validate a screening “portfolio” (short battery of tests) that could be used by community health workers, though the study is limited by minimal information about how the screen was used and did not report sensitivity or specificity.<sup>19</sup> Two studies, one in Malawi (the Malawi Developmental Assessment Tool) and another in Kenya (Kilifi Developmental Inventory), have developed developmental screening tools aimed at identifying children with neurodevelopmental disabilities including CP.<sup>21,22</sup> A Zimbabwean group described the use of the Neonatal Neurological Examination based on Prechtl’s general movements to identify neonates at risk for developmental disabilities including CP.<sup>43</sup>

Studies from the African continent published to date have reported a larger proportion of severely affected children or those with Gross Motor Functional Classification System IV

**Table Summary of Key Studies Evaluating Prevalence and Etiology of Cerebral Palsy in Africa**

Study	Country, Region, and Income Status	Design	Prevalence (per 1000)	Most common Etiologies
El-Tallawy et al <sup>41</sup>	Egypt North Africa Lower middle income	Prospective Community based Cross-sectional	2	Birth asphyxia (34.6%) Preterm birth (17.3%) Kernicterus (15.4%)
Ogunlesi and Ogundeyi <sup>3</sup>	Nigeria  West Africa Lower middle income	Retrospective  Case series Hospital based	Not measured	Birth asphyxia (57.6%)  Kernicterus (36.9%) CNS infections (21.7%)
Couper <sup>9</sup>	South Africa Southern Africa Upper middle income	Prospective Community based Cross-sectional	10	Not reported
Karumuna and Mgone <sup>15</sup>	Tanzania  Southern Africa Low income	Prospective  Hospital based Cross-sectional	Not measured	Birth asphyxia (36%)  Low birth weight (20%) CNS infections (15%)

CNS, central nervous system.

and V in their services compared with European and North American cohorts.<sup>4,9,15,17</sup>

## Management

Children with CP require multidisciplinary care for optimal management of their medical needs as well as support to maximize their developmental and educational potential.<sup>45</sup> Guidelines for the recommended investigation and management of specific aspects of CP are available through pediatric organizations such as the American Academy of Pediatrics.<sup>46</sup> However, little is known about either the availability of guidelines or the recommendations in practice across resource-poor countries in Africa. In the first instance, several studies identified barriers toward optimal care for children with CP in Africa. Limited access to health care facilities and specialists, as well as a lack of adaptive equipment such as wheelchairs and other ambulation aids, contributes to the treatment gap for children with CP.<sup>40</sup> In addition, high levels of social stigma toward children with neurologic disorders were reported as reasons for families failing to seek treatment even when it was available.<sup>41</sup>

Further contributing to difficulties in accessing health care facilities for children with CP is a lack of wheelchair-accessible transport.<sup>28,44</sup> This represents a particular challenge for families with older nonambulant children with CP. A multisite country study conducted by the African Child Policy Forum on the lives of children with disabilities in Africa noted that poverty powerfully affected the caregivers' ability to provide for the basic needs of their disabled children.<sup>44</sup> Caregivers found the financial costs attached to medical care, rehabilitation, assistive devices, and transportation to be a significant barrier to seeking care. Community-based rehabilitation programs were reported by parents as being effective in increasing access to education and assistive devices, but none of these have been evaluated systematically.

There are isolated reports on experience in specific aspects or approaches to management for children with CP in selected populations. These include a South African study in which an 8-week randomized controlled trial of a strength training program for adolescents with CP showed improved gait and body image perception.<sup>35</sup> Another study examined a video-based training model for providers working with children with CP.<sup>34</sup>

In relation to the medical complications of CP, spasticity occurs in most children with this condition and may significantly hamper mobility and fine motor function. Spasticity is defined as a velocity-dependent increase in muscle tone and is one of the core impairments experienced by children with CP.<sup>46</sup> Although effective use of botulinum toxin has been described in a group from Egypt,<sup>47</sup> there is no information regarding how accessible this treatment option is across the continent as a whole. There is limited access to several other supportive procedures reported in isolated centers across Africa. These include musculotendinous release for spastic hips<sup>48</sup> and gait analysis assessments.<sup>49-51</sup> Finally, a study on selective dorsal rhizotomy provided long-term evidence for

safety and efficacy of this treatment for spasticity in CP in a resource-limited setting in South Africa, although the sample size was small.<sup>31,53</sup>

Home- or community-based approaches to rehabilitation have been described as being well suited for low-resource settings as they often require relatively minimal infrastructure and resources to implement compared with alternative models, such as center-based interventions.<sup>54</sup> In addition, they have been described as providing opportunities for strengthening the parent-child bond.<sup>14</sup> There is a lack of information regarding any intervention studies or evidence-based clinical studies on the efficacy of home-based interventions from Africa. Most of the information on home-based studies has been done in high-income countries using varying interventions and presents conflicting results.<sup>54-56</sup>

## Comorbidities

Consistent with reports from North American and European groups,<sup>57,58</sup> epilepsy is one of the most common comorbidities of CP in Africa. In the cohort studied in Dar Es Salaam in Tanzania, the most common comorbidities were epilepsy (in 35% of children with CP), followed by deafness, speech impairment, and visual impairment. In the study by Ogunlesi et al conducted in Nigeria, the most common comorbidities were seizures (in 46.7%), speech impairment (43.5%), visual impairment (25%), and cognitive impairment (24%). Although not specifically identified as a comorbidity in this study, malnutrition was also identified in 80.4% of children with CP. A second Nigerian study by Lagunju and Fatunde<sup>38</sup> also identified epilepsy as a common comorbidity of CP, occurring in 38% of children.

Seizure-related morbidity and mortality are comorbidities that could potentially be reduced by appropriate anticonvulsant use in children with CP.<sup>59</sup> Of those living with epilepsy, 80% globally reside in resource-poor regions like Africa<sup>60</sup>; it is estimated that 23% of people with active epilepsy in Senegal and up to 100% of such patients in parts of Uganda, Togo, the Gambia, and Tanzania do not access the anticonvulsants they require.<sup>37,60</sup> Untreated epilepsy is likely to contribute to increased mortality as well as loss of developmental potential in children with CP and epilepsy in Africa.

Vision abnormalities have been identified in 58% of African children with CP in a study.<sup>39</sup> Strabismus occurred in 15%-34% of children in this group. This is a potentially a surgically correctible anomaly with resultant improvement in binocular alignment and fusion. Regular diagnosis and management of ocular anomalies in Africa is enhanced by several nongovernmental agencies, but access to intervention services remains suboptimal.<sup>39,40</sup>

Other common comorbidities reported in high-resource settings, such as poor feeding, orthopedic complications, and cognitive impairment, are likely to be even more common in African cohorts given the high percentage of more severe forms of CP and limited access to health care. However, there are minimal data on the prevalence or management of these comorbidities in Africa.

## Discussion

Despite concerns about underreporting of CP in many African communities, the prevalence estimates reported here were generally higher than the estimated 2-2.5 of 1000 in most studies conducted in the United States or Europe.<sup>61</sup> These discrepancies may in part be because of methodological differences in the way these studies were conducted, but it is likely that the prevalence of CP is indeed higher in Africa because of the level of perinatal complications such as birth asphyxia and neonatal infections. However, variations in data collection methodologies between studies make direct comparisons between countries difficult.

As noted earlier, several major contributors to CP in Africa are potentially preventable and include birth asphyxia, central nervous system infections, and kernicterus. Prematurity and low birth weight are proportionally much smaller contributors to CP in Africa than in high-resource settings, likely because of relatively poor survival among these infants. However, it should be noted that most studies on etiology lacked a control group and few discussed issues of confounding and bias. Further, CP is a term often used in African contexts to describe all motor disability syndromes, and the term is considered synonymous with birth asphyxia in many countries. Thus, in the absence of a control group, studies reporting high rates of CP due to birth asphyxia should be viewed with a certain amount of caution.

What is clear is that there is a lack of structured and consistent screening policy for developmental disabilities amongst infants and preschool children in Africa. Families that can afford privately funded health care have access to pediatricians directly and may have access to some screening. However, even within these relatively well-resourced communities, programs are not standardized and no consistent strategy exists for referral lines or interventions when problems are identified. For most populations in Africa, recognition of a child with a disability relies on the parents or caregivers identifying a problem and presenting to medical facilities. On a continent where access to even the most basic health services is often poor and the burden of severe diseases high, this situation leaves many children with disabilities unrecognized and hence without appropriate intervention.

There may be a greater proportion of children with more severe disability in resource-poor countries because of delayed presentation of a range of disorders as well as the lack of early intervention services. However, this hypothesis needs to be tested across population-based epidemiologic studies. The relatively increased severity spectrum seen in reports of African cohorts may also be because of a combination of selection bias (hospital cohorts vs community- or population-based studies), lack of access to medical services (resulting in only those severely affected children requiring medical input to survive being brought to medical attention), and powerful stigma against disabled children (resulting in children being “hidden” when medical attention is not considered essential). Children with disabilities and their families in African countries are frequently excluded from society because of stigmatization. Most of these children consequently confront many challenges socially, economically, and politically because they are often

denied the basics of health care, education, socialization, and recognition.

Despite reports from a number of groups describing aspects of care for children with CP on the African continent, there appears to be very little information on the coordinated management of multiple impairments and associated comorbidities which these children may be affected with. There is a shortage of trained health workers and rehabilitation personnel across the continent, with few personnel specifically trained in the tools necessary to care for and to manage a child with CP in rural areas. In addition, when infrastructure exists, it is constrained by a lack of cultural- and language-specific validated clinical assessment tools, low literacy rates, and limited facilities for rehabilitative services and equipment. Initiatives that have emerged in some countries in the last 10 years have included a combination of nongovernmental organization-driven and government-subsidized community- and centre-based programs. Promotion of the use of adaptable low-cost materials, community health care workers, development of parent support groups, and multidisciplinary outreach programs may serve as means to manage CP in Africa. These need to be evaluated for effectiveness in the African context.<sup>44,52,62</sup>

Limitations of a review such as this include the fact that there is likely to be much information on these topics that is not published in peer-reviewed English-language journals, such as conference abstracts, thesis dissertations, and other “gray literature.” In particular, the Francophone countries in West Africa represent a large subpopulation within the continent and are not well represented in this review because of the exclusion of non-English-language publications. In addition, owing to our emphasis on CP, we did not include studies on other neurologic disorders that might also have relevance for CP, for example, studies of community rehabilitation for children with visual impairment. Despite this, the available literature does highlight some useful points as described.

This review suggests several gaps of knowledge regarding CP in Africa. The lack of validated research criteria for identifying cases of CP in low-resource settings and the absence of standardized methodologies for assessing prevalence prevent meaningful comparisons between countries or assessing change in prevalence over time. Most studies are simple cross-sectional descriptive studies and there is a lack of appropriate control groups that would allow better assessment of risk factors. Large multicenter studies with a standardized methodology and appropriate control groups would help to address this gap. In addition, qualitative or mixed-methods studies assessing patient and family understanding of CP, access to resources, and barriers to care are virtually absent in Africa and would help in designing future interventions. Finally, there is a need for longitudinal studies that would assess outcomes over time for patients with CP as well as for randomized controlled trials of community-based treatment strategies that would be appropriate in an African setting.

## References

1. Bax M, Goldstein M, Rosenbaum P, et al: Proposed definition and classification of cerebral palsy. *Dev Med Child Neurol* 47:571-576, 2005

2. Shevell M, Dagenais L, Oskoui M: The epidemiology of cerebral palsy: New perspectives from a Canadian Registry. *Semin Pediatr Neurol* 20:60-64, 2013
3. Ogunlesi T, Ogundeyi M, Ogunfowora O, et al: Socio-clinical issues in cerebral palsy in Sagamu, Nigeria. *S Afr J Child Health* 3:120-124, 2008
4. Sathiakumar N, Yakubu AM: Cerebral palsy in Zaria, northern Nigeria—Is it preventable? *J Trop Pediatr* 33:263-265, 1987
5. Izuora GI, Iloje SO: A review of neurological disorders seen at the Paediatric Neurology Clinic of the University of Nigeria Teaching Hospital, Enugu. *Ann Trop Paediatr* 9:185-190, 1989
6. Nottidge VA, Okogbo M: Cerebral palsy in Ibadan, Nigeria. *Dev Med Child Neurol* 33:241-245, 1991
7. Christianson AL, Zwane ME, Manga P, et al: Children with intellectual disability in rural South Africa: Prevalence and associated disability. *J Intellect Disabil Res* 46:179-186, 2002
8. Arens LJ, Molteno CD: A comparative study of postnatally-acquired cerebral palsy in Cape Town. *Dev Med Child Neurol* 31:246-254, 1989
9. Couper J: Prevalence of childhood disability in rural KwaZulu-Natal. *S Afr Med J* 92:549-552, 2002
10. Rubin IL, Davis M: Etiology of developmental disabilities in Soweto, South Africa. *Am J Public Health* 76:1112-1114, 1986
11. Tekle-Haimanot R, Abebe M, Gebre-Mariam A, et al: Community-based study of neurological disorders in rural central Ethiopia. *Neuroepidemiology* 9:263-277, 1990
12. Tamrat G, Kebede Y, Alemu S, et al: The prevalence and characteristics of physical and sensory disabilities in northern Ethiopia. *Disabil Rehabil* 23:799-804, 2001
13. Biritwum RB, Devres JP, Ofosu-Amaah S, et al: Prevalence of children with disabilities in Central Region, Ghana. *West Afr J Med* 20:249-255, 2001
14. Commey JO, Richardson JE: Cerebral palsy in Ghanaian children. *West Afr J Med* 3:157-163, 1984
15. Karumuna JM, Mgone CS: Cerebral palsy in Dar Es Salaam. *Cent Afr J Med* 36:8-10, 1990
16. Makwabe CM, Mgone CS: The pattern an aetiology of cerebral palsy as seen in Dar es Salaam, Tanzania. *East Afr Med J* 61:896-899, 1984
17. El-Tallawy HN, Farghaly WM, Shehata GA, et al: Epidemiology of cerebral palsy in El-Kharga District-New Valley (Egypt). *Brain Dev* 33:406-411, 2011
18. Mung Ala-Odera V, Meehan R, Njuguna P, et al: Prevalence and risk factors of neurological disability and impairment in children living in rural Kenya. *Int J Epidemiol* 35:683-688, 2006
19. Wirz S, Edwards K, Flower J, et al: Field Testing of the ACCESS materials: A portfolio of materials to assist health workers to identify children with disabilities and offer simple advice to mothers. *Int J Rehabil* 28:293-302, 2005
20. Bower JH, Howlett W, Maro VP, et al: A screening instrument to measure the prevalence of neurological disability in resource-poor settings. *Neuroepidemiology* 32:313-320, 2009
21. Arubakar A, Holding P, Van Baar A, et al: Monitoring psychomotor development in a resource-limited setting: An evaluation of the Kilifi Developmental Inventory. *Ann Trop Paediatr* 28:217-226, 2008
22. Gladstone M, Lancaster GA, E. Umar, et al: The Malawi Developmental Assessment Tool (MDAT): The creation, validation, and reliability of a tool to assess child development in rural African settings. *PLoS Med* 7: e1000273. <http://dx.doi.org/10.1371/journal.pmed.1000273>
23. Gona J, Newton CR, Hartley S: Using participatory rural appraisal in the identification of children with disabilities in rural Kilifi, Kenya. *Rural Remote Health* 6:553, 2006
24. Mung'ala-Odera V, Meehan R, Njuguna P, et al: Validity and reliability of the "Ten Questions" questionnaire for detecting moderate to severe neurological impairment in children aged 6–9 years in rural Kenya. *Neuroepidemiology* 23:67-72, 2004
25. Mung'ala-Odera V, Newton CR: Identifying children with neurological impairment and disability in resource-poor countries. *Child Care Health Dev* 33:249-256, 2006
26. Maulik PK, Darmstadt GL: Childhood disability in low- and middle-income countries: Overview of screening, prevention, services, legislation, and epidemiology. *Pediatrics* 120:S1-S55, 2007
27. Geere JL, Gona J, Omondi FO, et al: Caring for children with physical disability in Kenya: Potential links between caregiving and carers' physical health. *Child Care Health Dev* 39:381-392, 2012
28. Hartley S, Ojwang P, Baguwemu A, et al: How do carers of disabled children cope? The Ugandan perspective. *Child Care Health Dev* 31:167-180, 2005
29. Hamzat TK, Mordi EL: Impact of caring for children with cerebral palsy on the general health of their caregivers in an African community. *Int J Rehabil Res* 30:191-194, 2007
30. Fatudimu MB, Hamzat TK, Akinyinka OO: Comparative quality of life of Nigerian caregivers of children with cerebral palsy. *Int J Ther Rehabil* 20:131-135, 2013
31. Langerak NG, Lamberts RP, Fieggen AG, et al: Selective dorsal rhizotomy: Long-term experience from Cape Town. *Childs Nerv Syst* 23:1003-1006, 2007
32. Lundgren-Lindquist B, Nordholm L: Community-based rehabilitation—A survey of the disabled in a village in Botswana. *Disabil Rehabil* 15:83-89, 1993
33. Iloje SO, Ejike-Orji E: Compliance by cerebral palsy patients attending a child neurology service in a developing country: A preliminary report. *West Afr J Med* 12:1-5, 1993
34. Holloway S, Lee L, McConkey R: Meeting the training needs of community-based service personnel in Africa through video-based training courses. *Disabil Rehabil* 21:448-454, 1999
35. Unger M, Faure M, Frieg A: Strength training in adolescent learners with cerebral palsy: A randomized controlled trial. *Clin Rehabil* 20:469-477, 2006
36. McConkey R, Mphole P: Training needs from developing countries: Experiences from Lesotho. *Int J Rehabil Res* 23:119-123, 2000
37. Edwards T, Scott AG, Munyoki G, et al: Active convulsive epilepsy in a rural district of Kenya: A study of prevalence and possible risk factors. *Lancet Neurol* 7:50-56, 2008
38. Lagunju IA, Fatunde OJ: Epilepsy in Nigerian children with cerebral palsy. *Dev Med Child Neurol* 49:158-159, 2007
39. Lagunju IA, Oluleye TS: Ocular abnormalities in children with cerebral palsy. *Afr J Med Sci* 36:71-75, 2007
40. Njambi L, Kariuki M, Masinde S: Ocular findings in children attending occupational therapy clinic at Kenyatta National Hospital, Nairobi, Kenya. *East Afr J Ophthalmol* 15:21-26, 2009
41. El Tallawy HNA, Farghaly WMA, Metwaly NA, et al: Door-to-Door Survey of Major Neurological Disorders in Al Kharga District, New Valley, Egypt: Methodological Aspects. *Neuroepidemiology* 35:185-190, 2010
42. Gladstone M: A review of the incidence and prevalence, types and aetiology of childhood cerebral palsy in resource-poor settings. *Ann Trop Paediatr* 30:181-196, 2010
43. Wolf MJ, Wolf B, Bijleveld C, et al: Neurodevelopmental outcome in babies with a low Apgar score from Zimbabwe. *Dev Med Child Neurol* 39:821-826, 1997
44. African Child Policy Forum. *The Lives of Children With Disabilities in Africa: A Glimpse into a Hidden World*. Addis Abbaba, Ethiopia: African Child Policy Forum, 2011
45. Novak I, McIntyre S, Morgan C, et al: A systematic review of interventions for children with cerebral palsy: State of the evidence. *Dev Med Child Neurol* 55:885-910, 2013
46. Cooley W: Providing a primary medical care home for children and youth with Cerebral Palsy. *Pediatrics* 114:1106-1113, 2004
47. Salem ME, El-Shakankiry HM, El-Kahky AM, et al: The effect of botulinum toxin type-A injection on spasticity, range of motion and gait patterns in children with spastic diplegic cerebral palsy: An Egyptian study. *Int J Rehabil Res* 27:275-281, 2004
48. Bishay SN: Short-term results of musculotendinous release for paralytic hip subluxation in children with spastic cerebral palsy. *Ann R Coll Surg Engl* 90:127-132, 2008
49. Langerak NG, Lamberts RP, Fieggen AG, et al: A prospective gait analysis study in patients with diplegic cerebral palsy 20 years after selective dorsal rhizotomy. *J Neurosurg Pediatr* 1:180-186, 2008
50. Hennessey MJ, Dixon SD, Simon SR: The development of gait: A study in African children aged one to five. *Child Dev* 55:844-853, 1984

51. Zverev Y, Adeloye A, Chisi J: Quantitative analysis of gait pattern in hemiparetic patients. *East Afr Med J* 79:420-422, 2002
52. Langerak NG, Tam N, Vaughan CL, et al: Gait status 17-26 years after selective dorsal rhizotomy. *Gait Posture* 35:244-249, 2012
53. Novak I, Cusick A, Lannin N: Occupational therapy home programs for cerebral palsy: Double-blind, randomized, controlled trial. *Pediatrics* 124: e606-e614, 2009
54. Gates PE, Banks D, Johnston TE, et al: Randomized controlled trial assessing participation and quality of life in a supported speed treadmill training exercise program vs. a strengthening program for children with cerebral palsy. *J Pediatr Rehabil Med* 5:75-88, 2012
55. Eugster-Buesch F, de Bruin ED, Boltshauser E, et al: Forced-use therapy for children with cerebral palsy in the community setting: A single-blinded randomized controlled pilot trial. *J Pediatr Rehabil Med* 5:65-74, 2012
56. Carlsson M, Hagberg G, Olsson I: Clinical and aetiological aspects of epilepsy in children with cerebral palsy. *Dev Med Child Neurol* 45:371-376, 2003
57. Sellier E, Uldall P, Calado E, et al: Epilepsy and cerebral palsy: Characteristics and trends in children born in 1976–1998. *Eur J Paediatr Neurol* 16:48-55, 2012
58. Birbeck G, Hesdorffer D: The geography of epilepsy: A fatal disease in resource-poor settings. *Neurology* 77:96-97, 2011
59. Mbuba CK, Ngugi AK, Newton CR, et al: The epilepsy treatment gap in developing countries: A systematic review of the magnitude, causes, and intervention strategies. *Epilepsia* 49:1491-1503, 2008
60. Durkin M: The epidemiology of developmental disabilities in low-income countries. *Ment Retard Dev Disabil Res Rev* 8:206-211, 2002
61. Gottlieb CA, Maenner MJ, Cappa C, et al: Child disability screening, nutrition, and early learning in 18 countries with low and middle incomes: Data for the third round of UNICEF's Multiple Indicator Cluster Survey (2005-06). *Lancet Neurol* 374:1831-1839, 2009
62. Engle PL, Black MM, Behrman JR: Strategies to avoid the loss of developmental potential in more than 200 million children in the developing world. *Lancet* 369:229-242, 2007