



Review Article

Open Access

Aflatoxins and Hepatitis B, C Viral Associated Hepatocarcinogenesis

Humaira Qureshi^{1,2}, Syed Shayan Ali^{1,2}, Mazhar Iqbal³, Anwar Ali Siddiqui¹, Naveed Ahmed Khan^{1*} and Saeed S Hamid²¹Department of Biological and Biomedical Sciences, Aga Khan University, Karachi, Pakistan²Department of Medicine, Aga Khan University, Karachi, Pakistan³Drug Discovery & Structural Biology, Health Biotechnology Division, National Institute for Biotechnology & Genetic Engineering (NIBGE), Faisalabad, Pakistan

Abstract

Hepatocellular carcinoma is a serious human disease with fatal consequences. The most distressing aspect of hepatocellular carcinoma is the limited improvement in mortality (mortality rate of more than 90%). At present, the underlying molecular mechanisms are not well understood and treatment options are often of limited efficacy. This review presents our current understanding of the burden of hepatocellular carcinoma on human health, pathogenesis and pathophysiology, and molecular mechanisms associated with the disease, as well as our knowledge of the physical barriers, cellular mechanisms and molecular elements that may be targets for therapeutic interventions and/or the development of preventative measures. As the proposed findings present a major risk to public health, it is hoped that robust intervention measures will be introduced for aflatoxins monitoring and reduction in diet.

Keywords: Hepatocarcinogenesis; Hepatocellular carcinoma; Aflatoxins; Mycotoxins; Risk factors; Pathogenesis

Introduction

Hepatocellular carcinoma (HCC) is the sixth most common cancer and the second leading cause of all cancer related deaths worldwide [1]. According to IARC (2012), 782,000 new cases of HCC are diagnosed per year with a mortality rate of 95% (Table 1) [2]. The overall incidence in developing countries is five times that of developed nations (513,000 vs. 110,000) [3]. In United States, the overall incidence rate of HCC has been estimated at 2.99 per 100,000 [4,5]. However, for Asian and African countries, figures are quite alarming, i.e. about 50-150 cases per 100,000 [3,6]. Another report suggests approximately 82% incidence rate in developing countries, particularly among South Asia with an annual incidence of 24 per 100,000 with much greater occurrence in the high-risk areas [7,8] (Figure 1 and Table 2). This is because these areas have higher risk factors, such as high prevalence of hepatitis B and C infections, behavioral factors like obesity, diabetes mellitus, betel nuts chewing, and an important but largely ignored factor of exposure to aflatoxins [8-10].

Aflatoxins are secondary metabolites primarily synthesized by food-borne *Aspergillus* species. Aflatoxins commonly contaminate maize, nuts, rice, wheat, and spices like chilies, black pepper, turmeric and other food items that are consumed in raw form. Products obtained from animals that have ingested contaminated feed are also recognized as indirect source. Aflatoxins being highly lipophilic are readily absorbed through the gastrointestinal tract. They may also enter the bloodstream directly by inhalation [11]. On a global scale, approximately 4.5 billion persons living in developing countries are frequently exposed to unchecked amounts of the toxin [12] (Table 3). Consumption of aflatoxin-contaminated food has been associated with HCC [13]. According to a study conducted in 2010, there are 550,000–600,000 new HCC cases reported and out of those about 25,200–155,000 may be attributable to aflatoxin exposure [14]. The regions where most of these cases occur are sub-Saharan Africa, China, and Southeast Asia; areas where the elevated prevalence of viral hepatitis, as well as higher occurrence of aflatoxin-contaminated foods, especially the maize have been reported [14]. For example in China, reported incidence of contamination in corn was 85%, while in Pakistan, reported incidence of aflatoxin contamination in maize was found to be 41.6% and 25% in red chillies, respectively [12,15] (Table 4). Both countries have significant prevalence of viral hepatitis infections as well [16].

There are many types of aflatoxins (B1, B2, G1, G2 and M1) but aflatoxin B1 (AFB1) has been classified as a Group 1 carcinogen (definitely carcinogenic to humans) and one of the most potent toxins associated with liver cancer [17]. AFB1 metabolites have the ability to bind with albumin and the measurement of aflatoxin-serum albumin adducts has been used to detect recent exposure in individuals [18-20]. AFB1 metabolites react with guanine residues in DNA resulting in (AFB1)-N7-guanine adducts. A Guanine to Thiamine transversion mutation in codon 249 of TP53 gene has been strongly associated with aflatoxin exposure and HCC cases [21]. The ability to modulate DNA activity by creating such lesions in the DNA may play a role in AFB1 induced carcinogenesis [22].

There is a significant synergetic effect between aflatoxin and Hepatitis B virus (HBV) infection [23]. Another study has associated aflatoxin exposure with advanced liver disease in chronic Hepatitis C patients [18]. This objective of this article is to emphasize the deleterious effects of aflatoxins-contaminated foods and to highlight the relationship between aflatoxins and viral associated hepatocellular carcinoma. As the proposed findings present a major risk to public

Region	World	More developed regions	Less developed regions
Cases	782	134	648
Deaths	746	123	622
Percentage (%)	95	92	96

Table 1: Incidence of HCC in the world. There is poor prognosis of HCC, with majority of patients expiring within 5 years. The majority of the cases are from less developed regions [2].

***Corresponding author:** Naveed Ahmed Khan, Department of Biological and Biomedical Sciences, Aga Khan University, Karachi – 74800, Pakistan, Tel: +92-21-3486-4541; E-mail: naveed5438@gmail.com

Received: July 18, 2014; **Accepted:** September 27, 2014; **Published:** September 29, 2014

Citation: Qureshi H, Ali SS, Iqbal M, Siddiqui AA, Khan NA, et al. (2014) Aflatoxins and Hepatitis B, C Viral Associated Hepatocarcinogenesis. J Cell Sci Ther 5: 179. doi: [10.4172/2157-7013.1000179](http://dx.doi.org/10.4172/2157-7013.1000179)

Copyright: © 2014 Qureshi H, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1: Number of incidence from 2000-2009.

at least 75% of HCC. In the US, Africa and Europe the prevalence of HBV among HCC cases was reported as 9%, 40% and 22%, respectively [16]. In Asia, proportion of HBsAg positive HCC cases was greater than 50% in China, Taiwan, Korea, Thailand, Vietnam, and Turkey. However, for Pakistan and India the prevalence is 30 and 45% respectively [16]. The annual risk of HCC is 0.5% for asymptomatic HBsAg carriers and 0.8% for patients with chronic hepatitis B [24]. Both HBV and HCV can directly lead to HCC reviewed in HBV is an oncogenic virus that partially contains double stranded DNA and it can directly integrate into the host genome, leading to changes in genomic function or chromosomal instability [25]. Of note, the integration of HBV DNA into the host genome is not required for its replication, however it does allow for the persistence of the viral genome within the host. Furthermore, several HBV factors have been implicated in hepatocarcinogenesis, including the HBx gene, the pre-S2/S gene and the HBV spliced protein. By contrast to HBV, HCV is a positive-stranded RNA virus. As it lacks reverse transcriptase, it cannot integrate into the host genome. Being a completely cytoplasmic replicating virus, its involvement in HCC occurs via indirect pathways through the effects of chronic inflammation and hepatocellular injury.

Year	Gender	ASR for HCC/ 100,000 in Karachi
1995 - 1997	Males	5.7
	Females	3.7
1998 - 2002	Males	5.3
	Females	4.0
Year	Gender	ASR for HCC/ 100,000 in Larkana
2000 - 2002	Males	10.5
	Females	2.0
Year	Gender	ASR for HCC/ 100,000 in Hyderabad
1998 - 2002	Males	4.4
	Females	1.2
Year	Gender	ASR for HCC/ 100,000 in Quetta
1998 - 1999	Males	12.3
	Females	3.1
Gender		ASR for HCC/ 100,000 in Karachi
1995 - 1997	Males	5.7
	Females	3.7
1998 - 2002	Males	5.3
	Females	4.0
Year		ASR for HCC/ 100,000 in Larkana
2000 - 2002	Males	10.5
	Females	2.0
Year		ASR for HCC/ 100,000 in Hyderabad
1998 - 2002	Males	4.4
	Females	1.2
Year		ASR for HCC/ 100,000 in Quetta
1998 - 1999	Males	12.3
	Females	3.1

Table 2: Year wise increase in HCC cases with respect to gender in different cities of Pakistan. The male to female ratio is 3.6:1.0 [7,61]

health, it is hoped that robust intervention measures will be introduced for aflatoxin monitoring and reduction in diet.

Synergism of Aflatoxin and Hepatitis B Viral Infection in HCC

HBV and HCC

According to WHO, HBV is the second known carcinogen affecting liver after tobacco and there are approximately 350 million people chronically infected with HBV worldwide and responsible for

Food Product	Country	Aflatoxin Levels in µg/kg	Maximum Allowed level according to commission of European Communities. (µg/kg)	Reference
Bread	Lebanon	1.12	2.0	[62]
Chocolate	Lebanon	4.33	2.0	[63]
Nuts	Lebanon	4.40	8.0	[63]
Rice, wheat, Oat, Barley, Maize	Malaysia	0.12-4.54	5.0	[64]
Peanuts	Malaysia	0.66-15.33	8.0	[64]
Rice	Iran	0.0029	5.0	[65]
Puffed Corn Snack	Iran	0.0012	8.0	[65]
Peanuts	Iran	0.00154	8.0	[65]
Bean	Iran	0.21-0.29	8.0	[66]
Maize	Iran	30	5.0	[67]
Peanuts	Iran	15	8.0	[67]
Rice	Pakistan	8.5	5.0	[68]
Corn	Pakistan	8	8.0	[68]
Corn Products	Pakistan	5.5	8.0	[68]
Dried Chili	Pakistan	Upto 90	5.0	[69]
Capsicum	Pakistan	Upto 25	5.0	[70]
Corn based products	Indonesia	5.8 - 12.4	8.0	[71]
Corn	Indonesia	119	8.0	[72]
Corn	Phillippines	49	8.0	[73]
Corn	Thailand	63	8.0	[73]
Maize	Taiwan	15	5.0	[67]
Peanuts	Taiwan	15	8.0	[67]
Rice	India	<15	5.0	[74]
Maize	India	30	5.0	[67]
Peanuts	India	30	8.0	[67]
Maize	China	40	5.0	[67]
Peanuts	China	49	8.0	[67]
Nuts and Product	Japan	0.3-128	8.0	[75]
Maize	Nepal	40	5.0	[67]
Peanuts	Nepal	40	8.0	[67]

Table 3: Aflatoxin levels in staple food products of different countries in Asia.

Sampling Region in Pakistan	Types of Mycotoxins studied	Types of Foodstuffs Analyzed	No. of Samples Analyzed	Aflatoxins Level µg/kg ^a	Percentage of positive samples ^b	References
Punjab	aflatoxin B ₁ , total and ochratoxin A	rice	68	12.08	28	[68]
		corn	105	5.47	14	
		corn products	102	7.85	20	
36 districts of Punjab	aflatoxin M ₁	milk sweet	232 138	0.252 0.58	32 78	[76]
Across Pakistan	aflatoxin B ₁ , B ₂ , G ₁ , G ₂	paddy rice	58	16.35	64	[40]
		parboiled rice	69	14.20	38	
		brown rice	84	9.85	33	
		white rice	93	7.10	42	
		broken rice	109	8.50	50	
Punjab	Aflatoxin M ₁	milk,	107	150.7	58	[77]
		yogurt,	96	90.4	47	
		white cheese,	119	147.8	15	
		cream,	150	102.6	11	
		Butter	74	69.7	52	
Faisalabad District Punjab	Aflatoxin M ₁	buffaloes milk,	55	0.013	15.8	[78]
		cows milk,	40	0.014	20	
		goats milk,	30	0.002	nil	
		sheep milk,	24	0.002	nil	
		camels milk	20	Nil	nil	
Punjab	Aflatoxin B ₁ , B ₂ , G ₁ , G ₂	Longi chillies	15	15.91	53	[79]
		Wonder hot chillies	12	5.77	8	
		Skyline 1 chillies	13	12.15	38	
Swat Valley NWFP	Aflatoxin B ₁ , ochratoxin A	Maize Kernels upper swat	18	14.94	66	[80]
		Maize Kernels lower swat	18	16.22	88	
Major districts of Punjab	Aflatoxin B ₁ Aflatoxin M ₁	halva	56	3.79	20	[81]
		pistachio	71	9.48	23	
		almonds	63	12.98	34	
		semolina	69	7.45	12	
		cardamom	34	2.34	11	
		raisins	46	5.34	7	
		halva puri	39	7.80	9	
		wheat powder	53	5.67	11	
Punjab 4 districts (Faisalabad, Lahore, Jhang and Toba Tek Singh)	aflatoxin B ₁ , B ₂ , G ₁ , G ₂ Ochratoxin A	<u>open market:</u>				[82]
		chilli sauce	26	10.4	12	
		crushed chilli	29	22.9	27	
		powdered chilli	34	27.5	35	
		<u>Restaurant:</u>				
		chilli sauce	24	11.3	8	
		crushed chilli	28	19.8	32	
		powdered chilli	29	21.1	41	
NWFP and Northern area of Pakistan	aflatoxin B ₁ , B ₂ , G ₁ , G ₂	<u>Dried fruits & nuts:</u>				[83]
		dried apricot,	20	4.55	5	
		dates,	20	2.5	nil	
		dried figs,	10	5.81	30	
		mulberries,	15	2.22	nil	
		dried raisins,	10	5.05	10	
		apricot kernels,	15	2.65	7	
		almonds with shells,	10	nil	nil	
		almonds without shells,	10	2.13	nil	
		walnuts with shells,	10	6.45	20	
		walnuts without shells,	10	3.43	30	
		peanut with shells,	10	5.10	10	
		peanuts without shells,	10	5.20	20	
		pistachios with shells,	10	2.10	nil	
		pistachios without shells,	10	6.34	20	
		pine nuts with shell,	10	3.25	nil	
Central area of punjab	Aflatoxin M ₁	Winter season & summer	48 & 56	0.073 & 0.028	27 & 23	[84]
		Raw milk	45 & 39	0.060 & 0.021	24 & 23	
		UHT milk	51 & 45	0.053 & 0.019	25 & 18	
		Yogurt	35 & 35	0.036 & 0.015	34 & 20	
		Butter	42 & 37	0.021 & 0.012	17 & 5	
		Ice cream				

Table 4: Natural Aflatoxin occurrence in Pakistani foodstuffs. (^aAverage concentration of total aflatoxins where multiple aflatoxins analyzed, otherwise aflatoxin B₁ or M₁; ^bWith respect to EU MRLs)

In this regard, several HCV proteins, including the core, envelope and nonstructural proteins may be involved in HCC development via oxidative stress, cell cycle control, and tumor formation.

Aflatoxin, HBV and HCC

Studies reveal that South East Asia and Africa show a significant overlap between AFB1 exposure and HBV prevalence [14]. According to a cohort study by Ming et al., even moderate levels of aflatoxin in diet almost tripled the chances of developing HCC in men infected with HBV. Another study reported 30 times greater risk of developing HCC in individuals co-exposed to both HBV and aflatoxin exposure [26]. In comparison to HCV, HBV infection appears to carry a higher risk in causing HCC with a 20-fold lower odds ratio for HCV. A mutation at 249ser-p53 has been identified as a marker to detect concurrent HBV infection and aflatoxin exposure. It was found to be present in 54% of the HBV positive HCC cases in Oidong, China [27]. While the same mutation was found to be absent in 42% matched HCC cases from Beijing, known for relatively low aflatoxin exposure [28,29]. Several epidemiological studies confirm that aflatoxin's cancer causing potency is much greater among HBV-positive than among HBV-negative individuals [23,27,30]. A strong testimony for the synergistic effect of aflatoxin in HBV-associated HCC comes from the fact that AFB1 is associated with a G-to-T transversion in codon 249 of the TP53 gene which changes arginine to serine. This mutation has been seen to interact with HBV X protein in altering cell proliferation and chromosome stability of liver cells [31].

Studies on animals have shown that injury to liver cells inflicted by HBV can increase the amount of cytochrome p450 in liver cells thereby inadvertently increasing aflatoxin metabolism and activation. This could also explain why children acutely infected with HBV show highest adduct levels, followed by chronic carriers as compared to uninfected children [32].

Increased risk of HCC with aflatoxin exposure and Hepatitis C Viral infection

HCV and HCC

An earlier report of the World Health Organization suggested that approximately 170 million people are chronically infected with HCV worldwide [33]. However, considerable regional variations in the prevalence of HCV infection in HCC patients is also reported. In the US, Africa and Europe, the prevalence of anti-HCV positive individuals among HCC cases was 22%, 69% and 45%, respectively. In Asia, majority of HCC cases are reported from Japan with 68% sero prevalence of anti-HCV antibodies. Prevalence of HCV infection in Pakistan is 6.5% with a high proportion (45%) of anti-HCV antibodies also detected among HCC cases [16,34]. While active HCV RNA infections reported in different regions of Pakistan ranged from 3.5-4.9% [35,36].

Aflatoxin, HCV and HCC

Even though aflatoxin and HBV synergism for HCC has been reported, similar synergistic effects of aflatoxin in chronic HCV infection remain to be settled. A conclusive evidence to this effect is hindered by the fact that chronic HCV infection usually occurs later in life and persists for a long time, whereas chronic HBV infection occurs much earlier, thus it is possible that the time of overlapped exposure for aflatoxin and HCV is less significant [14]. An attempt to link HCC prevalence and geographical overlap in populations exposed to aflatoxin, and populations with a high prevalence of HCV, remained inconclusive due to variations encountered in results. For example,

there was more prevalence of aflatoxin, HCV and HBV in Cameroon as compared to Gambia but the latter had higher HCC rates. Perhaps this was because of limited data available on HCV prevalence or other additional risk factors that were unaccounted for [37]. An association between HCV positive patients with Advanced Liver Disease (ALD; liver cirrhosis or HCC) and aflatoxin levels has also been reported. While ALD patients without anti-HCV positivity revealed no significant association with aflatoxin levels. Hence, it can be assumed that ALD does not cause increased aflatoxin activation. However, these studies did not look at dietary aflatoxin levels [18]. Therefore, investigations on associating dietary aflatoxin levels in HCV associated HCC are needed.

In another study in an HCV-endemic country, the rise in HCC cases was strongly linked with food and seeds contaminated with aflatoxin. The consumption of aflatoxin contaminated foods was assessed and correlated with liver biopsy grade and stage scores of chronic HCV patients. It was found that chronic HCV patients who had consumed all seven types of aflatoxin contaminated foods were positively correlated to grade 3 and stage 3 of liver disease, which is a strong precursor for development of HCC [38].

Sun et al., assessed the aflatoxin albumin adduct (AF-alb) levels for anti-HCV positive (n=36) and negative (n=406) subjects in a case control study in Taiwan. They found that anti-HCV positive subjects had less AF-alb detection rates (30.6%) as compared to anti-HCV negative subjects (41.6%). This could possibly be because of low sample size for anti-HCV positive subjects [39].

Generally, during chronic HCV infection about 10-20% of the infected population develops cirrhosis and out of those about 1-4% develop HCC. However, in Asian populations, a much higher percentage (approximately 20%) of chronic HCV infections are resulting in HCC (Figure 2). As increased prevalence of aflatoxin contaminated foods has been reported in these areas [14], this brings us to wonder whether AFB1 could be acting as a co-carcinogen. In Pakistan, rice is a major contributor to the dietary intake of aflatoxins. The total estimated amount of aflatoxin intake for average rice consumers ranged from 19.1 to 26.6 ng/kg body weight/day, which is much higher than the reference value of 1 ng/kg body weight/day [40].


Oncogene activation by HCV, could aflatoxins exacerbate it?

HCV enhances the transcription of c-myc oncogene which may contribute to the development of cancer [41]. The non-structural protein, NS4B is known to transform NIH3T3 cells in association with Ha-ras oncogene [42]. NS4B also causes upregulation of RAPI and FYN oncogenes in HeLa cells [43].

In a study in Taiwan, aflatoxin levels were determined in serum as well as plasma of the same patients and these samples were then tested for p53 mutations and p16 methylation. These two markers are indicative of cancer development, p53 being one of the most frequently mutated genes. A significant correlation between aflatoxin adducts, p53 mutation and p16 methylation was observed [44]. Another study has reported the ability of AFB1 epoxide form to cause mutations in human Ha-ras proto-oncogene in vitro [45].

Reactive Oxygen species (ROS) in HCV and role of aflatoxin as an agent of oxidative stress

During HCV infection, the production of ROS is enhanced as mitochondrial function is impaired, at the same time liver antioxidant system is affected leading to aggravated oxidative stress [46]. The envelope protein, E2, of HCV has been implicated in upregulation of collagen $\alpha(1)$ and oxidative stress in Huh7 cells [47]. Another viral



protein, core protein, has been observed to interact with heat shock protein Hsp60, a stress protein chaperon. This interaction increases ROS production, which upregulates TNF- α -induced apoptotic cell death in Huh7 cells [48].

Exposure to Aflatoxin G1 revealed induction of oxidative stress by increasing ROS generation and causing DNA double-strand breaks which was reduced upon treatment with anti-oxidants [49]. Interestingly, vitamin A, C and E exhibited a protective effect in fresh lymphocytes isolated from human blood by inhibiting ROS produced by AFB1 exposure [50]. As low as 0.05 ng/ml of AFB1 unregulated the production of ROS in polymorphonuclear leukocytes which are at the first line of defense against invading bacterial and fungal infections [51].

Immune system interference by HCV and role of aflatoxins as an immunomodulatory agent

The virus modulates a number of the host cell's defense processes to establish complete infection. Impairment of complement mediated pathway to hinder immune activation and thereby delaying resolution of infection was observed [52]. Down regulation of complement component C9 in patients as well as in cell culture was observed and it was attributed to the core protein of HCV [52]. Using transgenic mouse models, it has been observed that nonstructural protein 5A (NS5A) of HCV impairs antiviral response and weakens interferon immune response along with cytotoxic T lymphocyte response [53].

Numerous studies have attested that AFB1 exposure can affect the immune system in a variety of ways. Secretory Immunoglobulin A in saliva of Gambian children was noted to be lower when exposed to aflatoxin contaminated food [32]. At the same time these mycotoxins have led to increased susceptibility to bacterial and parasitic infections and adversely affect acquired immunity. Human monocytes when exposed to AFB1 in vitro showed reduced microbicidal activity against *Candida albicans* even at low doses of 0.05-0.1 $\mu\text{g}/\text{mL}$. Both decreased phagocytosis and decreased secretion of interleukin-1, interleukin-6 and tumor necrosis factor- α were observed [54].

Patients suffering from HIV were tested for aflatoxin exposure via aflatoxin-albumin adducts and it was observed that those with high aflatoxin-albumin adducts had significantly low T and B cells [55]. Hendrickse et al., observed that aflatoxin contaminated-heroin addicts showed rapid progression of human immunodeficiency virus and acquired immune deficiency syndromes [56]. They attributed

this accelerated rate of HIV progression to aflatoxin related immune suppression. This same synergy can be used to explain the faster rate of HIV progression in Africa when compared to Europe or the United States where aflatoxin exposure is comparatively controlled. Therefore, it is quite possible that accelerated rate of cancer development may be observed in those HCV patients who are consuming aflatoxin contaminated foods.

Conclusions

In order to improve food quality and to lower the aflatoxin levels in food items various practices need to be adopted including effective monitoring mechanisms from produce to products, fungal-resistant crops. At the consumer level, a change from aflatoxins-susceptible diet in highly endemic areas is another possibility. For example, recent reforms in China included changing the staple diet from maize to rice in the 1980s which actually led to reduced aflatoxin levels in sera and ultimately lower primary liver cancer cases [57]. Farmers can opt for crops that are resistant to the fungal infection or introduce fungal strains that are inefficient in producing aflatoxins. Irrigation techniques need to be improved along with drying and storage practices in order to keep the moisture content to a minimum. Testing of crops for aflatoxin contamination needs to be mandatory and cheap methods of doing so are needed. If crops have been contaminated then measures need to be introduced for decontamination and detoxification. These include food and feed processing, biochemical and microbial inactivation, chemical degradation and reduction in toxin bioavailability by selective chemisorption with clays. Physical methods may include thermal heating but due to toxin being heat stable, this has not been successful. Irradiation through radiolysis of water has shown some prospect. A last resort is always physical separation of contaminated parts by manual, mechanical or electronic means [38]. For example, in a study conducted in China, oltipraz, an antischistosomal drug, was administered to 234 healthy adults. It was noted that levels of aflatoxin m1, which is a phase 1 metabolites of AFB1, were reduced by 51% in urine samples of individuals receiving a high dose of oltipraz [58]. Inclusion of vitamins like A, C and E in the daily diet could also help nullify the toxic effects of aflatoxin exposure to some extent [50]. Chlorophyllin, which is a water-soluble derivative of chlorophyll, is reported to be anti-mutagenic. When chlorophyllin and aflatoxin were co-administered, this natural product absorbed the toxin before it could reach the liver and reduced AFB1-DNA adduct formation and hepatocarcinogenesis [59,60]. Although the aforementioned preventive measures for lowering aflatoxin levels in foods is necessary but until that becomes a reality, perhaps these drugs or vitamins could be introduced in areas where aflatoxin consumption is unavoidable in order to prevent the development of HCC.

Conflict of Interest

HQ and SSA mined the literature. MI and AAS provided gathered all data related with aflatoxins. NAK and SSH wrote the manuscript. HQ and AAS added and improved the overall manuscript structure and contents. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by the Aga Khan University.

References

1. IARC (2014) WHO. Press Release N 244.
2. IARC. GLOBOCAN (2012) Estimated cancer incidence, mortality and prevalence worldwide in 2012. International Agency for Research on Cancer, World Health Organization. Press Release N 223.

3. Parkin DM, Bray F, Ferlay J, Pisani P (2005) Global cancer statistics, 2002. *J Clin* 55: 74-108.
4. Lang K, Danchenko N, Gondek K, Shah S, Thompson D (2009) The burden of illness associated with hepatocellular carcinoma in the United States. *J Hepatol* 50: 89-99.
5. Yuen MF, Hou JL, Chutaputti A (2009) Asia Pacific Working Party on Prevention of Hepatocellular Carcinoma (2009) Hepatocellular carcinoma in the Asia Pacific region. *J Gastroenterol Hepatol* 24: 346-353.
6. Somi MH (2005) Hepatocellular Carcinoma. *Hepat Mon* 5: 65-67.
7. Butt AS, Abbas Z, Jafri W (2012) Hepatocellular carcinoma in Pakistan: where do we stand? *Hepat Mon* 12: 6023.
8. Schütte K, Bornschein J, Malfertheiner P (2009) Hepatocellular carcinoma--epidemiological trends and risk factors. *Dig Dis* 27: 80-92.
9. Ohishi W, Fujiwara S, Cologne JB, Suzuki G, Akahoshi M, et al. (2008) Risk factors for hepatocellular carcinoma in a Japanese population: a nested case-control study. *Cancer Epidemiol Biomarkers Prev* 17: 846-854.
10. Shariff MI, Cox IJ, Gomaa AI, Khan SA, Gedroyc W, et al. (2009) Hepatocellular carcinoma: current trends in worldwide epidemiology, risk factors, diagnosis and therapeutics. *Expert Rev Gastroenterol Hepatol* 3: 353-367.
11. Bbosa GS, Kitya D, Lubega A, Ogwal-Okeng J, Anokbonggo WW, et al. (2013) Review of the Biological and Health Effects of Aflatoxins on Body Organs and Body Systems. Review of the Biological and Health Effects of Aflatoxins on Body Organs and Body Systems.
12. Williams JH, Phillips TD, Jolly PE, Stiles JK, Jolly CM, et al. (2004) Human aflatoxicosis in developing countries: a review of toxicology, exposure, potential health consequences, and interventions. *Am J Clin Nutr* 80: 1106-1122.
13. Wild CP, Gong YY (2010) Mycotoxins and human disease: a largely ignored global health issue. *Carcinogenesis* 31: 71-82.
14. Liu Y, Wu F (2010) Global burden of aflatoxin-induced hepatocellular carcinoma: a risk assessment. *Environ Health Perspect* 118: 818-824.
15. Munir MA, Saleem M, Malik ZR, Ahmed M, Ali A (1989) Incidence of aflatoxin contamination in non-perishable food commodities. *J Pak Med Assoc* 39: 154-157.
16. Raza SA, Clifford GM, Franceschi S (2007) Worldwide variation in the relative importance of hepatitis B and hepatitis C viruses in hepatocellular carcinoma: a systematic review. *Br J Cancer* 96: 1127-1134.
17. Zaryabova V, Shalamanova T, Israel M (2013) Pilot study of extremely low frequency magnetic fields emitted by transformers in dwellings. *Social aspects. Electromagn Biol Med* 32: 209-217.
18. Chen CH, Wang MH, Wang JH, Hung CH, Hu TH, et al. (2007) Aflatoxin exposure and hepatitis C virus in advanced liver disease in a hepatitis C virus endemic area in Taiwan. *Am J Trop Med Hyg* 77: 747-752.
19. Sheabar FZ, Groopman JD, Qian GS, Wogan GN (1993) Quantitative analysis of aflatoxin-albumin adducts. *Carcinogenesis* 14: 1203-1208.
20. Wild CP, Hudson GJ, Sabbioni G, Chapot B, Hall AJ, et al. (1992) Dietary intake of aflatoxins and the level of albumin-bound aflatoxin in peripheral blood in The Gambia, West Africa. *Cancer Epidemiol Biomarkers Prev* 1: 229-234.
21. Gouas D, Shi H, Hainaut P (2009) The aflatoxin-induced TP53 mutation at codon 249 (R249S): biomarker of exposure, early detection and target for therapy. *Cancer Lett* 286: 29-37.
22. Bedard LL, Massey TE (2006) Aflatoxin B1-induced DNA damage and its repair. *Cancer Lett* 241: 174-183.
23. Qian GS, Ross RK, Yu MC (1994) A follow-up study of urinary markers of aflatoxin exposure and liver cancer risk in Shanghai, People's Republic of China. *Cancer Epidemiol Biomarkers Prev* 3:3-10.
24. WHO Hepatitis B Vaccines (2004) *Weekly Epidemiology Records* 49:425-427.
25. Fung J, Lai CL, Yuen MF (2009) Hepatitis B and C virus-related carcinogenesis. *Clin Microbiol Infect* 15: 964-970.
26. Groopman JD, Kensler TW, Wild CP (2008) Protective interventions to prevent aflatoxin-induced carcinogenesis in developing countries. *Annu Rev Public Health* 29: 187-203.
27. Kirk GD, Camus-Randon AM, Mendy M, Goedert JJ, Merle P, et al. (2000) Ser-249 p53 mutations in plasma DNA of patients with hepatocellular carcinoma from The Gambia. *J Natl Cancer Inst* 92: 148-153.
28. Kew MC (2003) Synergistic interaction between aflatoxin B1 and hepatitis B virus in hepatocarcinogenesis. *Liver Int* 23: 405-409.
29. Ming L, Thorgeirsson SS, Gail MH, Lu P, Harris CC, et al. (2002) Dominant role of hepatitis B virus and cofactor role of aflatoxin in hepatocarcinogenesis in Qidong, China. *Hepatology* 36: 1214-1220.
30. Ok HE, Kim HJ, Shim WB, Lee H, Bae DH, et al. (2007) Natural occurrence of aflatoxin B1 in marketed foods and risk estimates of dietary exposure in Koreans. *J Food Prot* 70: 2824-2828.
31. Jiang W, Wang XW, Unger T (2010) Cooperation of tumor-derived HBx mutants and p53-249(ser) mutant in regulating cell proliferation, anchorage-independent growth and aneuploidy in a telomerase-immortalized normal human hepatocyte-derived cell line. *Int J Cancer* 127:1011-1020.
32. Turner PC, Moore SE, Hall AJ, Prentice AM, Wild CP (2003) Modification of immune function through exposure to dietary aflatoxin in Gambian children. *Environ Health Perspect* 111: 217-220.
33. [No authors listed] (1999) Global surveillance and control of hepatitis C. Report of a WHO Consultation organized in collaboration with the Viral Hepatitis Prevention Board, Antwerp, Belgium. *J Viral Hepat* 6: 35-47.
34. Hamid S, Umar M, Alam A, Siddiqui A, Qureshi H, et al. (2004) PSG consensus statement on management of hepatitis C virus infection--2003. *J Pak Med Assoc* 54: 146-150.
35. Ali A, Ahmad H, Ali I, Khan S, Zaidi G, et al. (2010) Prevalence of active hepatitis C virus infection in district Mansehra Pakistan. *Virology* 7: 334.
36. Anwar MI, Rahman M, Hassan MU, Iqbal M (2013) Prevalence of active hepatitis C virus infections among general public of Lahore, Pakistan. *Virology* 10: 351.
37. Palliyaguru DL, Wu F (2013) Global geographical overlap of aflatoxin and hepatitis C: controlling risk factors for liver cancer worldwide. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* 30: 534-540.
38. Hifnawy M, Mangoud AM, Eissa MH, Nor Edin E, Mostafa Y, et al. (2004) The role of aflatoxin-contaminated food materials and HCV in developing hepatocellular carcinoma in Al-Sharkia Governorate, Egypt. *J Egypt Soc Parasitol* 34: 479-488.
39. Sun CA, Wu DM, Wang LY, Chen CJ, You SL, et al. (2002) Determinants of formation of aflatoxin-albumin adducts: a seven-township study in Taiwan. *Br J Cancer* 87: 966-970.
40. Iqbal SZ, Asi MR, Ariño A, Akram N, Zuber M (2012) Aflatoxin contamination in different fractions of rice from Pakistan and estimation of dietary intakes. *Mycotoxin Res* 28: 175-180.
41. Higgs MR, Lerat H, Pawlowsky JM (2013) Hepatitis C virus-induced activation of β -catenin promotes c-Myc expression and a cascade of pro-carcinogenic events. *Oncogene* 32: 4683-4693.
42. Park JS, Yang JM, Min MK (2000) Hepatitis C virus nonstructural protein NS4B transforms NIH3T3 cells in cooperation with the Ha-ras oncogene. *Biochem Biophys Res Commun* 267: 581-587.
43. Zheng Y, Ye LB, Liu J, Jing W, Timani KA, et al. (2005) Gene expression profiles of HeLa Cells impacted by hepatitis C virus non-structural protein NS4B. *J Biochem Mol Biol* 38: 151-160.
44. Zhang YJ, Rossner P Jr, Chen Y, Agrawal M, Wang Q, et al. (2006) Aflatoxin B1 and polycyclic aromatic hydrocarbon adducts, p53 mutations and p16 methylation in liver tissue and plasma of hepatocellular carcinoma patients. *Int J Cancer* 119: 985-991.
45. Riley J, Mandel HG, Sinha S, Judah DJ, Neal GE (1997) In vitro activation of the human Harvey-ras proto-oncogene by aflatoxin B1. *Carcinogenesis* 18: 905-910.
46. Fujinaga H, Tsutsumi T, Yotsuyanagi H, Moriya K, Koike K (2011) Hepatocarcinogenesis in hepatitis C: HCV shrewdly exacerbates oxidative stress by modulating both production and scavenging of reactive oxygen species. *Oncology* 81 Suppl 1: 11-17.
47. Ming-Ju H, Yih-Shou H, Tzy-Yen C, Hui-Ling C (2011) Hepatitis C virus E2 protein induce reactive oxygen species (ROS)-related fibrogenesis in the HSC-T6 hepatic stellate cell line. *J Cell Biochem* 112: 233-243.

48. Kang SM, Kim SJ, Kim JH, Lee W, Kim GW, et al. (2009) Interaction of hepatitis C virus core protein with Hsp60 triggers the production of reactive oxygen species and enhances TNF- α -mediated apoptosis. *Cancer Lett* 279: 230-237.
49. Shen H, Liu J, Wang Y, Lian H, Wang J, et al. (2013) Aflatoxin G1-induced oxidative stress causes DNA damage and triggers apoptosis through MAPK signaling pathway in A549 cells. *Food Chem Toxicol* 62: 661-669.
50. Alpsoy L, Yildirim A, Agar G (2009) The antioxidant effects of vitamin A, C, and E on aflatoxin B1-induced oxidative stress in human lymphocytes. *Toxicol Ind Health* 25: 121-127.
51. Ubagai T, Tansho S, Ito T, Ono Y (2008) Influences of aflatoxin B1 on reactive oxygen species generation and chemotaxis of human polymorphonuclear leukocytes. *Toxicol In Vitro* 22: 1115-1120.
52. Kim H, Meyer K, Di Bisceglie AM, Ray R (2013) Hepatitis C virus suppresses C9 complement synthesis and impairs membrane attack complex function. *J Virol* 87: 5858-5867.
53. Kriegs M, Bürckstümmer T, Himmelsbach K, Bruns M, Frelin L, et al. (2009) The hepatitis C virus non-structural NS5A protein impairs both the innate and adaptive hepatic immune response in vivo. *J Biol Chem* 284: 28343-28351.
54. Cusumano V, Rossano F, Merendino RA, Arena A, Costa GB, et al. (1996) Immunobiological activities of mould products: functional impairment of human monocytes exposed to aflatoxin B1. *Res Microbiol* 147: 385-391.
55. Jiang Y, Jolly PE, Preko P, Wang JS, Ellis WO, et al. (2008) Aflatoxin-related immune dysfunction in health and in human immunodeficiency virus disease. *Clin Dev Immunol*.
56. Hendrickse RG, Maxwell SM, Young R (1989) Aflatoxins and heroin. *BMJ* 299: 492-493.
57. Chen JG, Egner PA, Ng D, Jacobson LP, Muñoz A, et al. (2013) Reduced aflatoxin exposure presages decline in liver cancer mortality in an endemic region of China. *Cancer Prev Res (Phila)* 6: 1038-1045.
58. Wang JS, Shen X, He X, Zhu YR, Zhang BC, et al. (1999) Protective alterations in phase 1 and 2 metabolism of aflatoxin B1 by oltipraz in residents of Qidong, People's Republic of China. *J Natl Cancer Inst* 91: 347-354.
59. Dashwood R, Negishi T, Hayatsu H, Breinholt V, Hendricks J, et al. (1998) Chemopreventive properties of chlorophylls towards aflatoxin B1: a review of the antimutagenicity and anticarcinogenicity data in rainbow trout. *Mutat Res* 399: 245-253.
60. Jubert C, Mata J, Bench G, Dashwood R, Pereira C, et al. (2009) Effects of chlorophyll and chlorophyllin on low-dose aflatoxin B(1) pharmacokinetics in human volunteers. *Cancer Prev Res (Phila)* 2: 1015-1022.
61. Butt AS, Hamid S, Wadalawala AA (2013) Hepatocellular carcinoma in Native South Asian Pakistani population; trends, clinico-pathological characteristics & differences in viral marker negative & viral-hepatocellular carcinoma. *BMC Res Notes* 6:137.
62. Abbas Z (2013) Hepatocellular carcinoma in Pakistan. *J Coll Physicians Surg Pak* 23: 769-770.
63. Soubra L, Sarkis D, Hilan C, Verger P (2009) Occurrence of total aflatoxins, ochratoxin A and deoxynivalenol in foodstuffs available on the Lebanese market and their impact on dietary exposure of children and teenagers in Beirut. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* 26: 189-200.
64. Mohd-Redzwan S, Jamaluddin R, Abd-Mutalib MS, Ahmad Z (2013) A mini review on aflatoxin exposure in Malaysia: past, present and future. *Front Microbiol* 4: 334.
65. Yazdanpanah H1, Zarghi A, Shafaati AR, Foroutan SM, Aboul-Fathi F, et al. (2013) Analysis of aflatoxin b1 in Iranian foods using HPLC and a monolithic column and estimation of its dietary intake. *Iran J Pharm Res* 12: 83-89.
66. Fakoor Janati S, Beheshti HR, Khoshbakht Fahim N, Feizy J (2011) Aflatoxins and ochratoxinin A in bean from Iran. *Bull Environ Contam Toxicol* 87: 194-197.
67. Wu F, Stacy SL, Kensler TW (2013) Global risk assessment of aflatoxins in maize and peanuts: are regulatory standards adequately protective? *Toxicol Sci* 135: 251-259.
68. Majeed S, Iqbal M, Asi MR, Iqbal SZ (2013) Aflatoxins and Ochratoxin A contamination in rice, corn and corn products from Punjab, Pakistan. *J Cereal Sci* 58: 446-450.
69. Paterson RRM (2007) Aflatoxins contamination in chilli samples from Pakistan. *Food Control* 18: 817-82
70. Shamsuddin ZA, Khan MA, Khan BA, Ahmad MA, Ahmed A (1995) Contamination of red chilli with aflatoxin b1 in pakistan. *Mycotoxin Res* 11: 21-24.
71. Razzazi-Fazeli E, Noviandi CT, Porasuphatana S, Agus A, Böhm J (2004) A survey of aflatoxin B1 and total aflatoxin contamination in baby food, peanut and corn products sold at retail in Indonesia analysed by ELISA and HPLC. *Mycotoxin Res* 20: 51-58.
72. Ali N, Sardjono, Yamashita A, Yoshizawa T (1998) Natural co-occurrence of aflatoxins and Fusarium mycotoxins (fumonisins, deoxynivalenol, nivalenol and zearalenone) in corn from Indonesia. *Food Addit Contam* 15: 377-384.
73. Yamashita A, Yoshizawa T, Aiura Y, Sanchez PC, Dizon EI, et al. (1995) Fusarium mycotoxins (fumonisins, nivalenol, and zearalenone) and aflatoxins in corn from Southeast Asia. *Biosci Biotechnol Biochem* 59: 1804-1807.
74. Siruguri V, Kumar PU, Raghu P, Rao MV, Sesikeran B, et al. (2012) Aflatoxin contamination in stored rice variety PAU 201 collected from Punjab, India. *Indian J Med Res* 136: 89-97.
75. Tabata S, Kamimura H, Ibe A, Hashimoto H, Iida M, et al. (1993) Aflatoxin contamination in foods and foodstuffs in Tokyo: 1986-1990. *J AOAC Int* 76: 32-35.
76. Sadia A, Jabbar MA, Deng Yet, Hussain AE, Riffat S, al (2012) A survey of aflatoxin M1 in milk and sweets of Punjab, Pakistan. *Food Control* 26: 235-240.
77. Iqbal SZ, Asi MR (2013) Assessment of Aflatoxin M1 in Milk and Milk Products from Punjab, Pakistan. *Food Control* 30:235-239.
78. Hussain I, Anwar J, Asi MR, Munawar MA, Kashif M (2010) Aflatoxin M1 contamination in milk from five dairy species in Pakistan. *Food Control* 21:122-124.
79. Iqbal SZ, Paterson RRM, Bhatti IA, Asi MR (2011) Aflatoxin concentrations in chillies vary depending on variety. *Mycoscience* 52:296-299.
80. Shah HU, Simpson TJ, Alam S, Khattak KF, Perveen S (2010) Mould incidence and mycotoxin contamination in maize kernels from Swat Valley, North West Frontier Province of Pakistan. *Food Chem Toxicol* 48: 1111-1116.
81. Iqbal SZ, Asi MR, Jinab S (2013) Natural occurrence of aflatoxin B1 and aflatoxin M1 in "halva" and its ingredients. *Food Control* 34:404-407.
82. Iqbal SZ, Asi MR, Zuber M, Akhtar J, Saif MJ (2013) Natural occurrence of aflatoxins and ochratoxin A in commercial chilli and chilli sauce samples. *Food Control* 30: 621-625.
83. Lutfullah G, Hussain A (2011) Studies on contamination level of aflatoxins in some dried fruits and nuts of Pakistan. *Food Control* 22: 426-429.
84. Iqbal SZ, Asi MR, Jinab S (2013) Variation of aflatoxin M1 contamination in milk and milk products collected during winter and summer seasons. *Food Control* 34: 714-718.

Citation: Qureshi H, Ali SS, Iqbal M, Siddiqui AA, Khan NA, et al. (2014) Aflatoxins and Hepatitis B, C Viral Associated Hepatocarcinogenesis. *J Cell Sci Ther* 5: 179. doi:[10.4172/2157-7013.1000179](https://doi.org/10.4172/2157-7013.1000179)

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 350 Open Access Journals
- 30,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.editorialmanager.com/lifesciences>