



THE AGA KHAN UNIVERSITY

eCommons@AKU

Department of Emergency Medicine

Medical College, Pakistan

June 2013

A comparison of WHO guidelines issued in 1997 and 2009 for dengue fever - single centre experience.

Munawar Khursheed

Aga Khan University, munawar.khursheed@aku.edu

Uzma R. Khan

Aga Khan University, uzma.khan@aku.edu

Kiran Ejaz

Aga Khan University

Jabeen Fayyaz

Aga Khan University

Irum Qamar

Aga Khan University

See next page for additional authors

Follow this and additional works at: http://ecommons.aku.edu/pakistan_fhs_mc_emerg_med



Part of the [Infectious Disease Commons](#), and the [Parasitic Diseases Commons](#)

Recommended Citation

Khursheed, M., Khan, U., Ejaz, K., Fayyaz, J., Qamar, I., Razzak, J. (2013). A comparison of WHO guidelines issued in 1997 and 2009 for dengue fever - single centre experience.. *Journal of Pakistan Medical Association*, 63(6), 670-674.

Available at: http://ecommons.aku.edu/pakistan_fhs_mc_emerg_med/57

Authors

Munawar Khursheed, Uzma R. Khan, Kiran Ejaz, Jabeen Fayyaz, Irum Qamar, and J. Razzak

A comparison of WHO guidelines issued in 1997 and 2009 for dengue fever — single centre experience

Munawar Khurshheed, Uzma Rahim Khan, Kiran Ejaz, Jabeen Fayyaz, Irum Qamar, Junaid Abdul Razzak

Abstract

Objective: To compare the original (1997) and revised (2009) versions of World Health Organization guidelines for dengue patients

Methods: Adult patients with a positive dengue Immunoglobulin M serology, and a diagnosis of dengue were included in the study at Aga Khan University Hospital during a three-year period from January 2005 to December 2007. Data related to these dengue patients was collected from their medical records. Guidelines were then applied by the research assistant and correlation among these guidelines was computed. SPSS 19 was used for statistical analysis.

Results: A total of 612 patients were found with a diagnosis of dengue, but only 439 (71.73%) had a positive IgM. The median age of these 439 patients was 28 (interquartile range: 18) years and majority of them were males, 295 (67%). According to the 1997 guidelines, 383 (87%) patients were classified as having dengue, while according to the 2009 guidelines, all the 439 (100%) patients were classified with a dengue infection. Under WHO 1997, 21 (5.5%) cases were classified as dengue shock syndrome, while 2009 guidelines labelled 88 (20%) cases as severe dengue. There was a consensus on only 11 severe cases by both the guidelines, showing different results between the two.

Conclusion: By using 2009 guidelines, a physician would classify more dengue patients as having severe disease.

Keywords: Pakistan, Dengue, WHO guidelines. (JPMA 63: 670; 2013)

Introduction

Dengue infection is caused by Flavivirus and spreads through *Aedes aegypti* mosquito. The virus infects over 50 million people worldwide, resulting in over 24,000 deaths annually.^{1,2} Billions of people remain exposed to the disease across Africa, Eastern Mediterranean, Southeast Asia and Western Pacific region.³

Clinical presentation of patients with dengue infection varies from a self-limiting, non-specific acute febrile illness to a syndrome characterised by bleeding, severe intravascular volume depletion and shock.⁴ Given its vague presentation, early identification of severe infections can be challenging, causing delays in the institution of life-saving interventions. To assist clinicians in making triage decisions, the World Health Organisation (WHO) published a dengue infection triage and treatment guideline in 1997.⁵ It was based on the data from paediatric population of Bangkok and divided the clinical syndrome into dengue fever (DF), dengue haemorrhagic fever (DHF) and dengue shock syndrome (DSS).⁶⁻⁸ A revised version of the guideline was published in 2009 to address the low sensitivity in high-risk patients and was

more inclusive of adult patients as well.^{2,9,10} The new classification, while retaining the three-level severity grading, divided the infection into dengue without warning signs, dengue with warning signs and severe dengue.^{2,6,10}

Pakistan has experienced a major epidemic of dengue infection since 2005.¹¹ Hospitals, especially the emergency departments (ED), have seen a major surge in the dengue patient volume. In the beginning of the epidemic, triage decisions were based on the clinician's judgment, while WHO guidelines were slowly being accepted as the decision-making tool. Recently published studies from other settings have shown better accuracy of the revised guidelines.^{2,12,13} This study was carried out to compare the performance of 1997 and 2009 WHO guidelines based on the dengue Immunoglobulin (IgM) results in our setting.^{2,5}

Patients and Methods

This was a case series of Dengue patients recruited by retrospective chart review conducted at the Aga Khan University Hospital (AKUH) Karachi, Pakistan. All adult patients with a diagnosis of dengue who had a positive dengue IgM serology between January 2005 and December 2007 were included in the study. Cases with incomplete information such as signs, symptoms and

.....
Department of Emergency Medicine, Aga Khan University, Karachi, Pakistan.

Correspondence: Munawar Khurshheed. Email: munawar.khurshheed@aku.edu

laboratory figures were excluded. By using a standard tool, trained research assistants extracted data regarding clinical features, laboratory investigations, on-admission diagnosis in ED and diagnosis of hospital with serological test (IgM anti-dengue). In our current clinical setting, we send dengue IgM test on all suspicious patients. Anti-IgM becomes positive on the fifth day of dengue illness and can remain positive for 90 days.

No patient had polymerase chain reaction (PCR) test or culture of the virus. Each patient was classified/graded both according to 1997 (Dengue Fever, Dengue Haemorrhagic Fever and Dengue Shock Syndrome) and 2009 guidelines (dengue without warning signs, dengue with warning signs and severe dengue) (Table-1, Figure-1) by the research assistant using the standard definitions. The three categories of dengue used in each of the

Table-1: WHO guidelines (1999) for the treatment of dengue fever/ dengue hemorrhagic fever.

DF/DHF	Grade*	Symptoms	Laboratory picture
DF		Fever with two or more of the following signs: headache, retro-orbital pain, myalgia, arthralgia	Leukopenia occasionally. Thrombocytopenia may be present. No evidence of plasma loss
DHF	I	Above signs plus positive tourniquet test	Thrombocytopenia positive tourniquet < 100,000, Hct rise ≥ 20%
DHF	II	Above signs plus spontaneous bleeding	Thrombocytopenia < 100,000, Hct rise ≥ 20%
DHF	III	Above signs plus circulatory failure (weak pulse, hypotension restlessness)	Thrombocytopenia < 100,000, Hct rise > 20%
DHF	IV	Profound shock with undetectable blood pressure and pulse	Thrombocytopenia < 100,000, Hct rise ≥ 20%

DHF Grade III and IV are also called as Dengue Shock Syndrome (DSS).




Figure-1: Dengue, guidelines for diagnosis, treatment, prevention and control new edition 2009.

guidelines were referred to as minor, moderate and severe. Minor dengue was used for DF (1997 guidelines) and dengue without warning signs (2009 guidelines); moderate dengue for DHF (1997) and dengue with warning signs (2009); and severe dengue for DSS (1997) and severe dengue (2009).

Descriptive analysis was done using SPSS, version 19.0 and frequencies and percentages were calculated. Median with interquartile range (IQR) was reported for skewed data such as age and platelet count, while mean \pm standard deviation (SD) was used for normal data. Ethical approval was obtained from the institutional ethics review committee.

Results

A total of 612 patients were located with a diagnosis of dengue out of which 439 (71.73%) had a positive IgM. According to 1997 guidelines, 383 (87%) patients were classified as having dengue. On the other hand, 2009 guidelines classified 439 (100%) patients as dengue infected (Figure-2). Of the 56 (13%) cases which could not be labelled as dengue using the 1997 guidelines, 30 (54%) were classified as probable dengue without warning signs, 19 (34%) were classified as dengue with warning signs, and 7 (12%) as severe dengue using the 2009 guidelines.

The median age of the participants was 28 (IQR 18) years; 295 (67%) were males. All patients were discharged from the hospital except one patient who died in the hospital. Only 167 (38%) patients had three or more symptoms on history. Most (n=248; 56%) had no signs on examination (Table-2). All the patients presented with fever (100%), vomiting 281 (64%) and body ache 173 (39%). Rash (27%), petechiae (10%) and purpura (1%) were present in less number of patients.

Table-3 shows the comparison of both guidelines. The two guidelines classified approximately 50% of cases similarly as minor, moderate and severe dengue (Table-2). WHO 1997 had classified 21 (5.5%) cases as DSS while 2009 guidelines labelled 81 (21%) cases as severe dengue, with consensus on only 11 (52.4%) severe cases by both

Table-2: Clinical features of study population (n = 439).

Clinical Symptoms	n(%)
Fever	439 (100)
Vomiting	281(64)
Body ache	173(39)
Abdominal pain	152(35)
Bleeding other than nose and gum	75(17)
Headache	35(8)
Gum Bleed	32(7)
Nose Bleed	26(5)
Haematemesis	22(5)
Drowsiness	21(5)
Shortness of Breath	14(3)
Examination Findings	n(%)
Rash	120(27)
Ascites	44(10)
Petechie	45(10)
Hepatomegaly	19(4)
Pulmonary Effusion	16(4)
Splenomegaly	9(2)
Purpura	6(1)
Vital Examination Readings	Mean \pm SD
Temperature(oC)	38 \pm 3
Systolic Blood Pressure(mm Hg)	112 \pm 17
Diastolic Blood Pressure(mm Hg)	72 \pm 12
Pulse Pressure(beats/min)	40 \pm 14
Laboratory Parameters	Mean \pm SD
White Blood Count(x109/L)(n=431)	5 \pm 4
Haematocrit(gm/dl)(n=426)	41 \pm 7
Prothrombin Time(secs)(n=362)	13 \pm 8
International Normalized Ratio(n=359)	1 \pm 2
Activated Partial Thromboplastin Time(secs)(n=367)	37 \pm 20
Sodium(mEq/L)(n=406)	138 \pm 75
Bicarbonate(mEq/L)(n=332)	22 \pm 5
	Median
(IQR)	
Platelet(x109/L)(n=427)	45(67)
Blood Urea Nitrogen(mg/dL)(n=368)	9(6)
Creatinine(mg/dL)(n=409)	1(0.4)
Serum Glutamate-Oxaloacetate Transaminase(units/L)(n=388)	174(274)

the guidelines. The alarming result was that more than a quarter cases (29%) that were classified as moderate by the 1997 guideline were severe dengue according to the 2009 guidelines.

Table-3: Comparison of WHO guidelines for dengue cases (n=383).

Disease Severity as per 1997 Classification	Dengue Fever n(%)	Disease Severity as per 2009 Classification		Total n(%)
		Dengue Haemorrhagic Fever n(%)	Dengue Shock Syndrome n(%)	
Probable Dengue without warning signs	68 (45.3)	48(22.6)	4(19)	120(31.3)
Probable Dengue with warning signs	73(48.7)	103(48.6)	6(28.6)	182(45.7)
Severe dengue	9(6)	61(28.8)	11(52.4)	81(21.1)
Total	150(100)	212(100)	21(100)	383(100)




Figure-2: Flow chart of dengue patients recruitment (January 2005- December 2007).

Discussion

We not only found a difference between 1997 and 2009 guidelines in the accuracy of dengue diagnosis, but also in the assessment of the severity of the disease.^{2,5} There was a four-fold increase in the diagnosis of severe form of dengue using 2009 guidelines compared to the 1997 guidelines.

The diagnostic challenge is likely to be higher in our setting where other febrile illnesses are more common. Diseases such as typhoid fever or malaria contribute to a major bulk of acute clinical practice in Pakistan and, like dengue, they often present with non-specific fever, few signs or symptoms and a blood picture showing low white blood cell (WBC) and platelet counts.^{14,15} Atypical presentations such as abdominal pain, vomiting, diarrhoea, cough and headache resembling enteric fever have been reported in other studies from Pakistan, making clinical decision more difficult.¹⁶ In such endemic settings, concurrent infections such as dengue and malaria could mislead physician's initial impression. Coexistence of malaria and dengue have been reported to be in the range of 20% to as high as 80%.^{17,18} While malaria testing is more widely available, diagnostic tests for dengue (IgM) are either not available or not able to

detect dengue in the first few days of the disease onset, biasing the clinical diagnosis towards malaria.

Patients in our setting tend to start the use of antibiotic and other medicines either in consultation with a general physician or by themselves. In Pakistan, there are no strict prescription regulations and drugs are easily available over the counter. Patients visit the hospital only when their condition deteriorates, and this makes correct diagnosis challenging for the physicians.

Time also affects accuracy of diagnosis even when WHO guidelines are used. Leo et al reported an increase in the accuracy of diagnosis by the WHO guidelines from 14% to 32% and 61% to 79% from day 1 to day 7 of admission using the 1997 and 2009 guidelines respectively.¹³ A study from Karachi, showed that almost one-fourth of children with final diagnosis of DF were initially labelled as undifferentiated fever.¹⁹ Similarly, in Vietnam about a third of cases with DF were initially diagnosed as acute undifferentiated fever.²⁰

Revised guidelines were found to be better in diagnosing dengue cases overall, particularly those with severe form of illness. This is corroborated by work done in other settings, both among adults and children.^{12,21} DENCO, a

multi-country prospective study, found comparable results where 15% of patients with clinical shock were not correctly classified by 1997 classification as severe cases.²² One reason is, perhaps, paediatric focus of the 1997 guidelines which limited its application to adult population.^{7,9,23} The 2009 classification included more non-specific alarming signs such as mental status changes, abdominal pains and involvement of other organs such as liver.

The current study has several limitations. First, it is based on the retrospective data obtained from medical records. Not all features, for example, tourniquet test, were universally captured in the medical records. Second, we could not include dengue patients who were misdiagnosed and sent home and also those who were sent home because they had mild disease.²⁴ Third, we did not know if the individual physician applied the guidelines or was even aware of the guidelines.

Conclusion

By using the WHO guidelines 2009, a physician would end up classifying more dengue patients as having moderate or severe disease compared to the 1997 guideline. Widespread use of these guidelines needs to be encouraged among physicians.

Acknowledgements

We are thankful to Mr. Iqbal Azam for statistical help and Dr. Rehan Nasir and Dr. Munira Abdul Aziz for helping in data collection. Authors URK and JAR were partially supported through the "Johns Hopkins-Pakistan International Collaborative Trauma and Injury Research Training Program", Grant Number D43-TW007-292 from the Fogarty International Center of the United States National Institutes of Health.

References

- Gibbons RV, Vaughn DW. Dengue: an escalating problem. *BMJ* 2002; 324: 1563-6.
- TDR, WHO. Dengue: guidelines for diagnosis, treatment, prevention and control. New Edition. Geneva: WHO. 2009.
- Singhasivanon P, Jacobson J. Dengue is a major global health problem. Foreword. *J Clin Virol* 2009; 46 suppl 2: S1-2.
- Guzman MG, Vázquez S, Kouri G. Dengue: where are we today? *Malays J Med Sci* 2009; 16: 4-11.
- Dengue haemorrhagic fever: diagnosis, treatment, prevention and control. 2nd edition. Geneva: WHO . 1997.
- World Health Organization. Guidelines for treatment of dengue fever / dengue hemorrhagic fever in small hospitals WHO regional office for South-East Asia, New Dehli. *Dengue Bulletin* Volume 23. 1999. 1-23.
- Gupta P, Khare V, Tripathi S, Nag VL, Kumar R, Khan MY, et al. Assessment of World Health Organization definition of dengue hemorrhagic fever in North India. *J Infect Dev Ctries* 2010; 4: 150-5..
- Bandyopadhyay S, Lum LC, Kroeger A. Classifying dengue: a review of the difficulties in using the WHO case classification for dengue haemorrhagic fever. *Trop Med Int Health* 2006; 11: 1238-55.
- Rigau-Pérez JG. Severe dengue: the need for new case definitions. *Lancet Infect Dis* 2006; 6: 297-302.
- Research SPf, Diseases TIT, Diseases WDoCoNT, Epidemic WHO, Alert P. Dengue: Guidelines for diagnosis, treatment, prevention and control World Health Organization 2009.
- Khan E, Kisat M, Khan N, Nasir A, Ayub S, Hasan R. Demographic and clinical features of dengue fever in Pakistan from 2003-2007: a retrospective cross-sectional study. *PLoS One*. 2010; 5: e12505. Published Online First: 13 September 2010. doi: 10.1371/journal.pone.0012505 (accessed on ...)
- Basuki PS, Budiyo, Puspitasari D, Husada D, Darmowandowo W, Ismoedijanto, et al. Application of revised dengue classification criteria as a severity marker of dengue viral infection in Indonesia. *Southeast Asian J Trop Med Public Health* 2010; 41: 1088-94. .
- Leo YS, Thein TL, Fisher DA, Low JG, Oh HM, Narayanan RL, et al. Confirmed adult dengue deaths in Singapore: 5-year multi-center retrospective study. *BMC Infect Dis* 2011; 11: 123. Published Online First: 12 May 2011. doi: 10.1186/1471-2334-11-123.
- Potts JA, Rothman AL. Clinical and laboratory features that distinguish dengue from other febrile illnesses in endemic populations. *Trop Med Int Health* 2008; 13: 1328-40. Published Online First: 18 September 2008. doi: 10.1111/j.1365-3156.2008.02151.x.
- Shahid M, Razzak J, Jamali S, Ali SS, Ayubi TK. Initial diagnostic categories of patients presenting to four major emergency departments in Karachi. *J Coll Physicians Surg Pak* 2006; 16: 680.
- Sudjana P, Jusuf H. Concurrent dengue hemorrhagic fever and typhoid fever infection in adult: case report. *Southeast Asian J Trop Med Public Health* 1998; 29: 370-2.
- Ali N, Nadeem A, Anwar M, Tariq WU, Chotani RA. Dengue fever in malaria endemic areas. *J Coll Physicians Surg Pak* 2006; 16: 340-2.
- Abbasi A, Butt N, Sheikh QH, Bhutto AR, Munir SM, Ahmed SM. Clinical features, diagnostic techniques and management of dual dengue and malaria infection. *J Coll Physicians Surg Pak* 2009; 19: 25-9.
- Siddiqui FJ, Haider SR, Bhutta ZA. Endemic dengue fever: a seldom recognized hazard for Pakistani children. *J Infect Dev Ctries* 2009; 3: 306-12.
- Phuong H, de Vries PJ, Nga TTT, Giao PT, Hung LQ, Binh TQ, et al. Dengue as a cause of acute undifferentiated fever in Vietnam. *BMC Infect Dis*. 2006; 6: 123. Published Online First: 25 July 2006. doi:10.1186/1471-2334-6-123
- Balmaseda A, Hammond SN, Perez MA, Cuadra R, Solano S, Rocha J, et al. Short report: Assessment of the World Health Organization scheme for classification of dengue severity in Nicaragua. *Am J Trop Med Hyg* 2005; 73: 1059-62.
- Martínez -Torres E. Dengue. *Estudios avanzados*. 2008; 22: 33-52. <http://dx.doi.org/10.1590/S0103-40142008000300004>
- Rigau-Pérez J. Severe dengue: the need for new case definitions. *Lancet Infect Dis* 2006; 6: 297-302.
- Guzman MG, Kouri G. Dengue diagnosis, advances and challenges. *Int J Infect Dis* 2004; 8: 69-80.