



1-2016

An unusual presentation of gbs: case report and literature review

Sadaf Khattak

Pakistan Institute of Medical Sciences, Islamabad, Pakistan

Sumaira Nabi

Pakistan Institute of Medical Sciences, Islamabad, Pakistan

Irfan Khattak

Pakistan Institute of Medical Sciences, Islamabad,

Mazhar Badshah

Pakistan Institute of Medical Sciences, Islamabad, Pakistan

Follow this and additional works at: <http://ecommons.aku.edu/pjns>

 Part of the [Neurology Commons](#)

Recommended Citation

Khattak, Sadaf; Nabi, Sumaira; Khattak, Irfan; and Badshah, Mazhar (2016) "An unusual presentation of gbs: case report and literature review," *Pakistan Journal of Neurological Sciences (PJNS)*: Vol. 11: Iss. 1, Article 10.

Available at: <http://ecommons.aku.edu/pjns/vol11/iss1/10>

AN UNUSUAL PRESENTATION OF GBS: CASE REPORT AND LITERATURE REVIEW

Sadaf Khattak¹, Sumaira Nabi¹, Irfan Khattak², Mazhar Badshah³

¹ Senior Registrar

² Postgraduate resident of Neurology

³ Associate Professor

Correspondence to: Department of Neurology, Pakistan Institute of Medical Sciences, Islamabad, Pakistan. E-mail: sumairafn@gmail.com
Date of Submission: September 29, 2015 **Date of Revision:** December 05, 2015, **Date of Acceptance:** December 15, 2015

ABSTRACT

Guillain-Barre syndrome (GBS), also known as Landry paralysis is an acute idiopathic polyneuritis, believed to be immunologically mediated. It usually presents as a demyelinating neuropathy with ascending weakness, however, many clinical variants have been well documented in the medical literature, and variants involving the cranial nerves or pure motor involvement with axonal injury have also been described. We report a case of a 50 year old patient who initially presented to the ER with hemiparesis and cranial nerve palsies simulating a cerebrovascular event. Based on neurological examination, CSF analysis and needle EMG finding a diagnosis of GBS was made.

Introduction:

Guillain-Barre syndrome (GBS) was first reported by Landry in 1859 and later detailed by Guillain, Barré and Strohl, in 1916. The disease has become well-known internationally under the name of Guillain-Barré Syndrome¹. The disease is assumed to be autoimmune and operated by a preceding infection, most of the time respiratory or gastrointestinal infections². Generally infections by microorganisms such as Campylobacter jejuni, CMV, Mycoplasma pneumonia, or influenza virus exist several weeks prior to approximately two thirds of GBS cases². A small percentage of patients develop GBS after immunization with quadrivalent meningococcal polysaccharide conjugate vaccine (MCV4), following surgery or trauma, and bone-marrow transplantation³. It was associated with tumor necrosis factor alpha antagonist therapy in one report⁴. In another report it was associated with the use of isotretinoin in two patients, however, a definite causal link couldn't be identified⁵. GBS has also been linked to other autoimmune diseases and malignancy such as systemic lupus erythematosus, and Hodgkin lymphoma⁶. Although the classic description of GBS is that of a demyelinating neuropathy with ascending weakness, many clinical variants have been well documented in the medical literature, and variants involving the cranial nerves or pure motor involvement and axonal injury have also been described⁷. We describe a rare case of GBS presenting as hemiparesis.

CASE REPORT:

A 50 year old female known hypertensive for 3 years fairly controlled with medication, presented with h/o sudden onset,

generalized moderate intensity headache, and progressive weakness of right side of the body which was followed 2 days later by double vision (while looking towards left) and facial asymmetry. With the clinical suspicion of stroke her MRI brain was done at a local hospital and she was then referred to our hospital. On presentation to our hospital she had developed new onset left sided weakness 6 days later in addition to the persisting right sided weakness, facial asymmetry and diplopia. There was history of diarrhea 3 weeks prior to onset of weakness. She denied any history of fever, dysphagia, nasal regurgitation, urinary or fecal incontinence, backache, neck pain, photophobia, altered sensorium, arthralgias, oral ulcers or photosensitivity. Examination revealed an overweight female conscious, oriented time, place and person with a respiratory rate of 20/min without use of accessory muscles. Her pulse was regular 70 bpm and BP was 130/90 mmHg. Her higher mental functions were intact. Motor examination revealed normal bulk with hypotonia in all four extremities with power of 3/5 in both upper extremities and 2/5 in both lower extremities (both proximally and distally). She had generalized areflexia with mute planter response. There was facial diplegia (bilateral LMN type facial palsy) and bilateral 6th nerve paresis (left > right), rest of the cranial nerves were intact and symmetric. Cerebellar examination wasn't possible owing to patient's quadriplegia and pin prick and proprioception was intact. Fundoscopic examination revealed grade 1 hypertensive changes. Chest was clear to auscultation. Examination of CVS and GI system was unremarkable. Laboratory parameters showed normal complete blood picture and metabolic panel. Her ESR was 18, ANA, RA factor and hepatitis serology were negative. MRI brain

with contrast was normal. CSF showed cytoprotein dissociation with 5 cells, 77 mg/dl protein, and a normal glucose 65mg/dl(BSR134mg/dl). Nerve Conduction Studies/Electromyography was suggestive of a demyelinating polyneuropathy with secondary axonal degeneration. Left median, peroneal and tibial nerve showed markedly prolonged distal latency with decreased amplitude. Left median and ulnar showed mildly reduced velocity, low normal velocity was seen in left peroneal and normal in left tibial with mild dispersal of response in left tibial nerve. Left ulnar nerve showed normal motor and sensory response with absent response in left median nerve. F waves were absent in all the tested nerves. EMG of upper limb muscles showed reduced recruitment of normal duration and amplitude MUAPs while those of lower limbs showed reduced recruitment of mildly neuropathic MUAPs. On the basis of the above clinical and lab data a diagnosis of GBS was made and plasmapheresis was initiated alongwith supportive treatment. There wasn't much improvement in her power however her shortness of breath improved and she was discharged after 6 sessions of plasmapheresis and referred for rehabilitation. Unfortunately she was lost to follow ups.

DISCUSSION:

Guillain-Barré syndrome (GBS), also known as acute idiopathic polyneuritis, is a type of neuromuscular paralysis that has several variants⁴. In about 75% of cases it is preceded 1,2,7-8 weeks by an acute illness mostly respiratory or gastrointestinal. Campylobacter jejuni (*C. jejuni*) infection is, overall, the most common antecedent infection and has been reported in up to 32% of cases. *C. jejuni* GBS has marked motor axon degeneration, an elevated anti-GM1 antibody, and a delayed and often incomplete recovery. There appears to be an over-representation on of certain strains of *C. jejuni* suggesting that the lipopolysaccharides of these organisms share ganglioside-like epitopes with peripheral nerves. This molecular mimicry appears to confuse the immune system resulting in a mistaken attack against neural antigens⁹. Patients usually present with complaints of finger dysesthesias and muscle weakness of the lower extremities. The weakness may progress over hours to days to involve the arms, truncal muscles, cranial nerves and muscles of respiration. Variants of GBS may present as pure motor dysfunction (AMAN) or acute dysautonomia⁷. Half of GBS patients have some degree of cranial nerve dysfunction during their illness. Facial weakness is most common, especially if substantial limb weakness is present. Ophthalmoparesis is seen in 10-20% of patients with abducens palsy being most common which is usually bilateral. Acute motor axonal neuropathy (AMAN) is distinguished from AIDP by its involvement of exclusively motor nerves and an electrophysiologic pattern indicating axonal involvement. AMAN is seen commonly in northern China and Japan^{10,11},

usually associated with a preceding *Campylobacter jejuni* infection. The clinical features and recovery are very similar to those of AIDP¹². However, more AMAN patients require assisted ventilation because of impending respiratory failure. Acute motor-sensory axonal neuropathy (AMSAN) affects both sensory and motor fibers and is thus a more severe form of AMAN¹¹. Clinically, it resembles AMAN variant with the exception that sensory symptoms are more in AMSAN. There is marked axonal degeneration of both motor and sensory nerve fibers demonstrated by severely reduced or absent motor and sensory responses on nerve conduction studies and extensive denervation on follow-up EMG studies, resulting in delayed and incomplete recovery. Typical presentation of Miller Fischer Syndrome (MFS) is that of ophthalmoplegia with ataxia and areflexia¹³. About one-quarter of patients who present with MFS will develop some extremity weakness, clearly linking this disorder to GBS. A limited form of MFS presents as cerebellar ataxia and hyporeflexia without ophthalmoplegia. Antibodies against GQ1b are present in 85 to 90 percent of patients with MFS GBS. Elevated antibodies to GQ1b suggest an immune attack against GQ1b gangliosides which are concentrated in the Para nodal regions of extraocular nerves⁹. Certain features of MFS like ophthalmoplegia and ataxia along with encephalopathy and hyperreflexia characterize another variant of GBS, namely Bickerstaff encephalitis. It is also associated with anti-GQ1b antibodies and thus responds to IVIG or plasma exchange^{14,15}. The pharyngeal-cervical-brachial variant of GBS is characterized by acute muscular weakness of the oropharynx, neck, and shoulder girdle with swallowing abnormality^{16,17}. It represents a localized form of axonal GBS^{16,17}. Facial weakness may be seen. Motor strength and reflexes are typically preserved. This form overlaps with MFS¹⁷. Some patients with pharyngeal-cervical-brachial weakness have antibodies against GQ1b, or less often GD1a. It is thought by some experts that MFS, Bickerstaff encephalitis, and pharyngeal-cervical-brachial weakness with anti-GQ1b antibodies constitute overlapping expressions of the anti-GQ1b antibody syndrome¹⁸. Tatsumoto M et al showed in their study that isolated abducens nerve palsy can be categorized as a regional variant of Guillain-Barré syndrome or mild form of Fisher syndrome. They reviewed clinical profiles and laboratory findings for 100 cases of abducens nerve paresis. Tentative diagnoses made by the primary physicians on request of anti-ganglioside antibody testing were abducens nerve palsy (n=68), Fisher syndrome (n=14), acute ophthalmoparesis without ataxia (n=14). Serum anti-GQ1b antibody was positive in 25¹⁹. Yoon-Sik Jo, and Sang-Jun Nav have reported an unusual case of pure motor variant of GBS initially presenting with hemiparesis, (a 59 year-old-male presenting with left sided weakness 1 day prior to admission who developed right sided weakness 2 days following admission). Electrophysiological findings were consistent with

pure motor GBS²⁰. This patient presented with hemiparesis developing weakness of the other side few days later with NCS findings supporting diagnosis of GBS, similar to our patient however he didn't have facial diplegia or ophthalmoparesis as seen in our case report moreover the neurophysiology studies showed pure motor GBS while it was suggestive of demyelinating polyneuropathy with secondary axonal degeneration in our patient. Guillain-Barré syndrome (GBS) is generally diagnosed on clinical grounds⁷. Electromyography (EMG) and nerve conduction studies (NCS) can be very helpful in the diagnosis as well as for prognosis. Abnormalities in NCS that are consistent with demyelination are sensitive and represent specific findings for classic GBS(AIDP variant). Delayed distal latencies, slowed nerve conduction velocities, temporal dispersion of waveforms, conduction block, prolonged or absent F waves, and prolonged or absent H-reflexes are all findings that support demyelination. Needle EMG may be normal in acute nerve lesions, and it may take 3-4 weeks for fibrillation to develop. In the acute phase, the only needle EMG abnormality may be abnormal motor recruitment, with decreased recruitment and rapid firing motor units in weak muscles. Unfortunately, electrodiagnostic studies can be completely normal in early GBS and a normal study does not rule out the disease.^{21,22} Characteristic findings on CSF analysis include albuminocytologic dissociation, which is an elevation in CSF protein (>0.55 g/L) without an elevation in white blood cells. The increase in CSF protein is thought to reflect the widespread inflammation of the nerve roots⁷. Our patient initially presented with hemiparesis with cranial nerve palsies, a rare presentation of GBS her examination along with electrodiagnostic studies and CSF examination supported the diagnosis of GBS. Chauhan v et al have reported a case of GBS in India presenting as Bell's palsy with crossed hemiparesis who recovered over a month²³. Kim EJ and Yuki N from Australia have reported an unusual case of GBS who presented with left hemiparesis at the acute phase and right hemifacial weakness at the recovery phase of the hemiparesis²⁴. Acute hemiplegia as a clinical presentation of GBS could be confidently added to the category of atypical presentation of pediatric GBS which constitutes 11.2-24.3% of the whole pediatric GBS presentation reported previously. The management of patients with Guillain-Barré syndrome (GBS) can be intimidating.²⁵The unpredictable course and potential for rapidly producing life-threatening respiratory failure may prompt admission to an intensive care unit (ICU).Treatment of GBS though challenging, is mainly supportive and is centered at the anticipation and prevention of various complications. Immunomodulation with plasmapheresis and IV immunoglobulin (IVIG) helps to shorten the course of the disease.²⁶⁻²⁹ Patients with more severe disease and rapid progression of illness require extensive and prolonged rehabilitation and also show delayed and incomplete recovery.

REFERENCES:

1. Hughes RA, Comblath DR. Guillain-Barré syndrome. *Lancet*. 2005 Nov;366(9497):1653–66.
2. Vriesendorp FJ, Mishu B, Blaser MJ, Koski CL. Serum antibodies to GM1, GD1b, peripheral nerve myelin, and *Campylobacter jejuni* in patients with Guillain-Barre syndrome and controls: correlation and prognosis. *Ann Neurol*. 1993 Aug;34(2):130–5.
3. Ropper AH, Wijdicks EFM, Truax BT. Guillain-Barré syndrome, FA Davis, Philadelphia 1991. p.57
4. Shin IS, Baer AN, Kwon HJ, et al. Guillain-Barré and Miller Fisher syndromes occurring with tumor necrosis factor alpha antagonist therapy. *Arthritis Rheum* 2006; 54:1429.
5. Pritchard J, Appleton R, Howard R, Hughes RA. Guillain-Barré syndrome seen in users of isotretinoin. *BMJ* 2004; 328:1537.
6. Ropper AH. The Guillain-Barré syndrome. *N Engl J Med* 1992; 326:1130.
7. Andary TM, Oleszek LJ et al Guillain-Barre Syndrome Treatment & Management (online) 2014
8. (cited 20th May 2014). Available from (<http://emedicine.medscape.com/article/315632-overview>)
9. Kozminski, PM. Miller Fisher Variant of Guillain-Barré Syndrome: A Report of Case. *J Am Osteopath Assoc* 2008 Feb;108(2):51-2
10. Gorson, KC, and Ropper, AH. Guillain-Barre Syndrome (Acute Inflammatory Demyelinating Polyneuropathy) and Related Disorders. In: Katirji, B, Kaminski, HJ, Preston, DC, Ruff, RL, and Shapiro, BE, (eds.): *Neuromuscular Disorders in Clinical Practice*, Butterworth Heinemann, Boston, 2002, p. 544-566
11. McKhann GM, Comblath DR, Griffin JW, et al. Acute motor axonal neuropathy: a frequent cause of acute flaccid paralysis in China. *Ann Neurol* 1993; 33:333.
12. Feasby TE, Gilbert JJ, Brown WF, et al. An acute axonal form of Guillain-Barré polyneuropathy. *Brain* 1986; 109:1115.
13. Ho TW, Li CY, Comblath DR, et al. Patterns of recovery in the Guillain-Barre syndromes. *Neurology* 1997; 48:695.
14. FISHER M. An unusual variant of acute idiopathic polyneuritis (syndrome of ophthalmoplegia, ataxia and areflexia). *N Engl J Med* 1956; 255:57.
15. Winer JB. Bickerstaff's encephalitis and the Miller Fisher syndrome. *J NeurolNeurosurg Psychiatry* 2001; 71:433.
16. Odaka M, Yuki N, Hirata K. Anti-GQ1b IgG antibody syndrome: clinical and immunological range. *J NeurolNeurosurg Psychiatry* 2001; 70:50.

17. 16) Ropper AH. Unusual clinical variants and signs in Guillain-Barré syndrome. *Arch Neurol* 1986; 43:1150.
18. 17) Nagashima T, Koga M, Odaka M, et al. Continuous spectrum of pharyngeal-cervical-brachial variant of Guillain-Barré syndrome. *Arch Neurol* 2007; 64:1519.
19. 18) Shahrizaila N, Yuki N. Bickerstaff brainstem encephalitis and Fisher syndrome: anti-GQ1b antibody syndrome. *J NeurolNeurosurg Psychiatry* 2012.
20. 19) Tatsumoto M, Odaka M, Hirata K, et al. Isolated abducens nerve palsy as a regional variant of Guillain-Barré syndrome. *J NeurolSci* 2006;243:35-38.
21. 20) Jo S Yoon, Na J Sang. An Unusual Case of Pure Motor Variant of Guillain-Barre Syndrome Initially Presenting with Hemiparesis. *J. Korean Neurol Assoc.* 2006 Aug;24(4):385-388. Korean
22. 21) Albers JW, Kelly JJ Jr. Acquired inflammatory demyelinating polyneuropathies: clinical and electrodiagnostic features. *Muscle Nerve.* Jun 1989;12(6):435-51.
23. 22) Van den Bergh PY, Piéret F. Electrodiagnostic criteria for acute and chronic inflammatory demyelinating polyradiculoneuropathy. *Muscle Nerve.* Apr 2004;29(4):565-74.
24. 23) Chauhan V et al. Guillain-Barré syndrome presenting as Bell's palsy with crossed hemiparesis. *Int J Crit Illn Inj Sci.* 2015 Jan-Mar; 5(1): 62-63.
25. 24) Kim EJ, Yuki N. Hemiparetic Guillain-Barré syndrome. *J NeuroSci.* 2016; 361: 131-132
26. 25) Ropper AH, Samuels MA. Diseases of peripheral nerves. In: Davis JK, Sydor AM (editors) *Adams and Victor's Principles of Neurology.* 9th ed. Boston, USA: McGraw-Hill; 2009:1251-1325.
27. 26) Paul BS, Bhatia R, Prasad K, Padma MV, Tripathi M, Singh MB. Clinical predictors of mechanical ventilation in Guillain-Barre' syndrome. *Neurol India* 2012;60:150-3.
28. 27) Rajabally YA, Uncini A. Outcome and its predictors in Guillain-Barre' syndrome. *J NeurolNeurosurg Psychiatry.* 2011;10.1136.
29. 28) Doom PAV, Ruts L, Jacobs BC. Clinical features, pathogenesis and treatment of Guillain-Barre Syndrome. *Lancet Neurol* 2008; 7:939-50.
30. 29) Doom PAV. What's new in Guillain-Barre' syndrome in 2007-2008? *J PeripherNervSyst* 2009; 14:72-4.

Conflict of interest: Author declares no conflict of interest.

Funding disclosure: Nil

Author's contribution:

Dr. Sadaf Khattak: Study concept and design, protocol writing, data collection, data analysis, manuscript writing, manuscript review

Dr. Sumaira Nabi: data collection, data analysis, manuscript writing, manuscript review

Dr. Irfan Khattak: data collection, data analysis, manuscript writing, manuscript review

Dr. Mazhar Badshah: Study concept and design, data analysis, manuscript writing, manuscript review